.

INTERNATIONAL CRITICAL TABLES
OF
NUMERICAL DATA
PHYSICS, CHEMISTRY AND TECHNOLOGY

INTERNATIONAL CRITICAL TABLES

OF

NUMERICAL DATA,

PHYSICS, CHEMISTRY AND TECHNOLOGY

Prepared under the Auspices of the International
Research Council and the National
Academy of Sciences

.

.

PREFACE BY THE BOARD OF TRUSTEES

The publication of International Critical Tables at a price that would make possible a world-wide distribution required that the undertaking be financed by those appreciating its importance and in a position to make the necessary investment. Some 244 firms and individuals and two of the larger Foundations have provided the sum of $170,000 required for the compilation.

Many individuals have given freely of their time and effort in helping to obtain the funds necessary for the compilation of this work. In addition to those who have been responsible for assigned territory, there are a large number of others in industrial organizations which have supported the enterprise, and grateful acknowledgment is made of their interest and help, quite as much as if it were possible to give here the complete list of names. Indeed, it is impossible for the trustees to know of all those who at different stages of the work have rendered valuable assistance.

Special acknowledgment is due to the Carnegie Corporation of New York and to the International Education Board, whose appropriations in the support of this work were a large factor in making its successful completion possible.

It is appropriate to give here special recognition to those who assumed and carried out definite responsibility in the solicitation of funds, as well as to those whose financial support enabled the project to be made a reality.

LIST OF THOSE WHOSE SUPPORT HAS MADE POSSIBLE THE COMPLETION OF INTERNATIONAL CRITICAL TABLES

vii

The work of the trustees began with the appointment of Hugh K. Moore in 1920, with whom were later associated Julius Stieglitz, representing the American Chemical Society, and E. P. Hyde, representing the American Physical Society. After a substantial sum had been procured, the number was enlarged to include H. E. Howe and later George P. Adamson and Charles L. Reese. Mr. Hyde resigned to go abroad and was succeeded by Frank B. Jewett, who has lately been succeeded by Michael Pupin as representative of the American Physical Society. Upon relinquishing his active duties in the National Research Council, H. E. Howe was succeeded as Secretary of the Board of Trustees by W. M. Corse, but remained a member of the Board; and a little later Edward B. Craft was added to the Board.

The trustees have been obliged to place a maximum limit on the cost of this work, but they realize that other material which could not be included because of financial limitations should be made available and that International Critical Tables, if it is to render maximum service, should become an established institution, with supplements and revisions published from time to time, in order that these fundamental data may be made available as rapidly as the values are established through further research. An endowment therefore should be sought for International Critical Tables, and with the appearance of the completed set it is believed the enterprise will appeal to many of those able to make such an endowment a reality.

The trustees wish to express their gratitude to the many industrialists who have given of their time to become acquainted with this enterprise, for the courtesy which they have everywhere met, and for the widespread cooperation without which International Critical Tables could not have been brought into existence.

PREFACE BY THE
BOARD OF EDITORS

At the organization meeting of the International Union of Pure and Applied Chemistry, held in London in June 1919, the Union approved as one of its projects the compilation of International Critical Tables of Numerical Data of Physics, Chemistry, and Technology, and assigned to the United States of America the financial and editorial responsibility for the undertaking. The project was later given the patronage of the International Research Council at its Brussels meeting in 1923.

On behalf of the National Academy of Sciences, the National Research Council of the United States accepted the executive, editorial and financial responsibilities of the project, and with the cooperation of the American Chemical Society and the American Physical Society, created a Board of Trustees to take charge of the financial and business administration, and a Board of Editors to supervise and carry out the preparation of the text.

The first action of the Board of Editors, early in 1922, was to approve the appointment of Corresponding Editors in different parts of the world, particularly in all those countries in which conditions were such that they might be expected to take a really active part in the undertaking. In making these appointments, the Board first sought the advice of competent individuals in the several countries, and in accordance with the suggestions thus received, appointed ten Corresponding Editors and empowered them to arrange for Advisory Committees to assist in the work. In the case of certain countries, the Board was unsuccessful in its efforts to secure cooperation, usually either because of the receipt of no reply or an unfavorable reply, or through failure of the Corresponding Editor, after appointment, to perform his duties.

The general plan of preparation of the Tables was as follows: The subject matter was first divided into some 300 different sections. The Corresponding Editors were then asked to recommend for the several sections one or more persons who should either have some special knowledge of the subject matter of the section, or be otherwise qualified to pass critical judgment upon the available information on the subject. On the basis of the recommendations thus received, the Board of Editors selected the Cooperating Experts, to whom was intrusted the task of critically compiling, and displaying in suitable form, the available quantitative information upon the several topics. In making these selections, the Board consistently endeavored to secure the best man *available* in the light of all the information which it possessed. In certain special fields composed of closely related topics, the Board provided also for the appointment of Special Editors to supervise the work and to assist in the final arangement of the material.

In the course of its labors the Board of Editors has enjoyed the cooperation of numerous organizations and individuals whose advice, suggestions, and assistance, in many ways have greatly aided it in its complex and difficult task. It is especially indebted to the several Corresponding Editors and their Advisory Committees, who have generously contributed their time and thought to the success of the work; to the Special Editors; to the U. S. Bureau of Standards, the National Physical Laboratory of Great Britain and the Physical Society of France; to the International Commission in charge of Annual Tables; and to various organizations and individuals who made available unpublished data for the use of the Cooperating Experts.

PREFACE PAR LE
COMITÉ DES RÉDACTEURS

Lors de l'Assemblée d'organisation de l'Union internationale de Chimie pure et appliquée, qui eut lieu à Londres en Juin 1919, l'Union approuva comme l'un de ses projets l'élaboration de Tables critiques de valeurs numériques de physique, chimie et technologie, et elle chargea les Etats-Unis d'Amérique de la responsabilité financière et d'édition de l'entreprise. Le projet fut, plus tard, placé sous le patronage du Conseil international de Recherches, à son assemblée de Bruxelles en 1923.

Chargé de ces attributions, le Conseil national de Recherches des Etats-Unis, agissant en collaboration avec la Société chimique américaine et la Société physique américaine, nomma un Conseil d'Administration et un Comité des Rédacteurs.

La première activité que manifesta le Comité des Rédacteurs, au début de 1922, fut d'approuver la nomination de Rédacteurs-correspondants dans les différentes parties du monde, particulièrement dans tous les pays dont les conditions autorisaient l'espoir d'une collaboration active dans cette entreprise. Pour procéder à ces nominations, le Comité sollicita d'abord l'avis de personnalités compétentes dans les divers pays, et c'est en tenant compte des suggestions ainsi obtenues qu'il nomma dix Rédacteurs-correspondants et leur donna les pouvoirs nécessaires pour organiser des Comités-consultatifs dans le but d'aider à l'accomplissement du travail. Dans le cas de certains pays, les efforts du Comité en vue de s'assurer leur coopération furent vains, soit qu'il n'y eût pas de réponse ou que celle-ci fut défavorable, soit encore que le Rédacteur-correspondant, après sa nomination, eût manqué à ses engagements.

Le plan général de préparation de ces Tables fut le suivant: l'ensemble des matières à traiter fut d'abord divisé en quelque 300 différentes sections. Les Rédacteurs-correspondants furent alors priés de recommander, pour les différentes sections, une ou plusieurs personnes qui eussent des connaissances spéciales du sujet traité dans la section ou qui fussent qualifiées pour formuler un jugement critique sur les informations à disposition concernant le sujet. Sur la base des recommandations ainsi reçues, le Comité des Rédacteurs choisit les Experts-coopérants qui furent chargés de la compilation critique et de la disposition sous une forme convenable des informations quantitatives disponibles sur les différents sujets. En faisant cette sélection, le Comité s'efforça de s'assurer la collaboration de la personne qui, d'après les renseignements recueillis, était la plus qualifiée et qui se trouvait alors disponible. Dans certains domaines spéciaux, composés de sujets étroitement apparentés, le Comité se chargea aussi de nommer des rédacteurs spéciaux pour diriger le travail et pour aider à l'arrangement final de la matière.

Au cours de ses travaux, le Comité des Rédacteurs a eu le plaisir d'enregistrer la coopération de nombreuses organisations et de particuliers dont les conseils, les suggestions et l'aide lui ont été, en maintes circonstances, d'un grand secours dans l'accomplissement de sa tâche complexe et difficile. Il est spécialement reconnaissant aux nombreux Rédacteurs-correspondants et à leurs Comités-consultatifs qui ont généreusement donné leur temps et leur pensée pour assurer le succès de l'oeuvre; aux Rédacteurs spéciaux, au U. S. Bureau of Standards, au National Physical Laboratory of Great Britain et à la Société de Physique de France; à la Commission internationale chargée des Tables annuelles; ainsi qu'aux

x

VORWORT DER REDAKTIONS-KOMMISSION

An der geschäftlichen Sitzung der Internationalen Union für reine und angewandte Chemie in London, Juni 1919 billigte die Union, als eine ihrer Aufgaben, die Abfassung Internationaler kritischer Tafeln, numerischer Daten der Physik, Chemie und Technologie und betraute die Vereinigten Staaten von Amerika sowohl mit dem finanziellen als auch mit dem redaktionellen Teil dieser Aufgabe. Der Plan erhielt später die Förderung durch International Research Council an der Tagung in Brüssel 1923.

Entsprechend dieser Betrauung errichtete National Research Council der Vereinigten Staaten, zusammenwirkend mit American Chemical Society und American Physical Society vorgehend, eine geschäfts-führende Kommission und eine Redaktions-Kommission.

Die ersten Schritte, welche die Redaktions-Kommission zu Beginn des Jahres 1922 machte, war, sich korrespondierende Mitglieder in allen Teilen der Welt zu sichern, besonders in denjenigen in welchen die Bedingungen vorhanden waren, die eine lebhafte Beteiligung an dem Unternehmen erwarten liessen. Nach diesem nahm die Kommission zuerst den Rat massgebender Persönlichkeiten verschiedener Länder entgegen; in Übereinstimmung mit den so erhaltenen Vorschlägen, wurden zehn korrespondierende Mitglieder bestimmt, welche nun einen beratenden Ausschuss zu bilden hatten, um der Arbeit ihre Unterstützung zu zuwenden. In einigen Ländern gelang es der Kommission nicht Mitarbeiter zu erlangen, meistens deshalb weil keine, oder eine ablehnende Gegenäusserung erfolgte, oder, dass das korrespondierende Mitglied, nach der entsprechenden Zusage nicht vorging.

Die Grundlinien für die Bearbeitung der Tafeln waren die folgenden. Das Material wurde zuerst in etwa dreihundert verschiedene Abschnitte zerlegt. Die korrespondierenden Mitglieder wurden dann gebeten, für einige dieser Abschnitte, einen oder mehrere Mitarbeiter zu empfehlen, die entweder besondere Kenntnisse über den Gegenstand des Abschnittes besitzen, oder imstande waren, kritisch, vorhandenes Material durchzugehen. Auf Grund der so erhaltenen Empfehlungen, wählte die Redaktionskommission die Mitarbeiter aus, die mit der Aufgabe betraut wurden, kritisch die numerischen Daten des betreffenden Gegenstandes durchzuarbeiten und in entsprechender Form darzustellen. Bei dieser Auswahl war die Kommission ganz besonders bestrebt, nach den vorhandenen Mitteilungen, den besten *zur Verfügung* stehenden Mitarbeiter zu erhalten. In gewissen nahe verwandten Gebieten war man darauf bedacht, besondere Redaktions-mitglieder zu erhalten, um die Arbeit hier zu überwachen und tätigen Anteil der Schlussredaktion des Materials zu nehmen.

Im Laufe ihrer Bestrebungen konnte sich die Redaktion-Kommission der Mitarbeit zahlreicher Vereinigungen und einzelner Personen erfreuen, deren Ratschläge, Winke und Beihilfe ihnen bei der verwickelten und schweren Aufgabe von grossem Nutzen waren. Die Redaktionskommission ist besondren Dank ihren verschiedenen korrespondierenden Mitgliedern und dem beratenden Ausschuss schuldig, die in grossmütiger Weise ihre Zeit und Arbeit dem Erfolg dieser Tafeln gewidmet haben, ferner auch den Mitgliedern, die die Arbeit an den besonderen Kapiteln überwachten. Der Dank gebührt U. S. Bureau of Standards, National Physical Laboratory of Great Britain und Société de Physique de France, der Internationalen Kommission betraut mit der Herausgabe der Tables annuelles und den verschiedenen Ver-

PREFAZIONE DELL' UFFICIO DI REDAZIONE

Nella conferenza tenuta a Londra nel giugno 1919 per organizzare la Unione Internazionale della Chimica Pura ed Applicata venne, tra gli altri, formulato il progetto di compilare delle Tabelle Critiche Internazionali contenenti dati numerici di fisica, chimica e tecnologia, e venne affidata agli Stati Uniti la responsabilità finanziaria ed editoriale dell'impresa. Al progetto fu in seguito accordato il patronato del Consiglio Internazionale di Ricerche nella riunione del 1923 a Bruxelles.

In seguito all'incarico ricevuto, il Consiglio Nazionale di Ricerche degli Stati Uniti, d'accordo con la American Chemical Society e con la American Physical Society, nominò un Consiglio di Amministrazione ed un Ufficio Editoriale.

Come suo primo atto, l'Ufficio, nel 1922, nominò Redattori Corrispondenti in tutto il mondo, scegliendoli di preferenza nei Paesi dove poteva ritenersi che essi avrebbero preso parte attiva al lavoro. Le nomine furono fatte dopo aver sentito il parere di persone competenti. A questo modo furono scelti dieci Redattori Corrispondenti e ad essi venne data facoltà di nominare ciascuno un Comitato consultivo col compito di assisterli nel lavoro. In alcuni Paesi l'Ufficio non riuscì ad assicurarsi collaborazione di sorta, o perchè addirittura non gli fu possibile ottenere una risposta, o perchè la risposta fu negativa, o perchè il Redattore Corrispondente scelto, dopo essere stato nominato, mancò agli obblighi assunti.

Il piano generale di preparazione delle tabelle è stato il seguente. Si è diviso la materia in circa 300 capitoli differenti, e i Redattori Corrispondenti sono stati invitati a suggerire per ogni singolo capitolo il nome di una o più persone le quali o avessero una speciale competenza nell'argomento o potessero ritenersi capaci di vagliare criticamente tutto quello che si conosce al riguardo. In base alle proposte ricevute, l'Ufficio di Redazione scelse gli Esperti, e a questi affidò l'incarico di raccogliere, vagliare ed esporre in forma opportuna i dati quantitativi che si sono potuti riunire sui diversi argomenti.

Nel fare la scelta degli Esperti l'Ufficio cercò sempre di assicurarsi la collaborazione degli uomini che, in base alle informazioni avute, dovevano ritenersi i migliori di cui si potesse disporre. In certi campi speciali, comprendenti argomenti strettamente connessi, l'Ufficio nominò anche dei Redattori Speciali col compito di sorvegliare il lavoro e collaborare alla disposizione definitiva del materiale.

Nell'espletare il suo compito, l'Ufficio di Redazione ha potuto giovarsi della collaborazione di numerose organizzazioni e di numerose persone, le quali con consigli e suggerimenti vari sono state di grande aiuto nel portare a fine un lavoro che è stato certamente complesso e difficile. L'Ufficio è specialmente grato ai vari Redattori Corrispondenti e ai rispettivi Comitati Consultivi i quali hanno generosamente dato il loro tempo e la loro intelligenza al successo dell'opera, ai Redattori Speciali, al Bureau of Standards degli Stati Uniti, al National Physical Laboratory inglese e alla Société de Physique francese, alla Commissione Internazionale in carica per le Tabelle annuali e alle varie organizzazioni e persone che misero a disposizione degli Esperti dati inediti.

Infine i Membri dell'Ufficio desiderano manifestare l'alto apprezzamento che fanno dei contributi di tutti gli Esperti, il lavoro dei quali, compiuto in larga misura con entusiasmo e disinteressatamente, ha reso possibile queste tabella; ed in particolar modo

Finally, the members of the Board desire to record their appreciation of the work of all of the Cooperating Experts whose contributions, largely a labor of love, have made these tables possible; and in particular, of the work of the Editorial Staff, Messrs. Washburn, Dorsey, and West, to whom indeed the utility of this collection of tables should be largely accredited.

<div style="text-align:center">

George K. Burgess   S. C. Lind
Saul Dushman   C. E. Mendenhall
John Johnston   R. B. Moore.

</div>

organisations diverses et aux personnes qui ont procuré des données inédites à l'usage des Experts-coopérants.

Efin, les membres du Comité désirent exprimer leur appréciation pour le travail de tous les Experts-coopérants dont les contributions, pour une large part désintéressées, ont rendu possible l'élaboration de ces Tables, et en particulier pour le travail des Rédacteurs, MM. Washburn, Dorsey et West, auxquels nous sommes en grande partie redevables des services que rendra cette collection de Tables.

<div style="text-align:center">

George K. Burgess   S. C. Lind
Saul Dushman   C. E. Mendenhall
John Johnston   R. B. Moore.

</div>

INTRODUCTION

International Critical Tables is the result of the cooperative labors of a large number of specialists, each of whom has been charged with the responsibility for the critical compilation of the quantitative information available on his topic. The word "critical" in this connection means that the Cooperating Expert was requested to give in each instance the "best" value which he could derive from all the information available, together, where possible, with an indication of its probable reliability.

Through a cooperative arrangement with International Annual Tables, the Board of Editors has been able to place in the hands of each Cooperating Expert the literature references belonging to his topic for the years 1910–1923 inclusive, as compiled by the staff of International Annual Tables. For the period preceding 1910, each Cooperating Expert was directed to collect the necessary literature references from the various published handbooks, special treatises, works of reference, and other sources known to him as a specialist in the field. No attempt has been made to systematically cover the literature since 1923, although a certain amount of information published since then has been utilized.

In preparing the various sections, the Cooperating Experts were instructed,

1. To include in the bibliography only (a) the sources of the data upon which their reported values actually rest, and (b) the sources of available data of the same kind pertaining to those systems for which no numerical value is given. It is not intended to be a complete bibliography of the field.

2. To omit from the tables of numerical data all those systems for which the available data (a) were of slight scientific or practical interest, or (b) were so discordant as to be of little, if any, value.

3. To set forth the results of their work in the form of text, equations, tables, graphs, or charts, as seemed most appropriate under the circumstances, having regard to the necessity of space economy.

4. To give only selected samples illustrating types in the case of very large and heterogeneous fields, such as colloids, chemical kinetics, and certain classes of industrial materials

5. To restrict the accompanying explanatory text to the amount necessary for the intelligent use of the data. (Under this restriction, the Expert is given no opportunity to present a general discussion of his subject or of the methods by which he obtained the values given.)

In preparing the textual material for publication the Editors have been compelled, in the interest of economy of space, to enforce the restrictions imposed by sections 3 and 5 of the preceding paragraph and have freely rearranged and rewritten the text, whenever it was evident that a compression or an improvement in logical order could be so secured. With few exceptions, which are duly

INTRODUCTION

Les Tables critiques internationales sont le résultat du travail coopératif d'un grand nombre de spécialistes, chacun de ceux-ci ayant été chargé de la responsabilité de la compilation critique des informations disponibles sur son sujet. Le mot "critique" dans ce cas signifie que l'expert coopérant fut invité à donner dans chaque circonstance la "meilleure" valeur qu'il pouvait recueillir de toutes les informations disponibles, en ajoutant si possible une indication au sujet de la confiance probable qu'on pouvait avoir en elle.

Par le fait d'un arrangement coopératif avec les Tables annuelles internationales, le Comité des Rédacteurs a été en mesure de mettre à la disposition de chaque expert coopérant les références bibliographiques appartenant à son sujet de l'année 1910 à l'année 1923 inclusivement, celles-ci ayant été compilées par le Bureau des Tables annuelles internationales. Pour la période précédant 1910, chaque expert coopérant fut chargé de recuillir les références bibliographiques nécessaires en usant des manuels variés, des traités spéciaux, des ouvrages de références, et d'autres sources connues de lui en sa qualité de spécialiste du sujet traité. En ce qui concerne la littérature depuis 1923, aucune tentative n'a été faite pour la couvrir d'une façon systématique; un certain nombre d'informations postérieures à 1923 ont cependant été utilisées.

Pour la préparation des différentes sections, il fut recommandé aux experts coopérants:

1. D'inclure dans la bibliographie seulement (a) les sources de valeurs sur lesquelles reposent actuellement leurs valeurs reportées, et (b) les sources des données de même nature appartenant aux systèmes pour lesquels aucune valeur numérique n'est donnée. Le but poursuivi n'est pas de constituer une bibliographie complète du sujet.

2. De ne pas introduire dans les tables de valeurs numériques tous les systèmes pour lesquels les valeurs disponibles (a) sont de peu d'intérêt scientifique ou pratique, ou (b) sont par trop discordantes pour être d'une valeur quelconque, si toutefois elles en présentent une.

3. De disposer les résultats de leur travail sous la forme d'un texte, d'équations, de tables, de graphiques ou de cartes, en employant le moyen qui leur parut le mieux approprié suivant les circonstances, en ayant en vue la nécessité d'économiser de la place.

4. De ne donner que des exemples choisis, illustrant les types, dans le cas d'un champ très vaste et hétérogène, tel que: les colloïdes, la cinétique chimique et certaines classes de matières industrielles.

5. De restreindre le texte explicatif accompagnant les données au strict nécessarie pour la compréhension de celles-ci. (Vu cette restriction, l'expert n'a donc pas l'occasion de présenter une discussion générale de son sujet et des méthodes par lesquelles il a obtenu les valeurs données).

einigungen und Freunden, die noch nicht veröffentlichten Daten den Mitarbeiteren zur Verfügung stellten.

Schliesslich möchte die Redaktions-Kommission ihre Anerkennung den Mitarbeiteren ausdrücken, deren Arbeitsfreudigkeit diese Tafeln möglich machten, im besondrem aber auch der Mühewaltung des Redaktionsstabes der Herrn Washburn, Dorsey und West, denen man vorwiegend den Erfolg und die Nutzlichkeit dieses Tabellenwerkes schulden muss.

George K Burgess   S. C. Lind
Saul Dushman   C. E Mendenhall
John Johnston   R. B. Moore

ricordano l'opera dei dirigenti dell'Ufficio di Redazione, Sigg. Washburn, Dorsey, e West ai quali sopratutto si deve essere grati per l'utilità che si avrà dalla presente raccolta di tabelle.

George K. Burgess   S. C. Lind
Saul Dushman   C. E. Mendenhall
John Johnston   R B. Moore.

EINLEITUNG

Die Internationalen kritischen Tafeln stellen die Ergebnisse des Zusammenwirkens einer grossen Zahl von Mitarbeiteren mit besonderen Erfahrungen dar, die mit der Aufgabe betraut wurden, die erreichbaren Daten des entsprechenden Gebietes kritisch darzustellen. In dieser Verbindung bedeutet das Wort kritisch soviel, dass der Mitarbeiter gebeten wurde, in jedem einzelnem Fall die "besten" Werte zu geben, die er auf Grund aller zur Verfügung stehenden Literaturstellen, ableiten konnte, zugleich ferner, wenn möglich, alle Angaben mit dem Grade ihrer Zuverlässlichkeit zu vermerken.

Durch ein Übereinkommen mit der Redaktion der Tables annuelles konnte die Redaktionskommission jedem einzelnem Mitarbeiter, über seinen Gegenstand die Literatur der Jahre 1910 bis einschliesslich 1923 soweit übergeben, als sie durch die Redaktion der Tables annuelles ausgearbeitet worden ist. Für die Zeit vor 1910 wurde ein jeder Mitarbeiter gebeten, die notwendigen Literaturstellen und Daten aus den verschieden vorhandenen Handbüchern Spezial-und Nachschlagewerken und anderen, ihm als besonderem Kenner auf diesem Gebiete erreichbaren Quellen, zu sammeln. Es ist nicht versucht worden, die Literatur seit 1923 noch systematisch darzustellen, obwohl ein gewisser Teil davon noch Berücksichtigung finden konnte.

Bei der Bearbeitung der verschiedenen Abschnitte erhielt der Mitarbeiter folgende Anweisungen:

1. Als Literatur sind (a) nur diejenigen Stellen anzugeben, auf Grund deren die angegebenen Werte besonders folgerten, (b) die Quellen, über denselben Gegenstand, die aber keine numerischen Daten enthalten, die Verwendung gefunden haben.

2. Es sind in den Zahlenangaben der Tafeln alle diejenigen Systeme wegzulassen, deren vorliegende Daten, (a) von geringem wissenschaftlichen und praktischen Werte sind, order (b) die Daten sind so widersprechend, dass sie, wenn überhaupt, von geringem Werte sind.

3. Die Ergebnisse ihrer Arbeit sind in einer solchen Form darzustellen, dass durch den Text, die Gleichungen, Tabellen und Tafeln mit Rücksichtnahme auf Raumersparnis, der Zweck am besten erfüllt wird.

4. In sehr grossen, heterogenen Gebieten wie in denen der Kolloide, der chemischen Kinetik und in gewissen Fällen von technischer Bedeutung, sind nur ausgewählte Beispiele zu geben, die das Gebiet charakterisieren sollen.

5. Der erläuternde Text ist soweit zu beschränken, dass eine sachgemässe Verwertung der Tafeln noch möglich ist. (Bei dieser Einschränkung hat der Experte nicht die Gelegenheit allgemein seine Aufgabe, noch die Methode, darzustellen, nach welchen er seine Angaben erhalten hat.)

INTRODUZIONE

Le Tabelle Critiche internazionali sono il frutto della collaborazione di un gran numero di specialisti a ciascuno dei quali è stato affidato il compito di vagliare i dati disponibili sopra un determinato soggetto. La denominazione di tabelle "critiche" indica che l'esperto è stato incaricato di dare in ogni caso il valore "migliore," deducibile da tutte le notizie che si hanno a disposizione. Tutte le volte che è stato possibile l'esperto è stato incaricato anche di dare indicazioni sul grado di attendibilità dei valori numerici.

In seguito ad accordi intervenuti con le Tabelle annuali internazionali, l'ufficio di Redazione ha potuto fornire a ciascun esperto le indicazioni bibliografiche riferentisi agli anni dal 1910 al 1923 incluso, quali vengono compilate dalla direzione delle Tabelle internazionali. Per gli anni precedenti al 1910, gli esperti vennero consigliati a raccogliere la letteratura dai vari manuali, trattati speciali, lavori bibliografici e da altre fonti ad essi note data la qualità di ognuno di specialista in un determinato campo. Dei dati pubblicati dopo il 1923 si è tenuto conto solo in parte.

E' stato raccomandato agli esperti che, nel preparare le varie parti:

1. Includessero nella Bibliografia soltanto: (a) le fonti delle indicazioni sulle quali sono basati i valori riportati, e (b) le fonti delle indicazioni riguardanti i sistemi per i quali non viene dato nessun valore. Non si è riportare inteso una bibliografia completa del soggetto.

2. Omettessero nelle tabelle delle grandezze numeriche tutti quei sistemi per i quali i dati disponibili; (a) fossero di poco interesse scientifico o pratico, oppure (b) fossero così in disaccordo da essere di poco o di nessun valore.

3. Esponessero, a seconda dei casi, i risultati del loro lavoro in forma di testo, di equazioni, di tabelle, di grafici, o di tavole tenendo presente la necessità di economia di spazio.

4. Riportassero soltanto esempi tipici nei campi molto vasti ed eterogenei come colloidi, cinetica chimica ed alcune classi di prodotti industriali.

5. Limitassero il testo esplicativo a quel tanto sufficiente per un uso intelligente delle tabelle (data questa limitazione, all' esperto non è stato consentito di redigere una esposizione generale del suo soggetto o dei metodi con i quali egli ha ottenuto i valori che riporta).

Nel preparare il testo per la pubblicazione i Redattori sono stati obbligati, per economia di spazio, ad applicare le restrizioni imposte nei capoversi 3 e 5 del precedente paragrafo, ed hanno liberamente cambiato disposizione e forma al testo, ogni qualvolta era evidente che potesse derivarne un miglioramento. Salvo poche eccezioni, tutte indicate la forma definitiva del testo è stata sottoposta alla approvazione dell'Esperto.

noted, the final form of the rewritten text was submitted to the Expert and was accepted by him.

In preparing the numerical data for publication the Editors have made no change except in their arrangement and in their mode of presentation. In making such changes the Editors have been guided by the necessity of saving space. The numerical data are in all cases those submitted by the Expert, excepting that (a) a few additional values, all duly indicated, have been inserted, and (b) when an Expert has submitted a number of values for the same nominal quantity, these have been grouped so as to make a single entry with an indication of the range covered by the values submitted, whenever such grouping seemed justifiable. In these cases, the final manner of grouping was in every case where possible submitted to and accepted by the Expert. The exceptional cases are noted as they occur.

Owing to the method of publication, i e , one volume at a time, a strictly logical arrangement of subject matter is not always followed. Among such a large number of Cooperating Experts a few instances of greatly delayed reports, arising from illness, accident, or other unforeseen causes, are to be expected; and certain sections or parts of sections, therefore, may not appear in their logical places but will be found in a later volume. The whole set of volumes is very completely indexed, however, and the user who consults the index should have no difficulty in locating any information given.

Chemical compounds are arranged in the tables by formula according to a definite system, called the "Standard Arrangement." This system is based upon a set of key numbers for the chemical elements and is fully explained in Volume One.

In order to find a given substance in the longer tables it is therefore necessary to know its chemical formula, at least approximately. If only the name is known, the formula, for most organic compounds or minerals, may be found with the aid of the name indices in Volume One, p. 174 and 280.

Pour la préparation du texte destiné à la publication, les rédacteurs se sont vu obligés, afin d'économiser encore de la place, d'accentuer encore les restrictions imposées dans les sections 3 et 5 du paragraphe précédent et ils ont pris la liberté de ré-arranger et de ré-écrire le texte partout où il était évident qu'une compression ou une amélioration dans l'ordre logique pouvait ainsi être réalisée. A part de rares exceptions, qui sont du reste dûment notées, la forme définitive du texte ré-écrit fut soumise à l'expert et acceptée par lui.

En disposant les données numériques pour la publication, les rédacteurs n'ont fait aucune modification, excepté en ce qui concerne l'arrangement et le mode de présentation. En faisant ces changements, les rédacteurs ont été guidés par la nécessité d'épargner de la place.

Les données numériques sont dans tous les cas celles fournies par les experts, à l'exception (a) d'un petit nombre de valeurs, toutes dûment indiquées, qui ont été insérées, et (b) lorsqu'un expert à soumis un certain nombre de valeurs pour la même quantité nominale, ces valeurs ont été groupées de façon à constituer une entrée unique, avec une indication du range occupé par les valeurs fournies, toutes les fois qu'un tel groupement paraissait indiqué. Dans ces cas, la forme définitive du groupement fut, partout où cela était possible, soumise a l'expert et acceptée par lui. Les cas exceptionnels sont notés lorsqu'ils se présentent.

Etant donné le mode de publication par un volume à la fois, un arrangement strictement logique de la matière traitée n'est pas toujours possible. En effet, avec un tel nombre d'experts coopérants, il faut s'attendre à ce qu'il y ait quelques circonstances imprévues, telles que maladies, accidents ou autres causes, occasionnat un grand retard dans la remise des rapports; c'est pourquoi certaines sections ou parties de sections ne peuvent paraître à leur place logique mais se trouveront dans un volume suivant. Cependant, la série complète des volumes étant indexée d'une façon très détaillée, le lecteur qui consulte la table des matières n'aura aucune difficulté pour repérer toute information donnée.

Les composés chimiques sont disposés dans les tables suivant leurs formules et cela d'après un système défini appelé "arrangement type." Ce système est basé sur une suite de "nombres clés" pour les éléments chimiques, et il est expliqué d'une façon complète dans le volume I.

Afin de trouver une substance donnée dans les longues tables, il est nécessaire de connaître sa formule chimique au moins approximativement. Si le nom seul est connu, la formule peut être trouvée pour la plupart des composés organiques ou des minéraux au moyen des noms indices qui se trouvent dans le volume I, p. 174 et 280.

Bei der Zusammenstellung des Textes für die Veröffentlichung waren die Herausgeber gezwungen, im Interesse der Raumersparnis die unter 3 und 5 oben angegebenen Richtlinein besonders zu betonen. Sobald erkannt wurde, dass eine Zusammenziehung und eine Verbesserung in der logische Anordnung möglich sei, wurde der Text frei zusammengestellt und frisch geschrieben. Mit wenigen Ausnahmen, welche besonders bezeichnet sind, wurde die entgültige Form des neu geschriebenen Textes dem Experten vorgelegt und von ihm angenommen.

Bei der Vorbereitung des Zahlenmaterials für die Veröffentlichung änderten die Herausgeber nichts, ausgenommen war nur dessen Anordnung und die Form der Darstellung, wobei man sich von der Notwendigkeit, Raum zu sparen, leiten liess. Die Zahlenwerte sind in allen Fällen dieselben, welche vom Experten vorlagen, ausgenommen, (a) dass einige ergänzende, besonders bezeichnete Werte hinzugefügt wurden und (b), wenn der Experte für dieselbe quantitative Grösse mehrere Werte angegeben hat. Diese wurden dann, sobald ein solches Vorgehen gerechtfertigt war, zusammengestellt, so, dass nur eine Zahl, mit den Grenzen hingeschrieben werden konnte, welche durch die Werte gegeben sind. In so einem Falle wurde die Endform der Anordnung jedesmal dem Experten, wo möglich vorgelegt und von ihm angenommen. Die Ausnahmsfälle sind dorten wo sie vorgekommen bezeichnet.

Entsprechend der Publikationsmethode, der Herausgabe eines Bandes zu einer bestimmten möglichen Zeit, konnte eine genaue logische Anordnung eines bestimmten Kapitels nicht immer erreicht werden. Unter einer so grossen Zahl von Mitarbeiteren sind Fälle zu erwarten, wo sich einige Artikel stark verzögern werden, sei es durch Krankheit oder andere unvorhergesehene Ursachen. Deshalb werden gewisse Abschnitte oder deren Teile nicht an ihren richtigen Plätzen erscheinen, sonderen sie können in einem späteren Band gefunden werden. Die ganze Bänderfolge ist mit einem sehr vollständigem Verzeichnis versehen und der Leser, welcher das Verzeichnis benützt, wird keine Schwierigkeit haben, Vorhandenes aufzufinden.

Die chemischen Verbindungen sind in den Tafeln nach einem Formelsystem angeordnet, das als "Normalanordnung" (Standard Arrangement) bezeichnet wird. Dieses System, das im ersten Bande vollständig erklärt wird, beruht darauf, dass für die chemischen Elemente Schlüsselnummern gewählt werden.

Um im den längeren Tafeln eine gegebene Substanz aufzufinden, ist es notwendig, deren chemische Formel wenigstens annähernd zu kennen. Ist nur der Name bekannt, so kann die Formel der meisten organischen Verbindungen und der Minerale, mit Hilfe des englischen Namenverzeichnisses im Bande 1 Seite 174 und 280 gefunden werden.

Nell'allestire i dati numerici per la pubblicazione i Redattori hanno fatto cambiamenti solo nel modo di disporli e di presentarli. Nel fare questi cambiamenti i Redattori sono stati guidati dalla necessità di risparmiare spazio. I dati numerici sono in tutti i casi quelli forniti dall'Esperto; solo qualche volta sono stati aggiunti alcuni pochi valori, tutti bene indicati, e qualche altra, avendo l'Esperto riportato parecchi valori per una stessa grandezza, questi—allorchè è sembrato giustificato il farlo—sono stati raggruppati indicando un solo numero ed i limiti entro i quali oscillano i valori considerati. In questi casi, la disposizione finale, fu sempre, quando possibile, sottoposta all'approvazione dell' Esperto. Tutte le volte che è stato fatto diversamente, lo si è indicato.

Siccome le tabelle vengono pubblicate un volume alla volta, non sempre la disposizione della materia è fatta in modo strettamente logico.

Dato il numero grande di Esperti, è da aspettarsi che qualche rapporto sarà presentato con grande ritardo a causa di malattie o di incidenti imprevedibili. Certe parti perciò potranno comparire non nel posto che logicamente ad esse spetterebbe, ma in volumi posteriori. Tutti i volumi sono però muniti di indici accurati e il lettore, consultandoli, non avrà difficoltà a rintracciare una notizia qualunque.

I composti chimici sono disposti nelle tabelle in base alle formule seguendo un sistema chiamato "disposizione Standard." Questo sistema è fondato sopra una serie di numeri chiave assegnati agli elementi chimici ed è esaurientemente spiegato nel primo volume.

Per poter quindi trovare una data sostanza nelle tabelle più lunghe, è necessario conoscerne la formula chimica, almeno approssimativamente. Se si conosce solo il nome, la formula si può trovare (per la massima parte dei composti organici o minerali) con l'aiuto degli indici per nome contenuti nel 1° volume p. 174 e 280.

CONTENTS

MATIÈRES

INHALTSVERZEICHNIS

INDICE

COOPERATING EXPERTS—VOLUME ONE

ADAMS, LEASON H., B.S.: Physical Chemist, Geophysical Laboratory, Carnegie Institute of Washington, Washington, D. C.

ASTON, F. W, D.Sc , F.R.S , F.I.C : Fellow, Cavendish Laboratory, Trinity College, Cambridge, England.

AUGER, PIERRE: Agrégé des sciences physiques, Faculté des Sciences de Paris, France.

BANDWELL, D. C., Ph.D.: Associate Physical Chemist, Fixed Nitrogen Research Laboratory, U. S. Department of Agriculture, Washington, D. C.

BIGG, PHILIP, H., B.Sc.: Junior Assistant, The National Physical Laboratory, Teddington, Middlesex, England.

BRIGGS, LYMAN J., Ph D.: Chief, Division of Mechanics and Sound, Bureau of Standards, Washington, D. C.

CLARK, MANSFIELD, Ph.D.: Professor of Chemistry, Hygienic Laboratory, U. S. Public Health Service, Washington, D. C.

DRYDEN, HUGH L., Ph.D.: Chief, Aerodynamical Physics Section, Bureau of Standards, Washington, D. C.

DUSHMAN, SAUL, Ph.D.: Physicist, General Electric Co , Schenectady, N. Y.

FAIRCHILD, CHARLES O., M. S.: Physicist, Bureau of Standards, Washington, D. C.

FOOTE, HARRY W., Ph.D.: Professor of Chemistry, Yale University, New Haven, Connecticut.

FORSYTHE, WILLIAM E , Ph.D : Director, Nela Research Laboratory, Cleveland, Ohio.

GOULD, F. A., M.A.: Assistant, Metrology Department, The National Physical Laboratory, Teddington, Middlesex, England.

GUILLAUME, CHARLES-EDOUARD: Directeur du Bureau international des Poids et Mesures, Sèvres, France.

HANN, RAYMOND M., B.S.: Bureau of Chemistry, Washington, D. C.

HERSCHEL, WINSLOW H , A.B . Associate Physicist, Bureau of Standards, Washington, D C.

HEVESY, G., Ph.D.: Professor, Universitetets Institut for Teoretiskfysik, Copenhagen, Denmark

HOUGEN, OLAF A., Ph D.: Assistant Professor of Chemical Engineering, Univ. of Wisconsin, Madison, Wisconsin.

HUMPHREYS, WILLIAM J., Ph.D : Professor of Meteorological Physics, U. S. Weather Bureau, Washington, D. C.

JOY, ALFRED H., M.A.: Astronomer, Mount Wilson Observatory of the Carnegie Institution of Washington, Pasadena, California.

KANOLT, C. W., Ph. D : Physicist, Bureau of Standards, Washington, D. C.

KEENAN, GEORGE L , M.A.: Bureau of Chemistry, Washington, D. C.

KLEEMAN, RICHARD D., D Sc : Consulting and Research Physicist and Physical Chemist; Associate Professor of Physics, Union College, Schenectady, N. Y.

KRAMERS, H. A.: Institut for Teoretiskfysik, Copenhagen, Denmark.

LAMBERT, WALTER D., A.M : Mathematician, U. S. Coast and Geodetic Survey, Washington, D. C.

LIND, S. C., Ph D.: Assistant Director, Fixed Nitrogen Research Laboratory, U. S. Department of Agriculture, Washington, D C.

MEYER, STEFAN, Ph.D : O Ö Prof. d Physik, Universitat, Vienna, Austria.

MOORE, RICHARD B , Sc.D . General Manager, The Dorr Company, New York City, N Y.

MUELLER, EUGENE F.: Physicist; Chief, Heat Measurement Section, Bureau of Standards, Washington, D C.

PIENKOWSKY, ARTHUR T., Ph B : Associate Physicist, Bureau of Standards, Washington, D. C

RAGATZ, ROLAND A., M S : Instructor in Chemical Engineering, University of Wisconsin, Madison, Wisconsin.

RUDORF, G.: Air Ministry, Kingsway, London, England.

SCHLUNDT, HERMAN, Ph D : Chairman, Department of Chemistry, University of Missouri, Columbia, Missouri

SCHWEIDLER, EGON, Ph D O. Ö Professor der Physik, Physikalisches Institut der Universitat, Innsbruck, Austria.

SEARES, FREDERICK H , B.S.: Assistant Director, Mt. Wilson Observatory of the Carnegie Institution of Washington, Pasadena, California

SODDY, FREDERICK, M.A., F.R S.: Professor, Chemical Department, University Museum, Oxford, England.

SPENCER, HUGH M., Ph D : Instructor in Chemistry, Yale University, New Haven, Connecticut.

STOTT, VERNEY, B.A., F. Inst. P : Assistant, The National Physical Laboratory, Teddington, Middlesex, England.

STRÖMBERG, GUSTAF, Ph D.: Astronomer, Mt Wilson Observatory of the Carnegie Institution of Washington, Pasadena, California.

SWICK, CLARENCE E., C.E : Geodetic Mathematician, U S. Coast and Geodetic Survey, Washington, D. C.

VOLET, CHARLES: Licencié ès-Sciences, Adjoint au Bureau international des Poids et Mesures, Sèvres, France.

WALTON, C F., JR.: Research Chemist, Bureau of Chemistry, Washington, D. C.

WEIMARN, P. P., VON: Professor of Physical Chemistry; Mining Engineer; Head, Dispersoidological Department, Imperial Industrial Research Institute, Osaka, Japan.

WELLS, ROGER C., Ph.D.: Physical Chemist, U. S. Geological Survey, Washington, D. C.

WENSEL, H. T., Ph.D.: Associate Physicist, Bureau of Standards, Washington, D. C.

WHERRY, EDGAR T., Ph.D.: Senior Chemist, Bureau of Chemistry, Washington, D. C.

WHIPPLE, F. J. W., M.A , F. Inst. P.: Superintendent, Kew Observatory, Richmond, Surrey, England.

WOODWORTH, ROBERT S , Ph D.: Professor of Psychology, Columbia University, New York City, N. Y.

WYCKOFF, RALPH W. G.: Research Associate, Geophysical Laboratory, Carnegie Institution of Washington, Washington, D. C.

ZWAARDEMAKER, H., M.D.: Professor of Physiology, Univ. of Utrecht, Holland.

VOLUME I
ERRATA

PAGE	
xviii & xix	Ready Reference Tables. *Between* (c) and (d) *insert* Boiling points (inorganic) .. 162 To (d) *entry* add 165, 276. *Between* (d) and (e) *insert* Index to minerals 174. Index to C-Table 280
4	For Columbia read Colombia
7	Column 3, under *Mass* For 453 592 45 read 453 592 43 For 64 798 182 read 64 798 9182
8	Column 1, 1 bushel For 35 367 7048 1 read 36 367 7048 1
10	Column 1 For 1 mal = 10 s read 1 mål or maal = 9 843 or 10 s
12	Column 2 For 1 aln read 1 aln
	1 famn 1 famn
	1 stang = 16 1 stång = 10 or 16
	Column 3, under *Mass*
	For skålpund read skålpund
	1 ass 1 ass
	1 quintin 1 kvintin
	1 untz 1 uns
	1 nylast = 12 000 1 nyläst = 10 000 or 12 000
	Add 1 korn = 1/1000
	1 ort = 1/100
	Column 3, under *Capacity, dry*
	Delete 1 ort = 1/10
	For 1 junkfru read 1 jungfru or jumfru
	1 quarter 1 kvarter
	1 kappar 1 kappe
	1 fjerdingar 1 fj rding
	1 spanna 1 spann
13	Column 1, 1 saah For 1/18 read 1/8
17	Column 1, (v) For p 27 read p 38
	Accepted Basic Constants Regarding Uncertainty column *add* These values are rough estimates and those for e, e m₀ and h should probably be several times as great as the values given
18	Section A. These Derived Constants have been computed from the Accepted Basic Constants on p 17, and are vitiated by the errors in those values The greatest errors occur in ν ω and N α, which differ from the best experimental values by about 0 4 %, the computed value of ν ω being too small
	For r ω read ν ω
	Section B, log₁₀ λ For 4 808 7827 read 3 808 7827
23	Table 28, last line For 15 5951 read 14 5951. For 1 192 9882 read 1 143 3824.
26	Table 48. For 1 0000 lambert 0,000 0000
	1.0764 millilambert 0 031 9684
	read 3 1416 lambert 0 497 1499
	3 3816 millilambert 0 529 1183
34	Column 2, line 1 For 980 655 read 980 665
42	Column 1 For Synonical read Synodical
49	Column 1 For X read Xe
52	For F O Fairchild read C O Fairchild
62	Column 2. For above 20° read below 20°
66	Section (a), Phosphorus pentasulfide For 52° read 522°
91	High Vacuum Technique
	line 1, for *Amount* read *Mass*
	line 6, *after* molecules *add* striking 1 cm² sec⁻¹
	line 11, for *Q* = amount read *Q* = volume
102	The Gaseous State, viscosity column
	A, for 221 read 222
	Br, for 155 read 154.
	Air, for 284 2 read 180 8
106	Line 1. For Smithers read Smither.
	Line 3 For John C W Frazer read J C W Frazer
	Index No 6 For −76 read −59
	Index No 35 1 Delete 1 in density column for 4 182?
109	Index No 204, d₁⁰ Add 3 022
	Index No 205, d₁⁰ Add 4 49
	Index No 206, d₄⁰ Add 3 63.
110	Index No 259. Delete entry.
115	Index No. 500 For Attaite read Altaite
119	Index No. 766. Delete entry.
	Index No. 767 For 45 5 read 44 07.
122	Index No. 940 Add Eriochalcite.
	Index No. 1001. For Phosphochalite read Phosphochalcite
128	Index No. 1354. For Szomolnokite read Szomolnkite.
	Index No. 1355 For Siderotilate read Siderotilite.
129	Index No. 1394. For FeCO₃.H₂O read FeCO₃. For 133 855 read 115 84.
131	Index No 1507. For 2 818¹²⁄₁₂ read 4 13
133	Index No. 1631. For Crocotate read Crocoite.
134	Index No. 1683. Insert Tungstenite.
	Index No. 1726. For UO₃.CO₂ read UO₂ CO₃.
136	Index No 1819. For BN₃ read BN. For 38 8360 read 24 8280
139	Index No 1990. Insert Monazite.
143	Index No. 2236. For Hydrophyllite read Hydrophilite
149	Index No 2622. For d 29 6 read 29 88
152	Index No. 2807. Probably a decahydrate, v. Conroy, *54*, 17: 104; 98
153	Index No. 2877. For 3.55 read 2.55.

PAGE	
155	Index No. 2968. For 96 read 256 d.
	Index No 2994 For KC₁₀H₁₉O₂ read KC₁₀H₁₉O₃.
	Serial No 1 For 1 833 read 1 1833
165	Column 1 For Apjohnite read Apjohnite.
	Delete Arsenic siderite
174	Column 2, Automolite. For 1119 read 1911.
	Column 3 For Cerargytite read Cerargyrite
	Column 4. For Chrysotite read Chrysotile.
	For Cotinnite read Cotunnite
	Column 5 For Durfeidtite read Durfeldtite
	Column 6. For Eriochalite read Eriochalcite.
	For Ganophyllite read Ganophyllite.
175	Column 1. For Geoeronate read Geoeronite
	For Jeremejerite read Jeremejevite.
	Column 3 For Molybdophillite read Molybdophyllite
	After Molyssite insert Monazite, 1990
	Column 4 For Phosphochalite read Phosphochalcite
	Polianite For (Pyrosulite) read (Pyrolusite)
	Column 5, Scheelite For 2366 1 read 2366.
	After Spencerite insert Sperrylite, 1179
175	Column 5 For Szomolnokite read Szomolnikite
	For Tennamite read Tennantite
	Column 6. After Tsumebite insert Tungstenite, 1683.
	For Uru ilggite read Urvclgyite
176	Index No 20 Delete 2 53.
179	Index No. 232 Delete 1 7
180	Index No 263 For 1 617 read 0 6606
183	Index No 435 For NH(COCH₃)₂ read NH₂CONHCOCH₃.
187	Index No 686. For −18 read −5
188	Index No 725. For CH₃COC₃H₅ read CH₃CO₂C₃H₅.
	Index No 773 1 For Methy read Methyl
192	Index No 1012 For C₃H₅COₐH read C₃H₅CO₂H
193	Index No 1074 For Dimethy read Dimethyl
200	Index Nos 1466, 1468, and 1470. Data probably not for pure compounds o-Dihydrobenzene and 1, 3-cyclohexadiene are two names for the same compound.
211	Index No. 2328. Delete entry.
	Index No. 2330 For −126 4 read −126 3
	100 8 100 3
	0 764 0 7804?
220	Index Nos 2719, 2720, and 2721. For Creasyl read Tolyl
224	Index No 2942 For 2-Ethylhexane CH₃(C₃H₇)CHC₄H₉ read 3-Ethylhexane (C₂H₅)₂CHC₃H₇.
	Index No 2942 1 Delete entry.
227	Index No 3123 For C₉H₁₀O₄ read C₉H₁₀N₂O₄
	Index Nos 3150, 3151, and 3152 For Creasyl read Tolyl
234	Index No 3576 For C₅H₄C²H CHC₃H₅ read C₆H₅CH₂C:CCH₃.
235	Index No 3635 For Benzacetin read 5-Acetylamino-2-methoxy-benzene-1-carboxylic acid
238	Index No 3848 For (C₁₀H₁₆O) read (C₁₀H₁₆O)₂
	152 12 456 36
	295 310
	Index No. 3862. For [C₁₀H₁₆O]₂ read (C₁₀H₁₆O)₃
	[152.12]₂ 456 36
	264 285
242	Index No. 4078 Add 5-Acetylamino-2-ethoxybenzene-1-carboxylic acid
217	Index No 4394. 1 525 is the density for the monohydrate
253	Index No 4734. For Creasyl read Tolyl. For p-CH₃C₆H₄O₂CC₆H₄ read p-CH₃C₆H₄O₂CC₆H₄
254	Index Nos 4730, 4741, 4742, and 4744 For Creasyl read Tolyl .
258	Index Nos 4778, 4779, and 4780 For Creasyl read Tolyl.
	Index No. 5057. For C₁₀H₁₂N₂ read C₆H₁₂N₂. For 233.12 read 127 124
260	Index No 5152. For capronate read caproate.
262	Index No. 5291 For Chinosol read Quinosol
266	Index Nos. 5547, 5550 For Jelsemine read Gelsemine.
268	Index No. 5653. For Strychine read Strychnine.
269	Index No. 5711. For Gelsemine read Gelsemine.
270	Index No. 5779. For o-Cresol read o-Tritolyl.
271	Index No. 5902. Delete entry
272	Index No. 5928. For C₁₁H₃₂O₁₆ read C₁₁H₃₀O₁₆.
	612 25 610 23
	183 190
	Index No. 5967. For Octocosane read Octacosane.
274	Index No. 6054. For capronate read caproate.
	Index No 6082. For Filixic read Filicic.
	Index No. 6096. Delete entry
275	Index No 6110 For caprinate read caprate.
278	Serial No 910. Delete entry.
280	Column 3 After p-Acetylaminobenzoic acid insert 5-Acetylamino-2-ethoxybenzene-1-carboxylic acid, 4078.
	Column 4 After o-Acetylaminomethoxybenzene insert 5-Acetylamino-2-methoxybenzene-1-carboxylic acid, 3635.
	Column 4, Benzacetin Delete 3635.
282	Column 1. For Cerebrin, 5931, 6153 read Cerberin, 5931
285	Cerebrin, 6153

1

PAGE

286 Column 3. Delete 14 entries, beginning with o-Cresyl acetate 3150 and ending with p-Cresyl salicylate, 4743 Delete o-Cresol orthoacetate, 5779
Column 4. *After* m-Cymene *insert* p-Cymene, 3728 1

290 Column 4, Ergosterol *Delete* 5002
291 Column 3 *Delete* 2 Ethylhexane, 2942
3-Ethylhexane *For* 2942 1 *read* 2942
Ethyl hippurate *For* 4516 *read* 4077 1

292 Column 2, Fibric acid *For* 6896 *read* 6082
Fibixic acid Delete entry
Column 3. *For* Gelsemine, 5711 *read* Gelsemine, 5547
Gelsemine, 5711

295 Column 4, Jelsemine, 5547 and Jelsemine hydrochloride, 5550
Delete entries

300 Column 4, N-Phenylthiourethane *For* 321 *read* 3201
301 Column 4, Pyrene *For* 5206 *read* 5026
302 Column 2 *After* Quinoneoxime insert Quinosol, 5291
303 Column 1, Terephthalic acid Delete entry.
304 Column 2. *After* Toluylene-3, 5-diamine insert
o-Tolyl acetate, 3150
m-Tolyl acetate, 3151
p-Tolyl acetate, 3152
After p-Tolylantipyrene insert
m-Tolyl benzoate, 4736
p-Tolyl benzoate, 4734
After p-Tolyldimethylpyrazolone insert
o-Tolyl ether, 4778
m-Tolyl ether, 4779
p-Tolyl ether, 4780
After p-Tolyl isothiocyanate insert
o-Tolyl methyl ether, 2719
m-Tolyl methyl ether, 2720
p-Tolyl methyl ether, 2721
After p-Tolyl mustard oil insert
o-Tolyl salicylate, 4741
m-Tolyl salicylate, 4742
p-Tolyl salicylate, 4743

305 Column 3. *After* Tritluoglycerol insert o-Tritolyl orthoacetate, 5779.

306 Column 2, Xanthine. Delete entry
Property-Substance Tables, **—180.** *Delete* 2328.

307 **60:** *For* 1516 *read* 4077 1.
308 **96:** *Delete* 32908.
116: *After* 33347 insert 07.
310 **288:** *After* 4931 insert 32968.
261: *Delete* 3862.
288: *After* 2620 insert 3862.
292: *Delete* 3848
310: *After* 1385 insert 3846.
311 **98:** *Delete* 171
106: *Delete* 2328
116: *Delete* 2942.1.
116: *Delete* 67.
180: *For* 1516 *read* 4077 1

313 **0.760:** *Delete* 2328 and 2430
319 C₆H₁₂O₂ *For* caproate *read* caproate
C₆H₁₂O₂ *For* caprinate *read* caprate

322 SrC₂H₄(SO₃)₂·H₂O. *For* Strontium disulfonate *read* Strontium ethane disulfonate

331 C₁₁H₁₅O₂N *For* Glutaric anilide *read* Glutaranilide
353 (**111**), Sasahara *For* 329 read 210
358 Odoriferous Materials, Classification
For fragrance 1* read fragrance
Allyl — Alliaceous
empyreumatic — empyreumatic
tetri — tebri
nauseois — nauseous

360 Column 1, line 3 *For* 6.06 × 10³¹ *read* 6.06 × 10³³
Columns 1 and 2, table heading *For* Molecules per cc *read* Molecules per 0 01 cm³

PAGE

362 Column 2, line 8. *For* V = 0.0342a^{1/2} *read* V = 0.0342a^{1/4}.
Above Remarks Concerning the Nomenclature there should be a rule extending across entire page.
Column 2, Note 9. *For* DAI *read* DAI.
363 Series of Thorium, Thoron. *For* 0.574 *read* 0.0574
364 Column 2, Tables (b) and (c) *For* cm⁻² *read* cm³.
366 Chemical Effects of α-Particles, column 1, line 3 *After* α-particles insert in the time the M are reacting
equation, line 12. *For* ln *read* log₀

368 Column 1, line 4 *For* T_{1_2} *read* T (cf. p. 362, column 1, line 14).

372 Column 2, Literature
(⁴). *For* 181:1751 *read* 180:1750.
(¹⁰). 10:11 11:828
Lit. column. *For* (⁹⁹) *read* (⁹⁶).
Lit. column. *For* (⁹⁷) *read* (⁹⁸).
Column 2. *For* Japan (⁴²) *read* Japan (⁴¹).
Column 2, *Hokutolite* *For* (⁴²) *read* (⁴¹).
Column 2, *Skaldowskite* *read* Skiodowskite.
Column 2, *Torbernite* *For* (CCaPO₄aq) *read* (CCuPO₄aq)
Column 2, **Y** *For* Yttrotantalite *read* Yttrotantalite.

379 Oceanic Deposits Data from (¹³⁸) have been superseded by the author's later work (Joly, 3, **24**:094,12) and should read.

	No. speci-mens	R... mea...
Blue mud		
1240 fath	1	1.5
"Ooze"		
720 fath . . .	1	1.7
Globigerina ooze		
199 to 2493 fath .	1	3 3
3 of above sampl s		3 1
Radiolarian ooze		
2600 to 2750 fath	2	13 1
Red clay		
2350 fath ...	1	11 0

380 The Loetschberg Tunnel. *For* Aplete *read* Aphte.
Meteorites, Remarks *For* hexahydrite *read* hexahedrite
381 Column 1, line 2 of table *For* Anondoga *read* Onondaga
Column 2, (¹³⁸). *For* Joly, 3, **16**:190,18 *read* Joly, 3, **16**:190;08.
382 Lines 2 and 3 of table *For* Felixtowe *read* Felixstowe.
Line 16 of table *For* Frier *read* Friar.
Line 29 of table. *For* Charnokite *read* Charnockite
392 Characteristics of Members of Solar System
Sidereal rotation of Sun *For* 25 3 *read* 25 0
Number of satellites   Mars *For* 0 *read* 2
               Jupiter   7   9
               Saturn   9   10
Column 1, line 2 bottom. *For* 24 da *and* 30 da *read* 24 5 da *and* 30 6 da, respectively.
Column 2, Constant of notation *For* notation *read* nutation.
Column 2, Constant of aberration. Add this note. Astronomers now generally accept a value near 20 52, but the Paris conference value is used in the computation of the national ephemerides.
Column 2, Solar parallax Add this note. The direct determination (8 806″) is by far the most reliable, the one from the velocity of light is based upon the value for the constant of aberration adopted at the Paris conference of 1896, which is smaller than the value now generally accepted The two others are from the nature of the case somewhat uncertain
Column 2, Inclination of Moon's orbit to ecliptic. *For* about 5° *read* 5° 8′ 43″
394 Table 1, item 6 *For* meridonal *read* meridional
398 Column 1, Greenwich, g *For* 981 184 *read* 981 188
Column 1, Kew, g *For* 981 144 *read* 981 201

INTERNATIONAL CRITICAL TABLES

NATIONAL AND LOCAL SYSTEMS OF WEIGHTS AND MEASURES

Charles-Édouard Guillaume and Charles Volet

Plan.—Section A: International Metric System; list of countries in which its use was compulsory on January 1, 1925, list of those in which its use was either legally optional or partially compulsory on same date.

Section B: Other modern systems; the more important units at present in use or in use before adoption of metric system.

Section C: Weights and measures of antiquity.

Style and Abbreviations.—Only the singular number of the names of the units are used; ten meters will appear as 10 meter. Units of area and of volume will be written in the form centimeter² (= cm²) and centimeter³ (= cm³), respectively.

i.	Value given is only approximate.
h.	Units have changed from time to time.
m²	Square centimeter — centimètre carré = Quadrat-zentimeter — centimetro quadrato.
urrent	Units, other than metric, which are now in use; some of the units included in this class are practically obsolete. (*See* Local.)
ocal	Units of local or native origin or derivation which are in use, but which are embraced neither by the metric system nor by that of the central government. Applies mainly to colonial possessions (*See* Current.)
n³	Cubic meter = mètre cube — Kubikmeter — metro cubico.
↑ c.	International metric system compulsory since
↓o.	International metric system legally optional since
ilder	Units used before adoption of international metric system.
ilder =	The older units were those of . .
provincial	Units vary from one province or city to another
ince	. = . . Since the units have been the same as those of . .
i.	*Vide* = see.
rar.	Units are variable, not rigidly defined.

A. INTERNATIONAL METRIC SYSTEM

The decimal metric system, established in France by the Loi du 7 Avril, 1795, and represented by standards deposited in the Archives de France, became international on May 20, 1875, by the action of the Convention Internationale du Mètre. The new standards, of platinum-iridium, constructed at that time and serving as the basis of the international system, were copied from those of the Archives.

On January 1, 1925, the metric system was compulsory in:

Algeria	Greece	Peru
Allemagne	Guam	Poland
Argentina	Guatemala	Porto Rico
Austria	Haiti	Portugal and colonies
Autriche	Holland	Rumania
Belgium	Honduras	Russia
Bolivia	Hungary	Salvador
Brazil	Ireland	Schweden
Bulgaria	Italy & colonies	Schweiz
Chile	Japan	Serbie-Croatie-Slovénie
Colombia	Kolumbien	Seychelles Islands
Congo, Belgian	Kongo, Belgisch	Siam
Costa Rica	Kuba	Spain
Cuba	Luxemburg	Suddo
Czechoslovakia	Malta	Suisse
Denmark	Mauritius	Svézia
Deutschland	Mexico	Svizzera
Ecuador	Netherlands & colonies	Sweden
Equateur	Nicaragua	Switzerland
Espagne	Norway	Tchécoslovaquie
Filippine	Olanda	Tunis
Finland	Österreich	Ungarn
France	Panama	Ungheria
Germany	Pay-Bas & colonies	Uruguay
Gioppône	Philippine Islands	Venezuela
		Yugoslavia

On the same date, it was legally optional or partially compulsory in:

Canada	Great Britian	Irish Free State
China	India, British	Paraguay
Egypt		Turkey
Ethiopia	Ireland, Northern	United States of America

The fundamental units are: METER (m), which is the distance at 0°C between the axes of two lines ruled on the prototype deposited at the Bureau international des Poids et Mesures, Sèvres, France; KILOGRAM (kg), which is the mass of the prototype deposited at the same Bureau; and LITER (l), which is the volume of one kilogram of pure water at the temperature of its maximum density, under the pressure of one normal atmosphere.[1]

The primary units of the system are the *meter* (m), *micron* (μ) = 10^{-6} meter, *gram* (g) = 10^{-3} kilogram, *liter* (l), *are* (a) = area of a square with a side 10 meter long, and *stere* (s) = volume of a cube with an edge one meter long. The units of area [of volume], characterized by the adjective square [cubic], are *not* derived from a primary unit, but are each defined as the area [volume] of a square [cube] with side [edge] equal to the stated unit of length. The names of other secondary units are formed by attaching to the name of a primary unit certain prefixes of unvarying significance.

[1] Normal atmosphere, *v* p 18.

1

Secondary units.

LENGTH m = meter	
μ	micron* = 10⁻⁶ m
mm	millimeter = 10⁻³ m
cm	centimeter = 10⁻² m
dm	decimeter = 10⁻¹ m
dkm	dekameter = 10 m
hm	hectometer = 10² m
km	kilometer = 10³ m
Mm	myriameter = 10⁴ m
	megameter = 10⁶ m

* mμ millimicron = 10⁻⁹ m   μμ micromicron = 10⁻¹² m

MASS g = gram	
μg*	microgram = 10⁻⁶ g
mg	milligram = 10⁻³ g
cg	centigram = 10⁻² g
dg	decigram = 10⁻¹ g
dkg	dekagram = 10 g
hg	hectogram = 10² g
kg	kilogram = 10³ g
q	metric quintal · 10² kg = 10⁵ g
t	metric ton · 10³ kg = 10⁶ g
c	metric carat = 200 mg

* Symbol γ also used

CAPACITY l = liter = 1 000 027 dm³	
μl*	microliter = 10⁻⁶ l
ml	milliliter = 10⁻³ l
cl	centiliter = 10⁻² l
dl	deciliter = 10⁻¹ l
dkl	dekaliter = 10 l
hl	hectoliter = 10² l

* Symbol λ also used

AREA m² = square meter	
mm²	square millimeter = 10⁻⁶ m²
cm²	square centimeter = 10⁻⁴ m²
dm²	square decimeter = 10⁻² m²
a	are = 10² m²
ha	hectare = 10² a
km²	square kilometer = 10⁶ m²

VOLUME m³ = cubic meter	
mm³	cubic millimeter = 10⁻⁹ m³
cm³	cubic centimeter = 10⁻⁶ m³
dm³	cubic decimeter = 10⁻³ m³
km³	cubic kilometer = 10⁹ m³
ds	decistere = 0.1 s = 10⁻¹ m³
s	stere = 1 m³
dks	dekastere = 10 s = 10 m³

B. MODERN SYSTEMS

Abyssinia.—var.: current, ca.

Length

1 pic	= 0.686 m
1 farsang	= 5.07 km
1 berri	= ⅔ farsang

Mass

1 rottolo	= 311 g
Unit	Rottolo
1 drachm =	} 1/16
1 derime =	

1 wakea	= 1/12
1 mocha	= 1/10

Capacity, dry

1 madega	= 0.44 l
1 ardeb	= 10 or 24 madega

Capacity, liquid

1 kuba	= 1.016 l

Ägypten v. Egypt.
Äthiopien v. Ethiopia.
Algeria.—Since 1843 =
France. Older:

Length

1 pic (dzera à torky)	= 0.640 m
1 pic (dzera à rabry)	= 0.480 m
Unit	Pic
1 termin	= ⅛
1 rebia	= ¼
1 nus	= ½

Mass

1 ukkia	= 34.13 g
1 metical	= ca. 4.7 g
Unit	Ukkia
1 rottolo à thary	= 16
1 rottolo à khadhary	= 18
1 rottolo à kebyr	= 24
1 cantar	= 100 rottolo

Capacity, dry

1 caffiso	= 317.47 l
1 saah	= 58 l
1 tarri	= 1/16 caffiso

Capacity, liquid

1 khoull	= 16⅔ l or 16 l

Allemagne v. Germany.
Anam.—var.: ch., current:*

Length

1 thuoc moc	= 0.425 m
1 thuoc de ruong	= 0.470 m
1 thuoc vai	= 0.644 m
Unit	Thuoc
1 ly	= 0.001
1 phan	= 0.01
1 tat	= 0.1
1 tam }	= 5
1 ngu }	
1 truong	= 10
1 sao	= 15
1 chai vai }	= 30
1 that }	
1 mao	= 150
1 gon	= 300

Mass

1 dong	= 3.775 g
1 picul	= 60 kg
Unit	Dong
1 hao	= 0.001
1 li	= 0.01
1 fan	= 0.1
1 luong	= 10
1 neu	= 100
1 can	= 160
1 yen	= 1600
1 binh	= 8000
1 ta	= 16 000
1 quan	= 18 000

Area

1 ngu²	= 4.5156 m²
Unit	Ngu²
1 thuoc	= 6
1 sao	= 90

* By an ordinance of 1872, units were defined in terms of metric.

Unit	Ngu²
1 mau	= 900
1 quo	= 1800

Capacity

1 hao or shita	= 28.26 l
1 tao	= 2 hao

Angola.—m.c. 1910.
Arabia.—Provincial, current

Length

1 covid	= 0.482 m
1 guz	= 0.635 m
1 cassaba	= 3.84 m
1 farsakh	= 4.83 km
Unit	Farsakh
1 baryd	= 4
1 marhala	= 8

Mass

1 maund	= 1350 g
1 ratl	= ca. 460 g
Unit	Maund
1 coffilas	= 1/400
1 vakias }	= 1/40
1 tukeas }	
1 farzil }	= 10
1 farecella }	
1 bahar }	= 150
1 bokard }	

Capacity, dry

1 téman	= 85 l
Unit	Téman
1 mecmeda }	= 1/40
1 kella }	
1 mec dema	= 1/80

Capacity, liquid

1 nusfiah	= 0.79 l or = 0.95 l
Unit	Nusfiah
1 vakia	= 1/6
1 cuddy	= 4
1 zudda	= 8

Argentine Republic.—m.c. 1887; m.o. 1863. Older,* provincial:

Length

1 vara	= 0.8666 m
Unit	Vara
1 linéa	= 1/432
1 pulgada	= 1/36
1 pié	= ⅓
1 braza	= 2
1 cuadra	= 150
1 legua	= 6000

Mass

1 libra†	= 459.4 g
Unit	Libra
1 grano	= 1/9216
1 adarme	= 1/256
1 onza	= 1/16

* National system derived from old Spanish. Units given are those of province of Buenos Aires.

† 1 libra de farmacia = ¾ libra = 344.5 g.

Unit	Libra
1 arroba	= 25
1 quintal	= 100
1 tonelada	= 2000

Area

1 vara² = 0.75 m²

Capacity, dry

1 fanega = 137.1977 l

Unit	Fanega
1 cuartilla	= ¼
1 tonelada	= 7.5
1 lastre	= 15

Capacity, liquid

1 frasco = 2.375 l

Unit	Frasco
1 octava	= $\frac{1}{16}$
1 cuarta	= ¼
1 baril	= 32
1 cuarter	= 48
1 pipa	= 192

Austria.—m.c. 1876; m.o. 1873. Older:

Length

1 Fuss* = 0.316 08 m
1 Ell = 0.7792 m

Unit	Fuss
1 Punkt	= $\frac{1}{1728}$
1 Linie	= $\frac{1}{144}$
1 Zoll	= $\frac{1}{12}$
1 Klafter	= 6
1 Meile	= 24 000

Mass, (1) ordinary

1 Pfund = 560.01 g

Unit	Pfund
1 Pfennig } 1 Denat }	= $\frac{1}{512}$
1 Quentchen	= $\frac{1}{128}$
1 Loth	= $\frac{1}{32}$
1 Unze	= $\frac{1}{16}$
1 Vierding	= ¼
1 Mark	= ½
1 Stein	= 20
1 Zentner	= 100
1 Saum	= 275
1 Karch	= 400

Mass, (2) for drugs

1 Pfund apoth. = ¾ Pfund = 420.01 g

Unit	Pfund apoth.
1 Gran	= $\frac{1}{5760}$
1 Scrupel	= $\frac{1}{288}$
1 Drachme	= $\frac{1}{96}$
1 Unze	= $\frac{1}{12}$

Area

1 Joch = 1600 Klafter² = 57.557 a
1 Metze = ⅓ Joch

* Vienna.

Capacity, dry

1 Metze = 61.489 l

Unit	Metze
1 Probmetze	= $\frac{1}{1024}$
1 Becher	= $\frac{1}{128}$
1 Futtermassel	= $\frac{1}{32}$
1 Muthmassel	= $\frac{1}{16}$
1 Achtel	= ⅛
1 Viertel	= ¼
1 Muth	= 30

Capacity, liquid

1 Mass = 1.4151 l

Unit	Mass
1 Pfiff	= ¼
1 Seidel	= ½
1 Halbe	= ½
1 Viertel	= 10
1 Eimer	= 40
1 Fass	= 400
1 Dreiling	= 1200
1 Fuder	= 1280

Balearic Islands.—v. Spain. Local:

Length

1 canna = 1.564 m
1 palmos = ⅛ canna

Mass

1 rottolo = 408 g

Unit	Rottolo
1 libra major	= 3
1 corta	= 9
1 quartano	= 9
1 arroba	= 26
1 misura	= 36
1 cantaro barbaresco	= 100
1 cantaro	= 104
1 cargo	= 312

Capacity, dry

1 quartera = 71.97 l

Unit	Quartera
1 barcella	= ⅙
1 almude	= $\frac{1}{36}$

Capacity, liquid

1 quartin = 27.14 l

Unit	Quartin
1 quarte	= $\frac{1}{13}$
1 quarta	= $\frac{1}{26}$

Bavaria v. Germany.

Belgian Congo — m.c. 1911.

Belgium. —m.c. 1820; at first with the names· aune = m, litron = l, livre = kg, once = hg, lood = dg, wigtje = g, Older:

Length

1 perche = 0.497 m
1 pied = $\frac{1}{10}$ perche

Mass

1 livre = 489.5 g

Unit	Livre
1 loth	= $\frac{1}{32}$
1 once	= $\frac{1}{16}$
1 marc	= ½
1 stein	= 8
1 quintal	= 100
1 chariot	= 165
1 balle	= 200
1 schiffpfund	= 300
1 charge	= 400

Area

1 arpent = 400 perche² = 130.6 a

Birmanie v. British India, Rangoon.

Bolivia.—m.c. 1893; m.o. 1871. Older = Spain.

Brazil. m.c. 1862 Older *

Length

1 pé = 0.33 m

Unit	Pé
1 palmo	= ⅔
1 vara	= 3⅓
1 passo geometrico	= 5
1 braca	= 6⅔
1 legoa	= 20 000

Mass

1 libra = 459.05 g

Unit	Libra
1 onza	= $\frac{1}{16}$
1 marco	= ½
1 arroba†	= 32
1 quintal	= 128
1 tonelada	= 1728

Area

1 tarefa = 30 to 46 a
1 alqueire = 242 or 484 a

Capacity

1 almude = 31.944 l
1 alqueire = 40 to 320 l

Unit	Almude
1 canada	= $\frac{1}{12}$
1 pipa	= 15
1 tonel	= 30

Britain, British v. Great Britain.

British India.—m.o. 1920. Current: British and local. Local,‡ provincial·

BOMBAY.

Length

1 guz = 0.6858 m

Unit	Guz
1 tassoos	= $\frac{1}{24}$

* Those of Portugal, with notable local differences
† 1 arroba metrica = 15 kg
‡ Local or national measures are now defined by their equivalents in British units

Unit	Guz
1 hath } 1 covid } 1 cubit }	= ⅔

Mass

1 seer = 317.5147 g

Unit	Seer
1 tank	= $\frac{1}{12}$
1 pice } 1 parah }	= $\frac{1}{30}$ or $\frac{1}{18}$
1 maund	= 40
1 candy	= 800

Area

Unit	Are
1 ground	= 2.03
1 biggah	= 24.68
1 kani	= 30.75
1 cawnie	= 54
1 chahar	= 2902

Capacity

1 parah = 110.1 l

Unit	Parah
1 tipree	= $\frac{1}{128}$
1 seer	= $\frac{1}{64}$
1 adoulie	= $\frac{1}{16}$
1 candy	= 8
1 garce	= 80

CALCUTTA.

Length

1 guz* = 0.9144 m

Unit	Guz
1 jaob } 1 jow }	= $\frac{1}{144}$
1 unglee	= ⅛
1 moot	= $\frac{1}{12}$
1 span	= ¼
1 covid } 1 haut }	= ½
1 danda	= 2
1 niranga	= 10
1 coss	= 2000

Mass

1 seer = 933.04 g

Unit	Seer
1 ruttee	= $\frac{1}{7680}$
1 masha	= $\frac{1}{960}$
1 tolah } 1 sicca }	= $\frac{1}{80}$
1 chittack	= $\frac{1}{16}$
1 pouah	= ¼
1 raik	= ½
1 pally } 1 dhurra }	= 5
1 maund (bazar)	= 40

Area

1 guz² = 0.836126 m²

Unit	Guz²
1 chattack	= 5
1 cottah	= 80
1 biggah	= 1600
1 tenab	= 2500

* Old guz = 0.915 m.

British India.—*Cont'd.*

Capacity

1 pally = 5.0 to 5.5 l

Unit	Pally
1 chattack	= $\frac{1}{80}$
1 khoonke	= $\frac{1}{64}$
1 kunk	= $\frac{1}{16}$
1 raik	= $\frac{1}{4}$
1 soally	= 20
1 khahoon	= 320

CEYLON.

Length

1 covid = 0.464 m

Mass

1 candy }
1 bahar } = 226.8 kg

Capacity

1 ammonam = 203.4 l

Unit	Ammonam
1 parrah	= $\frac{1}{4}$
1 seer	= $\frac{1}{2\cdot88}$

MADRAS.

Length

1 covid = 0.472 m

Mass

1 seer = 283.495 g

1 cafh = 1 230.147 mg

Unit	Cafh
1 fanam	= 80
1 pagoda	= 2880

Unit	Seer
1 pagoda	= $\frac{1}{10}$
1 pollam	= $\frac{1}{8}$
1 varahan	
1 powe	= $\frac{1}{4}$
1 vis	= 5
1 maund	= 40
1 candy	= 800

Area

1 cawnie = 53.11 a

1 maoney = $\frac{1}{24}$ cawnie

Capacity

1 puddy = 1.533 l

Unit	Puddy
1 olluck	= $\frac{1}{8}$
1 measure	= 1
1 mareal	= 8
1 parah	= 40
1 garce	= 3200

RANGOON.

Length

1 sandong = 0.5588 m

Unit	Sandong
1 palgat	= $\frac{1}{2\cdot2}$
1 taim	
1 cubit	= $\frac{9}{11}$
1 lan	= 4
1 bamboo	
1 dha	= 7
1 oke thapal	= 140
1 dain	= 7000

Mass

1 tical = 16.32 g

Unit	Tical
1 ruay	= $\frac{1}{64}$
1 pai	= $\frac{1}{16}$
1 moo	= $\frac{1}{4}$
1 mat	= $\frac{1}{2}$
1 cattie	= $33\frac{1}{3}$
1 viss	= 100
1 candy	= 15 000

Capacity

1 byee = 0.505 l

Unit	Byee
1 lamany	= $\frac{1}{8}$
1 zalay	= $\frac{1}{2}$
1 zayoot	= 2
1 seit	= 4
1 kwai	= 8

STRAITS SETTLEMENTS.

Mass

1 kati = 604.79 g

Unit	Kati
1 tahil	= $\frac{1}{16}$
1 pikul	= 100
1 bhara	= 300
1 koyan	= 4000

Capacity

1 gantang* = 4 545.96 l

Unit	Gantang
1 para	= 10
1 koyan	= 800

Bulgaria. m.c. 1892.

Burma *v.* British India.

Cambodia *v.* Indo-China.

Canada.—m.o. 1871. Current = British,† French names are:

Length

1 pouce	= 1 inch
1 chainon	= 1 link
1 pied	= 1 foot
1 verge	= 1 yard
1 perche	= 1 rod, pole
1 chaine	= 1 chain‡

Mass

1 livre	= 1 pound av.
1 cent	
1 quintal	= 1 hundred weight
1 tonneau	= 1 short ton

Area

1 arpent = 34.196 a

Capacity

1 pinte	= 1 quart
1 chopine	= 1 pint
1 boisseau	= 8 gallons
1 minot	= 39.025 l

*Gantang = British gallon

† Old French measures have been used, but only minot and arpent are now in use.

‡ Gunther's.

Ceylon *v.* British India.

Chile.—m.c. 1848. Older were from Spanish; legal values:

Length

1 bara = 0.836 m

Unit	Bara
1 linea	= $\frac{1}{432}$
1 pulgada	= $\frac{1}{36}$
1 pié	= $\frac{1}{3}$
1 cuadra	= 150
1 legua	= 5400

Mass

1 libra = 460.093 g

Unit	Libra
1 granos	= $\frac{1}{9216}$
1 adarme	= $\frac{1}{256}$
1 castellano	= $\frac{1}{100}$
1 onza	= $\frac{1}{16}$
1 arroba	= 25
1 quintale	= 100

Area

1 bara² = 0.698 896 m²

Capacity, dry

1 almude	= 8.083 l
1 fanega	= 12 almude

Capacity, liquid

1 cuartillo	= 1.111 l
1 arroba	= 32 cuartillo

China.—m.o. 1903 with the following names:

Length

kilometer	= sin li
hectometer	= sin yin
dekameter	= sin tchang
meter	= sin tchi
decimeter	= sin tshwen
centimeter	= sin fen
millimeter	= sin li

Area

hectare	= sin khing
are	= sin meou
centare	= sin li

Capacity

kiloliter	= sin ping
hectoliter	= sin chi
dekaliter	= sin teou
liter	= sin cheng
deciliter	= sin ho
centiliter	= sin cho
milliliter	= sin tshwo

Great diversity in national system; since 1908, defined by metric equivalents. (The orthography here employed is arbitrary; there is diversity in provincial pronunciation.)

Length

1 tchi = 0.32 m

Unit	Tchi
1 hoé	= 10⁻⁶
1 su	= 10⁻⁴

Unit	Tchi
1 hao	= 10^{-4}
1 li	= 10^{-3}
1 fen	= 10^{-2}
1 tsouen	= 10^{-1}
1 pou	= 5
1 tchang	= 10
1 yin	
1 van	= 100
1 fen	= 120
1 kyo	= 300
1 li	= 1800
1 poû	= 18 000
1 thsan	= 144 000
1 tou	= 450 000

Mass

1 liang = 37.301 g

Unit	Liang
1 hao	= 0.0001
1 li	= 0.001
1 fen	= 0.01
1 tsien	= 0.1
1 km	
1 tchin	= 16
1 kwan	= 480
1 tan	= 1600
1 shih	= 1920

Area

1 meou = 6000 tchi² = 614.4 m²

Unit	Meou
1 hao	= $\frac{1}{10000}$
1 pou²	
1 kung	= $\frac{1}{240}$
1 lvi	= $\frac{1}{160}$
1 fen	= $\frac{1}{10}$
1 kish	= $\frac{1}{4}$
1 king	= 10
1 ching	= 100

Volume

1 tchi³ = 32.768 dm³

1 ma	
1 fang	= 100 tchi³

Capacity

1 cheng = 1 035.44 l

Unit	Cheng
1 quei	= 0.0001
1 ço	= 0.001
1 chao	= 0.01
1 yo	= 0.5
1 khô	= 0.1
1 to	= 10
1 hou	= 50
1 chet	
1 sei	= 100
1 ping	= 500

Capacity, liquid

Liquids are measured by weight.

Chypre, Cipro *v.* Cyprus.

Cochin-China *v.* Indo-China.

Columbia.—m.c. 1854, but following, derived from metric system, are current:

Length

Unit	
1 vara	= 0.8 m
Unit	Vara
1 pulgada	= $\frac{1}{37}$
1 cuarta	= $\frac{1}{4}$
1 cuadra	= 100
1 legua	= 6250

Mass

Unit	
1 libra	= 500 g
Unit	Libra
1 onza	= $\frac{1}{16}$
1 arroba	= 25
1 quintal	= 100
1 saco	= 125
1 carga	= 250
1 tonelada	= 2000

Area

1 vara²	= 0.64 m²
1 fanegada	= 10 000 vara²

Cirénaïque v. Tripoli.
Congo, Belgian.—m c. 1911.
Costa Rica, Guatemala, Honduras, Nicaragua, Salvador.—m.c. 1912 by a joint convention; in partial use at earlier dates. Older (modified Spanish, English, and local):

Length

1 vara	= 0.8393 m (Costa Rica)
	= 0.8359 m (Guatemala)
	= 0.8128 m (Honduras)
Unit	Vara
1 cuarta	= $\frac{1}{4}$
1 tercia	= $\frac{1}{3}$
1 mecate	= 24

Mass

1 caja	= 16 kg
1 fanega	= 92 kg
1 carga	= 161 kg

Area

1 manzana	= 10 000 vara²
	= 6960.5 m² (Costa Rica)
	= 6987.4 m² (Guatemala)
	= 6987.4 m² (Nicaragua)
1 caballeria	= 64 manzana

Capacity

1 botella	= 0.63 to 0.67 l
1 cajuela	= 16.6 l

Cuartillo is very variable.
Cuba.—m.c. 1858, but others (old Spanish, American, and local) are current:

Mass

1 tonelada	= 1015.65 kg
1 tercio	= 72.22 kg

Area

1 caballeria Cubana	= 1342.02 a
1 cordele	= $\frac{1}{324}$ caballeria

Capacity

1 bocoy	= 136.27 l
1 barrile	= $\frac{1}{4}$ bocoy

Cyprus.—British system. Accepted equivalents:

Length

1 pic	= 2 foot
	= 0.6096 m

Mass

1 oke	$\{$ = 2.8 pound av
	= 1270.06 g
1 moosa *	= 50 700 g
Unit	Oke
1 drachme	= $\frac{1}{400}$
1 rottolo	= 0.44
1 stone	= 5
1 kantar	= 44
1 kantar (Aleppo)	= 180
1 ton	= 800

Area

1 donum	$\{$ = 1600 yard²
	= 13.378 a
1 scala	= 1 donum

Capacity

1 oke	= 1.278 55 l
1 cass	= 4.73 l
1 kile†	= 36.368 l
1 medimno	= 75.05 l
1 kartos	= 4 oke
1 kouza	= 8 oke
1 gomari	= 128 oke

Cyrenaica v. Tripoli.
Czechoslovakia.—m.c. 1876 ‡ Local:

Length

1 latro	= 1.917 m
BOHEMIA.	
1 stopa§	= 0.296 m
1 sah	= 1.778 m
1 mile	= 7.003 km
PRAGUE	
1 loket	= 0.593 m
MORAVIA.	
1 stopa§	= 0.284 m
1 loket	= 0.594 m
SILESIA.	
1 loket	= 0.579 m
1 mile	= 6.483 km

Area

BOHEMIA.	
1 merice	= 19.99 a
1 korec	$\}$
1 strych	= 28.78 a
1 mira	
Unit	Korec
1 jitro	= 2
1 lan	= 60

* Moosa = hundredweight
† Kile = bushel.
‡ Old Vienna (v Austria) and some local measures were still in use when the state was established
§ Stopa = strevic.

Capacity

1 merice*	= 70.6 l
1 korec $\}$	
1 strych $\}$	= 93.592 l

Denmark.—m.c. 1912; m.o. 1910. Older:

Length

1 fod	= 0.313 857 m
Unit	Fod
1 linie	= $\frac{1}{144}$
1 tomme	= $\frac{1}{12}$
1 aln	= 2
1 faon, favn	= 6
1 ruthe	= 10
1 mil	= 24 000

Mass

1 pund	= 500 g
Unit	Pund
1 es	= $\frac{1}{9152}$
1 ort	= $\frac{1}{512}$
1 quintin	= $\frac{1}{128}$
1 loth	= $\frac{1}{32}$
1 unze	= $\frac{1}{16}$
1 mark	= $\frac{1}{2}$
1 bismerpund	= 12
1 lispund	= 16
1 wog	$\}$ = 36
1 waag	
1 quintal	$\}$ = 100
1 centner	
1 skippund	= 320
1 skyplast	= 5200
1 quint	= 0.1
1 ort	= 0.01
1 kvint	= 0.001

Area

1 tondeland	= 55.162 a
1 tonde	= 283.69 a
Unit	Tonde
1 penge	= $\frac{1}{3}\frac{1}{8}$
1 album	= $\frac{1}{9}\frac{1}{6}$
1 fjerdingkar	= $\frac{1}{3}\frac{1}{2}$
1 skiepper	= $\frac{1}{4}$
1 pflug	= 32

Capacity, dry

1 korntonde	= 139.12 l
Unit	Korntonde
1 pott	= $\frac{1}{144}$
1 achtel	= $\frac{1}{64}$
1 viertel	= $\frac{1}{32}$
1 skieppe $\}$	= $\frac{1}{8}$
1 ottingkar $\}$	
1 fjerdingkar	= $\frac{1}{4}$
1 last	= 22

Capacity, liquid

1 pott	= 0.9661 l
Unit	Pott
1 paegel	= $\frac{1}{4}$
1 kande	= 2
1 stubchen	= 4

* Moravian.

Unit	Pott
1 viertel	= 8
1 fod³	= 32
1 anker*	= 40
1 ohm*	= 160
1 oxhoft*	= 240
1 pipe*	= 480
1 fuder*	= 960

Deutschland v. Germany.
Dutch East Indies.—Same as Netherlands. Old Dutch and local measures are also used. Latter very variable; recently they have been legally defined by their metric equivalents. Current:

Length

1 depa	= 1.70 m
Unit	Depa
1 hasta	= $\frac{1}{4}$
1 kilan	= $\frac{1}{8}$

Mass. (1) *Ordinary*

1 pikol $\}$	= 61.761 3025 kg
1 pecul $\}$	
Unit	Pikol
1 thail	= $\frac{1}{1600}$
1 catti $\}$	= $\frac{1}{100}$
1 kabi $\}$	
1 kulack	= 0.0725
1 amat	= 2
1 small bahar	= 3
1 large bahar	= 4.5
1 timbang	= 5
1 kojang (Batavia)	= 1667.555 kg
1 kojang (Semarang)	= 1729.316 kg
1 kojang (Soerabaya)	= 1852.839 kg

Mass. (2) *For precious metals*

1 thail	= 54.000 g
Unit	Thail
1 wang	= $\frac{1}{8}$
1 tali	= $\frac{1}{6}$
1 soekoe	= $\frac{1}{4}$
1 reaal	= $\frac{1}{2}$

Mass. (3) *For opium*

1 thail	= 38.601 g
Unit	Thail
1 tji	= 0.1
1 tjembang Mata $\}$	= 0.001
1 hoen $\}$	

Area

1 bahoe $\}$	= 70.965 a
1 bouw $\}$	
1 lieue² †	= 55.0632 km

Volume

1 kojang	= 1.976 362 m³
1 toembak	= 6.684 m³

Capacity, dry

1 kojang	= 2011.2679 l
1 pikol	= $\frac{1}{30}$ kojang

* Variable.
† Geographic.

Dutch East Indies.—Cont'd.

Capacity, liquid

(Legal equivalents)

Unit	Liter
1 takar*	= 25.770
1 kit*	= 15.159
1 koelak*	= 3 709
1 kan†	= 1.5751
1 mutsje†	= 0.1516
1 pintje*	= 0.0758

Ecuador.—m.c. 1865, but the British and, more generally the old Spanish, measures are currently used.

Egypt.—m.o. 1873; m.c. in government use, 1891. Current:‡

Length

| 1 diraa baladi | = 0 58 m |
| 1 kassabah | = 3 55 m |

Unit	Diraa
1 kirat	= $\frac{1}{24}$
1 abdat	= $\frac{1}{6}$
1 kadam	= $\frac{1}{3}$
1 pic	= 1
1 gasab	= 4
1 mil hachmi	= 1000
1 farsakh	= 3000

Mass

| 1 oke | = 1248 g |

Unit	Oke
1 kirat	= $\frac{1}{6400}$
1 dirhem	= $\frac{1}{400}$
1 miskal	= $\frac{3}{800}$
1 okieh	= 0 03
1 rotoli	= 0 36
1 kantar	= 36
1 helm	= 200

Area

| 1 feddan | = 42.008 a |

Unit	Feddan
1 sahme	= $\frac{1}{500}$
1 kirat kamel	= $\frac{1}{24}$
1 feddan masri	= 1

Capacity

| 1 keddah | = 2.0625 l |

Unit	Keddah
1 kirat	= $\frac{1}{32}$
1 khanoubah	= $\frac{1}{8}$
1 toumnah	= $\frac{1}{6}$
1 robhah	= $\frac{1}{3}$
1 nisf keddah	= $\frac{1}{2}$
1 malouah	= 2
1 rob	} = 4
1 roubouh	}
1 keila	= 8
1 ardeb	= 96
1 daribah	= 768

* For oil
† For various products
‡ In national system, units and their interrelations were very variable, but since 1891, have been defined by their metric equivalents.

England v. **Great Britain.**
Equateur v. **Ecuador.**
Eritrea.—m.o. Local, provincial·

Length

1 cubi	= 0.32 m
1 emmet }	
1 derah }	= 0.46 m

Mass

1 rotolo	= 448 g
1 okia	= $\frac{1}{16}$ rotolo
1 gisla	= 163 kg

Capacity

| 1 messé | = 1.50 l |

Unit	Messé
1 cababo	= 4
1 tunica	= 12
1 ghebeta	= 16
1 entelam	= 128

Espagne v. **Spain**
Esthonia.—Russian and local Current:

Length

| 1 archine (Russian) | = 0.7112 m |
| 1 elle (Livonian) | = 0.6096 m |

Unit	Archine
1 elle (Kunnar)	= 0.75
1 faden	= 3

Mass

| 1 pfund | = 430 g |

Unit	Pfund
1 quent	= $\frac{1}{128}$
1 loth	= $\frac{1}{32}$
1 bespfund	= 20
1 centner	= 120
1 tonne	= 240
1 schiffspfund	= 400

Area

	Reval
1 lofstelle	= 18 55 a
1 tonnland	= 54 627 a

	Livonian
1 lofstelle	= 37 1 a
1 tonnland	= 51.94 a

Capacity

| 1 hulmit | = 11.48 l |

Unit	Hulmit
1 lof (Reval)	= 3
1 lof (Livonian)	= 6
1 tonne (Livonian)	= 12

Etablissements des Détroits v. **British India.**
Etats-Unis v. **United States.**
Ethiopia.—var. Current

Length

(Approximate only)

Unit	cm
1 tat	= 2.5
1 gat	= 8
1 sinzér	= 16
1 kend	= 49

Mass

1 kasm	= 3.90 g
1 neter	= 336 g
1 farasula*	= 13.478 kg
1 farasula†	= 16.85 kg
1 farasula‡	= 17.072 kg

Unit	Kasm
1 mutagalla	= 2
1 alada	= 4
1 wogiet	= 8

Capacity

| 1 menelik | = 1 l (approximate) |

Filippine v. **Philippine.**
Finland.—m.c. 1892; m.o. 1887. Older (Russian and local):

Area

| 1 tunnland | = 46 54 a |

Capacity

1 tunna	= 163.49 l
1 kannor	= $\frac{1}{30}$ tunna
1 ottingar	= 15.71 l
1 sextingkar	= $\frac{1}{2}$ ottingar

France.—m.c. 1794. Other legal units.

Length

| 1 mille marin | = 1852 m |

Volume

| 1 tonneau de jauge | = 2.83 m³ |
| 1 tonneau de mer | = 1.44 m³ |

Old measures derived from the system of Charlemagne are:

Length

| 1 toise§ | = 1.949 0365 m‖ |
| 1 toise§ | = 1.949 090 m¶ |

Unit	Toise
1 ligne	= $\frac{1}{864}$
1 pouce	= $\frac{1}{72}$
1 pied	= $\frac{1}{6}$
1 aune	= 0.6064
1 lieue	= 2280.3
1 mille marin	= 950.13
1 lieue marine	= 2850.4

Mass

| 1 livre** | = 489.505 85 g |

Unit	Livre
1 grain	= $\frac{1}{9216}$
1 scruple	= $\frac{1}{384}$
1 gros	}
1 drachme	} = $\frac{1}{128}$
1 once	= $\frac{1}{16}$
1 marc ††	= $\frac{1}{2}$

* For ivory.
† For coffee.
‡ For rubber.
§ Toise de Perou at 16 25°C
‖ Equivalent made legal in 1799
¶ By measurement, in 1887, by J R. Benoit
** One livre de Charlemagne = 367.128 g
†† 1 Marc de la Rochelle = 244 75 g
   1 Marc de Limoges = 240 93 g
   1 Marc de Tours = 237 87 g
   1 Marc de Troyes et Paris = 260 05 g

Unit	Livre
1 quintal	= 100
1 millier	= 1000

Unit	Livre (Ch)
1 sol	= $\frac{1}{20}$
1 denier	= $\frac{1}{240}$
1 obole	= $\frac{1}{480}$
1 grain	= $\frac{1}{5760}$

Area

| 1 pied² | = 0.10552 m² |

Unit	Pied²
1 toise²	= 36
1 perche de Paris	= 324
1 perche des Eaux et Forêts	= 484
1 arpent de Paris	= 32 400
1 arpent des Eaux et Forêts	= 48 400

Capacity, dry

| 1 boisseau | = 1.862 78 l* |

Unit	Boisseau
1 litron	= $\frac{1}{16}$
1 quart	= $\frac{1}{4}$
1 minot	= 3
1 mine	= 6
1 setier	= 12
1 muid	= 144

Capacity, liquid

1 muid	= 274.239 l†
1 muid	= 268.241 l‡
1 pinte	= 0.931 389 l§

Unit	Pinte
1 roquille	= $\frac{1}{32}$
1 posson	= $\frac{1}{8}$
1 demi-setier	= $\frac{1}{4}$
1 chopine	= $\frac{1}{2}$
1 pot	= 2
1 velte	= 8
1 quarteau	= 72
1 feuillette	= 144
1 muid	= 288

Francia, Isola di v. **Mauritius.**
Frankreich v. **France.**

Germany.—m.c. 1872. Since the beginning of the nineteenth century, the other units and their interrelations have been fairly definite, but before that there was great diversity. *Length:* fundamental unit was Fuss (foot), its value, depending upon the state, varied from 0.280 to 0.320 m. The one most extensively used was the Rheinlandischer Fuss (Rhenish foot) = 0.313 857 m. *Mass:* fundamental unit was Pfund

* From 1 muid = 268.241 l by relation 144 boisseau = 1 muid (see Capacity, Liquid).
† Legal value.
‡ Derived from concrete standards
§ From 1 muid = 268.241 l by relation 288 pinte = 1 muid.

(pound), its value generally varied little from 467 g; during transition period preceding 1872 the accepted equivalents were Pfund = 30 Loth = 300 Zent = 3000 Korn; Centner = 100 Pfund. Older:

BAVARIA.

Length

| 1 Fuss | = 0.291 86 m |
| 1 Elle | = 0.833 01 m |

Unit	Fuss
1 Linie	= $\frac{1}{144}$
1 Zoll	= $\frac{1}{12}$
1 Ruthe	= 10
1 Chausseemeile	= 25 406

Mass

| 1 Pfund | = 560 g |

Unit	Pfund
1 Gran	= $\frac{1}{7680}$
1 Pfennig	= $\frac{1}{512}$
1 Quint	= $\frac{1}{128}$
1 Loth	= $\frac{1}{32}$
1 Unze	= $\frac{1}{16}$
1 Zentner	= 100

Area

1 Morgen	
1 Tagwerk	= 34 072 a
1 Juchert	
	= 400 Ruthe²

Capacity, dry

| 1 Metzen | = 37.0596 l |

Unit	Metzen
1 Dreissiger	= $\frac{1}{33}$
1 Mässel	= $\frac{1}{8}$
1 Scheffel	= 6

Capacity, liquid

| 1 Masskanne | = 1.069 03 l |

Unit	Masskanne
1 Zoll³	= $\frac{1}{43}$
1 Eimer	= 60 or 64
1 Fass	= 1600

PRUSSIA.

Length

| 1 Fuss | = 0 313 857 m |

Unit	Fuss
1 Linie	= $\frac{1}{144}$
1 Zoll	= $\frac{1}{12}$
1 Ruthe	= 12
1 Meile	= 24 000
1 Elle	= 25.5 Zoll

Mass

| 1 Pfund | = 467.711 g |

Unit	Pfund
1 Quentchen	= $\frac{1}{60}$
1 Loth	= $\frac{1}{32}$
1 Stein	= 22
1 Centner	= 110
1 Schiffspfund	= 330

Area

| 1 Morgen | = 25.532 24 a |
| 1 Morgen | = 180 Ruthe² |

Capacity, dry

| 1 Metze | = 3.435 89 l |

Unit	Metze
1 Quart	= $\frac{1}{4}$
1 Zoll³	= $\frac{1}{192}$
1 Scheffel	= 16

Capacity, liquid

| 1 Quart | = 64 Zoll³ |
| 1 Quart | = 1.145 03 l |

Unit	Quart
1 Anker	= 30
1 Eimer	= 60
1 Ohm	= 120
1 Oxhoft	= 180
1 Fuder	= 720

WÜRTTEMBERG.

Length

| 1 Fuss | = 0 286 49 m |

Unit	Fuss
1 Linie	= 0 01
1 Zoll	= 0.1
1 Elle	= 2 144
1 Ruthe	= 10
1 Meile	= 26 000

Mass

| 1 Pfund | = 467.728 g |
| 1 Apotheker-Pfund | = 357 647 g |

Unit	Pfund
1 Quentlein	= $\frac{1}{96}$
1 Loth	= $\frac{1}{32}$
1 Mark	= $\frac{1}{2}$
1 Zentner	= 104

Area

1 Ruthe²	= 8 207 66 m²
1 Morgen	= 384 Ruthe²
1 Juchart	
1 Tagwerk	= 576 Ruthe²

Capacity, dry

| 1 Simri | = 942.125 Zoll³ |
| | = 22.1533 l |

Unit	Simri
1 Viertelein	= $\frac{1}{128}$
1 Erklein	= $\frac{1}{32}$
1 Vierling	= $\frac{1}{4}$
1 Scheffel	= 8

Capacity, liquid

| 1 Maass | = 78.125 Zoll³ |
| | = 1.837 05 l |

Unit	Maass
1 Schoppe	= $\frac{1}{4}$
1 Imi	= 10
1 Eimer	= 160
1 Fuder	= 960

Gioppóne v. Japan.

Great Britain, Irish Free State, and Northern Ireland.—m.o. 1864. Since 1898, the national measures are convertible to metric by the legally sanctioned factors given below. National fundamental units defined thus: *Length*: The yard is distance at 62°F between axes of two lines traced on gold plugs set in a bronze bar preserved at the Standards Department of the Board of Trade. *Mass*: The pound avoirdupois is the mass of a certain platinum standard, similarly preserved. *Capacity*: The gallon is the volume of 10 pounds avoirdupois of pure water, as weighed in air against brass weights, the water and air being at the temperature of 62°F and the barometer at 30 inches. In official comparisons, the density of brass is taken as 8.143 g/cm³. Some of the units in the following tables are not in current use.

Length

1 yard* (yd.)	= 0 914 3992 m
1 foot (ft.)	= $\frac{1}{3}$ yd.
	= 30.479 97 cm
1 inch (in.)	= $\frac{1}{36}$ yd.
	= 2.539 998 cm

Unit	Inch
1 mil	= 0.001
1 point	= $\frac{1}{72}$
1 line	= $\frac{1}{12}$
1 barleycorn	= $\frac{1}{3}$
1 nail	= 2 25
1 palm	= 3
1 hand	= 4
1 span	= 9
1 quarter	= 9
1 foot	= 12
1 cubit	= 18
1 pace	= 30
1 yard	= 36
1 ell	= 45

Unit	Foot
1 fathom	= 6
1 pole	
1 rod (rd.)	= 16.5
1 perch	
1 rope	= 20
1 chain†	= 66
1 skein	= 360
1 furlong	= 660
1 cable length	= 720
1 mile (statute)	= 5280
1 mile (nautical)	
1 knot	= 6080
1 league	= 15 840

Mass

1 pound avoirdupois (lb. av.)	
	= 453.592 45 g
	= 7 000 grain
1 grain (gr.)	= 64.798 182 mg

(Three systems: avoirdupois, troy, apothecary.)

* This is the present legal equivalent of the imperial yard; recent comparisons by the National Physical Laboratory show that the yard as defined by the Weights and Measures Act of 1878 = 0 914 3987 m.

† Gunther's chain, divided into 100 link.

Avoirdupois (av.)
(General use)

Unit	Pound
1 dram (dm.)	= $\frac{1}{256}$
1 ounce (oz.)	= $\frac{1}{16}$
1 clove or customary stone	= 8
1 stone (legal)	= 14
1 quarter	= 28
1 cental	= 100
1 hundred-weight (cwt.)	= 112
1 wey	
1 load	= 252*
1 ton	= 2240

Troy (t.)
(For precious metals)

Unit	Grain
1 pennyweight (dwt.)	= 24
1 ounce (oz.)	= 480
1 pound (lb.)	= 5760

Apothecary (ap.)
(For dispensing drugs)

Unit	Grain
1 scruple (s.)	= 20
1 drachm (dr.)	= 60
1 ounce (oz.)	= 480
1 pound (lb.)	= 5760

Area

1 inch² (sq. in.)	= 6.451 5808 cm²
1 foot² (sq. ft.)	= 929.0289 cm²
1 yard² (sq. yd.)	= 0.836 1259 m²
1 acre (A.)	= 4046.849 m²

Unit	Foot²
1 inch²	= $\frac{1}{144}$
1 yard²	= 9

Unit	Yard²
1 pole² (sq. po.)	
1 rod²	= 30.25
1 perch²	
1 chain²† (ch.)	= 484
1 rood	= 1210
1 acre (A.)	= 4840

Unit	Acre
1 mile² (sq. mi.)	= 640

Volume

1 yard³ (cu. yd.)	= 0.764 552 85 m³
1 foot³ (cu. ft.)	= 28 316.77 cm³
1 inch³ (cu. in.)	= 16.387 0253 cm³

Unit	Foot³
1 inch³	= $\frac{1}{1728}$
1 yard³	= 27

* Variable
† Gunther's chain.

Great Britain. *Cont'd.*

Unit	Foot³
1 register	
ton	= 100
1 rod	= 1000

Capacity, dry

1 gallon (gal.) = 4.545 9631 l
1 bushel (bu.) = 8 gallon
= 35.367 7048 l

Unit	Gallon
1 quartern	= ½
1 peck	= 2
1 bucket	= 4
1 bushel	= 8
1 firkin	= 9
1 kilderkin	18
1 barrel	= 36
1 hogshead	63
1 puncheon	= 84
1 butt	= 126

Unit	Bushel
1 strike	2
1 sack	
1 bag	= 3
1 coomb	= 4
1 quarter	= 8
1 seam	8
1 chaldron	32*
1 wey	
1 load	= 40*
1 last	80*

Capacity, Liquid

1 gallon (gal.) = 4.545 9631 l

Unit	Gallon
1 gill	
1 quartern	= 1/32
1 noggin	
1 pint	= 1/8
1 quart	= 1/4
1 pottle	= 1/2

Greece. m.c. 1922; m.o. 1836. Older.

Length

1 piki varies 0.640 to 0.670 m
1 pic = 1 piki
1 small piki of Constantinople = 0.648 m
1 large piki of Constantinople = 0.669 m
1 piki (masonry) = 0.750 m

Mass

1 dramme = 3.2 g
1 livre (Venetian) = 150 g
1 mna = 1.5 kg
1 mine (royal) = 1.5 kg
1 oka† = 1.280 kg
1 oka = 1.250 to 1.333 kg
1 stater = 56.32 kg
1 talanton = 150 kg

Area

1 stemma = 10 a

* Variable.
† 0.85331 royal mine.

Capacity

1 oka = 1.333 to 1.340 l
1 baril = 74.236 l

Grossbritannien *v.* Great Britain.

Guam.—Metric is compulsory

Guatemala *v.* Costa Rica.

Guinea. m.c. 1910. Older = Portugal, England, and local:

Length

1 pik = 0.578 m
1 jacktan = 3.658 m

Mass

1 benda = 64.2 g
1 kantar = 977 kg
1 gammell = ½ kantar

Unit	Benda
1 akey	= 1/8
1 mediatabla	= 3½
1 aguirage	= 1/6
1 quinto	= 3½
1 piso	
1 uzan	= 1/8
1 seron	= 1/16
1 benda (offa)	= ½

Haiti. m.c. 1921. Older = British, old French, and Spanish; legal equivalents during transition period:

Length

1 toise = 1.9488 m
1 aune = 1.188 m

Area

1 carreau = 1292.3 m

Volume

1 baril = 0.1 m³
1 corde = 3.84 m³
1 toise = 8 m³

Holland *v.* Netherlands.
Honduras *v.* Costa Rica.
Hungary.—m.c. 1876. Older = old Vienna·

Length

1 mertfold	
1 meile	= 8.3536 km
1 marok	
1 faust	= 0.105 36 m

Area

1 hold = 43.16 a
1 joch = 43.16 a
1 meile² = 6978 ha

Volume

1 eimer = 54.30 l
1 halbe
1 itcze = 1/64 eimer
1 metzen
1 ako = 62.53 l

Iceland.—m.c. 1907. Older (analogous to Danish) were defined by their metric equivalents.

Length

1 fet = 0.313 85 m
1 sjomila = 1855 m

Unit	Fet
1 lina	= 1/144
1 þumlungur	= 1/12
1 alin	= 2
1 faðmur	= 6
1 mila a landi	= 24.000

Mass

1 pund = 0.5 kg

Unit	Pund
1 mark	= 2
1 fisk	= 8
1 fierding	= 40
1 hespund	= 64
1 tunna smjors	= 224
1 skippund	
1 batt	= 320

Area

1 ferfaðmur = 3.546 m²
1 fermila = 56.7383 km²

Unit	Ferfaðmur
1 ferþumlungur	= 5 1/84
1 ferfet	= 3/6
1 feralin	= 1/9
1 tundagslatta	= 900
1 engjateigur	= 1600

Capacity

1 pottar = 3½ fet³
= 0.9661 l

Unit	Pottar
1 kornskeppa	= 18
1 anker	= 39
1 almenn turma	= 120
1 öltunna	= 136
1 korntunna	= 144

India *v.* British India; *v.* Indo-China.
Indies, East *v.* British India; *v.* Dutch East Indies.
Indo-China, British *v.* British India.
Indo-China, French:
COCHIN CHINA. m.c. 1911, with the names:

Length

1 môt thuoc = 1 m

Mass

1 môt cân tây = 1 kg
1 môt đông cân tây = 1 g
1 picul = 60 kg

Capacity

1 vuông môt bat tây = 1 l
1 vuông môt gia = 40 l
CAMBODIA.—m.c. 1914, with the names:

Length

1 muoi mètre = 1 m

Mass

1 pram rôi = 1 kg
1 muoi gramme = 1 g
1 hoesep = 60 kg

Capacity

1 muoi litre = 1 l
1 sĕsep litre = 40 l

Irish Free State *v.* Great Britain.
Islande *v.* Iceland.
Italian colonies.—Metric compulsory.
Italy. m.c. 1861; adopted in Milan as early as 1803, with the following names:

Length

metro = m
palmo = dm
dito = cm
atomo = mm

Mass

libbra nuova = kg
oncia = hg
grosso = dkg
denar = g
grano = dg

Capacity

soma = hl
mina = dkl
pinta = l
coppo = dl
Older, provincial:

Length

1 piede liprando = 0.513 77 m

Unit	Piede lip.
1 punto	= 1/144
1 oncia	= 1/12
1 canna	= 4
1 trabucco	= 6
1 miglio	= 4333⅓

Mass

1 libbra = 307 to 398 g

Unit	Libbra
1 grano	= 1/6912
1 denaro	= 1/288
1 ottavo	= 1/96
1 oncia	= 1/12
1 rubbo	= 25
1 cantaro	= 150

Area

1 quadrao
1 giornata = 38 a
1 tavola = 1/100 giornata

Capacity, dry

1 mine = varies 12 to 120 l

Capacity, liquid

1 barile da vino = 45.6 l
1 barile da olio = 33.4 l

Japan.—m.o. 1893. Before 1891, great diversity; since 1891, fundamental units defined by metric equivalents.

Length

1 shaku*	= $\frac{10}{33}$ m
	= 0.303 0303 m

Unit	Shaku
1 shi	= 10⁻⁵
1 mô	= 10⁻⁴
1 rin	= 10⁻³
1 bu	= 10⁻²
1 sun	= 10⁻¹
1 yabiki	= 2.5
1 hiro	= 5
1 ken	= 6
1 jô	= 10
1 chô	= 360
1 ri†	= 12 960

Mass

1 kwan	= $1_1^{.5}$ kg
	= 3.75 kg

Unit	Kwan
1 shi	= 10⁻⁶
1 mô	= 10⁻⁵
1 rin	= 10⁻⁴
1 fun	= 10⁻⁴
1 candareen	= 10⁻⁴
1 mommé	= 10⁻³
1 niyo	= 0.001
1 hyaku-mé	= 0.10
1 kin	= 0 16
1 ninsoku-ichi-nin	= 7
1 kiyak-kin	= 16
1 karus hari-ichi-da	= 18
1 komma-ichi-da	= 40

Area
(Land Measure)

1 bu	= $\frac{100}{30\ 25}$ m²
	= 3.305 785 12 m²

Unit	Bu
1 gô	= 0 1
1 tsubo	= 1
1 sé	= 30
1 tan	= 300
1 chô	= 3000
1 ri²	= 46 656

Capacity

1 shô	= $1\frac{4\,9\,1}{3\,3\,1}$ l
	= 1.803 9068 l
	= 64827 bu³

Unit	Shô
1 shaku	= 10⁻²
1 gô	= 10⁻¹
1 to	= 10
1 koku	= 100

Kanada r. Canada.
Kolumbien r. Columbia.
Kongo r. Congo.

---

* The old shaku (kujirajaku) = 1.25 shaku is legal for fabrics
† One n mann (kai-ri) = nautical ri

Kuba r. Cuba.
Latvia.—m.o. Russian and local measures since 1845. Old measures were those of Holland.

Length

1 elle	= 0.537 m
1 quartier	= $\frac{1}{4}$ elle
1 meile	= 7 verste (Russian)
	= 7.468 km

Mass

1 pfund = 419 g

For secondary units, see Esthonia

Area

1 kapp	1.4864 a

Unit	Kapp
1 pourvete	25
1 loofstelle	25
1 tonnstelle	35

Volume

1 faden = 4 077 s

Capacity

1 stoof	= 1 2752 l

Unit	Stoof
1 kanne	2
1 kulmet	9
1 anker	30
1 poure }	54
1 loof }	
1 tonne	108

Lettonie r. Latvia.
Luxemburg. m.c. 1820 Previously used a local unit
1 malter = 191 l.
Malacca. —

Length

1 asta	= 0 457 m
1 depa	= 4 asta
1 jumba	= 8 asta

Mass

1 catty	= 0 61 kg

Unit	Catty
1 miam	$\frac{1}{3\,0}$
1 buncal	$\frac{1}{2\,0}$
1 tampang	1
1 bedur	2
1 kip	15
1 pecul	100
1 bahar	300

Area

1 jumba²	13.38 m²
1 orlong	= 400 jumba²
	53 52 a

Capacity

1 chupa	ca. 1 l
1 gantang	= 4 chupa

Malaysia r. British India; r. Dutch East Indies.
Malta.—m.c. 1914. Older, British and local (old Sicilian):

Length

1 foot	= 0.2836 m
1 canna	= 2.088 m
1 palmo	= $\frac{1}{8}$ canna

Mass

1 rottolo	= 1.75 lb. av.
	= 0.793 79 kg

Unit	Rottolo
1 parto	= $\frac{1}{480}$
1 ounce	= $\frac{1}{30}$
1 cantaro	= 100

Capacity

1 caffiso	= 20 457 l
1 barıl	= 43.162 l
1 salma	= 290.944 l

Marokko r. Morocco
Mauritius and Seychelles Islands. m.c. Older = old French, British, and the following:

Capacity

1 cash	= 227.11 l
1 velt	= $\frac{1}{50}$ cash

Mexico. m c. 1896; m o 1857. Older (from Spanish, Castillian), legally defined, during transition period, in terms of metric equivalents:

Length

1 vara	= 0 838 m

Unit	Vara
1 linea	$\frac{1}{432}$
1 pulgada	$\frac{1}{36}$
1 pie	$\frac{1}{3}$
1 legua	5000

Mass

1 libra	= 160 246 34 g

Unit	Libra
1 tomin	$\frac{1}{768}$
1 adarme	$\frac{1}{256}$
1 ochava	$\frac{1}{128}$
1 onza	$\frac{1}{16}$
1 arroba	25
1 quintal	100
1 tercio	160

Area

1 fanega	356 628 a

Unit	Fanega
1 caballeria	= 12
1 labor	18
1 sitio	= 192 28

Capacity, dry

1 cuartillo	= 1.8918 l

Unit	Cuartillo
1 almud	4
1 fanega	48
1 carga	96

Capacity, liquid

1 cuartillo	= 0.456 264 l
1 cuartillo for oil	= 0.506 162 l
1 jarra	= 18 cuartillos

Morocco.—m.o.; local, var.:

Length

1 cubit }	= 0.533 m
1 canna }	
1 pic	= 0.61 m
1 tonni	= $\frac{1}{4}$ pic

Mass

1 rotal }	= 507.5 g
1 artal }	
1 gerbe	= 3 kg
1 kula	= 22 rotal
1 kantar	= 100 rotal

Capacity

1 sahh }	= 56 l
1 fanega }	
1 mudd }	= 14 l
1 almude }	

Mozambique r. Portuguese East Africa.

Netherlands. —m.c. 1820, with the names:

Length

streep	= mm
duim	= cm
palm	= dm
elle	= m
roede	= dkm
mijle	= km

Mass

korrel	= dg
wigtje	= g
lood	= dkg
once	= hg
pond	= kg

Capacity, dry

maatje	= dl
kop	= l
schepel	= dkl
mudde	= hl
zak	= hl
last	= 30 hl

Capacity, liquid

vingerhoed	= cl
maatje	= dl
kan	= l
dekaliter	= dkl
vat	= hl

Old national system is more or less current in some of the old colonies:

Length
(Amsterdam)

1 roeden	= 3.079 77 m
1 elle	= 0.687 813 m
1 voeten	= 0.283 0594 m
1 duime	= 25.733 mm
1 lyne	= 2.144 mm

Mass

1 pond	= 492.16772 g
1 pond*	= 494.090 32 g

* Amsterdam.

Netherlands.—Cont'd.

1 pond (Apothecary)
 = ⅚ pond
 = 369.126 g

Unit	Pond
1 mark	= ½
1 unze	= 1/16
1 drachme	= 1 1/8
1 engel	= 1/320
1 vierling	= 1 1/1280
1 grein	= 1/7680

Area

1 morgen = 81.244 346 a

Capacity, dry

1 schepel = 27.26 l

Unit	Schepel
1 kop	= 1/32
1 vierd	= ¼
1 zak	= 3
1 mud	= 4
1 last	= 108

Capacity, liquid

1 mingelen = 1.200 to 1.237 l

Unit	Mingelen
1 vat	= 768
1 oxhooft	= 192
1 aam	= 128
1 anker	= 32
1 streekan	= 16
1 stoop	= 2
1 pint	= ½
1 mutsje	= ⅛

Nicaragua r. Costa Rica.
Niederlande r. Netherlands.
Northern Ireland r. Great Britain.
Norway.—m.c. 1882; m.o. 1879. Older differed very little from Danish; legal equivalents:

Length

1 fod = 0.3137 m

Mass

1 skaalpund = 0.4981 kg

Area

1 mal = 10 a

Capacity, dry

1 korntonde = 138.97 l

Capacity, liquid

1 pot = 0.9651 l

Oceania.—British measures.
Olanda r. Netherlands.
Österreich r. Austria.
Paësi Bäesi r. Netherlands
Panama.— Metric compulsory.
Paraguay.—Metric almost exclusively used. m.o. 1899. Older = Spain; legal equivalents:

Length

1 vara (old)	= 0.838 56 m
1 cuerda	=83⅓ vara = 69.88 m
1 cordel	
1 vara	= 0.866 m

Unit	Vara
1 piede	= ¼
1 pouce	= 1/30
1 ligne	= 1/432
1 cuadra	= 100
1 lieue	= 5000

Mass

1 libra (old)	= 460.08 g
1 libra	= 459 g

Unit	Libra
1 once	= 1/16
1 arrobe	= 25
1 quintal	= 100
1 tonne	= 2000

Area

1 liño (old)	= 48.832 a
1 liño	= 100 vara²
1 liño	= 75 m²

Capacity, dry

1 fanega	= 288 l
1 almude	= 1/12 fanega

Capacity, liquid

1 frasco = 3.029 l

Unit	Frasco
1 cuarta	= ¼
1 baril	= 32
1 pipe	= 192

Pays-Bas r. Netherlands.
Persia.—Metric is in process of adoption. By 1924 the following assimilation had occurred: 1 zar = 1 m, 1 dram = 1 g, 1 ralte = 1 l. National measures, provincial, var.; even today, in retail commerce, cereal grains are used as weights:

Length

1 guerze (common)	= 0.63 to 0.97 m
	= 1 monk-elzer
1 zar	= 1.04 m

Unit	Zar
1 gireh	= 1/16
1 ouroub	= ⅛
1 charac	= ¼
1 gez	
1 guerze	= 1
1 farsakh	
1 parasang	= 6000

Mass

1 miskal = 4.60 g

Unit	Miskal
1 una	= 3 ¼
1 gandum	
1 grain	= 1/96
1 abbas	= 1/24

Unit	Miskal
1 nakhod	
1 carat	= 1/24
1 dung	= ⅙
1 dartung	= 0.22
1 dirhem	= 2
1 sir	= 16
1 pinar	= 20
1 danar	= 40
1 abbassi	= 80
1 rottel	= 100
1 tcheirek	= 160
1 saddirham	= 320
1 batman (Tauris)	= 640
1 batman (Shirez)	= 1280
1 batman	= 600 to 1000
1 karvar	= 100 batman

Area

1 jerib = 1082 m² to 1153 m²
 = 1000 to 1066 zar²

Capacity

1 chenica	= 1.32 l
Unit	Chenica
1 sextario	= 0 25
1 capichas	= 2
1 sabbitha	= 5.5
1 colluthun	= 6 25
1 legana	= 30
1 artaba	= 50

Peru.—m c. 1869. Older (from Spanish, Castilian):

Length

1 vara = 0.835 98 m

Mass

1 libra = 460.09 g

Unit	Libra
1 arroba	= 25
1 quintal	= 100
1 fanega	= 140

Area

1 topo	= 27.06 a
1 fanegada	= 64.596 a

Philippine Islands. –m.c. 1860. Older = Spain. Local:

Mass

1 catty = about 600 g

Unit	Catty
1 punto	= ⅓
1 chinanta	= 10
1 lachsa	= 48
1 caban	= 97
1 pecul	= 100

Area

1 balita = 27.95 a

Unit	Balita
1 loan	= 0.1
1 quignon	= 10

Capacity

1 kaban	= 99.90 l
1 chupa	= 3.75 cm³
1 ganta	= 1/25 kaban
1 apatan	= ¼ chupa

Poland.—Metric in process of adoption; in some provinces it has been in use since 1872. Russian system legalized in 1849, without displacing national measurements. Since 1819 these have been defined by their metric equivalents. National:

Length

1 stopa = 0.288 m

Unit	Stopa
1 linja	= 1 1/12
1 cal	= 1/12
1 lokiec	= 2
1 sazen	= 6
1 pret	= 15

Old measures

1 pied (Warsaw)	= 0.2978 m
1 pied (Cracow)	= 0.3564 m
1 aune	= 0.620 m

Mass

1 funt = 405.504 g

Unit	Funt
1 gran	= 1/9216
1 skrupul	= 1/384
1 drachma	= 1/128
1 lut	= 1/32
1 uneja	= 1/16
1 kamian	= 25
1 centnar	= 100

Old measures

1 funt	= 404 g
1 centner	= 16 funt
1 stein	= 3.2 funt

Area

1 pret²	= 18.6624 m²
1 morga	= 300 pret²
1 wloka	= 9000 pret²

Capacity

1 kwarta = 1 l

Unit	Kwarta
1 kwarterka	= ¼
1 garniec	= 4
1 ewierc	= 32
1 korzec	= 128

Porto Rico.—m.c. 1860. Older = Spain:

Area

1 cuerdo = 2250 vara²
 = 15.72 a

Portugal.—m.c. 1872; m.o. 1852. Older:*

Length

1 pe	= 0.3285 m
1 estadio	= 258 m
1 milha	= 8 estadio
1 legoa	= 24 estadio

* In some of the older colonies the old Portuguese system, more or less modified, is still in use.

Unit	Pe
1 linha	= 1¼₄
1 pollegada	= ₁½
1 palmo	= ⅜
1 covada	= 2
1 vara	= 1⅒

Mass

1 libra*	= 459 g

Unit	Libra
1 grao	= ₉₂₁₆
1 escrupulo	= ₃₁₄
1 outava	= 1½₈
1 onca	= 1⅙
1 marco ⎫ 1 meio ⎬	= ½
arratel	= 1
1 arroba	= 32
1 quintal	= 128

Area

1 vara²	= 1.2 m²

Unit	Vara²
1 ferrado	= 605
1 geira	= 4840

Capacity, dry

1 fanga	= 54 l

Unit	Fanga
1 outava	= ₃½₂
1 quarto	= ₁⅙
1 meio	= ½
1 alqueira	= ¼
1 moio	= 15

Capacity, liquid

1 almude	= 16.5 l

Unit	Almude
1 quartillo	= ₄⅛
1 meio	= ₂¼
1 canada	= ₁½
1 alqueira	= ⅙
1 bota ⎫ 1 pipa ⎬	= 26
1 tonelada	= 52

Portuguese Colonies.—Metric compulsory.

Portuguese East Africa (Mozambique).—m.c. 1910. Older, mainly of Portugal; one bahar is considered equivalent to 109 kg.

Prussia v. Germany.

Rumania.—m.c. 1884; m.o. 1866. In old Bessarabia, Russian measures replaced by metric in 1922. Older:

Length

1 halibiu	= 0.701 m
1 endere	= 0.662 m
1 stringene	= 1.96 m

Mass

1 cantar	= ca. 56 kg
1 oke	= ₁¼ cantar

*For drugs 1 libra = ⅓ libra = 344.25 g.

Capacity

1 dimerla	= 24.6 l

Unit	Dimerla
1 oke	= ₁⅙
1 mirze	= 8
1 kilo	= 16

Capacity, liquid

1 viacka	= 14.15 l
1 oke	= 0.1 viacka

Russia.—m.o. 1900. Definitions of fundamental national units: *Length* Archine is distance at 17°C between the axes of two lines drawn on the platinum-iridium prototype marked "H 1894." *Mass* Fount is mass of the platinum-iridium prototype marked "H 1894." *Capacity, liquid* Vedro is volume of 30 founts of pure water at 16²₃°C. *Capacity, dry* Garnetz is ⁴₁₅ vedro.

Length

1 archine	= 0 711 200 m
1 totchka	= 0.254 0000 mm

Unit	Totchka
1 ligne	= 10
1 paletz	= 50
1 sotka	= 84
1 dume	= 100
1 verchoc	= 175
1 foute	= 1200
1 archine	= 2800

Unit	Archine
1 sagène	= 3
1 verste	= 1500

Mass (1) Ordinary

1 fount	= 409.51241 g
1 doli	= 44.434 9403 mg

Unit	Doli
1 sol ⎫ 1 zolotnik ⎬	= 96
1 lote	= 288
1 once	= 576
1 lana	= 768
1 fount	= 9216

Unit	Fount
1 poud	= 40
1 berkovets	= 400
1 tonne marine	= 2400

Mass (2) For drugs

Unit	Doli
1 grain	= 1.4
1 scrupule	= 28
1 drachme	= 84
1 once	= 672
1 livre	= 8064

Area

1 archine²	= 0.505 8054 m²
1 ligne²	= 6.451 600 mm²

Unit	Ligne³
1 dulme³	= 100
1 verchoc³	= 306.25
1 foute³	= 14 400
1 archine³	= 78 400

Unit	Archine³
1 sagène³	= 9
1 décatine	= 21 600
1 verste³	= 2 250 000

Volume

1 archine³	= 0.359 7288 m³
1 ligne³	= 16.387 06 mm³

Unit	Ligne³
1 dulme³	= 1000
1 verchoc³	= 5359.375
1 foute³	= 1 728 000
1 archine³	= 21 952 000

Unit	Archine³
1 sagène³	= 27
1 tonne marine	= 7 871 72
1 last marin	= 15.713 44

Capacity, dry

1 garnetz	= 3.279 842 l
1 tchast	= 0.109 328 07 l

Unit	Tchast
1 pologarnetz	= 15
1 garnetz	= 30
1 lof	= 592

Unit	Garnetz
1 tchetverik	= 8
1 polouosmina	= 16
1 osmina	= 32
1 tchetvert	= 64

Capacity, liquid

1 vedro	= 12.299 41 l
1 tcharka	= 0.122 9941 l

Unit	Tcharka
1 chkalik	= 0.5
1 bottle (vodka)	= 5
1 bottle (wine)	= 6.25
1 krouchka	= 10
1 shtoff	= 12.5
1 vedro	= 100

Unit	Vedro
1 stekar	= 1.5
1 anker	= 3
1 pipe	= 36
1 fass ⎫ 1 botchka ⎬	= 40

Salvador v. Costa Rica.

Schottland v. Great Britain.

Schweden v. Sweden.

Schweiz v. Switzerland.

Scotland, Scozia v. Great Britain.

Serbie-Croatie-Slovénie v. Yugoslavia.

Seychelles Islands v. Mauritius.

Siam.—m.c. 1923; m.o. 1889. Older now defined by metric equivalents; those of transition period:

Length

1 wah	= 2 m

Unit	Wah
1 anukabiet	= ₇¼₈
1 kabiet	= ₄¼₄
1 niou	= ₀⅙
1 keup	= ⅛
1 sawk ⎫ 1 sock ⎬	= ¼
1 ken	= ½
1 sen	= 20
1 roeneng	= 2000
1 yote	= 8000

Mass

1 tchang*	= 1200 g

Unit	Tchang
1 klom	= ₁₀¼₄₀
1 klam	= ₅₁½₂₀
1 pai	= ₂₅⅘₀
1 sompay ⎫ 1 grani ⎬	= ₁₇⅘₀
1 fuang	= ₆¼₀
1 salung	= ₃⅙₀
1 baht	= ₈¹₀
1 tamlung	= ₂⅒
1 doon	= 20
1 hap	= 50
1 bara	= 400

Area

1 wah²	= 4 m²
1 ngan	= 100 wah²
1 rai	= 400 wah²

Capacity

1 tanan†	= 1 l

Unit	Tanan
1 niou	= ₁⅘₀
1 chai meu	= ₂¹₄
1 kam meu	= ⅝
1 laang ⎫ 1 chang awn ⎬	= ½
1 kanahn	= 1
1 sat	= 20
1 tang	= 40
1 tamlaum	= 400
1 seste	= 800
1 ban	= 1600
1 kwien ⎫ 1 koyan ⎬	= 2000 or 3200
1 cohi	= 32 000

Siria v. Syria.

Somaliland.—m.o.; local, vary with material and province:

Length

1 top	= 3.92 m
1 cubito	= ½ top

Mass

1 rottolo	= 448 g

* Previously, 1 tchang = 600 to 1300 g.

† Previously, 1 tanan = 0.9 to 1.2 liter.

Somaliland. -- *Cont'd.*

Unit	Rottolo
1 okia	$= \frac{1}{16}$
1 frasla	$= 36$
1 gisla	$= 360$

Area

1 darat $= 80$ a

Capacity, dry

1 chela $= 1.359$ l

Unit	Chela
1 tabla	$= 15$
1 gisla	$= 120$

Capacity, liquid

1 caba $= 0.453$ l

Soudan *v.* Sudan.

South Africa *v.* Union of South Africa

Spain.--m c. 1860. Older,* var., provincial; Castilian:

Length

1 vara $= 0.835$ 905 m

(Other vara comprised between 0.768 m and 0.912 m)

Unit	Vara
1 punto	$= \frac{1}{6912}$
1 linea	$= \frac{1}{576}$
1 diedo	$= \frac{1}{48}$
1 pulgada	$= \frac{1}{36}$
1 sesma	$= \frac{1}{6}$
1 palma	$= \frac{1}{4}$
1 pie	$= \frac{1}{3}$
1 codos	$= \frac{1}{2}$
1 passo	$= 1\frac{2}{3}$
1 estado	$= 2$
1 estadal	$= 4$
1 milla†	$= 1666\frac{2}{3}$
1 legua	$= 5000$ or 8000

Mass

1 libra $= 460$ 093 g

(Other libra comprised between 350 g and 575 g)

Unit	Libra
1 grano	$= \frac{1}{9216}$
1 arienzo	$= \frac{1}{2304}$
1 tomin	$= \frac{1}{768}$
1 dinero	$= \frac{1}{384}$
1 adarme \}	
1 dracma /	$= \frac{1}{256}$
1 ochava \}	
1 caracter /	$= \frac{1}{128}$
1 escrupulo	$= \frac{3}{64}$
1 onza	$= \frac{1}{16}$
1 marco	$= \frac{1}{2}$
1 arroba	$= 25$
1 barril	$= 50$
1 quintal	$= 100$
1 quintalmacho	$= 150$
1 tonelada	$= 2000$

* Old national system, more or less modified, is still in use in the old Spanish colonies.

† Milla $= 5000$ pie.

Area

1 vara² $= 0.698$ 7372 m²

Unit	Vara²
1 cuartilla	$= 25$
1 calemin	$= 768$
1 aranzada	$= 6400$
1 fanega \}	
1 fanegada /	$= 9216$
1 yugada	$= 460$ 800

Capacity, dry

1 fanega $= 55.501$ l

Unit	Fanega
1 ochavillo	$= \frac{1}{768}$
1 racion	$= \frac{1}{192}$
1 cuartillo	$= \frac{1}{48}$
1 medio	$= \frac{1}{24}$
1 calemin	$= \frac{1}{12}$
1 almude	$= \frac{1}{12}$
1 cuartilla	$= \frac{1}{4}$
1 cahiz	$= 12$

Capacity, liquid

(Arroba was defined as volume of 34 libra of river water. The arroba for oil was volume of 25 libra of oil)

1 arroba (wine) $= 16.133$ l

1 arroba (oil) $= 12.563$ l

Unit	Arroba
1 copas	$= \frac{1}{128}$
1 quarterone \}	
1 panilla* /	$= \frac{1}{100}$
1 libra \}	
1 cuartillo /	$= \frac{1}{32}$
1 azumbre	$= \frac{1}{8}$
1 cuartilla*	$= \frac{1}{4}$
1 cantara	$= 1$
1 moio	$= 16$
1 pipa	$= 27$
1 bota	$= 30$

Stati Uniti *v.* United States.

Straits Settlements *v.* British India.

Sud-Africaine, Union *v.* Union of South Africa.

Sudan. Egyptian in use.

Suède *v.* Sweden.

Suisse *v.* Switzerland.

Svézia *v.* Sweden.

Svizzera *v.* Switzerland.

Sweden. m c. 1889; m.o. 1879. Older:

Length

1 fot $= 0.296$ 90 m

Unit	Fot†
1 linie	$= \frac{1}{144}$
1 tum	$= \frac{1}{12}$
1 alm	$= 2$
1 famn	$= 6$
1 stang	$= 16$
1 ref	$= 100$ or 160
1 mil	$= 18$ 000

* Ools

† The fot is also divided into decimals.

Mass

1 skålpund $= 425.076$ g

Unit	Skålpund
1 as	$= \frac{1}{8848}$
1 quintin	$= \frac{1}{128}$
1 lod	$= \frac{1}{32}$
1 untz	$= \frac{1}{16}$
1 lispund	$= 20$
1 sten	$= 32$
1 centner	$= 100$ or 120
1 waag	$= 165$
1 skeppund	$= 400$
1 nylåst	$= 12$ 000

Area

1 fot²	$= 0.088$ 149 61 m²
1 kappland $\begin{cases} \\ \end{cases}$	$= 1.542$ 618 17 a
	$= 1750$ fot²
1 ref²	$= 8.814$ 961 a
1 tunland $\begin{cases} \\ \end{cases}$	$= 49.363$ 781 6 a
	$= 56$ 000 fot²

Capacity, dry

1 kanna $= 2.617$ l

Unit	Kanna
1 ort	$= \frac{1}{32}$
1 junkfra	$= \frac{1}{32}$
1 quarter	$= \frac{1}{8}$
1 stop	$= \frac{1}{2}$
1 kappar	$= 1$
1 fjerdingar	$= 7$
1 spanna	$= 28$
1 tunna	$= 56$
1 koltunna	$= 63$
1 kollast	$= 756$

Capacity, liquid

1 kanna	$= 0.1$ fot³
	$= 2.617$ 162 l

Unit	Kanna
1 jungfrur \}	
1 jungfer /	$= \frac{1}{32}$
1 quarter	$= \frac{1}{8}$
1 stop	$= \frac{1}{2}$
1 ankar	$= 15$
1 eimer	$= 30$
1 am \}	
1 ohm /	$= 60$
1 oxhufud \}	
1 oxhoft /	$= 90$
1 pipe	$= 180$
1 fuder	$= 360$

Switzerland.--m.c. 1877; m.o. 1868. Older, var.; during transition were fixed as follows:

Length

1 pied \}	
1 fuss /	$= 30$ cm

Unit	Pied
1 ligne \}	
1 linie /	$= \frac{1}{144}$
1 pouce \}	
1 zoll /	$= \frac{1}{12}$
1 aune \}	
1 elle /	$= 2$
1 toise \}	
1 ruthe /	$= 6$

Unit	Pied
1 perche	$= 16$
1 lieue	$= 16$ 000

Mass (1) Ordinary

1 livre $= 500$ g

Unit	Livre
1 loth	$= \frac{1}{32}$
1 once	$= \frac{1}{16}$

Mass (2) For medicine

1 livre $= 375$ g

Unit	Livre
1 grain	$= \frac{1}{5760}$
1 scruple	$= \frac{1}{288}$
1 drachme	$= \frac{1}{96}$
1 once	$= \frac{1}{12}$

Syria.—m.o.; current:

Length

1 pic $= 0.582$ m

Mass

1 rottolo $= 1785$ g

Unit	Rottolo
1 drachme \}	
1 pesi /	$= \frac{1}{800}$
1 metecali	$= \frac{1}{400}$
1 miteal	$= \frac{1}{400}$
1 once	$= \frac{1}{10}$
1 zurbo	$= 27.5$
1 cola	$= 35$
1 cantar	$= 100$

Capacity

1 rotl $= 3.2$ l

Unit	Rotl
1 makuk	$= 250$
1 garava	$= 450$

Tchéco-Slovaquie *v.* Czechoslovakia.

Tonkin.—Same as Anam (*q v.*)

Tripoli and Cyrenaïca.—m.o., current defined by metric equivalents:

Length

1 pik	$= 0.68$ m
	$= 1$ handaze
1 palmo	$= \frac{1}{3}$ pik
1 draa	$= 0.46$ m

Mass

1 rottolo	$= 512.8$ g
1 oka $\begin{cases} \\ \end{cases}$	$= 2.5$ rottolo
	$= 1282$ g
1 metical	$= 4.76$ g

Unit	Rottolo
1 kharouba	$= \frac{1}{2560}$
1 dram	$= \frac{1}{160}$
1 termino	$= \frac{1}{128}$
1 uckin	$= \frac{1}{16}$
1 mattaro	$= 42$
1 cantar	$= 100$

Area

1 pik² $= 0.4624$ m²

Unit	Pik*
1 denum	= 1600
1 jabia	= 1800

Capacity, dry

1 orba	= 7.6 l

Unit	Orba
1 nufsorbah	= ½
1 temen	= 4
1 ueba	= 16

(Measured by weight)

1 oka	= 1282 g
1 marta	= 11 to 14 oka
1 kele	= 2 marta

Capacity, liquid

1 barile	= 64.8 l
1 bozzo	= ¼ barile

(Measured by weight)

1 oka	= 1282 g

Unit	Oka
1 gorraf	= 9.75
1 giarra	= 58.5

Tschechoslovak *v.* Czechoslovakia.

Tunis. —m.c. 1895. Current.

Length

1 pic arabe	= 48.8 cm
1 pic ture	= 63.7 cm
1 pic endazé	= 67.3 cm

The pic used depends upon the object measured.

Mass

1 uckir	= 31.495 g

Unit	Uckir
1 rottolo attari	= 16
1 rottolo sucki	= 18
1 rottolo khaddari	= 20
1 cantaro	= 100

Capacity

1 cafisso	= 496 l
1 millerole (Marseilles)	= ca. 64 l

Unit	Cafisso
1 saah	= 1 ½ 9
1 whiba	= 1 ⅙

Turkestan.

Length

1 hasch	= 0.7112 m

Unit	Hasch
1 archine*	
1 altschin	= 1

Mass

1 batman	= 125 kg to 128 kg

Unit	Batman
1 sir	= ⅛
1 tscharik	= 6¼
1 mimtscha	= 2 ⅙

Turkey. —m.o.; current, var.:

*Russian.

Length

1 archine	= 64 to 76 cm
1 archine (for architecture)	= 75.77 cm
1 nul	= 1 km

Unit	Archine
1 nocktat	= 3 ⅟₅₆
1 hatt	= 2 ⅛
1 parmack	= 2¼
1 ouromb	= ⅞
1 pic	= 1

Mass

1 oka	= 1283 g

Unit	Oka
1 karat	= 6 ⅟₀ ₀
1 denke	= 1 ⅟₀ ₀
1 dirhem	
1 drachme	= ⅟₀ ₀
1 miskal	= ⅟₀ ₀
1 cequi	= ⅓
1 yusdrum	
1 rottel	= 0 44
1 batman	= 6
1 kantar	= 44
1 tcheki	= 176 to 195

Area

1 deunum	= 1600 archine²
	= 913 m²
1 djeil	= 100 a

Capacity

1 kile	= 32 to 43 l
1 zira³	= 0.435 m³

Unit	Kile
1 chinik	= ¼
1 fortin	= 4

Ungarn, Ungheria *v.* Hungary.

Union of South Africa. —Metric, British, and old Dutch:

Length

1 elle	= 0.685 m

Mass

1 bundle	= 3175 g

Area

1 morgen	= 85.5 a

Capacity

1 gantang	= 9.2 l
1 balli	= 5 gantang
1 muid	= 109.1 l
1 legger	= 516 l

Unit	Legger
1 kanne	= 3 ⅛ ₈
1 ahm	= ¼

United States of America. —m.o. 1866; m.c. for certain governmental purposes. Fundamental units of national system are defined in terms of metric units. For less common and obsolescent units, see Great Britain.

Length

1 yard (yd.)	= ⅜⅜⅜ m
	= 0.914 401 83 m
1 foot (ft.)	= ⅓ yd.
	= 30.480 061 cm
1 inch (in.)	= ⅟₃₆ yd.
	= 2.540 005 08 cm

Unit	Inch
1 mil	= 0.001
1 hand	= 4
1 span	= 9
1 foot	= 12
1 yard	= 36

Unit	Foot
1 fathom	= 6
1 rod	
1 pole	= 16.5
1 perch	
1 chain* (Gunther's)	= 66
1 chain* (engineer's)	= 100
1 bolt	= 120
1 furlong	= 660
1 cable length	= 720
1 mile (statute)	= 5280
1 mile (nautical)†	= 6080.20
1 league (statute)	= 3 st. mile
1 league (nautical)	= 3 n. mile

Mass

1 pound avoirdupois (lb. av.)	= 453.592 4277 g
	= 7000 grain (gr)
1 grain	= 64.798 918 21 mg

(Three systems: avoirdupois, troy, apothecary.)

Avoirdupois (av.)

(General use)

Unit	Pound
1 dram (dr.)	= 2 ⅟₅₆
1 ounce (oz.)	= ⅟₁₆
1 hundred-weight (cwt.) (long)	= 112
1 ton (short) (sh. tn.)	= 2000
1 ton (long) (l. tn.)	= 2240

Troy (t.)

(For precious metals)

Unit	Grain
1 pennyweight (dwt.)	= 24
1 ounce (oz)	= 480
1 pound (lb.)	= 5760

Apothecary (ap.)

(For dispensing drugs)

Unit	Grain
1 scruple (s. or ℈)	= 20
1 dram (dr. or ℨ)	= 60
1 ounce (oz. or ℥)	= 480
1 pound (lb.)	= 5760

* 1 link = 0.01 chain.
† 1 nautical mile = 1853.249 m

Area

1 inch² (sq. in.)	= 6.451 6258 cm²
1 foot² (sq. ft.)	= 929.0341 cm²
1 yard² (sq. yd.)	= 0.836 130 71 m²
1 acre (A.)	= 4046.873 m²

Unit	Foot²
1 inch²	= ⅟₁₄₄
1 yard²	= 9

Unit	Yard²
1 rod² (sq. rd.)	= 30.25
1 perch	
1 chain²*	= 484
1 rood	= 1210
1 acre (A.)	= 4840

Unit	Acre
1 mile² (sq. mi.)	= 640
1 township†	= 23 040

Volume

1 yard³ (cu. yd.)	= 0.764 559 45 m³
1 foot³ (cu. ft.)	= 28 317.0 cm³
1 inch⁴ (cu. in.)	= 16.387 162 cm³

Unit	Foot³
1 inch³	= 1 ⅟₇₂₈
1 board foot (bd. ft.)	= ⅟₁₂
1 yard³	= 27
1 shipping ton	= 40
1 register ton	= 100
1 cord (cd.)	= 128

Capacity, dry

1 bushel (bu.)	= 2150.42 inch³
	= 35.238 329 l

Unit	Bushel
1 pint (pt.)	= 6 ¼
1 quart (qt.)	= 3 ½
1 peck (pk.)	= ¼
1 barrel‡ (bbl.)	= 3.281
1 chaldron§	= 36
1 firkin	= 9 gallon

Capacity, liquid

1 gallon (gal.)	= 231 inch³
	= 3.785 332 l
1 minim (min. or ♏)	= 6 ⅟₄₀ gal.
	= 0.061 6102 ml

Unit	Minim
1 fluid dram (fl. dr.)	= 60
1 fluid ounce (fl. oz.)	= 480
1 gill (gi.)	= 1920

* Gunther's chain.
† 36 mile².
‡ For dry commodities, except cranberries, barrel = 7056 inch³; cranberry barrel = 5826 inch³; lime barrel contains 180 lb. av. or 280 lb. av.; by custom, flour barrel = 196 lb. av.
§ Variable.

United States.— Cont'd.

Unit	Gallon
1 gill (gi.)	= $\frac{1}{32}$
1 pint (pt.)	= $\frac{1}{8}$
1 quart (qt.)	= $\frac{1}{4}$
1 barrel*	= 31.5
1 hogshead	= 63

Uruguay.—m.c. 1894; m.o. 1866. Older = Spain (Castilian), more or less modified.

Venezuela.—m c. 1914; m o. 1857. Older = Spain (Castilian), more or less modified, and the following of Granada:

Length

1 vara	= 0.8 m
1 meile	= 6280 vara

Mass

1 libra	= 1 kg
1 bag	= 62.5 kg

Vereinigte Staaten r United States.

Württemberg r Germany

Yugoslavia. m c. 1883 Older:

C. SYSTEMS OF ANTIQUITY

Our knowledge of the measures of antiquity is derived from the texts and monuments which have persisted to modern times, and some actual standards which have come down to us. The latter enable us to establish quite exact equivalence between the measures which they represent and ours. But most frequently such equivalence is only very roughly known, or is actually unknown In this section are given only the more important or the best studied of these systems. The values given must not be taken too literally. Indeed, especially in antiquity, systems do not succeed one another; they evolve. Several may coexist among a single people; it is generally impossible to fix the dates at which these systems were used. The ancients had no capacity measures, such as ours; they weighed liquids and grains in terms of standards forming a second system of weights

Arabian System.

Length

1 foot	= 0.320 m

Unit	Foot
1 assbaa (finger)	= $\frac{1}{16}$
1 cabda (palm)	= $\frac{1}{4}$
1 cubit (new)	= $1\frac{1}{2}$
1 cubit †	= 2
1 orgye (pace)	= 6
1 qasab	= 12
1 seir	= 600
1 ghalva	= 720
1 mille	= 6000
1 parasang	= 18 000
1 barid } 1 veredus }	= 72 000
1 marhala	= 144 000

* Wine barrel
† Hachemic.

Length

1 linija	= 21.95 mm
1 palaz	= 36.34 mm
1 archine	= 660 mm to 712 mm
1 khvat	= 1.896 m
1 stopa	= $\frac{1}{6}$ kvat

Mass

1 oka	= 1280 g

Unit	Oka
1 dratmm	= $\frac{1}{100}$
1 satljk	= $\frac{1}{4}$
1 litra	= $\frac{1}{4}$
1 akov	= 40
1 tovar	= 100

Area

1 stopa²	= 998.56 cm²

Unit	m²
1 dunum	= 700
1 motyka	= 800
1 raliza	= 2500
1 dan oranja	= 3597
1 lanaz	= 5760 = 1600 khvat²

Capacity

(Liquids are measured by weight.)

Mass

(So-called system of the Prophet)

1 rotl	= 340 g

Unit	Rotl
1 dirhem	= $\frac{1}{130}$
1 nevat	= $\frac{1}{34}$
1 nasch	= $\frac{1}{6}$
1 oukia	= $\frac{1}{3}$
1 man } 1 mine }	= 2
1 ocque	= 4
1 qanthar	= 100
1 kikkar	= 125

Area

1 feddan	= 14 400 cubit²† = 59 a

Unit	Feddan
1 achir } 1 qasaba }	= $\frac{1}{100}$
1 qamha	= $\frac{1}{36}$
1 habbah	= $\frac{1}{72}$
1 cafiz	= $\frac{1}{40}$
1 qirat	= $\frac{1}{24}$
1 daneq	= $\frac{1}{6}$
1 djarib	= $\frac{1}{4}$

Capacity

(Measured by weight)

1 cafiz	= 32.64 kg

Unit	Cafiz
1 mudd	= $\frac{1}{8}$
1 kiladja } 1 caphite } 1 kist }	= $\frac{1}{24}$
1 sâa	= $\frac{1}{12}$
1 makuk	= $\frac{1}{8}$
1 ferk	= $\frac{1}{4}$
1 woebe } 1 khoull }	= 1
1 modius	= $1\frac{1}{3}$
1 artabe } 1 amphora }	= 2
1 gariba } 1 den }	= 8

Assyro-Chaldean-Persian System.

Length

1 foot	= 0.320 m

Unit	Foot
1 finger	= $\frac{1}{16}$
1 palm	= $\frac{1}{4}$
1 zereth	= 1
1 cubit	= 2
1 pace	= 6
1 qasab } 1 cane }	= 12
1 chebel	= 80
1 stadion } 1 ghalva }	= 720
1 mille	= 5400
1 parasang	= 20 000
1 schoeme	= 21 600
1 stathmos } 1 mansion }	= 80 000

Mass

1 talent	= 32.6 kg

(Talent divided into 50, 60 or 100 mina)

1 drachma = 0.01 mina

Area

1 gar	= 14.7 m² = 144 foot²

Unit	Gar
1 dizaine	= 10
1 gan	= 100
1 gur	= 1000

Capacity

(Measured by weight)

1 amphora	= 32.6 kg

Unit	Amphora
1 cados	= $\frac{1}{12}$
1 makuk	= $\frac{1}{8}$
1 woebe } 1 modius }	= $\frac{1}{2}$
1 small artaba	= $1\frac{1}{2}$
1 large artaba	= 2
1 large amphora	= 3
1 gariba	= 8

Egypt: System of the Pharaohs.

Length

1 pied	= 0.349 m

Unit	Pied
1 doigt, finger } 1 theb }	= $\frac{1}{16}$
1 palme } 1 choryos }	= $\frac{1}{4}$
1 dichas	= $\frac{1}{2}$
1 spithame	= $\frac{3}{4}$
1 pied royal } 1 zereth }	= 1
1 pigon	= $1\frac{1}{4}$
1 coudée royale } 1 derah }	= $1\frac{1}{2}$
1 coudée longue	= 2
1 pas	= $2\frac{1}{3}$
1 xilon	= $4\frac{1}{2}$
1 orgye	= 6
1 canne	= $11\frac{2}{3}$
1 senus	= 150
1 stade	= 500 or 600
1 mille	= 5000
1 atour vulgaire	= 15 000
1 schoeme	= 18 000
1 parasange	= 20 000
1 atour royal	= 30 000

Mass

1 mine	= 850 g

Unit	Mine
1 gerah	= $\frac{1}{1200}$
1 sicle	= $\frac{1}{60}$
1 kikkar } 1 talent }	= 50

Area

1 pekeis	= 27.405 m²

Unit	Pekeis
1 coudée²	= $\frac{1}{100}$
1 sû	= 6.25
1 dizaine	= 10
1 rema	= 50
1 aurure } 1 aroure }	= 100
1 setta	= 1000

Capacity
(Measured by weight)

1 khar = 34 kg

Unit	Khar
1 outen	= 1/80
1 man } 1 mine }	= 1/40
1 hecte	= 1/6
1 apt	= 1/4
1 keramion	= 1
1 metretes d'Héron	= 1 1/4
1 artabe des septante	= 1 1/2
1 artabe } 1 letech }	= 4 3/32

Greek System.
Length

1 pous* = 0.308 56 m

Unit	Pous
1 daktylos (finger)	= 1/16
1 condylos	= 1/8
1 palestra (palm)	= 1/4
1 dichas	= 1/2
1 spithame (span)	= 3/4
1 cubit†	= 1 1/2
1 Grecian cubit	= 2
1 bema (pace)	= 2 1/2
1 orgyia	= 6
1 amma (corde)	= 60
1 plethron	= 100
1 stadion	= 600
1 mille	= 4500
1 kiloorgyia	= 6000

Mass

1 mina = 425 g

Unit	Mina
1 chalque	= 1/4800
1 obol	= 1/600
1 diobol	= 1/300
1 drachma	= 0.01
1 tetradrachma	= 0.04
1 talent	= 60

Area

1 pous² = 0.095 209 m²

Unit	Pous²
1 dekapode²	= 100
1 plethron²	= 10 000

Capacity
(Measured by weight)

1 chenica = 816 g

Unit	Chenica
1 cyanthos	= 1/24
1 oxybaphon	= 1/16
1 cotyle	= 1/4
1 sexte	= 1/2

* The Olympic foot of Egyptian origin.
† Lapidary.

Unit	Chenica
1 maris	= 2
1 choüs	= 3
1 hemiektos	= 4
1 hektos } 1 modius }	= 8
1 metretes	= 36
1 medimnos	= 48

Hebrew System.
Length

1 sacred cubit = 0.640 m
1 cubit* = 0.555 m

Unit	Cubit*
1 finger	= 1/24
1 palm	= 1/4
1 zereth	= 1/2

Mass (Sacred system)

1 mina = 850 g

Unit	Mina
1 obol } 1 gerah }	= 1/1200
1 rabah	= 1/240
1 bekah	= 1/120
1 shekel	= 1/60
1 talent†	= 50

Mass (Talmudist or Rabbinical system)

1 mina = 354.2 g

Unit	Mina
1 pondiuscule	= 1/1200
1 mehah } 1 gerah } 1 obol }	= 1/600
1 zuzah } 1 drachma } 1 shekel } 1 tetradrachma }	= 1/100
1 talent	= 60

Capacity, dry
(Measured by weight)

1 ephah { (old) = 29.376 kg
{ (new) = 21.120 kg

Unit	Ephah
1 log	= 1/72
1 cab	= 1/18
1 gomor	= 0.1
1 sath } 1 modius }	= 0.3
1 cor	= 10

Capacity, liquid
(Measured by weight)

1 bath (old) = 29.376 kg
1 bath (new) = 21.120 kg

Unit	Bath
1 log	= 1/72
1 hin	= 1/6
1 cor	= 10

* Talmudist
† Of Moses.

Hindu System.
Length

1 hasta = 0.457 m

Unit	Hasta
1 angula (finger)	= 1/24
1 vitasti (span)	= 1/2
1 cubit	= 1
1 dhanush } 1 orgyla }	4
1 crosa	= 8000
1 gavyuti	= 16 000
1 yodjana	= 32 000

Mass

1 retti } 1 ratica }	0.117 g
1 pala	= 47 g

Unit	Retti
1 yava	= 0.1
1 masha	= 2, 5, 6, or 8
1 tank-sala	= 24
1 kona	= 48
1 tola	= 80
1 karsha	= 96
1 dharana	= { 32 (silver) { 3200 (gold)
1 pala	= 320

Unit	Pala
1 tuba	= 100
1 bara	= 200
1 bara	= 2000
1 achita	= 20 000

Capacity
(Measured by weight)

1 drona = 13.2 kg

Unit	Drona
1 pala } 1 musti }	= 1/256
1 cudava	= 1/32
1 prastha	= 1/16
1 adhaka	= 1/4
1 cumbha (small)	= 2
1 shari	= 16
1 cumbha	= 20
1 bahu	= 200

Persian System v. Assyrio-Chaldean-Persian.
Roman System.
Length

1 pes (common or Drusian) (foot) = 0.3196 m
1 legal pes (1st) = 0.2962 m
1 legal pes (2nd) = 0.2967 m

Unit	Pes
1 digitus (finger)	= 1/16
1 uncia (inch)	= 1/12
1 cubitus (cubit)	= 1 1/2
1 passus (pace)	= 5

---

1 decempeda (perch)	= 10
1 actus (chain)	= 120
1 millarium (mile)	= 5000

Mass

1 podium = 326 g

Unit	Podium
1 scrupulus	= 1/288
1 denier*	= 1/96
1 denier†	= 1/90
1 denarius	= 1/84
1 solidus } 1 sextula }	= 1/72
1 miliaresum	= 1/60
1 sicilium	= 1/48
1 duella	= 1/36
1 semuncia	= 1/24
1 ounce	= 1/12
1 mina	= 1 3/4
1 centum-podium	= 100

Area

1 common pes² = 0.102 14 m²
1 legal pes³ (1st) = 0.087 73 m²
1 legal pes² (2nd) = 0.088 03 m²

Unit	Pes²
1 decempeda¹	= 100
1 actus (small)	= 400
1 clima	= 3600
1 versum	= 10 000
1 actus	= 14 400
1 jugerum	= 28 800
1 heredium	= 57 600
1 centuria	= 5 760 000
1 saltus	= 23 040 000

Capacity, dry

1 sextarius = 544 g

Unit	Sextarius
1 modius	= 16
1 quadrantal	= 48
1 pes'‡ (of water)	= 48

Capacity, liquid
(Measured by weight)

1 sextarius } 1 sextus }	= 544 g

Unit	Sextarius
1 cyathus	= 1/12
1 acetabulum	= 1/8
1 quartus	= 1/4
1 hemina	= 1/2
1 congius	= 6
1 urna	= 24
1 amphora	= 48
1 culeus } 1 dolium }	= 960

* Silver
† Neronian.
‡ Legal pes (2).

SYMBOLS, BASIC CONSTANTS, CONVERSION DATA, DIMENSIONS, DEFINITIONS

BASES OF DATA CONTAINED IN I. C. T.

When many experts are cooperating in the assembling of data, it is essential that the same values for the fundamental constants and for the necessary conversion factors shall be employed by all. Consequently, at the very beginning of the work, the Editors compiled a set of accepted, or I. C. T., values for such constants and factors; and the Experts were instructed to base all their data upon these values. In the few cases in which it was not feasible to follow these instructions, the data were to be accompanied by a statement of the actual basis upon which they rest.

In compiling this list, and in choosing the accepted values of such of the quantities as were independently chosen, the Editors secured and utilized the advice of the United States Bureau of Standards, the National Physical Laboratory of Great Britain, and the Société Française de Physique. Acknowledgments are also due to Dr. F. E. Fowle, of the Smithsonian Institution, for his valued assistance in preparing the initial table of fundamental constants, and to Professors T. W. Richards and G. P. Baxter for their recommendations concerning the table of atomic weights.

The list so prepared comprised (1) a table of atomic weights (p. 43), (2) a set of nine basic constants (p. 17) (the estimated uncertainties were added at a later date), (3) twenty-one derived constants (computed directly from the nine basic constants), five conventional constants, and two experimental constants (p. 18) and (4) certain conversion factors selected from Tables 1 to 79 (p. 20-32). Although the accepted values were close approximations to the best values at that time available, it was not claimed that they were such best values.

SYMBOLS AND ABBREVIATIONS

Except as the contrary is definitely stated, the following symbols and abbreviations will always be used in the sense here indicated. Other symbols will be defined in the sections in which they are used. For those quantities which are included in the list of symbols approved by the International Association of Chemical Societies (*i*, **119**: 502; 21), the symbols so approved have, in general, been used; in some cases, this has necessitated the use of the same symbol to represent two distinct quantities; the context will serve to indicate which interpretation is correct. For explanations of the several technical terms, consult Selected Technical Terms, p. 34.

Å	Angstrom unit	ap	Apothecaries
A.	Acre	Av.	Average
A_n	Normal atmosphere	av.	Avoirdupois
A_{45}	Atmosphere, 45° latitude	*a*	Van der Waal's pressure
A	Atomic weight. Maximum		constant. Capillary con-
	work of a thermodynamic		stant
	system		
a	Are	BTU	British Thermal Unit
(a)	Based on Int. ohm and Int	bbl	Barrel
	ampere as defined by sil-	bd.	Board
	ver voltameter. (*See*	bu.	Bushel
	Int. elec. units, p. 27)	*b*	Van der Waal's volume
abs.	Absolute		constant

C	Centigrade	fir.	Firkin
CTU	Centigrade thermal unit	fl.	Fluid
C	Concentration. Molecular	fps	Foot-pound-second system
	heat		of units
C_1, C_2	Radiation constants of	fpse	Fps electrostatic system
	black body. (*See* defini-	fpsm	Fps electromagnetic system
	tion of black body)	ft.	Foot
C_j	Intensity coefficient. (*See*	ft.²	Square foot
	definition of black body)	ft.³	Cubic foot
C_p, C_v	Molecular heat at constant	fur.	Furlong
	pressure, at constant		
	volume	G	Gravitation constant
c	Velocity of light in vacuo	g	Gram
c	Carat. Centi-	gal.	Gallon
ca	Candle	gi.	Gill
ca.	circa ~ about, approxi-	gr.	Grain
	mately	*g*	Acceleration due to gravity
cal	Calorie (gram)	g_s	Standard gravity
cd.	Cord		
cf.	Confer = compare	IP	Horse-power
cgs	Centimeter-gram-second	*H*	Atomic weight of hydrogen
	system of units	h	Planck's constant of action
cgse	Cgs electrostatic system	h	Hecto-
cgsm	Cgs electromagnetic system	ha	Hectare
ch.	Chain	hhd.	Hogshead
cm	Centimeter	h p.	Horse-power
cm²	Square centimeter	hr	Hour
cm³	Cubic centimeter	*h*	Height
c p.	Candle power		
cu.	Cubic	Int.	International
cu ft.	Cubic foot	I. C. T.	International Critical
cwt.	Hundredweight		Tables
c	Specific heat ~ heat capac-	*I*	Electric current
	ity of the substance	ibid.	Ibidem = in the same place
c_p, c_v	Specific heat at constant	i.e.	Id est = that is
	pressure, at constant vol-	in.	Inch
	ume	in.³	Cubic inch
D	Density	*J*	Radiance
d	Derivative. Deci-	J_λ	Intensity of monochromatic
da	Day		radiance of wave-length λ
deg	Thermometric degree, abso-	J_m	Value of J_λ for λ = $λ_m$
	lute C unless contrary is		
	indicated	k	Karat. Kelvin, or absolute
dk	Deka-		C, scale of temperature
dm³	Cubic decimeter	*K*	Constant of chemical equi-
dr	Dram		librium
dwt	Pennyweight	k	Kilo-
d	Density. Diameter	kg	Kilogram
d_c	Critical density	km	Kilometer
$d_{t_1}^{t_2}$	Specific gravity at temper-	km²	Square kilometer
	rature t_2, with reference	*k*	Velocity coefficient of
	to water at temperature		chemical reaction
	t_1	k_0	Boltzmann's gas constant
E	Electromotive force	*L*	Latent heat per mole
E_0	Mean translational energy	l	Liter
	of molecule of ideal gas	l.	Long
	at 0°C	lat.	Latitude
e	Electronic charge	lb.	Pound
e	Base of natural system of	li.	Link
	logarithms = 2 71828 +	liq.	Liquid
e.g.	Exempli gratia = for	long.	Longitude
	example	*l*	Length. Latent heat per
em	Cgsm unit of quantity of		gram
	electricity		
emf	Electromotive force	*M*	Molecular weight
equiv	Electrochemical equivalent	M [α]	Molecular rotatory power
es	Cgse unit of quantity of	M [ω]	Molecular magnetic rota-
	electricity		tory power
etc.	Et cetera = and so forth	m_e	Mass of electron at low
et seq	Et sequentes = and the		velocity
	following	m	Meter. Milli-
e_0	Ratio of E_0 to T_0	m²	Square meter
		max.	Maximum
F	Faraday	mg	Milligram
F	Fahrenheit	mi.	Mile
fath.	Fathom	min	Minute

Min.	Minim, Minimum	T_a	Ice point, absolute C			Minim
ml	Milliliter	T	Temperature on absolute C scale	s	Susceptibility (magnetic). Electrical (volume) conductivity	Apothecaries' ounce
mmf	Magnetomotive force					Apothecaries' dram
mμ	Millimicron. Millimicro-	T_c	Critical temperature, absolute C	Λ	Equivalent conductivity (electrical)	Apothecaries' scruple
m	Mass					Degree (arc or temperature)
m_H	Mass of a hydrogen atom	t	Metric ton	λ	Wave-length. λ5890 = spectral line of wave-length = 5890Å	Minute of arc (sexagesimal)
		t.	Troy			Second of arc (sexagesimal)
N	Numeric	tn.	Ton	$λ_m$	Wave-length of maximum monochromatic radiance of black-body at stated temperature	Percent = per hundred
N_0	Avogadro's number	t	Time. Temperature C (above ice point)			Per thousand = 0.1 %
N_∞	Rydberg's universal series constant	t_c	Critical temperature C (above ice point)	μ	Permeability (magnetic) Micron, Micro-, Molecular conductivity (electrical)	Dimensional expressions are inclosed in [ ]. In text, [ ] is used to inclose a second reading. (E.g., Length [diameter] of the bar is 10 cm [1 cm] = length of bar is 10 cm, diameter of bar is 1 cm)
n	Refractive index	U. S.	United States of America			
n_a, n_k	Transport number for anion, kation	V	Volume	μμ	Micromicron Micromicro-	
n_0	Loschmidt's number	V_0	Volume per gram-mole of ideal gas at 0°C and A_D	ν	Frequency	$A < B$ [$A > B$] denotes that A is less than [greater than] B
O	Atomic weight of oxygen	v.	Vide = see	$ν_{\infty}$	Rydberg's fundamental frequency	
oz.	Ounce	(v)	Based on Int ohm and Int. volt as defined by standard cell. (See Int elec. units, p 27)	π	Ratio of circumference of a circle to its diameter	Negative of <; $A < B$ denotes that A is not less than B
P	Pressure			σ	Stefan's constant (radiation)	
pk	Peck	v	Volume	ψ	Fluidity. Angle	Combination of < and =; $A \leq B$ denotes that A is equal to or less than, B
pt.	Pint	v_c, v_r	Critical volume, reduced volume	ψ	Luminous flux	
p	Pressure			Ω	Ohm	Is not equal to
p_c, p_r	Critical pressure, reduced pressure	W	Electrical resistance	[Ω]	Relative molecular magnetic rotatory power with reference to water	Identically equal to; used in defining symbols, etc.
		wt.	Weight			
Q	Quantity	w	Wien's displacement constant	ω	Solid angle	Approximately (or essentially) equal to
q	Quantal			[ω]	Specific magnetic rotatory power	
qt.	Quart	yd.	Yard			Infinity
q.v.	Quod vide = which see	yr	Year			
R	Réaumur	Z	Atomic number			
R	Gas constant per mole of ideal gas. Electrical resistance.	α	Degree of dissociation. Angle of optical rotation			
rd.	Rod	[α]	Specific rotatory power			
r	Radius	β	Specific heat constant			
r_G	Specific refractivity (Gladstone and Dale)	γ	Surface tension. Ratio of c_p/c_v Gamma (magnetic unit)			
r_L	Specific refraction (Lorentz and Lorenz)	Δ	Diffusion coefficient			
r_1	Radius of first Bohr ring, hydrogen	ε	Dielectric constant Electrode potential			
S.E.	Siemens unit	$ε_h$, $ε_0$	Electrode potential above that of normal hydrogen, of normal calomel, electrode			
S	Entropy					
s	Stere	η	Viscosity			
s.	Scruple	θ	Angle (plane). Temperature C above ice point			
sec	Second (mean solar unless contrary is stated)					
'	Short					
sh.	Square					
sq	Square foot					
sq ft.						

[1] In every computation it is tacitly assumed that the values employed are exact. If but three digits are employed, it is assumed that all others are zero; if a computing machine is used, the assumption is carried out to the extreme limit of the machine; if logarithms are used, it is carried to the limit within which the logarithms are interpolated. To adopt an accepted or a conventional value, and to give as its logarithm an abbreviated value, is to introduce an ambiguity of a magnitude determined by the degree of abbreviation of the logarithm. But the sole object in adopting accepted or conventional values is to avoid ambiguity.

FUNDAMENTAL CONSTANTS

By an *accepted, conventional,* or *defined* value, is meant one which is to be regarded as exactly correct for purposes of computation.[1] Thus, errors from computational approximations are avoided and do not enter into consideration in any future revision of the computed result for a discovered difference between the true and the accepted value. When the computation involves several accepted values, it is especially important that each shall be regarded as exactly correct, for only then can the result be independently revised (without complete recalculation) for changes in the values of each. For this reason the logarithms of the several accepted values are given to the full precision of Vega's seven-place table. The degree of uncertainty in the value accepted is indicated by the number of significant figures retained in the value itself, not by the logarithm.

ACCEPTED BASIC CONSTANTS Units: cgs, °C, liter, A_D, absolute electric

	Quantity	Value	Uncertainty	Log_{10} (value)
c	Velocity of light	2 9986 × 10^10 cm sec^-1	0 0003	10.476 9185
G	Gravitation constant	6 66 × 10^-8 cm^3 g^-1 sec^-2	0 01	8.823 4742
e	Electronic charge	4 774 × 10^-10 es	0 005	10.678 8824
e	Electronic charge	*1 592 × 10^-20 em		20.201 9639
e/m_0	Electronic ratio	5 305 × 10^17 es g^-1	0 010	17.724 6854
e/m_0	Electronic ratio	*1 769 × 10^7 emg^-1		7.247 7669
F	Faraday	9 6500 × 10^3 coulombs	0 0010	4.984 5273
F	Faraday	*2 893 65 × 10^14 es		14.461 4458
V_0	Volume 1 mole at 0°C, A_D	†22 4115 × 10^3 cm^3 mole^-1	0 002	4.350 4709
h	Planck's constant	6 554 × 10^-27 erg sec	0 001	27.816 5064
T_0	Ice point, absolute	273 1 deg C	+0.15 to −0 05	2.436 3217
O	Atomic weight of oxygen	16 000 (by definition)	(definition)	1.204 1200

* This value is derived from the preceding one, which is the value actually accepted.
† Derived from volume at 0°C, A_D = 22.412 liters/g-mole on assumption log_{10} (A_D/A_{bs}) = 0.000 0214, liter = 1000.027 cm^3.

ACCEPTED CONSTANTS:—CONVENTIONAL AND NON-BASIC Units: cgs, °C, liter, A_n absolute electric, international angstrom

Quantity		Value	Log_{10} (value)
		A. Derived Constants	
R	Gas constant	$8\ 315 \times 10^7$ erg deg^{-1} mole^{-1}	7.919 8658
R	Gas constant	$0\ 082\ 06$ liter atm deg^{-1} mole^{-1}	$\bar{2}$.914 1375
R	Gas constant	$1\ 9869$ cal$_{15}$ deg^{-1} mole^{-1}	0.298 1703
N_0	Avogadro's number	$6\ 061 \times 10^{23}$ mole^{-1}	23.782 5634
n_0	Loschmidt's number	$2\ 705 \times 10^{19}$ cm^{-3} (at 0°C, A_n)	19.432 0925
k_0	Molecular gas constant	$1\ 372 \times 10^{-16}$ erg deg^{-1}	$\bar{16}$.137 3024
E_0	Translational energy of molecules, 0°C	$5\ 620 \times 10^{-14}$ erg	$\bar{14}$.749 7154
e_0	Ratio of E_0 to T_0	$2\ 058 \times 10^{-16}$ erg deg^{-1}	$\bar{16}$.313 3937
m_H	Mass of hydrogen atom	1.663×10^{-24} g	$\bar{24}$.220 7679
m_0	Electronic mass	$8\ 999 \times 10^{-28}$ g	$\bar{28}$.954 1970
r_1	Radius 1st Bohr ring of hydrogen	0.5305×10^{-8} cm	$\bar{9}$.724 6912
h/e	Photo-electric constant	$1\ 373 \times 10^{-17}$ erg sec es^{-1}	$\bar{17}$.137 6240
h/e	Photo-electric constant	$*4.117 \times 10^{-15}$ volt sec	$\bar{15}$.614 5425
hc/e	Photo-electric constant	$4\ 117 \times 10^{-7}$ erg cm es^{-1}	7.614 5425
hc/e	Photo-electric constant	$1\ 2344 \times 10^4$ volt Å	4.091 4610
β	Specific heat constant	$4\ 778 \times 10^{-11}$ sec deg	$\bar{11}$.679 2040
σ	Stefan's constant	$5\ 709 \times 10^{-5}$ erg cm^{-2} sec^{-1} deg^{-4}	$\bar{5}$.756 5416
C_1	Radiation constant, first	$3\ 703 \times 10^{-5}$ erg cm^2 sec^{-1}	$\bar{5}$.568 5233
C_2	Radiation constant, second	$1\ 433$ cm deg	0.156 1225
w	Wien's displacement constant	$0\ 2885$ cm deg	$\bar{1}$.460 1933
C_i	Intensity coefficient	$1\ 301 \times 10^{-4}$ erg cm^{-3} sec^{-1} deg^{-5}	$\bar{4}$.114 2762
v_∞	Rydberg frequency	$3\ 2775 \times 10^{15}$ sec^{-1}	15.515 5372
N_∞	Rydberg wave number	$1\ 0930 \times 10^5$ cm^{-1}	5.038 6187
		B. Conventional Constants	
A_n	Normal atmosphere	$1\ 0132\ 50 \times 10^6$ dyne cm^{-2}	6 005 7166
A_{45}	Atmosphere, latitude 45°	$1\ 0132\ 00 \times 10^6$ dyne cm^{-2}	6.005 6952
Å	Wave-length of red Cd line is	$6438\ 4696$ Å	4.808 7827
g_s	Standard gravity	$980\ 665$ cm sec^{-2}	2 991 5207
	Aberration constant	$20\ 47''$	1 311 1178
		C. Experimental Constants	
Grating space in calcite		$3\ 028$ Å	0.481 1559
H	Atomic weight of hydrogen	$1\ 0077$	0.003 3313
‡Liter		$1000\ 027$ cm^3	3.000 0117
‡Gram calorie (20°C)		4.181 joule	0.621 2802
‡Gram calorie (15°C)		4.185 joule	0.621 6955
‡Gram calorie (mean)		4.186 joule	0.621 7992
‡British Thermal Unit (39°F)		$1060\ 4$ joule	3.025 4697
‡British Thermal Unit (mean)		$1054\ 8$ joule	3 023 1701
‡British Thermal Unit (60°F)		1054.6 joule	3.023 0878
‡International ohm		$1\ 000\ 52$ ohm	0 000 2259
‡International ampere (v)§		$0\ 999\ 90$ ampere	0.999 9566
‡International ampere (a)§		$0\ 999\ 93$ ampere	0 999 9696

* This value is derived from the preceding one, which is the value actually accepted.

‡ In the original list, this quantity was included solely in the list of conversion factors; its value, however, is an independently selected, *accepted* constant, and, consequently, is treated as exact in all computations

§ (v) = Based on Int ohm and Weston normal cell = 1 018300 Int volts at 20°C, (a) = based on deposit of 1 11800 mg of silver per Int ampere second.

CONVERSION FACTORS AND DIMENSIONAL FORMULAE

N. ERNEST DORSEY

In the following tables are given the factors by which values expressed in other units must be multiplied in order to obtain their equivalents in units of the centimeter-gram-second (cgs) system. To convert in the reverse direction, divide by the factor given. The dimensional formula in the cgs, or any similarly constructed, system is given in the title of each table.

Conversion Factors.—With few exceptions,[1] the values given are based exclusively upon legal definitions, conventional constants, and the I. C. T. accepted values (p. 16). Consequently, they are computable to as extreme a precision as may be desired. They have been computed by means of Vega's seven-place logarithms, and it is hoped that their logarithms as given are correct to a unit in the last digit. Obviously, those factors which involve the accepted value of an experimentally determined constant will be in error by an amount determined by the error in the accepted value; but quantities converted by means of the logarithms given will retain their same relative precision, however great this may be, within the limit set by the seven-place table, and may at any time be as exactly corrected for a revision of the accepted value. This would not be true if an abbreviated logarithm were used, unless the exact value of the abbreviated logarithm itself were given. The latter would be equivalent merely to the adoption of another accepted value for the experimental constant involved;

[1] The exceptions are (1) astronomical unit of distance, (2) parsec, (3) sidereal second, (4) certain units of luminous intensity, (5) international electrical units prior to 1911, and (6) the data for hydrometers

and the new value so fixed would, in general, be expressible only by an indefinite number of digits. The former procedure is to be preferred.

Frequently, the same factor applies to more than one type of physical quantity; if the units of the several types have distinctive names, separate tables are given, otherwise, not. In general, the tables are arranged in the order of increasing complexity of the dimensional formulae. Some quantities for which conversion factors are seldom required, and a few dimensionless quantities have been grouped together in Table 78. The dimensional formulae of the more inportant electric and magnetic units, and the numerical relations connecting these units in the three systems most frequently used, are assembled in Table 77. To find the conversion factor for a given quantity, consult the index below.

Dimensions.—Two types of dimensional equations need to be considered, viz.: (1) Those in which the dimensions are expressed in terms of the quantities directly involved in the phenomenon under consideration, and (2) those in which the dimensions are expressed in terms of certain fundamental units.

As an illustration of the first we may consider the force of repulsion between two point charges (e, e') of electricity situated at a distance, r, apart in a medium of dielectric constant ϵ. If this force is denoted by f, then $f = ee'/\epsilon r^2$, and we may write $[e^2] = [f\epsilon l^2]$, $[e] = [e^2 f^{-1}l^{-2}]$, etc., where $[\ ]$ denotes that we are concerned with dimensions only; $[l]$ denotes the dimension of length, $[f]$ that of force, etc. These dimensional equations are true whatever be the system of units employed. As they involve quantities, such as force, which can be expressed in terms of other units that are usually considered more fundamental, such dimensional equations will be referred to as "unreduced," in order to distinguish them from those of the second class in which the dimensions are expressed solely in terms of a small number of fundamental units.

It is evident that the dimensions of a quantity in terms of fundamental units can be assigned only in relation to a specific system of units and to a specific method of derivation. For example, (1) if the unit of volume is defined as the volume occupied by a unit mass of water when at its greatest density under a pressure of one atmosphere, then the volume so defined will be independent of the units of length and time, and will vary directly as the unit of mass; we will have $[v] = [m]$. (2) If the unit of volume is defined as the volume occupied by a mass of water (when at its greatest density, etc.) which is equal to the mass of a specified block of platinum, then the volume so defined will not change as we change our units of length, of mass, and of time: that is $[v] = [v]$. In this case $[v]$ is an independent unit and must be so regarded in all dimensional equations. (3) If the unit of volume is defined as the volume of a cube of which the edge is equal to the unit of length then $[v] = [l^3]$. A unit may be defined in any desired unambiguous manner and, in general, the dimensions of the unit will vary from definition to definition.

Dimensional equations of the second type stand in marked contrast to those of the former, in being far less general and in implying the acceptance of a very exactly defined system of units. This, however, is the type of equation which is commonly in mind when dimensional equations are mentioned, and is probably the one which is the more generally useful; the unreduced dimensional expressions (the first type), however, are often simpler, convey more detailed information, and in many cases are to be preferred. For these reasons, unreduced dimensional expressions are to be found in explanations of technical terms (p. 34); they are followed by others, the final one in each case being the fully reduced dimensions on the centimeter, gram, second, degree centigrade absolute, electrostatic system. Wherever necessary, this system of units will be denoted by the symbol cgse in order to distinguish it from the corresponding electromagnetic system, which will be denoted by cgsm. In the conversion tables, dimensional formulae only of the cgse and of the cgsm systems are given. In the cgse system, the fundamental units and their symbols are those of length $[l]$ the centimeter, of mass $[m]$ the gram, of time $[t]$ the mean solar second, of temperature $[T]$ the absolute centigrade degree, and of dielectric constant $[\epsilon]$, that of a vacuum. The fundamental units in the cgsm system differ from those in the cgse system only by the replacement of dielectric constant by magnetic permeability $[\mu]$, the unit being the permeability of a vacuum.

It should be realized that dimensional expressions give no positive information regarding the ultimate nature of the quantity to which they refer; e.g., energy and torque have the same dimensions, but differ vastly in their nature.

Symbols.—(U. S.) before a logarithm denotes that it is based upon the U. S. yard; for explanation of other symbols, see Symbols and Abbreviations, p. 16.

INDEX TO TABLES OF CONVERSION FACTORS

INDEX TO TABLES OF CONVERSION FACTORS.—*Continued*

CONVERSION FACTORS

1. Length [l] (*see also* p. 1)

Unit		Value		Log_{10} (value)
1 angström unit	=	1 0000 × 10⁻⁸ cm		$\overline{8}$.000 0000
1 micron	=	1 0000 × 10⁻⁴ cm		$\overline{4}$.000 0000
1 mil	=	2.5400 × 10⁻³ cm		$\overline{3}$.404 8346
1 inch	=	2.5400 cm	(U. S.)	0.404 8346
1 foot	=	30.480 cm	(U. S.)	1.484 0158
1 yard (U. S.)	=	91.44018 cm		1.961 1371
1 yard (British)	=	91 43992 cm		1.961 1350
1 mile, statute	=	1.6093 km	(U. S.)	0.206 6497
1 light year	=	9.4627 × 10¹² km		12.976 0131
1 astronomical unit	=	1.495 × 10⁸ km		8.174 6712
1 parsec	=	3 084 × 10¹³ km		13.489 09

2. Length⁻¹; Absorptivity; Coefficient of Absorption* [l^{-1}]

1 angstrom⁻¹	=	1.0000 × 10⁸ cm⁻¹		$\overline{8}$.000 0000
1 micron⁻¹	=	1 0000 × 10⁴ cm⁻¹		4.000 0000
1 mil⁻¹	=	393 70 cm⁻¹		2 595 1654
1 inch⁻¹	=	0 39370 cm⁻¹	(U. S.)	$\overline{1}$ 595 1654
1 foot⁻¹	=	3 2808 × 10⁻² cm⁻¹	(U. S.)	$\overline{2}$.515 9842
1 mile⁻¹	=	0 62137 km⁻¹		$\overline{1}$.793 3503

* Coefficient of transmission (τ) is so defined that − log₁₀ τ = coefficient of absorption

3. Mass [m]; Weight (*see also* p. 1)

1 grain	=	64.799 mg		1 811 5677
1 carat (metric)	=	200 000 mg		2 301 0300
1 ounce (avoirdupois)	=	28.350 g		1 452 5458
1 ounce (apothecary) or (troy)	=	31.103 g		1 492 8090
1 pound (avoirdupois)	=	453 59243 g		2.656 6658
1 pound (apothecary) or (troy)	=	373.2417 g		2.571 9902
1 ton, short (2000 pounds)	=	907 185 kg		2 957 6958
1 ton, long (2240 pounds)	=	1016 047 kg		3 006 9138
1 slug (g_s)	=	14 594 kg		1.164 1707
1 gram mole	=	M. W.† g·		
1 molecule/M. W.†	=	1.6498 × 10⁻²⁴ g		$\overline{24}$.217 4366
1 assay ton	=	29 1667 g		1 464 8868

† M. W denotes the molecular weight of the substance

4. Mass⁻¹ [m^{-1}]

1 grain⁻¹	=	1 5432 × 10⁻² mg⁻¹		2 188 4323
1 ounce⁻¹ (avoirdupois)	=	3.5274 × 10⁻² g⁻¹		$\overline{2}$ 547 4542
1 ounce⁻¹ (troy)	=	3 2151 × 10⁻² g⁻¹		$\overline{2}$.507 1910
1 pound⁻¹ (avoirdupois)	=	2 2046 × 10⁻³ g⁻¹		$\overline{3}$.343 3342
1 ton⁻¹ (2000 pounds)	=	11 0231 × 10⁻⁴ kg⁻¹		$\overline{3}$.042 3042
1 ton⁻¹ (2240 pounds)	=	9 8421 × 10⁻⁴ kg⁻¹		$\overline{4}$.993 0862
1 (gram mole)⁻¹	=	†(M. W.)⁻¹ g⁻¹		

† M. W. denotes the molecular weight of the substance

5. Time [t]

1 second, mean solar	=	1 00273791 sidereal sec		0.001 1874
1 second, sidereal	=	0 997270 sec (mean solar)		$\overline{1}$ 998 8126
1 hour (tropical, mean solar)	=	3 6000 × 10³ sec (mean solar)		3.556 3025
1 day (tropical, mean solar)	=	8 6400 × 10⁴ sec (mean solar)		4.936 5137
1 day (sidereal)	=	8 6164 × 10⁴ sec (mean solar)		4.935 3263
1 year (tropical, mean solar)	=	31 5569 × 10⁶ sec (mean solar)		7.499 0946
1 year (tropical, mean solar)	=	365 2422 day (mean solar)		2 562 5809

CONVERSION FACTORS.—*Continued*

6. Time^{-1}; Frequency; "Velocity" of a Process $[t^{-1}]$

1 second^{-1} (sidereal)	=	1 002738 sec^{-1} (mean solar)	0.001 1874
1 minute^{-1} (mean solar)	=	1 66667 × 10^{-2} sec^{-1} (mean solar)	$\bar{2}$.221 8487
1 hour^{-1} (mean solar)	=	2.77778 × 10^{-4} sec^{-1} (mean solar)	$\bar{4}$.443 6975
1 day^{-1} (mean solar)	=	1 15741 × 10^{-5} sec^{-1} (mean solar)	$\bar{5}$.063 4863
1 year^{-1} (mean solar)	=	3 16888 × 10^{-8} sec^{-1} (mean solar)	$\bar{8}$.500 9054
1 year^{-1} (mean solar)	=	2.73791 × 10^{-3} day^{-1} (mean solar)	$\bar{3}$.437 4191
1 electron-volt, quantum^{-1}	=	2 4292 × 10^{14} sec^{-1} (mean solar)	14.385 4575
1 joule per mole, N_0^{-1} quantum^{-1}	=	2 5173 × 10^{9} sec^{-1} (mean solar)	9.400 9301
1 velocity of light, (angström unit)$^{-1}$	=	2 9986 × 10^{18} sec^{-1} (mean solar)	18.476 9185
1 velocity of light, millimicron^{-1}	=	2 9986 × 10^{17} sec^{-1} (mean solar)	17.476 9185
1 velocity of light, micron^{-1}	=	2 9986 × 10^{14} sec^{-1} (mean solar)	14.476 9185
1 velocity of light, millimeter^{-1}	=	2 9986 × 10^{11} sec^{-1} (mean solar)	11.476 9185
1 velocity of light, meter^{-1}	=	2 9986 × 10^{8} sec^{-1} (mean solar)	8.476 9185

7. Angle $[\theta]$

1 radian	=	57.29578 degree	1.758 1226
1 circumference	=	6 28319 radian	0 798 1799
1 quadrant	=	1.57080 radian	0.196 1199
1 degree	=	1 74533 × 10^{-2} radian	$\bar{2}$.241 8774
1 minute	=	2 90888 × 10^{-4} radian	$\bar{4}$.463 7261
1 second	=	4 84814 × 10^{-6} radian	$\bar{6}$.685 5749

8. Angle^{-1} $[\theta^{-1}]$

1 circumference^{-1}	=	0 159155 radian^{-1}	$\bar{1}$.201 8201
1 degree^{-1}	=	57 29578 radian^{-1}	1.758 1226
1 minute^{-1}	=	3.43775 × 10^{3} radian^{-1}	3.536 2739
1 second^{-1}	=	2 06265 × 10^{5} radian^{-1}	5.314 4251

9. Solid Angle $[\omega]$

Entire space	=	12 5664 steradian	1.099 2099
1 hemisphere	=	6 2832 steradian	0.798 1799
1 square degree	=	3 0462 × 10^{-4} steradian	$\bar{4}$.483 7548

10. Solid Angle^{-1} $[\omega^{-1}]$

Entire space^{-1}	=	7 9577 × 10^{-2} steradian^{-1}	$\bar{2}$.900 7901
1 hemisphere^{-1}	=	1 5916 × 10^{-1} steradian^{-1}	$\bar{1}$.201 8201
1 square degree^{-1}	=	3.2828 × 10^{3} steradian^{-1}	3.516 2452

11. Temperature $[T]$ (*See also* Thermometry, p. 52)

Fahrenheit.	$x°$ F	$= (\frac{5}{9})(x - 32)°C$
Réaumur	$x°$ R	$= (\frac{5}{4})x°C$
Absolute (Centigrade).	$x°$ K	$= (x - T_0)°C$
Absolute (Fahrenheit)	$x°$ Rankine	$= (\frac{5}{9})(x - 491.58)°C$

12. Degree^{-1} (Thermometric); Expansivity; Curie's Constant (magnetic) $[T^{-1}]$

1 per degree F	=	1.8000 per degree C	0 255 2725
1 per degree R	=	0 8000 per degree C	1 903 0900
1 per degree K	=	1 000 per degree C	0.000 0000

13. Luminous Flux $[\psi]$

By definition, the total luminous flux emitted by a point source of one spherical candle power is 4π *lumen*.

14. Dielectric Constant; Electrical Inductivity $[\epsilon]$; $[\mu^{-1}l^{-2}t^{2}]$

Specific inductive capacity is of zero dimensions It is numerically equal to the dielectric constant expressed in cgse or in fpse units.

1 cgsm unit	=	8 9916 × 10^{20} cgse unit	20 953 8370
1 fpse unit	=	1 0000 cgse unit	0.000 0000
1 fpsm unit	=	1 0764 × 10^{-3} cgsm unit	3.031 9684
1 fpsm unit	=	9 6784 × 10^{17} cgse unit	17.985 8054

15. Magnetic Permeability; Susceptibility $[\epsilon^{-1}l^{-2}t^{2}]$; $[\mu]$

1 cgse unit	=	8 9916 × 10^{20} cgsm unit	20.953 8370
1 fpsm unit	=	1.0000 cgsm unit	0.000 0000
1 fpse unit	=	1.0764 × 10^{-3} cgse unit	3.031 9684
1 fpse unit	=	9 6784 × 10^{17} cgsm unit	17.985 8054

INTERNATIONAL CRITICAL TABLES

CONVERSION FACTORS.—*Continued*

16. Area $[l^2]$

1 circular millimeter	7.8540 × 10⁻³ cm²	3.895 0899
1 circular mil	5.0671 × 10⁻⁶ cm²	(U. S.) $\bar{6}$.704 7591
1 square inch	6.4516 cm²	(U. S.) 0.809 6692
1 square foot	9.2903 × 10² cm²	(U. S.) 2.968 0316
1 square yard	8.3613 × 10³ cm²	(U. S.) 3.922 2742
1 square mile	2.5900 km²	(U. S.) 0.413 2995
1 are	1.0000 × 10² m²	2.000 0000
1 hectare	1.0000 × 10⁴ m²	4.000 0000
1 acre	4.0469 × 10³ m²	3.607 1196

17. Area⁻¹ $[l^{-2}]$

1 (circular millimeter)⁻¹	127.324 cm⁻²	2.104 9101
1 millimeter⁻²	100.0000 cm⁻²	2.000 0000
1 meter⁻²	0.0001 cm⁻²	$\bar{4}$.000 0000
1 (circular mil)⁻¹	1.9735 × 10⁵ cm⁻²	(U. S.) 5.295 2409
1 inch⁻²	0.15500 cm⁻²	(U. S.) $\bar{1}$.190 3308
1 foot⁻²	1.0764 × 10⁻³ cm⁻²	(U. S.) $\bar{3}$.031 9684
1 yard⁻²	1.19599 × 10⁻⁴ cm⁻²	(U. S.) $\bar{4}$.077 7258
1 mile⁻²	0.38610 km⁻²	(U. S.) $\bar{1}$.586 7005

18. Volume $[l^3]$ or $[v]$

1 liter	1000.027 cm³	3.000 0117
1 cubic inch	16.387 cm³	(U. S.) 1.214 5038
1 cubic foot	2.8317 × 10⁴ cm³	(U. S.) 4.452 0474
1 cubic yard	7.6456 × 10⁵ cm³	(U. S.) 5.883 4112
1 gallon (U. S.)	3.7854 × 10³ cm³	3.578 1157
1 gallon (British)	4.5461 × 10³ cm³	3.657 6376
1 bushel (U. S.)	3.5239 × 10⁴ cm³	4.547 0271
1 bushel (British)	3.6369 × 10⁴ cm³	4.560 7276
1 quart, dry (U. S.)	1101.23 cm³	3.041 8771
1 quart, liquid (U. S.)	946.358 cm³	2.976 0557
1 quart (British)	1136.521 cm³	3.055 5776
1 fluid ounce (U. S.)	29.5737 cm³	1.470 9057
1 fluid ounce (British)	28.4130 cm³	1.453 5176

19. Volume⁻¹ $[l^{-3}]$ or $[v^{-1}]$

1 liter⁻¹	9.9997 × 10⁻⁴ cm⁻³	$\bar{4}$.999 9883
1 inch⁻³	6.1023 × 10⁻² cm⁻³	(U. S.) $\bar{2}$.785 4962
1 foot⁻³	3.5314 × 10⁻⁵ cm⁻³	(U. S.) $\bar{5}$.547 9526
1 yard⁻³	1.3079 m⁻³	(U. S.) 0.116 5888
1 gallon⁻¹ (U. S.)	2.6417 × 10⁻⁴ cm⁻³	$\bar{4}$.421 8843
1 gallon⁻¹ (British)	2.1997 × 10⁻⁴ cm⁻³	$\bar{4}$.342 3624
1 quart⁻¹, dry (U. S.)	9.0808 × 10⁻⁴ cm⁻³	$\bar{4}$.958 1229
1 quart⁻¹, liquid (U. S.)	1.0567 × 10⁻³ cm⁻³	$\bar{3}$.023 9443
1 quart⁻¹ (British)	8.7988 × 10⁻⁴ cm⁻³	$\bar{4}$.944 4224
1 (fluid ounce)⁻¹ (U. S.)	3.3814 × 10⁻² cm⁻³	$\bar{2}$.529 0943
1 (fluid ounce)⁻¹ (British)	3.5195 × 10⁻² cm⁻³	$\bar{2}$.546 4824

20. Length Degree⁻¹ $[lT^{-1}]$

1 inch per °F	4.5720 cm per °C	0.660 1071
1 foot per °F	54.864 cm per °C	1.739 2883
1 meter per °C	100.00 cm per °C	2.000 0000

21. Mass⁻¹ Degree⁻¹ $[m^{-1}T^{-1}]$

1 per gram °F	1.8000 per gram °C	0.255 2725
1 per pound °F	3.9683 × 10⁻³ per gram °C	$\bar{3}$.598 6067
1 per pound °C	2.2046 × 10⁻³ per gram °C	$\bar{3}$.343 3342

22. Area⁻¹ Time⁻¹ $[l^{-2}t^{-1}]$

1 foot⁻² second⁻¹	3.8750 cm⁻² hr⁻¹	(U. S.) 0.588 2709
1 foot⁻² second⁻¹	1.0764 × 10⁻³ cm⁻² sec⁻¹	(U. S.) 3.031 9684
1 mile⁻² second⁻¹	1.2184 × 10⁻³ cm⁻² yr⁻¹	(U. S.) $\bar{3}$.085 7951
1 meter⁻² second⁻¹	3.600 × 10⁻¹ cm⁻² hr⁻¹	$\bar{1}$.556 3025

CONVERSION FACTORS.—*Continued*

23. Velocity [lt⁻¹]

1 foot per second	=	30 4801	cm sec⁻¹	(U. S.) 1.484 0158
1 foot per minute	=	0 5080	cm sec⁻¹	(U. S.) 1.705 8645
1 mile per hour	=	44 7041	cm sec⁻¹	(U. S.) 1 650 3472
1 mile per minute	=	2.6822 × 10³ cm sec⁻¹		(U. S.) 3 428 4984
1 meter per minute	=	1 6667	cm sec⁻¹	0.221 8487
1 kilometer per hour	=	27 7778	cm sec⁻¹	1.443 6975
Velocity of light	=	2 9986 × 10¹⁰ cm sec⁻¹		10.476 9185

24. Acceleration [lt⁻²]

1 foot per second²	=	30.480	cm sec⁻²	(U. S.) 1.484 0158
1 mile per hour second	=	44 704	cm sec⁻²	(U S.) 1.650 3472
1 mile per hour minute	=	0 74507	cm sec⁻²	(U. S.) 1.872 1959
1 meter per second²	=	100 000	cm sec⁻²	2.000 0000
1 kilometer per hour second	=	27 778	cm sec⁻²	1 443 6975
Gravity, standard	=	980 665	cm sec⁻²	2 991 5207
Gravity, standard	=	32 171	ft sec⁻²	(U. S.) 1 507 5049

25. Angular Velocity [θt⁻¹]

1 revolution per day	=	7 2722 × 10⁻⁵ radian sec⁻¹	5 861 6062
1 revolution per minute	=	1 0472 × 10⁻¹ radian sec⁻¹	1 020 0286
1 revolution per second	=	6 2832 radian sec⁻¹	0.798 1799
1 degree per second	=	1 7453 × 10⁻² radian sec⁻¹	2.241 8774

26. Angular Acceleration [θt⁻²]

1 revolution per second²	=	6 2832 radian sec⁻²	0.798 1799
1 revolution per minute²	=	1.7453 × 10⁻³ radian sec⁻²	3.241 8773
1 revolution per minute second	=	0 10420 radian sec⁻²	1 020 0286

27. Twist; Rotatory Power [θl⁻¹]

1 degree per inch	=	6 8714 × 10⁻³ radian cm⁻¹	(U. S.) 3.837 0428
1 degree per foot	=	5 7261 × 10⁻⁴ radian cm⁻¹	(U. S.) 4.757 8616
1 degree per centimeter	=	1 7453 × 10⁻² radian cm⁻¹	2 241 8774
1 minute per centimeter	=	2 9089 × 10⁻⁴ radian cm⁻¹	4 463 7261

28. Density; Volume Concentration; Solubility (Non-gases) [ml⁻³] or [mv⁻¹] (*See also* Hydrometer Tables, p. 31)

1 gram per milliliter*	=	0 999973 g cm⁻³	1 999 9883
1 pound per inch³	=	27 680 g cm⁻³	(U. S.) 1 442 1621
1 pound per foot³	=	0 016018 g cm⁻³	(U. S.) 2 204 6183
1 pound per gallon (U. S.)	=	0 119826 g cm⁻³	1 078 5502
1 pound per gallon (British)	=	0 099776 g cm⁻³	2 999 0282
1 slug per foot³ (gₛ)	=	0 5154 g cm⁻³	(U. S.) 1 712 1233
Mercury† at 0°C	=	15 5951 g cm⁻³	1 192 9882

* Numerically equal to specific gravity t° 4°   † Internationally accepted conventional value to be used in expressing pressures in terms of columns of mercury.

29. Mass Concentration [m₁m₂⁻¹]

(This quantity involves two distinct units of mass; when the two units are the same, the concentration is called the "titer," or is denoted as a per cent.)

1 gram per ton (2000 pound)	=	1 1023 mg per kilogram	0.042 3042
1 gram per ton (2240 pound)	=	0 9842 mg per kilogram	1.993 0862
1 milligram per assay ton	=	*34.286 mg per kilogram	1 535 1132
1 ounce (av.) per ton (2000 lb.)	=	31 2500 mg per kilogram	1.494 8500
1 ounce (av) per ton (2240 lb.)	=	27 9018 mg per kilogram	1.445 6320
1 pound (av) per ton (2000 lb.)	=	500 000 mg per kilogram	2 698 9700
1 pound (av.) per ton (2240 lb.)	=	446 429 mg per kilogram	2 649 7520
1 gram per ton (metric)	=	1 0000 mg per kilogram	0.000 0000
1 karat†	=	41 667 mg per gram	1 619 7888

* Equals one troy ounce per 2000 lb. av   † 1 of gold to 24 of mixture

30. Force [mlt⁻²]

1 gram weight (g.)	=	980 665 dyne	2 991 5207
1 poundal	=	1.3825 × 10⁴ dyne	(U. S.) 4.140 6816
1 pound weight (g.)	=	4.4482 × 10⁵ dyne	5 648 1864
1 ton weight (2000 lb.) (g.)	=	8.8964 × 10⁸ dyne	8.949 2164
1 ton weight (2240 lb.) (g.)	=	9 9640 × 10⁸ dyne	8 998 4344

CONVERSION FACTORS.—Continued

31. Force^{-1} $[m^{-1}l^{-1}t^2]$

1 (gram weight)$^{-1}$ (g.)	=	1 0917 × 10^{-3} dyne^{-1}	$\bar{3}$.008 4793
1 poundal^{-1}	=	7 2330 × 10^{-5} dyne^{-1}	$\bar{5}$.859 3184
1 (pound weight)$^{-1}$ (g.)	=	2.2481 × 10^{-6} dyne^{-1}	$\bar{6}$.351 8136

32. Torque; Moment of a Force $[ml^2t^{-2}]$

1 pound-foot (g.)	=	1 3558 × 10^7 dyne cm	(U. S.) 7 132 2022
1 pound-inch (g.)	=	1 1298 × 10^6 dyne cm	(U. S.) 6.053 0210
1 kilogram-meter (g.)	=	9.8066 × 10^7 dyne cm	7.991 5207
1 poundal-foot	=	4 2140 × 10^5 dyne cm	(U. S.) 5.624 6974

33. Stress; Pressure; Tension; Young's Modulus; Modulus of Rigidity; Modulus of Compression; Bulk Modulus; Coefficient of Skin Friction $[ml^{-1}t^{-2}]$

1 barye	=	1 0000	dyne cm^{-2}	0.000 0000
1 bar	=	*1 0000 × 10^6 dyne cm^{-2}		6.000 0000
1 gram weight per cm^2 (g.)	=	980 665	dyne cm^{-2}	2 991 5207
1 kilogram weight per m^2 (g.)	=	98 0665	dyne cm^{-2}	1.991 5207
1 kilogram weight per mm^2 (g.)	=	9 8066 × 10^7 dyne cm^{-2}		7.991 5207
1 pound weight per in.^{2} (g.)	=	6 8947 × 10^4 dyne cm^{-2}		(U. S.) 4.838 5173
1 pound weight per ft.^{2} (g.)	=	4 7880 × 10^2 dyne cm^{-2}		(U. S.)2.680 1548
1 ton (2000 lb.) weight per in.^{2} (g.)	=	1.3789 × 10^8 dyne cm^{-2}		(U. S.) 8.139 5473
1 ton (2240 lb.) weight per in.^{2} (g.)	=	1 5444 × 10^8 dyne cm^{-2}		(U. S.) 8 188 7653
1 ton (2000 lb.) weight per ft ^{2} (g.)	=	9.5760 × 10^5 dyne cm^{-2}		(U. S.) 5.981 1848
1 ton (2240 lb.) weight per ft.^{2} (g.)	=	10 7251 × 10^5 dyne cm^{-2}		(U. S.) 6 030 4028
1 centimeter of water at 4°C (g.)	=	9.80638 × 10^2 dyne cm^{-2}		2.991 5090
1 inch of water at 4°C (g.)	=	2.49082 × 10^3 dyne cm^{-2}		(U. S.) 3.396 3436
1 centimeter of mercury at 0°C (g.)	=	1 33322 × 10^4 dyne cm^{-2}		4 124 9031
1 inch of mercury at 0°C (g.)	=	3.38639 × 10^4 dyne cm^{-2}		.(U. S.) 4 529 7377
1 normal atmosphere (g.)	=	1 01325 × 10^6 dyne cm^{-2}		6 005 7166

* This value accords with the only internationally accepted use of this term, but "bar" has also been used to denote a pressure of one dyne per cm^2.

34. Stress^{-1}; Compressibility $[m^{-1}lt^2]$

1 centimeter^2 per gram weight (g.)	=	1 0197 × 10^{-3} cm^2 dyne^{-1}	$\bar{3}$ 008 4793
1 centimeter^2 per kilogram weight (g.)	=	1 0197 × 10^{-6} cm^2 dyne^{-1}	$\bar{6}$ 008 4793
1 millimeter^2 per kilogram weight (g.)	=	1 0197 × 10^{-8} cm^2 dyne^{-1}	$\bar{8}$.008 4793
1 inch^2 per pound weight (g.)	=	1 4504 × 10^{-5} cm^2 dyne^{-1}	(U. S.) $\bar{5}$.161 4827
1 inch^2 per ton weight (2000 lb.) (g.)	=	7 2519 × 10^{-9} cm^2 dyne^{-1}	(U. S.) $\bar{9}$.860 4527
1 inch^2 per ton weight (2240 lb.) (g.)	=	6.4749 × 10^{-9} cm^2 dyne^{-1}	(U. S.) 9 811 2347
1 foot^2 per pound weight (g.)	=	2.0886 × 10^{-3} cm^2 dyne^{-1}	(U. S.) $\bar{3}$ 319 8452
1 (centimeter of water at 4°C)$^{-1}$ (g.)	=	1 0197 × 10^{-3} cm^2 dyne^{-1}	$\bar{3}$ 008 4910
1 (inch of water at 4°C)$^{-1}$ (g.)	=	4.0147 × 10^{-4} cm^2 dyne^{-1}	(U. S.) $\bar{4}$ 603 6564
1 (centimeter of mercury at 0°C)$^{-1}$ (g.)	=	7.5006 × 10^{-5} cm^2 dyne^{-1}	$\bar{5}$.875 0969
1 (inch of mercury at 0°C)$^{-1}$ (g.)	=	2 9530 × 10^{-5} cm^2 dyne^{-1}	(U. S.) $\bar{5}$ 470 2623
1 (normal atmosphere)$^{-1}$ (g.)	=	9.8692 × 10^{-7} cm^2 dyne^{-1}	$\bar{7}$ 994 2834

35. Work; Energy; Heat $[ml^2t^{-2}]$

1 centimeter-dyne	=	1 0000	erg	0 000 0000
1 joule (absolute)	=	1 0000 × 10^7 erg		7.000 0000
1 joule (International) (v)	=	1 00032	joule (abs.)	0 000 1390
1 meter-kilogram (g.)	=	9 80665	joule (abs.)	0.991 5207
1 foot-pound (g.)	=	1 35582	joule (abs.)	(U. S.) 0.132 2022
1 liter-atmosphere (normal) (g.)	=	101.328	joule (abs.)	2.005 7283
1 liter-atmosphere (45° lat.)	=	*101 323	joule (abs.)	2.005 7067
1 cubic centimeter-atmosphere (normal) (g.)	=	0 101325	joule (abs.)	$\bar{1}$.005 7166
1 horse-power hour (HP hr.) (g.)	=	2 6845 × 10^6 joule (abs.)		(U. S.) 6.428 8674
1 horse-power hour (electrical, U. S., British)	=	2 6856 × 10^6 joule (abs.)		6 429 0413
1 cheval-vapeur heure (g.)	=	2 6478 × 10^6 joule (abs.)		6 422 8845
1 kilowatt-hour (abs.)	=	3.6000 × 10^6 joule (abs.)		6 556 3025
1 International volt (v) faraday	=	9.6541 × 10^4 joule (abs.)		4 984 7097
1 International volt (v) electronic charge	=	1 5927 × 10^{-19} joule (abs.)		$\overline{19}$.202 1463
1 gram calorie (20°C)	=	4 181	joule (abs.)	0.621 2802
1 gram calorie (15°C)	=	4 185	joule (abs.)	0.621 6955
1 gram calorie (mean)	=	4 186	joule (abs.)	0.621 7992
1 British Thermal Unit (39°F)	=	1060 4	joule (abs.)	3 025 4697
1 British Thermal Unit (mean)	=	1054 8	joule (abs.)	3.023 1701
1 British Thermal Unit (60°F)	=	1054 6	joule (abs.)	3.023 0878
1 Centigrade Thermal Unit (15°C)	=	1 8983 × 10^3 joule (abs.)		3.278 3613

* g_n = 980 616 cm sec^{-1}.

CONVERSION FACTORS.—Continued

36. Power $[ml^2t^{-3}]$

1 watt (absolute)	=	1.0000 $\times$ 10⁷ erg sec⁻¹	7.000 0000
1 watt (International) (v)	=	1.00032 watt (abs.)	0.000 1390
1 meter-kilogram per second (g_s)	=	9.80665 watt (abs.)	0.991 5207
1 foot-pound per second (g_s)	=	1.35582 watt (abs.)	(U. S.) 0.132 2022
1 horsepower, electrical (U. S., British)	=	*746 00 watt (abs.)	2.872 7388
1 horsepower, electrical (Continental Europe)	=	*736.00 watt (abs.)	2.866 0778
1 horsepower (HP) (g_s)	=	†745.70 watt (abs.)	2.872 5649
1 cheval-vapeur (g_s)	=	735.499 watt (abs.)	2.866 5820

* Defined in terms of the watt, commonly used in rating electrical machinery.   † Defined as 550 ft. lb per sec

37. Action $[ml^2t^{-1}]$

1 Planck's quantum	=	6.554 $\times$ 10⁻²⁷ erg sec	27.816 5064
1 volt electronic-charge second	=	2.4292 $\times$ 10¹⁴ quanta	14.385 4575
1 volt faraday second	=	1 4724 $\times$ 10³⁸ quanta	38.168 0209
1 joule second	=	1 5258 $\times$ 10³³ quanta	33.183 4936
1 calorie (15°C) second	=	6.3854 $\times$ 10³³ quanta	33.805 1891
1 joule second/N_0*	=	2.5173 $\times$ 10⁹ quanta	9.400 9302
1 calorie (15°C) second/N_0*	=	1 0535 $\times$ 10¹⁰ quanta	10.022 6257

* N_0 denotes Avogadro's number, the number of molecules per gram mole

38. Fluidity $[m^{-1}lt]$ (See also 39)

1 rhe	=	1.0000 poise⁻¹	0.000 0000

39. Viscosity $[ml^{-1}t^{-1}]$

1 poise	=	1 000 gram cm⁻¹ sec⁻¹	0.000 0000
1 gram weight sec cm⁻² (g_s)	=	980 665 poise	2.991 5207
1 pound weight sec inch⁻² (g_s)	=	6.895 $\times$ 10⁴ poise	(U. S.) 4.838 5173
1 pound weight sec foot⁻² (g_s)	=	4.788 $\times$ 10² poise	(U. S.) 2.680 1548

40. Kinematic Viscosity $[l^2t^{-1}]$

1 poise centimeter² gram⁻¹	=	1.000 cm² sec⁻¹	0.000 0000
1 poise inch² gram⁻¹	=	16.387 cm² sec⁻¹	1.214 5038
1 inch² second⁻¹	=	6.451 cm² sec⁻¹	(U. S.) 0.809 6692
1 poise foot² pound⁻¹	=	62.43 cm² sec⁻¹	(U. S.) 1.795 3817

41. Diffusivity; Diffusion, Coefficient of $[l^2t^{-1}]$

All quantities of the thing diffusing are to be expressed in terms of the same units. Heat diffusivity is numerically equal to heat conductivity divided by the product of the density times the heat capacity (per unit of mass); all must be expressed in the same system of units.

1 liter centimeter⁻¹ day⁻¹	=	1 1574 $\times$ 10⁻² cm² sec⁻¹	2.063 4980
1 centimeter² day⁻¹	=	1 1574 $\times$ 10⁻⁵ cm² sec⁻¹	5 063 4863
1 inch² sec⁻¹	=	6 4516 cm² sec⁻¹	(U. S.) 0.809 6692

42. Surface Tension $[mt^{-2}]$ (See also Capillary Constant, Table 43)

1 milligram weight per mm (g_s)	=	9 80665 dyne cm⁻¹	0.991 5207
1 milligram weight per inch (g_s)	=	0 38609 dyne cm⁻¹	(U. S.) 1.586 6861
1 erg per centimeter²	=	1 00000 dyne cm⁻¹	0.000 0000
1 erg per millimeter²	=	100 00000 dyne cm⁻¹	2.000 0000

43. (Capillary Constant)² $[l^2]$

The term "Capillary Constant" is used in two different senses; viz, either to denote $a_1 = \sqrt{\gamma/\rho g}$, or to denote $a_2 = \sqrt{2\gamma/\rho g}$. English authors generally follow the former practice, and German authors the latter; neither use the subscript. γ denotes the surface tension, g the acceleration of gravity, and ρ the positive difference in the densities of the adjacent fluids

1 inch²	=	6 451 cm²	0 809 6692
1 millimeter² (a_1^2) (g_s)	=	*9 807 dyne cm⁻¹ per (g cm⁻²)	0.991 5207
1 millimeter² (a_2^2) (g_s)	=	*4 903 dyne cm⁻¹ per (g cm⁻²)	0 690 4907
1 inch² (a_1^2) (g_s)	=	*6 327 $\times$ 10² dyne cm⁻¹ per (g cm⁻²)	(U. S.) 3.801 1899
1 inch² (a_2^2) (g_s)	=	*3 163 $\times$ 10² dyne cm⁻¹ per (g cm⁻²)	(U. S.) 3 500 1599

* To convert a^2, when referred to ρ_s, to surface tension in dynes per cm, multiply a^2 by the factor given in this table and by the difference in the densities (gram per cm³) of the adjacent fluids, if a^2 is referred to g, multiply the resulting product by g/g_s

44. Thermal Conductivity $[T^{-1}mlt^{-3}]$

The dimensions practically employed in expressing this property are (Heat Area⁻¹ Time⁻¹ per Degree Length⁻¹). Other conversion factors may be obtained by combining those of Tables 35 (Heat), 22 (Area⁻¹ Time⁻¹) and 20 (Length Degree⁻¹).

1 calorie (15°) cm⁻² sec⁻¹ (°C, cm⁻¹)⁻¹	=	4.185 joules (abs.) cm⁻² sec⁻¹ (°C, cm⁻¹)⁻¹	0.621 6955
1 calorie (20°) cm⁻² sec⁻¹ (°C, cm⁻¹)⁻¹	=	4.181 joules (abs.) cm⁻² sec⁻¹ (°C, cm⁻¹)⁻¹	0.621 2802

CONVERSION FACTORS.—*Continued*

44. Thermal Conductivity [$T^{-1}mll^{-3}$].—*Continued*

1 British Thermal Unit (39°F) ft.$^{-2}$ sec^{-1} (°F, in.$^{-1}$)$^{-1}$ =	5.218 joules (abs.) cm^{-2} sec^{-1} (°C, cm^{-1})$^{-1}$	0.717 5452
1 British Thermal Unit (mean) ft.$^{-2}$ sec^{-1} (°F, in.$^{-1}$)$^{-1}$ =	5.191 joules (abs.) cm^{-2} sec^{-1} (°C, cm^{-1})$^{-1}$	0.715 2456
1 British Thermal Unit (60°F) ft.$^{-2}$ sec^{-1} (°F, in^{-1})$^{-1}$ =	5 190 joules (abs.) cm^{-2} sec^{-1} (°C, cm^{-1})$^{-1}$	0 715 1633

45. Intensity of Radiation [ml^{-3}] or [$ml^{-1}t^{-2}$]

The dimensions depend upon the point of view, when the receptor is considered, they are [Energy, Area^{-1}, Time^{-1}]; when the radiation itself is considered they are [Energy, Volume^{-1}] Conversion from one to the other involves the velocity of propagation, if this is the velocity of light in vacuo, the factors are as given below, if the velocity is a cm sec^{-1}, the factors given must be multiplied by $v/(2.9986 \times 10^{10})$ For other units, combine these factors with those of Tables 19 (Volume^{-1}), 22 (Area^{-1} Time^{-1}), and 45 (Energy)

1 erg cm^{-3}	=	2 9986 $\times$ 10^{10} erg cm^{-2} sec^{-1}	10 476 9185
1 foot-pound ft^{-3} (g_v)	=	1 4357 $\times$ 10^{13} erg cm^{-2} sec^{-1}	(U. S.) 13.157 0733

46. Luminous Intensity of a Source in a Given Direction [$\psi \omega^{-1}$]

By definition of the lumen, a source of one spherical candle power emits 4π (= 12.566) lumens. (*See also* Photometric Standards, in another section (consult index))

1 candle, International	=	1 0000 Int. lumen per steradian	0.000 0000
1 pentane candle	=	1 0 Int. candle	
1 Hefner unit	=	0 9$_0$ Int. candle	
1 Carcel unit	=	9 6 Int. candle ⎫ Approximate	
1 bougie decimale	=	1 0 Int. candle	
1 English sperm candle	=	1 0 Int. candle	

47. Illumination of a Surface [ψl^{-2}]

1 lux	=	1 000 lumen meter^{-2}	0 000 0000
1 meter-candle	=	1 000 lumen meter^{-2}	0 000 0000
1 phot	=	1 000 $\times$ 10^4 lumen meter^{-2}	4 000 0000
1 foot-candle	=	10 764 lumen meter^{-2}	(U. S.) 1 031 9684
1 lumen foot^{-2}	=	10 764 lumen meter^{-2}	(U. S.) 1 031 9684

48. Surface Brightness [$\psi l^{-2}\omega^{-1}$]

1 lumen centimeter^{-2} steradian^{-1}	=	1 0000 lambert	0 000 0000
1 lumen foot^{-2} steradian^{-1}	=	1 0764 millilambert	(U. S.) 0 031 9684
1 candle centimeter^{-2}	=	3 1416 $\times$ 10^3 millilambert	3 497 1499
1 candle inch^{-2}	=	4 8695 $\times$ 10^2 millilambert	(U S.) 2 687 4807

49. Electrical Quantity; Charge; Total Electric Displacement; Flux of Induction [$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$]; [$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}$]

1 absolute coulomb	=	1 00010 Int. coulomb (v)	0.000 0434
1 absolute coulomb	=	1 00007 Int. coulomb (a)	0 000 0304
1 International coulomb (v)	=	0 99990 abs. coulomb	$\bar{1}$.999 9566
1 International coulomb (a)	=	0 99993 abs. coulomb	1.999 9696
1 cgsm unit	=	10 0000 abs. coulomb	1.000 0000
1 cgsm unit	=	*2 9986 $\times$ 10^{10} cgse unit	10 476 9185
1 cgse unit	=	3 3349 $\times$ 10^{-10} abs. coulomb	$\bar{10}$.523 0815
1 fpsm unit	=	1 1758 $\times$ 10^2 cgsm unit	2 070 3408
1 fpse unit	=	3 5839 $\times$ 10^3 cgse unit	3.554 3566
1 fpse unit	=	1 1952 $\times$ 10^{-6} abs. coulomb	6 077 4381
1 ampere-hour (abs.)	=	3 6000 $\times$ 10^4 abs. coulomb	3 556 3025
1 electronic charge	=	1 5921 $\times$ 10^{-19} abs. coulomb	$\bar{19}$ 201 9639
1 electronic charge	=	4 774 $\times$ 10^{-10} cgse unit	$\bar{10}$ 678 8824
1 faraday	=	9 6500 $\times$ 10^4 abs. coulomb	4 984 5273
1 faraday	=	9 6510 $\times$ 10^4 Int. coulomb (v)	4.984 5707
1 faraday	=	9 6507 $\times$ 10^4 Int. coulomb (a)	4.984 5577
1 faraday	=	2 89365 $\times$ 10^{14} cgse unit	14 461 4458

* Value of c; experimental value = 2 9979 $\times$ 10^{10} (Rosa and Dorsey, *Bull. U. S. Bur. Standards*, **3**: 433, 07)

50. Electrical Quantity^{-1}; Charge^{-1}; Total Electric Displacement^{-1}; Flux of Induction^{-1} [$\epsilon^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{3}{2}}t$]; [$\mu^{\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}$]

1 absolute coulomb^{-1}	=	0 99990 Int. coulomb^{-1} (v)	1 999 9566
1 absolute coulomb^{-1}	=	0 99993 Int. coulomb^{-1} (a)	$\bar{1}$ 999 9696
1 cgsm unit^{-1}	=	0 1000 abs. coulomb^{-1}	$\bar{1}$.000 0000
1 cgse unit^{-1}	=	2 9986 $\times$ 10^9 abs. coulomb^{-1}	9 476 9185
1 ampere-hour^{-1}	=	2 7778 $\times$ 10^{-4} abs. coulomb^{-1}	$\bar{4}$.443 6975
1 faraday^{-1}	=	1 0363 $\times$ 10^{-5} abs. coulomb^{-1}	$\bar{5}$.015 4727
1 electronic charge^{-1}	=	6 281 $\times$ 10^{18} abs. coulomb^{-1}	18 798 0361

CONVERSION FACTORS.—*Continued*

51. Electrical Current $[\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}]$; $[\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}]$

absolute ampere	=	1 00010	Int. ampere (v)	0.000 0434
absolute ampere	=	1 00007	Int. ampere (a)	0.000 0304
International ampere (v)	=	0 99990	abs. ampere	$\bar{1}$.999 9566
International ampere (a)	=	0 99993	abs. ampere	$\bar{1}$.999 9696
cgsm unit	=	10 0000	abs. ampere	1 000 0000
cgse unit	=	3 3349 $\times$ 10⁻¹⁰	abs. ampere	$\overline{10}$.523 0815
faraday second⁻¹	=	9 6500 $\times$ 10⁴	abs. ampere	4.984 5273
International ampere (U. S. before 1911)	=	0 99916	Int. ampere (v)	$\bar{1}$.999 6353
International ampere (England before 1906)	=	0 99870	Int. ampere (v)	$\bar{1}$.999 4358
International ampere (England 1906–8)	=	0 99894	Int. ampere (v)	$\bar{1}$.999 5399
International ampere (England 1909–10)	=	0 99990	Int. ampere (v)	$\bar{1}$.999 9566
International ampere (France before 1911)	=	0 9998	Int. ampere (v)	$\bar{1}$.999 9131
International ampere (Germany before 1911)	=	0 99968	Int. ampere (v)	$\bar{1}$.999 8610

52. Electrical Potential $[\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}]$

absolute volt	=	0 99958	Int. volt (v)	1.999 8176
absolute volt	=	0 99955	Int. volt (a)	1.999 8046
International volt (v)	=	1 00042	abs. volt	0.000 1824
International volt (a)	=	1 00045	abs. volt	0.000 1954
cgsm unit	=	1 0000 $\times$ 10⁻⁸	abs. volt	$\bar{8}$.000 0000
cgse unit	=	299 86	abs. volt	2.476 9185
International volt (U. S. before 1911)	=	0 99916	Int. volt (v)	$\bar{1}$.999 6353
International volt (England before 1906)	=	0 99870	Int. volt (v)	$\bar{1}$.999 4358
International volt (England 1906–8)	=	0 99894	Int. volt (v)	$\bar{1}$.999 5399
International volt (England 1909–10)	=	0 99990	Int. volt (v)	$\bar{1}$.999 9566
International volt (Germany and France, before 1911)	=	0 99968	Int. volt (v)	$\bar{1}$ 999 8610

53. Electrical Field Strength; Potential Gradient; Dielectric Strength $[\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}]$

cgsm centimeter⁻¹	=	1 0000 $\times$ 10⁻⁸	abs. volt cm⁻¹	$\bar{8}$ 000 0000
cgsm inch⁻¹	=	3 9370 $\times$ 10⁻⁹	abs. volt cm⁻¹	(U. S.) $\bar{9}$ 595 1654
cgse centimeter⁻¹	=	2 9986 $\times$ 10²	abs. volt cm⁻¹	2.476 9185
cgse inch⁻¹	=	1 1805 $\times$ 10²	abs. volt cm⁻¹	(U. S.) 2 072 0839
volt inch⁻¹	=	3 9370 $\times$ 10⁻¹	volt cm⁻¹	(U. S.) $\bar{1}$ 595 1654

54. Electrical Resistance; Surface Resistivity $[\epsilon^{-1}l^{-1}t]$; $[\mu l t^{-1}]$

1 absolute ohm	=	0 99948	Int. ohm	$\bar{1}$ 999 7741
1 International ohm	=	1 00052	abs. ohm	0.000 2259
1 cgsm unit	=	1 0000 $\times$ 10⁻⁹	abs. ohm	$\bar{9}$.000 0000
1 cgse unit	=	8 9916 $\times$ 10¹¹	abs. ohm	11.953 8370
1 International ohm (France before 1911)	=	0 9999	Int. ohm	$\bar{1}$.999 9566
1 Board of Trade unit (England 1903)	=	0 99984	Int. ohm	$\bar{1}$.999 9306
1 B. A. unit	=	0 98660	Int. ohm	$\bar{1}$ 994 1420
1 "Legal ohm" of 1884 (England)	=	0 99718	Int. ohm	1 998 7727
1 Siemens unit	=	0 94073	Int. ohm	$\bar{1}$.973 4667

55. Electrical Inductance $[\epsilon^{-1}l^{-1}t^2]$; $[\mu l]$

1 absolute henry	=	0 99948	Int. henry	1 999 7741
1 International henry	=	1.00052	abs. henry	0.000 2259
1 cgsm unit*	=	1 0000 $\times$ 10⁻⁹	abs. henry	$\bar{9}$.000 0000
1 cgse unit	=	8 9916 $\times$ 10¹¹	abs. henry	11.953 8370

* Occasionally called a centimeter

56. Electrical Capacity $[\epsilon l]$; $[\mu^{-1}l^{-1}t^2]$

1 absolute farad	=	1 00052	Int. farad	0.000 2259
1 International farad	=	0 99948	abs. farad	$\bar{1}$.999 7741
1 cgsm unit	=	1 0000 $\times$ 10⁹	abs. farad	9.000 0000
1 cgse unit*	=	1 1121 $\times$ 10⁻¹²	abs. farad	$\overline{12}$.046 1630
1 cgsm unit	=	8 9916 $\times$ 10²⁰	cgse unit	20.953 8370
1 absolute farad	=	8 9916 $\times$ 10¹¹	cgse unit	11.953 8370

* Frequently called a centimeter

57. Electrical Volume Resistivity $[\epsilon^{-1}t]$; $[\mu l^2 t^{-1}]$

1 absolute ohm-centimeter	=	0.99948	Int. ohm-cm	1.999 7741
1 International ohm-centimeter	=	1.00052	abs. ohm-cm	0.000 2259
1 cgsm unit	=	9.9948 $\times$ 10⁻¹⁰	Int. ohm-cm	$\overline{10}$.999 7741
1 cgse unit	=	8.9869 $\times$ 10¹¹	Int. ohm-cm	11.953 6111

CONVERSION FACTORS.—*Continued*

57. Electrical Volume Resistivity $[\epsilon^{-1}t]$; $[\mu l^2 t^{-1}]$.—*Continued*

1 microhm-centimeter	=	1.0000 × 10⁻⁸	ohm-cm		$\bar{6}$.000 0000
1 microhm-inch	=	2.5400	microhm-cm	(U. S.)	0.404 8346
1 ohm-inch	=	2.5400 × 10⁶	microhm-cm	(U. S.)	6.404 8346
1 ohm (meter, millimeter²)	=	100.0000	microhm-cm		2.000 0000
1 ohm (meter, millimeter)	=	78.540	microhm-cm		1.895 0899
1 ohm (mil, foot)	=	1.6624 × 10⁻¹	microhm-cm	(U. S.)	$\bar{1}$.220 7433
International Annealed Copper Standard (20°C)	=	1.7241	microhm-cm		0.236 5720

58. Volume Conductivity $[\epsilon t^{-1}]$; $[\mu^{-1} l^{-2} t]$

1 absolute *ohm⁻¹-centimeter⁻¹	=	1 00052	Int.* ohm⁻¹ cm⁻¹		0.000 2259
1 International ohm⁻¹-centimeter⁻¹	=	0 99948	abs. ohm⁻¹ cm⁻¹		$\bar{1}$.999 7741
1 cgsm unit	=	1 00052 × 10⁹	Int. ohm⁻¹ cm⁻¹		9.000 2259
1 cgse unit	=	1.11273 × 10⁻¹²	Int. ohm⁻¹ cm⁻¹		$\overline{12}$.046 3889
1 microhm⁻¹-centimeter⁻¹	=	1.0000 × 10⁶	ohm⁻¹ cm⁻¹		6.000 0000
1 microhm⁻¹-inch⁻¹	=	3 9370 × 10⁻¹	microhm⁻¹ cm⁻¹	(U. S.)	$\bar{1}$.595 1654
1 ohm⁻¹-inch⁻¹	=	3 9370 × 10⁻⁷	microhm⁻¹ cm⁻¹	(U. S.)	$\bar{7}$.595 1654
1 ohm⁻¹ (meter, millimeter²)⁻¹	=	1.000 × 10⁻²	microhm⁻¹ cm⁻¹		$\bar{2}$.000 0000
1 ohm⁻¹ (meter, millimeter)⁻¹	=	1.2732 × 10⁻²	microhm⁻¹ cm⁻¹		$\bar{2}$.104 9101
1 ohm⁻¹ (mil, foot)⁻¹	=	6 0153	microhm⁻¹ cm⁻¹	(U. S.)	0.779 2567
International Annealed Copper Standard (20°C)	=	0 5800	microhm⁻¹ cm⁻¹		$\bar{1}$.763 4280
100% conductivity (20°C)	=	0 5800	microhm⁻¹ cm⁻¹		$\bar{1}$.763 4280

* "Mho" is occasionally used instead of ohm⁻¹.

59. Electrical Mass Resistivity $[\epsilon^{-1} m l^{-1} t]$; $[\mu m l^{-1} t^{-1}]$

1 absolute ohm (meter, gram)	=	0 99948	Int. ohm (meter, gram)		1.999 7741
1 International ohm (meter, gram)	=	1.00052	abs. ohm (meter, gram)		0 000 2259
1 cgsm unit	=	9 9948 × 10⁻⁶	Int. ohm (meter, gram)		$\bar{6}$.999 7741
1 cgse unit	=	8 9869 × 10¹⁵	Int. ohm (meter, gram)		15.953 6111
1 ohm (mile, pound)	=	1 7513 × 10⁻⁴	ohm (meter, gram)	(U. S.)	$\bar{4}$.243 3663
1 ohm (centimeter, gram)	=	1.0000 × 10⁴	ohm (meter, gram)		4.000 0000
1 ohm (centimeter, gram)	=	D^*	ohm-cm		
†International Annealed Copper Standard at 20°C	=	0 15328	ohm (meter, gram)		$\bar{1}$.185 4738

* D represents the density in grams per centimeter³ † Density = 8.89 grams per centimeter³. *See* Table 61

60. Electrical Mass Conductivity $[\epsilon m^{-1} l t^{-1}]$; $[\mu^{-1} m^{-1} l t]$

1 absolute ohm⁻¹ (meter, gram)	=	1.00052	Int. ohm⁻¹ (meter, gram)		0.000 2259
1 International ohm⁻¹ (meter, gram)	=	0 99948	abs. ohm⁻¹ (meter, gram)		$\bar{1}$.999 7741
1 cgsm unit⁻¹	=	1 00052 × 10⁵	Int. ohm⁻¹ (meter, gram)		5.000 2259
1 cgse unit⁻¹	=	1 1127 × 10⁻¹⁶	Int. ohm⁻¹ (meter, gram)		$\overline{16}$ 046 3889
1 ohm⁻¹ (mile, pound)	=	5.7100 × 10⁻³	ohm⁻¹ (meter, gram)		$\bar{3}$.756 6337
1 ohm⁻¹ (centimeter, gram)	=	1.0000 × 10⁻⁴	ohm⁻¹ (meter, gram)		$\bar{4}$.000 0000
1 ohm⁻¹ (centimeter, gram)	=	*D^{-1}	(ohm-centimeter)⁻¹		

* D^{-1} = reciprocal of the density in grams per centimeter³.

61. Constants of Annealed Copper as Accepted at Various Times
Data taken from U. S. Bur. Standards Circular No. 31

Temperature °C	England (Eng. Stds. Com. 1904)	Germany (Old "Normal Kupfer" density = 8 91)	Germany (Old "Normal Kupfer" assuming density 8.89)	Lindeck, Matthiessen, assuming density 8.89	A. I. E. E. before 1907 (Matthiessen value)	A. I. E. E. 1907 to 1910	Bureau Standards and A. I. E. E. 1911	Inter. Annealed Copper Standard 1913
			Resistivity in ohms (meter, grams)					
0	0.141362	0 139590	0 139277	0.141571	**0.141729**	**0.14172₈**	0.141068	0.141332
15	0 150137	**0.148502**	**0.148164**	**0.149974**	0.150141	0.150658	0 150034	0.150290
15.6	**0.1508**							
20	0 153463	0.151470	0 151130	0.152851	0 153022	0 153634	**0.153022**	**0.15328**
25	0 156488	0 154440	0 154098	0 155765	0 155938	0 156610	0.156010	0.156262
			Temperature coefficient of resistance (mass constant)					
0	**0.00428**	0.004255	0 004255	$\frac{1}{R_t} = \frac{1}{R_o}(1 - 3\ 8701t \times 10^{-3}$	**0.0042**	0 003951	0.004277	0.004268
15	0.004022	**0.004**	**0.004**	$+ 9.009t^2 \times 10^{-6})$	0 003951	0.004019	0.004009
20	0 003943	0 003922	0 003922			0.003875	**0.00394**	**0.00393**
25	0 003866	0 003846	0.003846			0.003801	0.003864	0.003854
			Density					
	8.89 15.6°	8.91	(8 89)	(8 89)	8.89	8.89	8.89 20°	8.89 20°

CONVERSION FACTORS.—*Continued*

62. Ionic Mobility $[\epsilon^{\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}t]$; $[\mu^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}t]$

ntimeter² second⁻¹ per cgse unit of potential	=	3.3349 × 10⁻² cm² sec⁻¹ volt⁻¹ (abs.)		$\bar{3}$.523 0815
ch² second⁻¹ per cgse unit of potential	=	2.1515 × 10⁻² cm² sec⁻¹ volt⁻¹ (abs.)	(U. S.)	$\bar{2}$.332 7507
gh² second⁻¹ volt⁻¹ (absolute)	=	6 4516 cm² sec⁻¹ volt⁻¹ (abs.)	(U. S.)	0.809 6692

63. Thermoelectric Power $[\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}T^{-1}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}T^{-1}]$

gm unit of potential per °C	=	1.0000 × 10⁻² microvolt per °C (abs.)		$\bar{2}$.000 0000
gm unit of potential per °F	=	1 8000 × 10⁻² microvolt per °C (abs.)		$\bar{2}$.255 2725
se unit of potential per °C	=	2 9986 × 10⁸ microvolt per °C (abs.)		8 476 9185
se unit of potential per °F	=	5 3975 × 10⁸ microvolt per °C (abs.)		8 732 1910
icrovolt per °F	=	1 8000 microvolt per °C		0.255 2725

64. Peltier Coefficient $[\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}]$

joule per ampere-hour (absolute)	=	2.7778 × 10⁻¹ joule cm⁻¹		$\bar{3}$.443 6975
joule per ampere-hour (absolute)	=	9 2636 × 10⁻¹⁴ joule es⁻¹		$\bar{14}$.966 7790
joule per coulomb	=	10 000 joule cm⁻¹		1.000 0000
joule per faraday	=	1 0363 × 10⁻⁴ joule cm⁻¹		$\bar{4}$.015 4727
joule per electron	=	6 2811 × 10¹⁹ joule cm⁻¹		19.798 0361
calorie (15°C) per ampere-hour	=	1.1625 × 10⁻² joule cm⁻¹		$\bar{2}$.065 3930
calorie (15°C) per coulomb	=	41 850 joule cm⁻¹		1.621 6055
millivolt	=	1 0000 × 10⁻² joule cm⁻¹		$\bar{2}$.000 0000

65. Thomson Effect, Coefficient of; Specific Heat of Electricity $[\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}T^{-1}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}T^{-1}]$

joule coulomb⁻¹ per °F	=	1 8000 joule coulomb⁻¹ per °C		0.255 2725
joule es⁻¹ per °F	=	5 3975 × 10⁹ joule coulomb⁻¹ per °C		9.732 1910
joule cm⁻¹ per °F	=	0 1800 joule coulomb⁻¹ per °C		$\bar{1}$.255 2725
joule es⁻¹ per °C	=	2 9986 × 10⁹ joule coulomb⁻¹ per °C		9.476 9185
joule faraday⁻¹ per °C	=	1.0363 × 10⁻⁵ joule coulomb⁻¹ per °C		$\bar{5}$.015 4727
joule electron⁻¹ per °C	=	6 2811 × 10¹⁸ joule coulomb⁻¹ per °C		18.798 0361
volt per °C	=	1 0000 joule coulomb⁻¹ per °C		0.000 0000

66. Piezoelectric Constant $[\epsilon^{\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}t]$; $[\mu^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t^2]$

em per kilogram weight (g.)	=	3 0577 × 10⁴ es per dyne		4.485 3978
em per pound weight (g.)	=	6 7411 × 10⁴ es per dyne		4.828 7321
es per kilogram weight (g.)	=	1 0197 × 10⁻⁶ es per dyne		$\bar{6}$.008 4793
es per pound weight (g.)	=	2 2481 × 10⁻⁶ es per dyne		$\bar{6}$.351 8136
coulomb per kilogram weight (g.)	=	3 0577 × 10³ es per dyne		3.485 3978
faraday per kilogram weight (g.)	=	2 9507 × 10⁸ es per dyne		8.469 9251
electron per kilogram weight (g.)	=	4 868 × 10⁻¹⁶ es per dyne		$\bar{16}$.687 3617

67. Magnetic Field Intensity; Magnetic Potential Gradient; Magnetizing Force $[\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-2}]$; $[\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}]$

1 gauss, absolute	=	1.00010 Int. gauss (v)		0.000 0434
1 gauss, absolute	=	1 00007 Int. gauss (a)		0 000 0304
1 International gauss (v)	=	0 99990 abs. gauss		$\bar{1}$.999 9566
1 International gauss (a)	=	0 99993 abs. gauss		$\bar{1}$.999 9696
1 cgsm unit	=	1.0000 abs. gauss		0.000 0000
1 cgse unit	=	3 3349 × 10⁻¹¹ abs. gauss		$\bar{11}$.523 0815
1 gilbert per centimeter	=	1 0000 gauss		0.000 0000
1 ampere-turn per centimeter	=	1 2566 gauss		0.099 2099
1 ampere-turn per inch	=	0 49474 gauss	(U. S.)	$\bar{1}$.694 3753
1 gamma, γ	=	1 0000 × 10⁻⁵ gauss		$\bar{5}$ 000 0000

68. (Magnetic Field Intensity)⁻¹; Coefficient of Leduc Effect $[\epsilon^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t^2]$; $[\mu^{\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}t]$

1 gauss⁻¹ (absolute)	=	0 99990 Int. gauss⁻¹ (v)		1.999 9566
1 International gauss⁻¹ (v)	=	1.00010 gauss⁻¹ (abs.)		0.000 0434
1 cgsm unit⁻¹	=	1 0000 gauss⁻¹ (abs.)		0.000 0000
1 cgse unit⁻¹	=	2.9986 × 10¹⁰ gauss⁻¹ (abs.)		10.476 9185
1 centimeter per gilbert	=	1 0000 gauss⁻¹		0.000 0000
1 centimeter per ampere-turn	=	7 9577 × 10⁻¹ gauss⁻¹		$\bar{1}$.900 7901
1 inch per ampere-turn	=	2 0213 gauss⁻¹		0.305 6246

69. Magnetomotive Force; Magnetic Potential $[\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}]$; $[\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}]$

1 gilbert, absolute	=	1 00010	Int. gilbert (v)	0.000 0434
1 gilbert, absolute	=	1.00007	Int. gilbert (a)	0.000 0304
1 International gilbert (v)	=	0.99990	abs. gilbert	$\bar{1}$.999 9566
1 International gilbert (a)	=	0.99993	abs. gilbert	$\bar{1}$.999 9696
1 cgsm unit	=	1 00000	abs. gilbert	0.000 0000
1 cgse unit	=	3 3349 $\times 10^{-11}$ abs. gilbert		$\bar{11}$.523 0815
1 ampere-turn	=	1 2566	gilbert	0.099 2099

70. Magnetic Induction; Intensity of Magnetization $[\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}]$
Units of Magnetization are not named

1 maxwell per centimeter², absolute	=	0.99958	Int. maxwell per cm² (v)	$\bar{1}$.999 8176
1 maxwell per centimeter², absolute	=	0 99955	Int. maxwell per cm² (a)	$\bar{1}$.999 8046
1 International maxwell per centimeter² (v)	=	1 00042	abs. maxwell per cm²	0.000 1824
1 International maxwell per centimeter² (a)	=	1 00045	abs. maxwell per cm²	0.000 1954
1 maxwell per inch²	=	0 15500	maxwell per cm²	(U. S.) $\bar{1}$.190 3308
1 cgsm unit	=	1 00000	abs. maxwell per cm²	0.000 0000
1 cgse unit	=	2 9986 $\times 10^{10}$ abs. maxwell per cm²		10 476 9185
1 line per centimeter²	=	1 00000	maxwell per cm²	0.000 0000
1 line per inch²	=	0.15500	maxwell per cm²	(U. S.) $\bar{1}$ 190 3308

71. Flux of Magnetic Induction; Magnetic Flux; Pole Strength; Quantity of Magnetism $[\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}]$; $[\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}]$
Units of Pole Strength and Quantity of Magnetism are not named

1 maxwell, absolute	=	0 99958	Int. maxwell (v)	1 999 8176
1 maxwell, absolute	=	0 99955	Int. maxwell (a)	$\bar{1}$ 999 8046
1 International maxwell (v)	=	1.00042	abs. maxwell	0.000 1824
1 International maxwell (a)	=	1 00045	abs. maxwell	0 000 1954
1 cgsm unit	=	1 0000	abs. maxwell	0.000 0000
1 cgse unit	=	2 9986 $\times 10^{10}$ abs. maxwell		10.476 9185
1 line	=	1 0000	abs. maxwell	0 000 0000
1 volt-second	=	1 0000 $\times 10^8$	maxwell	8 000 0000

72. Magnetic Reluctance $[\epsilon l t^{-2}]$; $[\mu^{-1}l^{-1}]$

1 oersted, absolute	=	1 00052	Int. oersted	0.000 2259
1 International oersted	=	0 99948	abs. oersted	$\bar{1}$.999 7741
1 cgsm unit	=	1 0000	abs. oersted	0.000 0000
1 cgse unit	=	1 1122 $\times 10^{-21}$ abs. oersted		$\bar{21}$ 046 1630

73. Hall Effect, Coefficient of $[\epsilon^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t^3]$; $[\mu^{\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}]$

1 volt centimeter per ampere gauss (absolute)	=	1.0000 $\times 10^9$ cgsm unit	9.000 0000
1 volt inch per ampere gauss (absolute)	=	2.5400 $\times 10^9$ cgsm unit	(U. S.) 9.404 8346
1 cgse unit	=	2.6962 $\times 10^{31}$ cgsm unit	31.430 7555

74. Ettinghausen Effect, Coefficient of $[\epsilon^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t^4T]$; $[\mu m^{-\frac{1}{2}}l^{\frac{1}{2}}t^2T]$

1°C centimeter per ampere gauss (absolute)	=	10 000 °C cm per cgsm unit	1 000 0000
1°F inch per ampere gauss (absolute)	=	45.720 °C cm per cgsm unit	1.660 1071
1°C centimeter per cgse unit	=	8 9916 $\times 10^{20}$ °C cm per cgsm unit	20 953 8370

75. Nernst Effect, Coefficient of $[\epsilon^{-\frac{1}{2}}tT^{-1}]$; $[\mu l^2 t^{-1}T^{-1}]$

1 volt per gauss °C (absolute)	=	1.0000 $\times 10^8$ cgsm unit per °C	8.000 0000
1 volt per gauss °F (absolute)	=	1.8000 $\times 10^8$ cgsm unit per °C	8.255 2725
1 cgse unit per °C	=	8.9916 $\times 10^{20}$ cgsm unit per °C	20.953 8370

76. Verdet's Constant $[\epsilon^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t^2\theta]$; $[\mu^{\frac{1}{2}}m^{-\frac{1}{2}}l^{-\frac{1}{2}}t\theta]$

1 minute per gilbert	=	1.0000 minute per cgsm unit	0.000 0000
1 minute per ampere-turn	=	1.2566 minute per cgsm unit	0.099 2099
1 radian per gilbert	=	3 4377 $\times 10^3$ minute per cgsm unit	3.536 2739

77. Fundamental Electric and Magnetic Units

Name of quantity	Cgse units	1 *Cgsm unit equals Practical units (abs.)	Cgse system	Cgsm system	‡Practical system
			Dimensions		
Electric:					
Capacity	c^2	10^9 farad	d	$\mu^{-1}l^{-1}t^2$	$lE^{-1}t$
Charge, quantity	c	10 coulomb	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}$	It

CONVERSION FACTORS.—*Continued*

77. Fundamental Electric and Magnetic Units.—*(Continued)*

Conductivity (mass)	c^2	10^9 ohm^{-1} (cm, g)	$\epsilon m^{-1}l^{3}t^{-1}$	$\mu^{-1}m^{-1}lt$	$R^{-1}m^{-1}l^2$
Conductivity (surface)	c^2	10^9 ohm^{-1}	$\epsilon l t^{-1}$	$\mu^{-1}l^{-1}lt$	R^{-1}
Conductivity (volume)	c^2	10^9 ohm^{-1} cm^{-1}	ϵt^{-1}	$\mu^{-1}l^{-2}t$	$R^{-1}l^{-1}$
Current	c	10 ampere	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-2}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}$	I
Dielectric constant	c^2	†10^9 ohm^{-1} per (cm sec^{-1})	ϵ	$\mu^{-1}l^{-2}t^2$	†$IE^{-1}l^{-1}t$
Displacement (local)	c	10 coulomb per cm^2	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{3}{2}}$	$Il^{-2}t$
Displacement (integral)	c	10 coulomb	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}$	It
Electromotive force	c^{-1}	10^{-8} volt	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-2}$	E
Field strength	c^{-1}	10^{-8} volt cm^{-1}	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}$	El^{-1}
Inductance	c^{-2}	10^{-9} henry	$\epsilon^{-1}l^{-1}t^{2}$	μl	Rt
Inductivity	c^2	† 10^9 ohm^{-1} per (cm sec^{-1})	ϵ	$\mu^{-1}l^{-2}t^2$	†$IE^{-1}l^{-1}t$
Ionic mobility	c	10^5 cm sec^{-1} per (volt cm^{-1})	$\epsilon^{\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}$	$\mu^{-\frac{1}{2}}m^{-\frac{1}{2}}l^{\frac{1}{2}}t$	$E^{-1}l^{2}t^{-1}$
Polarization capacity	c^2	10^9 farad cm^{-2}	ϵl^{-1}	$\mu^{-1}l^{-3}t^2$	$IE^{-1}l^{-2}t$
Potential	c^{-1}	10^{-8} volt	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-2}$	E
Resistance	c^{-2}	10^{-9} ohm	$\epsilon^{-1}l^{-1}t$	$\mu l t^{-1}$	R
Resistivity (mass)	c^{-2}	10^{-9} ohm (cm, g)	$\epsilon^{-1}ml^{-3}t$	$\mu m l^{-1}t^{-1}$	Rml^{-2}
Resistivity (surface)	c^{-2}	10^{-9} ohm	$\epsilon^{-1}l^{-1}t$	$\mu l t^{-1}$	R
Resistivity (volume)	c^{-2}	10^{-9} ohm-cm	$\epsilon^{-1}t$	$\mu l^2 t^{-1}$	Rl
Specific heat of electricity (Thomson)	c^{-1}	10^{-8} volt deg^{-1}	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}T^{-1}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-2}T^{-1}$	ET^{-1}
Specific inductive capacity	1	1	zero	zero	zero
Magnetic:					
Field intensity	c	1 gauss	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	Il^{-1}
Flux of induction (integral)	c^{-1}	1 maxwell	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	Et
Induction (local)	c^{-1}	1 maxwell cm^{-2}	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	$El^{-2}t$
Intensity of magnetization (volume)	c^{-1}	1	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	$El^{-2}t$
Magnetic flux (integral)	c^{-1}	1 maxwell	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	Et
Magnetizing force	c	1 gauss	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{-\frac{1}{2}}t^{-1}$	Il^{-1}
Magnetomotive force	c	1 gilbert	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}$	I
Permeability	c^{-2}	1 maxwell cm^{-2} per gauss	$\epsilon^{-1}l^{-2}t^2$	μ	$I^{-1}El^{-1}t$
Pole strength	c^{-1}	1	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	Et
Potential	c	1 gilbert	$\epsilon^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-2}$	$\mu^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{1}{2}}t^{-1}$	I
Quantity	c^{-1}	1	$\epsilon^{-\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}$	$\mu^{\frac{1}{2}}m^{\frac{1}{2}}l^{\frac{3}{2}}t^{-1}$	Et
Reluctance	c^2	1 oersted	$\epsilon l t^{-2}$	$\mu^{-1}l^{-1}$	$IE^{-1}l^{-1}$
Susceptibility	c^{-2}	$\frac{1}{4\pi}$ maxwell cm^{-2} per gauss	$\epsilon^{-1}l^{-2}t^2$	μ	$I^{-1}El^{-1}t$

* For the purposes of International Critical Tables, c has been taken as 2.99086×10^{10} cm per sec, $\log_{10} c = 10.476\ 9185$, $\log_{10} c^{-1} = 11.523\ 0815$. This is the accepted value for the velocity of light in vacuo. The best directly determined value of the ratio of the two electrical units of quantity gives $c = 1.9979 \times 10^{10}$ cm per sec. (Rosa and Dorsey, *Bull. U. S. Bur. Standards*, **3**: 433; 07.)

† In practice this unit is not used; the quantity given in essentially every instance is the dimensionless "specific inductive capacity," which is numerically equal to the dielectric constant expressed in cgse units.

‡ In this column are given the dimensions in terms of the practical electrical units, as these generally enter into the actual determinations of the several quantities. As three basic electrical units are employed, alternative expressions are possible. T = thermometric degree, R = potential, I = current, E = resistance.

78. Indicated Conversion Factors

a = area, C = electrical capacity, T = thermometric degree, l = density, E = electrical potential, e = electric charge, F = electrical field intensity, h = heat, m = mass, Q = quantity of magnetism, R = electrical resistance, t = time, v = volume, ϵ = dielectric constant, η = viscosity, θ = plane angle.

Name of quantity	Dimensions	Tables
Electricity		
Electric displacement	ϵF	14, 53
Polarization capacity	Ca^{-1}	56, 17
Pyroelectric constant	$ea^{-1}T^{-1}$	19, 17, 12
Specific inductive capacity	zero	
Surface density of charge	ea^{-1}	49, 17
Thermoelectric power	ET^{-1}	52, 12
Volume density of charge	ev^{-1}	49, 19
Heat, capacity	$hm^{-1}T^{-1}$	35, 21
Latent	hm^{-1}	35, 4
Reaction	hm^{-1}	35, 4
Superficial latent	ha^{-1}	35, 17
Transformation	hm^{-1}	35, 4

Name of quantity	Dimensions	Tables
Radiation, index of absorption	zero	
Intensity of	$ha^{-1}t^{-1}$	35, 22
Kerr's constant (magneto-optic)	$\theta Q^{-1}a$	7, 71, 16
Reflectivity	zero	
Refraction, index of	zero	
Solubility, gases in liquids	zero	
Viscosity, kinematic	ηd^{-1}	39, 28

79. Hydrometer Scales

Unless the hydrometer is used in the liquid and at the temperature for which it is graduated, corrections must be applied for the changed capillary depression and for the expansion (or contraction) of the instrument. (The following table does not include all scales which have been used.)

T = temperature at which the instrument is to be used; r = reading of instrument; the specific gravity is with reference to water at temperature T unless another temperature is indicated in the last column.

79. Hydrometer Scales.—*Continued*

Hydrometer	T	Specific gravity Dense	Specific gravity Light	Remarks
A. P. I. = American Petroleum Institute.	60°F = 15.56°C		$\dfrac{141.5}{131.5 + r}$	Petroleum
Balling	17.5°C	$\dfrac{200}{200 - r}$	$\dfrac{200}{200 + r}$	
Bates	60°F = 15.56°C	$\dfrac{1000 + 2.78r}{1000}$		
Baumé	10°R = 12.5°C	$\dfrac{145.88}{145.88 - r}$	$\dfrac{145.88}{135.88 + r}$	
Baumé	15°C	$\dfrac{146.3}{146.3 - r}$	$\dfrac{146.3}{136.3 + r}$	
Baumé	17.5°C	$\dfrac{146.78}{146.78 - r}$	$\dfrac{146.78}{136.78 + r}$	
Baumé	15°C	$\dfrac{144.3}{144.3 - r}$		"Rational"
Baumé	15°C	$\dfrac{144.3}{144.3 - r}$		"Rational" (water at 4°C)
Baumé-Lunge	12.5°C	$\dfrac{144.32}{144.32 - r}$	$\dfrac{144.32}{144.32 + r}$	"Rational"
Baumé	15°C	$\dfrac{144.32}{144.32 - r}$	$\dfrac{144.32}{144.32 + r}$	French (water at 4°C)
Baumé	60°F = 15.56°C	$\dfrac{145}{115 - r}$	$\dfrac{140}{130 + r}$	American
Beck	12.5°C	$\dfrac{170}{170 - r}$	$\dfrac{170}{170 + r}$	
Brix	12.5°R = 15.625°C	$\dfrac{400}{400 - r}$	$\dfrac{400}{400 + r}$	
Cartier	12.5°C	$\dfrac{136.8}{126.1 - r}$	$\dfrac{136.8}{126.1 + r}$	
Fischer	12.5°R = 15.625°C	$\dfrac{400}{100 - r}$	$\dfrac{400}{400 + r}$	
Fleischer		$\dfrac{1000 + 10r}{1000}$		
Gay-Lussac		$\dfrac{100}{100 - r}$	$\dfrac{100}{100 + r}$	
Gerlach, or "new"	17.5°C	$\dfrac{146.78}{146.78 - r}$		
Holland, or "old"	12.5°C	$\dfrac{141}{144 - r}$		
Stoppani	12.5°R = 15.625°C	$\dfrac{166}{166 - r}$		
Twaddell	60°F = 15.56°C	$\dfrac{1000 + 5r}{1000}$		British (water at 4°C)

TECHNICAL EFFLUX VISCOMETERS: INTERPRETATION AND INTERCONVERSION OF READINGS

WINSLOW H. HERSCHEL

Since changes are made from time to time in the standardization or method of operation of these instruments, and many old instruments are still in use, it is believed that in general the determination of kinematic viscosity from the readings of the instruments, and direct interconversions between instruments, when used at the same temperature, may be made by the use of Fig. 1, with as great precision (about 5%) as the data will warrant. It is assumed that the instruments are used in the normal manner. For the Saybolt instruments, a higher precision is occasionally justified, and may be obtained by the use of Table 2.

If the instruments are used at different temperatures, appropriate temperature corrections must be applied. For lubricating oils, the viscosity at one temperature may be estimated from that at another by the approximate empirical rule, applicable between 100° and 212°F (37.8° and 100°C), that the logarithmic viscosity-temperature graphs are straight and meet at a point, temperatures being expressed in degrees Fahrenheit. (For other temperatures see ([1, 7, 8])). The location of the point of intersection for several classes of oils is given in Table 1.

TABLE 1.—COORDINATES OF POINTS OF INTERSECTION OF LOGARITHMIC GRAPHS([5])

η_0 = viscosity in poises; t_0 = temperature in °F

Class of oils	$\log_{10} \eta_0$	η_0	$\log_{10} t_0$	t_0
Paraffin base	$\bar{3}.58$	0.0038	2.77	589
Naphthene base	$\bar{3}.88$	.0076	2.57	371
Mixed base	$\bar{3}.43$	.0027	2.78	605
Fatty oils	$\bar{3}.75$	.0056	2.82	661

In estimating the viscometer reading at a given temperature for a certain type of instrument, from an observed reading at another temperature with another type of instrument, the following steps may be taken.

1. Determine the kinematic viscosity corresponding to the observed reading by means of Fig. 1.

2. Multiply by the density (g/cm³) so as to obtain the absolute viscosity (η) in poises; find the logarithm of the absolute viscosity and the logarithm of the temperature (t) of test (°F).

3. Plot the observed η, t and the η_0, t_0 of the point of intersection, as given in Table 1, on logarithmic paper. Or plot the corresponding logarithms on equispaced coordinate paper. In either case, these two points locate a straight graph upon which the viscosity at the desired temperature will be found.

4. Divide the absolute viscosity at the desired temperature by the density at that temperature to get the kinematic viscosity. From this, determine, by means of Fig. 1, the corresponding time of flow on the desired viscometer.

It will be noted that the density under (2) and (4) must be the density at the temperature under consideration, and not the density at 60°F (15.6°C), which is generally the standard for such density determinations.

If an instrument is used in an irregular manner, appropriate corrections must be applied ([2, 3, 6, 9]).

TABLE 2.—SAYBOLT UNIVERSAL AND SAYBOLT FUROL VISCOMETERS
Units: Time (t), sec; kinematic viscosity = (η/d), poise/(g per cm³).

Saybolt Universal		Saybolt Furol	
t	η/d	t	η/d
32	0.0115	25	0.486
40	0.0417	26	0.512
50	0.0740	27	0.537
60	0.103	28	0.562
70	0.130	29	0.586
80	0.156	30	0.610
90	0.181	35	0.730
100	0.206	40	0.846
125	0.266	45	0.960
150	0.324	50	1.072
175	0.381	60	1.292
200	0.437	70	1.507
225	0.492	80	1.724
250	0.548	90	1.939
275	0.603	100	2.155
300	0.658		

For higher viscosities the kinematic viscosity is equal to $0.00220t$ for the Saybolt Universal, or to $0.0216t$ for the Saybolt Furol.

LITERATURE
(For a key to the periodicals see end of volume)

([1]) Fortsch and Wilson, *45*, **16**: 789, 24. ([2]) Gans, *252*, **6**: 218; 99. ([3]) Herschel, *52*, No. **100**; 17. ([4]) Herschel, *244*, **10**: 31; 22. ([5]) Herschel, *45*, **14**: 715; 22. ([6]) Holde, Examination of hydrocarbon oils, 1917. ([7]) Lane and Dean, *45*, **16**: 905; 24. ([8]) MacCoull, *255*, **7**: No. 6; 21. ([9]) Ubbelohde, Tabellen zum Engler' schen Viskosimeter, 1907.

Poises/gram per cm³

Kinematic Viscosity

B   Barbey (Fluidity)
E° Engler (Degrees)
ES Engler (Seconds)
RA Redwood Admiralty
RS Redwood Standard
SF Saybolt Furol
SU Saybolt Universal

For Curves 1,2,4,5 and 6, T=
time in seconds.

For Curve 3, T= 10×Engler degrees
51×Engler degrees = time in seconds

For Curve 7, Time = Barbey fluidity
(= rate of flow, cm³/hr).

FIG. 1.—Conversion diagram for viscosimeters at a common temperature (4).

SELECTED TECHNICAL TERMS

N. Ernest Dorsey

In this section are given the definitions of numerous units, and very brief explanations of such technical terms as occur in many sections of the I. C. T. or are for other reasons more suitably considered here than elsewhere. Other terms will be explained where they occur in the body of the work. Symbolical explanations will be given wherever they appear to be satisfactory. In many cases, dimensional formulae (see p. 18) are given; these are enclosed in [ ]. Symbols are enclosed in ( ). The sequence will be: Name, symbol or symbols, dimensional formula, definition or explanation; but the symbol or formula, or both may be omitted. For the explanation of the symbols employed in the formulae and explanations, see p. 16.

Aberration, Constant of. $[\theta]$. $\tan (V\text{-}v)/c$. V, v = maximum and minimum velocity of earth in its orbit, c = velocity of light in vacuo.

Absolute. (abs.). 1. An adjective, descriptive of a system of units which is based upon the smallest possible number of independent units. In this connection, every specification of a definite substance or of a vacuum is to be regarded as the introduction of an independent unit. 2. **Absolute zero.** The temperature at which the pressure of a fixed mass of an ideal gas, maintained at a constant volume, becomes zero. 3. **Absolute temperature.** The temperature reckoned from the absolute zero.

Absorption. When the absorption of radiation by a substance is such that $J = J_o e^{-kl}$, J, J_o = intensity, l = length of path, k is the **coefficient of absorption.** k/d = coefficient of **mass absorption.** Writing $k = (4\pi k'n)/\lambda$, n = index of refraction, λ = wave length in vacuo, k' = **index of absorption.** (Some call $k'n$ the index.)

Absorptivity. Ratio of radiant energy absorbed to that absorbed, under same conditions, by a black body.

Action, Planck's constant of. See Planck.

Ampere. Unit of electric current. **Abs. ampere** = 0.1 cgs unit. **Int. ampere** is that unvarying electric current which, when passed through a solution of silver nitrate in water, in accordance with certain specifications, deposits silver at the rate of 0.00111800 gram per second.

Ampere-turn. Unit of mmf. Difference in magnetic potential between the faces of a coil of one turn carrying one ampere.

Ångstrom unit. (Å). [l]. 10^{-10} meters. **International Ångstrom** defined as such a length that wave-length of red cadmium line in air at 15°C, A_n, is exactly 6438.4696 Int. Å; it = 10^{-10} m within experimental error.

Anomalistic. -Anom. year [month] = time between successive passages of earth [moon] through perihelion [perigee].

Aphelion. Point of planet's orbit farthest from sun.

Apogee. Point of moon's orbit farthest from earth.

Aries, First point of. Designation of position of vernal equinox (see Celestial sphere); not at present in constellation Aries.

Assay ton. [m]. $29\frac{1}{6}$ grams; as many mg as there are troy ounces in short ton.

Astronomical unit of length. Mean distance (q v.) earth to sun; 149.50 × 10⁶ km.

Astronomical unit of mass. Mass of sun.

Astronomical unit of time. Mean solar day.

Atmosphere. [force area⁻¹], $[m/lt^2]$. 1. **Normal atmosphere** (A_n) defined as pressure exerted by vertical column of liquid 76 cm long, density 13.5951 grams per cm³, acceleration of gravity being 980.665 cm sec⁻². 2. **Atmosphere at 45°** (A_{45}) differs from A_n only in use of acceleration of gravity at sea level

and lat. 45° instead of 980.655 cm sec⁻². 3. **British atmosphere** is based on 30 inches instead of 76 cm.

Avogadro's number. -(N_0). $[m^{-1}]$. Number of molecules in a mole.

Bar. —[force/area], $[m/lt^2]$. Internationally accepted unit of pressure; = 10⁶ dyne/cm². Has also been used to denote one dyne/cm² (cf. Barye).

Barye.—[force/area], $[m/lt^2]$. The cgs unit of pressure, one dyne/cm². (In accordance with recommendation of special committee of International Congress of Physicists, Paris, 1900, and with the usage of the International Bureau of Weights and Measures.) (cf. Bar).

B. A. unit.—A unit of electrical resistance based on certain coils prepared in 1863-1864 by British Association for Advancement of Science.

Black Body.—One which absorbs all radiant energy incident upon it. Its radiance of wave-length λ is J_λ $d\lambda$; the **intensity,** J_λ = $C_1\lambda^{-5}[e^{C_2/\lambda T} - 1]^{-1}$, T = absolute temperature, C_1, C_2 are **radiation constants. Total radiance** (J) is $\int J_\lambda$ $d\lambda$ taken over all wave-lengths. $J = \sigma T^4$, σ = **Stefan,** or **Stefan-Boltzmann** constant of total radiation. For each T there is a wave-length (λ_m) for which J_λ $(=J_m)$ is a maximum; $J_m = C_j T^5$, C_j = **intensity coefficient;** $\lambda_m = w/T$, w = **Wien's displacement constant.**

Board of Trade unit. 1. A unit of electrical resistance based upon certain coils preserved by British Board of Trade. 2. (B.T.u.). Unit of **work.** Generally used in England as equivalent of one kilowatt-hour. (To be distinguished from British thermal unit (BTU).)

Boltzmann's molecular gas constant.—(k_o). $[ml^2/t^2T]$. Gas constant (q.r.) per molecule.

Bougie decimale.—$[\psi\omega^{-1}]$. An old unit of luminous intensity, 0.05 Violle unit.

Brightness.—$[\psi/l^2\omega]$. Luminous intensity per unit of apparent area of the luminous surface; if emission follows Lambert's law, brightness is independent of direction of line of sight, otherwise it is not; in latter case, line of sight is assumed to be normal to the surface unless the contrary is stated.

British Thermal Unit. (BTU). [energy], $[ml^2/t^2]$. Heat per pound, per °F of rise, required to produce small rise in temperature of water under pressure A_n; varies with temperature, which must be stated. **"Mean" BTU** = $\frac{1}{180}$ of heat required to raise one lb. of water from 32°F to 212°F, pressure A_n. (To be distinguished from Board of Trade unit (B.T.u.).)

Bulk modulus. —[stress], $[m/lt^2]$. Hydrostatic pressure divided by resulting decrease in volume per unit volume. Also called **volume elasticity, cubical elasticity, resistance to compression, modulus of compression** (cf. compressibility).

Calorie.—[Heat], $[ml^2/t^2]$. 1. Heat per unit of mass, per °C of rise, required to produce small rise in temperature of water under pressure A_n; varies with temperature, which must be stated. If unit of mass is gram, it is called small calorie, gram calorie, or calorie; symbol is cal. If unit of mass is kilogram, it is called large calorie, kilogram calorie, or Calorie; symbol, Cal. (2) **Mean calorie** = $\frac{1}{100}$ of heat required to raise unit mass of water from 0°C to 100°C, pressure A_n.

Candle. —(ca). $[\psi\omega^{-1}]$. Basic photometric unit of luminous-intensity. A value determined by international agreement, and maintained at certain national laboratories by means of incandescent electric lamps is known as the "International candle."

Candle per square centimeter.—$[\psi/l^2\omega]$. Brightness of surface which, in direction considered, has a luminous intensity of one

candle per cm² of apparent area; τ lamberts. Similarly: Candle per sq. in., etc.

Candlepower.—(c.p.). Luminous intensity in terms of candles.

Capacity, heat.—1. Of a substance, is heat per unit of mass, per degree of rise, required to produce a very small rise in temperature, also called **specific heat**, and **thermal capacity.** 2. Of a body, is heat, per degree of rise, required to heat the body.

Capacity, electrical.—Of body A with reference to body B is $Q/(V_A - V_B)$, all other bodies in the field being insulated and uncharged; Q = charge on A; V_A, V_B = potential of A, B.

Capacity, polarization.—Of one electrode with reference to another is its electrical capacity per unit of area.

Capillary constant. -(a). [l]. 1. British usage: $a_1^2 = \gamma/(d_1 - d_2)g$; γ = surface tension, g = acceleration of gravity, $(d_1 - d_2)$ = positive difference in the densities of the fluids separated by the surface. 2. German usage: $a_2^2 = 2\gamma/(d_1 - d_2)g$. (The subscripts to the a are usually omitted.)

Carat fine. -See Karat.

Carcel unit.—A superseded unit of luminous intensity; approximately = 9.6 Int. candles.

Celestial sphere.—Sphere, concentric with earth, serving to locate angular positions of celestial bodies; its intersection with plane of earth's orbit [equator] is called **ecliptic [celestial equator];** intersections of ecliptic and equator are called **equinoxes;** motion of equinoxes with reference to stars is called **precession of equinoxes,** it is resultant of an oscillatory and a nearly uniform motion, a fictitious equinox possessing only the latter motion is called **mean equinox.** The mean equinox through which sun passes in spring of northern terrestrial hemisphere is called **mean vernal equinox,** and is point from which **celestial longitude** (along the ecliptic) and **mean right ascension** (R. A.) (along the equator) are measured—positive to the east. Intersections of the sphere and the axis of rotation of earth are called **celestial poles;** that of the sphere and its diameter perpendicular to plane of ecliptic called **poles of the ecliptic.** Declinations are measured from equator along great circles passing through the poles—positive towards north; **celestial latitudes,** from ecliptic along great circles passing through poles of ecliptic—positive towards north. The pole of the sphere has a motion compounded of a nearly uniform progressive motion and a rotation about a point having the former motion; that point is called **mean pole,** its motion is the **precession of the pole,** the rotation of the true pole about the mean pole is called the **nutation** of the pole; mean (angular) distance between mean pole and true pole is called **constant of nutation.**

Centi-.—Prefix denoting $\frac{1}{100}$.

Centigrade.—(C). Thermometric system in which freezing point of water is called 0° and its boiling point is called 100°; pressure = A_n.

Centigrade thermal unit. -(CTU). [energy]. $[ml^2/t^2]$. Differs from British Thermal Unit only in the substitution of Centigrade for Fahrenheit scale.

Centimeter.—(cm). 1. The cgs unit of length, 0.01 meter. 2. Often used to denote cgse unit of electrical capacity. 3. Occasionally used to denote cgsm unit of electrical inductance.

Centimeter-dyne.—[work], $[ml^2/t^2]$. One erg.

Centimeter of water [of mercury, etc.] at $t°$. -[force/area], $[m/lt^2]$ Denotes pressure exerted by a vertical column of water [of mercury, etc.] one cm long, temperature $t°$, at a place where acceleration of gravity is g, (= 980.665 cm/sec²).

Cheval-vapeur.—[work/time], $[ml^2/t^3]$. 1. Primary definition, 75 meter-kilograms per second. Also called **force de cheval, continental horsepower, Pferdekraft.** 2. For electrical purposes, generally regarded as exactly 736 watts; may be called **continental electrical horsepower.**

Circular inch.—(cir. in.). $[l^2]$. Area of a circle one inch in diameter. Similarly for **circular mil** (cir. mil), **circular millimeter** (cir. mm), etc.

Compressibility.—$[lt^2/m]$. Reciprocal of bulk modulus.

Compression, modulus of. -$[m/lt^2]$. See Bulk modulus.

Concentration.—1. The amount per unit of volume; may be called **volume concentration.** If amount is measured by mass, the symbol is C. 2. The mass of the material per unit of mass of the mixture containing it; may be called **mass concentration.** If both masses are expressed in terms of the same unit, this concentration is generally called the **titer** of the mixture.

Conductance.—Reciprocal of resistance.

Conductance, Specific. -See Conductivity, electrical.

Conductivity, Electrical.—Reciprocal of electrical resistivity (q.v.). 1. (κ) **Volume conductivity** = reciprocal of volume resistivity; specific conductance. 2. **Mass conductivity** = κ/d; d = density. 3. **Equivalent conductivity** (Λ) is κ/c; c = equivalents of solute per unit volume of solution. 4. **Molecular conductivity** (μ) is κ/m; m = moles of solute per unit volume of solution.

Conductivity, Thermal. —$[(\text{heat}/\text{area-time})/(T/l)]$; $[ml/Tt^3]$.

$dQ/dt = -k\,lxdy\dfrac{d\theta}{dz}$; k = thermal conductivity, dQ = amount of heat through $dxdy$, in direction dz, in time dt, $d\theta$ = increase in temperature in distance dz

Coulomb.—The quantity of electricity transferred in one second by a current of one ampere.

Critical. 1. Any point, line, or region serving to locate a well marked **transition** may be described as **critical.** 2. As regards **condensation** of vapors, the temperature corresponding to the isotherm above which liquefaction is impossible is called the **critical temperature;** the vapor pressure at which the two phases are in equilibrium at the critical temperature is the **critical pressure;** volume of unit mass at the critical temperature and temperature is the **critical volume.** These three values are called the **critical constants.**

Cubic. (cu.), (³). Used in conjunction with name of unit of length to form name of a related unit of volume; e.g., **cubic meter** (cu. m) (m³) is name of a unit of volume equivalent to volume of a cube with edges one meter long.

Cubic centimeter atmosphere. - See Liter-atmosphere.

Curie. Internationally defined as amount of radon (radium emanation) which can exist in equilibrium with one gram of radium.

Current. —(I). The current of x through a surface S is $I = dx/dt$, where dx is the amount of x which passes through S in time dt. The density of the current through S at a given point is σ. = dI/dS, where dI is the current at that point through an element of S of area dS. The value of σ varies with the orientation of dS, and for a certain orientation it is a maximum. The normal, in the direction of the flux, to the element so oriented is the **direction of the current;** and this maximum value of σ is called the **density,** or the **intensity,** of the current at that point.

Dalton. -$[m]$. A unit of mass, $\frac{1}{16}$ mass of atom of oxygen. Approximately 1.650×10^{-24} grams.

Day.—(da). $[t]$. 1. **Solar day** = interval between successive transits of sun across same meridian. It is not of uniform length. 2. **Mean solar day** = average length of all the solar days in a tropical year. This is the basis of all our time measurements and is what is meant by day unless the contrary is definitely indicated. 3. **Sidereal day** = interval between successive transits of true vernal equinox. 4. The day defined by successive transits of same fixed star is not used in astronomical computations, and appears to have no name.

Deci-. -Prefix denoting $\frac{1}{10}$.

Declination.—1. Of celestial objects. See Celestial sphere. 2. **Magnetic declination** = angular deviation of horizontal com-

ponent of earth's magnetic field from northerly measured geographic meridian; easterly deviations, positive.

Degree.—1. (°), (deg). Unit of difference in **temperature**; size depends upon thermometric scale employed. 2. (°). Unit of angle, $\frac{1}{360}$ of complete circumference. 3. (°). Hydrometer **degree** is an arbitrary unit of difference in specific gravity; its value depends upon type of hydrometer (*see* p. 31).

Deka-.—Prefix denoting 10.

Demal.—A concentration of one g-equivalent per dm³.

Density.—1. Volume density $= dQ/dv$, $dQ =$ amount of the physical quantity considered which is contained in the element of volume dv. 2. **Density of a substance**, (d), (D), is dm/dv, $m =$ mass. When, on a particular scale of operation, the density varies from point to point, it may be that on a larger scale it will not; then the density on the larger scale may properly be called the **apparent density** (sometimes called **bulk density**) when operations on the smaller scale are being considered. 3. **Surface density** $= dQ/ds$, $ds =$ element of area of surface over which dQ is distributed.

Dielectric constant.—(ϵ). $[t^2/\mu l^2]$, $[\epsilon]$. The force (f) of repulsion between two point charges (e, e') of electricity at a distance (r) apart in a uniform medium of great extent is $f = ee'/\epsilon r^2$; ϵ depends upon the nature of the medium, and is called its dielectric constant.

Diffusion, Coefficient of.—*See* Diffusivity.

Diffusivity.—1. (Δ). $\left[\dfrac{\text{quantity}}{\text{area time}} \quad \dfrac{\text{vol. concen.}}{\text{distance}}\right]$, $[l^2/t]$. dQ/dt

$= -\Delta(dc/dx)dydz$. $dQ =$ amount of Q passing through area $dydz$ in direction of x in time dt, $dc/dx =$ rate of increase, in direction of x, of volume concentration of Q. Also called **coefficient of diffusion.** 2. **Heat diffusivity.** $\left[\dfrac{\text{heat}}{\text{area} \times \text{time}}\right.$

$+ \dfrac{\text{specific heat} \times \text{density} \times \text{temp.}}{\text{distance}}\right]$, $\left[\dfrac{\text{heat conductivity}}{\text{density} \times \text{specific heat}}\right]$,

$[l^2/t]$. $dQ/dt = -\Delta_t cd(dT/dx)dydz$, $\Delta_t =$ **heat diffusivity**, $c =$ specific heat, $d =$ density, $T =$ temperature. $\Delta_t cd =$ **thermal conductivity.** Δ_t also called **temperature conductivity.**

Displacement constant, Wien's.—*See* Black body.

Displacement, Electric.—*See* Induction, electrostatic.

Draconic month.—*See* Nodical month.

Dyne.—$[ml/t^2]$. The cgs unit of force. The force which, when acting continuously upon a mass of one gram and not opposed by another, will impart to the mass a uniform acceleration of one cm per sec.²

Dyne-centimeter.—[force length], $[ml^2/t^2]$. The torque of one dyne acting on a lever-arm of one cm.

Ecliptic.—*See* Celestial sphere.

Elastic modulus.—Ratio of stress to resulting elastic strain. There are as many types of moduli as there are types of strain. 2. Occasionally used to denote **Young's modulus.**

Elasticity.—1. Cubical; *see* Bulk modulus. 2. **Longitudinal**; *see* Young's modulus. 3. **Shear**; *see* Rigidity. 4. **Torsional**; *see* Rigidity. 5. **Modulus of**; *see* Elastic modulus.

Electric displacement, field strength, etc.—*See* corresponding nouns.

Electromagnetic unit of quantity of electricity.—*See* Quantity of electricity.

Electromotive force.—(E), (emf). *See* Potential.

Electron.—Negative electrons are very small negatively charged particles observed under many, very diverse conditions. All appear to be alike in every way, including amount of charge carried. They appear to be one of the basic elements of which atoms are made.

Electronic charge.—(e). A quantity of electricity, of either sign, which is numerically equal to the electric charge carried by an electron.

Electronic mass.—(m_e). The mass of a negative electron when moving with a velocity much less than that of light.

Electronic ratio.—(e/m_e). Ratio of electronic charge to electronic mass.

Electrostatic unit of quantity of electricity.—*See* Quantity of electricity.

Elongation.—Distance of an oscillating, or of a revolving, body from a point of reference; *e.g.*, the distance of an electron from the nucleus about which it revolves.

Emissivity.—Ratio of radiance of the body to that of a black body at same temperature. If radiation of only one wave-length is considered, it is **monochromatic emissivity**; if all wave-lengths, it is **total emissivity.** The ratio of the radiances (or of the emissivities) of two non-black bodies is called **relative emissivity** of first with respect to second.

English sperm candle.—*See* Sperm candle.

Equation of time.—*See* Time.

Equator.—1. The intersection of surface of the **earth**, or other rotating spheroid, with the plane through its center perpendicular to its axis of rotation. 2. The intersection of the surface of a spheroid with a plane through its center and perpendicular to any diameter chosen as axis. 3. **Celestial equator.** *See* Celestial sphere.

Equinox.—*See* Celestial sphere.

Equivalent.—(equiv). Electrochemical equivalent (briefly equivalent) of an ion—actual or potential—is its formula weight divided by its valence.

Erg.—[force · distance], $[ml^2/t^2]$. Work done by a force of one dyne while acting through a distance of one centimeter in its own direction.

Erg-second.—[work · time], $[ml^2/t]$. The action produced by one dyne acting through one cm in one sec.

Expansion, coefficient of.—*See* Expansivity.

Expansivity.—$[T^{-1}]$. 1. **Volume expansivity** $= dv/(vdT)$. 2. **Linear expansivity** $= dl/(ldT)$. $v, l, T =$ volume, length, temperature; $dv[dl]$ is change in $v[l]$ produced by change dT in temperature.

Fahrenheit.—(F). A thermometric system in which 32° denotes the freezing, and 212°, the boiling point of water under pressure of A_n.

Farad.—Capacity of electrical condenser which is charged to a potential difference of one volt by one coulomb.

Faraday.—(F). A subsidiary unit, the electrical charge carried in electrolysis by one gram-equivalent.

Field.—The field of a physical quantity is the region of space within which phenomena characteristic of the quantity exist. The strength, or intensity, of the field at any point is measured by the magnitude at that point of some chosen, characteristic phenomenon, and the complete designation of the field includes an indication of this phenomenon; *e.g.*, electrical field of force. As force is the phenomenon most frequently chosen, and in other cases the context indicates what is intended, the explicit designation of the chosen phenomenon is quite frequently omitted.

Field intensity.—The strength, or intensity, of a field of force at any point is df/dm, where df is the mechanical force experienced by dm, a vanishingly small amount of m placed at that point. For an **electrical field**, m is positive electricity; for a **magnetic field** it is a north magnetic pole; for a **gravitational field** it is mass. Magnetic field strength is frequently called **magnetizing force.**

Fluidity.—(φ). Reciprocal of viscosity. Also called **coefficient of fluidity.**

Flux.—1. Flux (ψ) of **vector** (V) through surface S is $\psi = \int_S V_n dS$; $V_n =$ component of V normal to dS, integral is to be taken over S. 2. Flux of a **quantity** Q through surface is $\psi = dQ/dt$,

dQ = amount of Q which passes through S in time dt. 3. From point source. If $V = I/r^2$, where r = distance from source and I is a constant independent of direction, I is called intensity of the source, and $\psi = I\omega$; ω = solid angle subtended, at the source, by S (cf. Intensity, luminous).

Flux, Luminous.—(ψ). Flux of radiant energy expressed in terms of its power to produce luminous sensation in the human eye.

Flux, Magnetic.—Flux of magnetic induction.

Foot-candle.—[ψ/l^2]. Unit of illumination, one lumen per square foot.

Foot-lambert.—[$\psi/l^2\omega$]. Unit of brightness; see Lambert.

Foot-pound.—[ml^2/t^2]. Work required to raise one pound a vertical distance of one foot, where g = 980.665 cm/sec^2 (cf. meter-kilogram).

Foot-poundal.—[ml^2/t^2]. Work done by force of one poundal (q.v.) acting through a distance of one foot.

Force.—[ml/t^2]. That which imparts acceleration to material bodies.

Force, Electromotive.—See Potential.

Force, Magnetizing.—See Field intensity.

Force, Magnetomotive.—See Potential.

Force de cheval.—See Cheval-vapeur.

Frequency.—(ν). [N/t]. Number per unit of time. In case of vibrations, waves, etc., the frequency is the number of complete vibrations, of complete waves, etc., per unit of time.

Gamma.—(γ). [$\sqrt{m/\mu l^2}$], [$\sqrt{m l_c/t^4}$]. A unit of magnetic field intensity; 0.000 01 gauss.

Gas constant.—1. (R). [work/mass-degree], [$l^2/t^2 T$]. The coefficient R in the ideal gas equation $pv = RTm$; p = pressure, v = volume of the mass m at absolute temperature T. 2. (R). [work/mole-degree]. **Gas constant per mole** obtained by expressing m in moles. 3. (k). [work/molecule-degree], [$ml^2/t^2 T$]. **Boltzmann's molecular gas constant:** obtained by expressing m in terms of number of molecules.

Gas, Ideal.—One which strictly satisfies the equation ($pv = RTm$) and other relations deduced from the classical kinetic theory of gases on the assumption that the molecules are infinitely small and devoid of mutual attraction.

Gauss.—[$\sqrt{m/\mu l^2}$], [$\sqrt{m l_c/t^4}$]. The cgsm unit of magnetic field intensity.

Gaussian gravitation constant.—The square root of the intensity of the gravitational field of force of the sun at a point whose distance from the sun is the astronomical unit of length (cf. Gravitation constant).

Geepound.—See Slug.

Gilbert.—[$\sqrt{ml/\mu t^2}$], [$\sqrt{\epsilon ml^3/t^4}$]. Electromagnetic unit of magnetic potential, of magnetomotive force. Unless contrary is indicated, it is the cgsm unit. In precise work, the International gilbert, based upon the Int. elec. units, should be distinguished from the absolute, or cgsm, gilbert.

Grade.—[θ]. Unit of plane angle, $\frac{1}{400}$ of complete circumference.

Gram atom.—See Mole.

Gram calorie.—See Calorie.

Gram equivalent.—See Mole.

Gram formula weight.—See Mole.

Gram weight.—See Weight.

Gravitation constant.—(G). [l^3/mt^2]. The coefficient G occurring in the equation $f = G\ (mm')/r^2$; f = force of gravitational attraction between two point masses (m, m') in vacuo, r = distance between m and m' (cf. Gaussian gravitation constant).

Gravity, Acceleration of.—(g), (g_s). [l/t^2]. Unless the contrary is indicated, this expression refers specifically to the earth, and denotes the resultant acceleration downward experienced by a freely falling body placed at the point considered. It includes centrifugal effects arising from the rotation of the

earth, as well as the effects of gravitational attraction (cf. Gravity, standard).

Gravity, Specific.—See Specific gravity.

Gravity, Standard.—(g_s). [l/t^2]. Standard gravity is the value adopted by the International Committee on Weights and Measures as the "accepted" value of the acceleration of gravity to which all measurements involving this quantity are to be referred. Thus a pressure of x cm of mercury at t°C is to be understood as denoting the pressure exerted by x cm of mercury at t°C at a place where the acceleration of gravity is g_s. The accepted value is g_s = 980.665 cm/sec^2 (= 32.174 ft./sec^2).

Heat. —1. By the heat of a process is meant the amount of heat evolved, per unit quantity of material involved, during the isothermal process, the process proceeding in the direction indicated. The quantity of material may be expressed in terms of mass, of moles, of equivalents, etc., as may seem desirable. 2. By the latent heat of a transformation is meant the amount of heat absorbed per unit quantity of material transformed, the transformation proceeding in the direction indicated. Latent heat of transformation of A to B = −(heat of transformation of A to B) = heat of transformation of B to A.

Heat diffusivity.—See Diffusivity.

Heat, Specific.—See Capacity, and Specific heat.

Hecto-.—Prefix denoting 100.

Hefner unit.—A superseded unit of luminous intensity; approximately = 0.9 Int. candles.

Henry.—[μl], [t^2/d]. Unit of electromagnetic inductance. Defined as that inductance for which an induced electromotive force of one volt is produced when the inducing current is changed at the uniform rate of one ampere per second.

Horsepower.—(h.p.). [work/time], [ml^2/t^3]. 1. (HP) **Primary definition** of the term is work done at the rate of 550 foot-pounds per second. 2. For electrical purposes it is regarded as exactly = 746 watts, which is frequently called the **electrical horsepower.** 3. **Continental horsepower.** See Cheval-vapeur.

Humidity. —1. **Absolute humidity** of a gas is the actual amount of water vapor per unit volume of the gas. Usually expressed in terms of the actual pressure of the water vapor present. 2. **Relative humidity** of a gas = ratio of the pressure of water vapor present to the pressure of water vapor which is in equilibrium with water at the same temperature. 3. **Dew-point** of a gas is the temperature at which the pressure of water vapor in equilibrium with water is equal to the actual pressure of the water vapor contained in the gas. If the temperature of the gas be varied while its absolute humidity remains unchanged, then the dew-point is that temperature at which the relative humidity is 100%. 4. If the bulb of a thermometer be encased in a fabric which is kept wet with water (wet-bulb), the thermometer will record a lower temperature than if the bulb were dry (dry-bulb). If the circulation over the wet bulb is sufficiently rapid, the difference in the temperatures depends solely upon the total pressure of the gas, its absolute humidity, and its temperature. Hence the humidity of the atmosphere, or of any other very large volume of gas, can be readily determined by the use of wet- and dry-bulb thermometers.

Hydrometer.—An instrument which, by the extent of its submergence, indicates the specific gravity of the liquid in which it floats. Frequently, its readings are expressed in degrees (°). Various systems of graduations are in use, see p. 31.

Hygrometric.—Pertaining to humidity of atmosphere.

Hypsometry.—The art of measuring the elevation above sea-level. More specifically, the use of the boiling-point of water for such measurements.

Ice point.—(T_0). Temperature at which water freezes when under the pressure of one normal atmosphere.

Ideal gas.—See Gas, ideal.

Illumination.—$[\psi/l^2]$. The illumination at a point of a surface is the surface density of the luminous flux incident at that point.

Inch of water [of mercury, etc.] at $t°$.—Analogous to cm of water (q.v.)

Index of absorption. —See Absorption.

Index of refraction. See Refraction.

Inductance.—The electrical inductance of circuit A with reference to circuit B is ψ_A/I_B; ψ_A = flux of magnetic induction through A as a result of the current I_B in B. A and B may be the same circuit.

Induction.—1. That modification which is acquired by a medium when it becomes the seat of a field of force, and which is evidenced by the fact that its boundaries with other media exhibit distinctive properties which they do not possess in the absence of the field. 2. The distinctive **properties** mentioned in (1); as in magnetization by induction, induced electric charges, etc. 3. **Electrostatic induction.** $[\sqrt{m/\mu l^3}]$, $[\sqrt{\epsilon m/l l^2}]$. ϵF, ϵ = dielectric constant, F = intensity of electrostatic field of force. **Electric displacement** = $\epsilon F/4\pi$. 4. **Magnetic induction** (B). $[\sqrt{\mu m/l l^2}]$, $[\sqrt{m/\epsilon l^3}]$. $B = \mu H$, μ = magnetic permeability, H = intensity of magnetic field of force. 5. **Electromagnetic induction** is the phenomenon which is characterized by the appearance, in every circuit, of a cyclical emf which is proportional to the rate of change of the flux of magnetic induction through that circuit.

Intensity coefficient.—See Black body.

Intensity, Field.—See Field intensity.

Intensity, luminous. 1 Of a point source in a given direction = amount of luminous flux, per unit of solid angle, which the source emits in the direction considered. 2. Of a point of an extended source = brightness of that point of the source; also called intrinsic brightness. 3. Of an extended source, in a given direction, is its intensity at a point so distant in the stated direction that the source may be regarded as a point. For nearer points the apparent intensity will depend upon the distance, and is defined as the intensity of that point source which at the same distance will produce the same illumination (cf. flux).

Intensity of magnetization.—See Magnetization.

Intensity of radiation. 1 The intensity of the radiation emitted in a specified direction by a body is the amount of radiant energy emitted in that direction, per unit of time, per unit of area, and per unit of solid angle of emission. For spectral, or monochromatic, intensity, See Radiance. 2. Of received radiation, See Irradiation. 3. Of radiation in transit. The amount of radiant power per unit area which passes through an element of area which is normal to the direction of propagation; this equals the volume density of radiant energy at the point considered.

International electrical units.—A system of electrical and magnetic units based upon the ohm, the ampere, and secondarily upon the volt, all as realized by certain concrete standards which have been internationally agreed upon, and upon the cgs units for such other quantities as may be involved. The concrete standards have been so chosen as to make the international system nearly identical with the practical system; as now defined, the outstanding discrepancy in no case exceeds 52 parts in 100 000. In distinguishing between the two systems, the units of the practical system are described as absolute, those of the other, as international. The introduction of the volt as a secondary unit defined by a concrete standard (Weston normal cell = 1.018300 Int. volts at 20°C) introduces confusion when measurements of high precision are to be recorded. In these Tables, values based upon the Int. ohm and the Int. ampere (as defined by the silver voltameter) are denoted by (a). Those based on the Int. ohm and the Int. volt (as defined by the standard cell) are denoted by (v).

Irradiation.—The radiant power, per unit of area, incident upon a surface.

Joule.—$[ml^2/t^2]$. 1. Absolute joule = 10^7 ergs. 2. International joule = work expended per second by an Int. ampere in an Int. ohm.

Karat.—(K). Denotes the "fineness of gold" in terms of parts (by weight) of gold per 24 parts of the alloy. Twenty-four g of an n karat alloy contains n g of gold, the alloy is "n carats fine."

Kelvin.—(K). Name applied to the absolute centigrade scale of temperature.

Kilo-.—Prefix denoting 1000.

Kilogram calorie.—See Calorie.

Kilogram-meter.—A torque equivalent to that of one kilogram weight acting on a lever-arm one meter long.

Kilowatt-hour.—Work expended by one kilowatt in one hour. In Great Britain it is quite generally called **Board of Trade unit** (B.T.u.).

Kinematic viscosity.—$[l^2/t]$. Ratio of viscosity to density.

Lambert. $[\psi/l^2\omega]$. The brightness of a surface which, radiating in accordance with Lambert's law, emits a total luminous flux of one lumen per cm². For such a surface, brightness is independent of direction of the line of sight and equals $1/\pi$ lumen, per steradian, per cm² = $1/\pi$ candles per cm². If the total emission is one lumen per sq. ft., the brightness is called one **foot-lambert.**

Lambert's law.— $I = I_o\cos\theta$; $I_o[I]$ = intensity of radiation emitted in direction normal [at angle θ with normal] to the surface. In many cases this law does not express the facts.

Latent heat.—(l, L). See Heat.

Latitude.—(lat.). 1. The angular distance of a point from the equator of a spheroid, measured along a great circle passing through the poles. 2. **Celestial latitude.** See Celestial sphere.

Legal ohm.—A unit of resistance; so designated by the International Conference of 1884, and defined as the resistance of a column of mercury 1 mm² in cross-section and 106 cm in length at the temperature of melting ice. It was never legalized.

Light-year.—Distance traveled by light in free space in one year.

Line.—Unit of flux of magnetic induction = one maxwell.

Liter-atmosphere.—The amount of external work done when a volume is increased by one liter against an external pressure of one atmosphere.

Longitude. (long.). 1. The longitude of a point is the angle which its axial plane makes with a fiducial one. For the **earth**, angles measured from the fiducial plane towards the west are usually considered positive. 2. **Celestial or astronomical** longitude. See Celestial sphere.

Loschmidt's number.—(n_o). $[l^{-3}]$. Number of molecules per unit volume of an ideal gas at 0°C and pressure A_n.

Lumen. $[\psi]$. Fundamental unit of luminous flux. A uniform point source of one candle emits 4π lumens.

Luminous flux.—See Flux, luminous.

Luminous intensity.—See Intensity, luminous.

Lunar month.—The time which elapses between successive new moons. Also called synodical month.

Lux.—A unit of illumination, one lumen per square meter.

Magnetic flux.—See Flux, magnetic.

Magnetic induction.—See Induction.

Magnetic moment.—See Moment.

Magnetization, Intensity of.—Magnetic moment per unit of volume (cf. moment).

Magnetomotive force.—(mmf). See Potential.

Magnitude.—The **magnitude**, or **apparent magnitude**, (*m*) of a star is primarily an indication of the amount of light the earth receives from it. The value to be assigned to the latter depends upon the characteristics of the perceptive apparatus: visual, photovisual, photographic, and radiometric magnitudes are to be distinguished. Certain stars near the north pole have been chosen as standards; the numerical magnitudes assigned to them are such as represent satisfactorily the range covered by early naked-eye estimates, and satisfy the equation $m = 2.5$ $(\log_{10} I_0 - \log_{10} I)$, I = intensity of light from a star of magnitude *m*, and I_0 = that from one of magnitude zero. For Vega, $m = 0.2$; a star of $m = 6$ is near the limit of naked-eye visibility. The **absolute magnitude** *M* is internationally defined as the apparent magnitude the star would have if its distance were 0.1 parsec; $M = m + 5 + 5 \log_{10}\pi$, π = parallax expressed in ".

Mass, Engineers' unit of.—*See* Slug.

Maxwell.—The cgsm unit of flux of magnetic induction.

Mean distance.—In astronomical parlance, the mean distance of a planet from the sun denotes the mean of the greatest and the least distance from the sun to the path of the planet. Similarly in other cases.

Mean spherical candlepower.—Average candlepower of a source, in all directions.

Mega-.—Prefix = 1 000 000.

Megmho.—Conductance of one reciprocal microhm.

Meter-candle.—The illumination of an element of surface one meter distant from a uniform source of one candle situated upon the normal to the center of the element. One lux.

Meter-kilogram. -$[ml^2/t^2]$. Work required to raise one kilogram a vertical distance of one meter at a place where the acceleration of gravity is 980 665 cm/sec.^{2}

Mho.—An electrical conductance of one reciprocal ohm.

Micro-.—Prefix denoting $1/10^6$.

Microhm.—10^{-6} ohm.

Micromicro-.—Prefix denoting $1/10^{12}$

Micron.—(*μ*). Unit of length = $1/10^6$ m = 0.001 mm.

Mil.- 0.001 in. (*cf.* Circular inch).

Milli-.—Prefix = 0.001.

Millimicro-.—Prefix = 0.000 000 001.

Minute.—1. (min). Time, 1_{1440} of a day. 2. ('). Unit of angle, 1_{60} degree. 3. (`). **Centesimal minute** = unit of angle = 0.01 grade.

Modulus.—1. *See* Elastic modulus. 2. For the several elastic moduli—bulk, compression, elasticity, rigidity, torsion, Young's - *see* distinguishing name.

Mohs.—An arbitrary scale of hardness based upon a selected list of 10 native minerals.

Mole.—A variable, derived unit of mass; its mass is numerically equal to the molecular weight of the substance measured. The expressions gram-mole, kilogram-mole, etc. are used to designate the basic unit of mass employed. Similarly derived units based upon the atomic weight, the formula weight, or the equivalent are called the **gram-atom, gram-formula weight** or **gram-equivalent** when the gram is the basic unit, and correspondingly in other cases.

Molecular.— For molecular properties, *see* appropriate properties.

Molecular volume.—Volume occupied by one mole. Molecular weight divided by density.

Molecular weight.—(*M*). The sum of the atomic weights of all the atoms contained in a molecule.

Moment.—1. Of force (*F*) about a point = Fl, l = perpendicular distance from the point to the line of *F*. 2. Of a couple = product of either force times perpendicular distance between them. 3. Of a magnet = moment of couple acting upon it when it is at right angles to a magnetic field of unit intensity. 4. Of inertia about an axis = sum of the products

of each element of mass times the square of its distance from the axis.

Month.—1. Period of time determined by motion of moon. *See* lunar, synodical, tropical, sidereal, anomalistic, nodical, draconic. 2. **Solar month** = 1_{12} of tropical year. 3. **Calendar month** = conventional subdivision of year.

Myria-.—Prefix = 10 000.

Node.—1. A point of a **standing wave** where the displacement is independent of the time. 2. In **astronomy**, the points where an orbital, or other, plane cuts the ecliptic; the **rising node** is the one at which the passage across the plane of the ecliptic is from south to north.

Nodical month.—Time required by the moon to pass from one rising node to the next. Also called **draconic month**.

Noon.—*See* Time.

Normal.—1. The normal to a **surface** is a line drawn perpendicular to the surface at the point considered. 2. Any **line** perpendicular to another may be said to be normal to it. 3. A **concentration** of one gram-equivalent per liter.

Normal atmosphere. (*A_n*). *See* Atmosphere.

Numeric. (*N*). A pure number A dimensionless quantity.

Nutation. *See* Celestial sphere.

Oersted. The cgsm unit of magnetic reluctance.

Ohm. (Ω). A unit of electrical resistance. 1. **Absolute ohm** = 10^9 cgsm units. 2. **International ohm** is the resistance, at the temperature of melting ice, offered to an unvarying electric current by a column of mercury, of constant sectional area, having a mass of 14.4521 grams and a length, at the temperature mentioned, of 106 300 cm.

Ohm-centimeter.—Unit of electrical volume resistivity. The resistivity of a material of which a uniform bar one cm^2 in sectional area has a longitudinal resistance of one ohm per cm of length. Frequently called one ohm per centimeter cube.

Ohm (cm, gram). Unit of electrical mass resistivity. The resistivity of a material of which a bar, having such a uniform section that its mass per linear cm is one gram, has a longitudinal resistance of one ohm per cm of length.

Ohm (meter, mm).—Unit of electrical volume resistivity. The resistivity of a material of which a circular cylinder one mm in diameter has a longitudinal resistance of one ohm per meter.

Ohm (meter, mm²).—Unit of electrical volume resistivity. The resistivity of a material of which a circular cylinder one square mm in sectional area has a longitudinal resistance of one ohm per meter.

Ohm (mil, ft.). -Analogous to ohm (meter, mm). Cylinder one mil in diameter, resistance of one ohm per foot.

Ohm (mile, pound).—Analogous to ohm (cm, gram).

Ohm-inch.—Analogous to ohm-centimeter.

Parallax. -1. The annual parallax of a star is defined as the maximum angle subtended by one astronomical unit of length at the distance of the star from the sun. 2. The **equatorial horizontal parallax** of a member of the solar system is the maximum angle subtended by the equatorial radius of the earth at the distance of the earth from the member considered.

Parsec. -The distance of a star for which the annual parallax is one second of arc.

Pentane candle.—A superseded unit of luminous intensity = one Int. candle.

Percent. (%). The number of units of the constituent in 100 units of the mixture containing it. If units of volume are used, the ratio is called **volume percent**; if units of mass, it is called **mass percent, weight percent**, or simply **percent**. (% must be distinguished from ‰ which is frequently used to denote per thousand.)

Perigee.—That point of the moon's orbit which is nearest to the earth (*cf.* apogee).

Perihelion.—That point of a planet's, or comet's, orbit which is nearest to the sun (cf aphelion).

Permeability.—(μ.) The force (f) of repulsion between two rigidly magnetized poles (m, m') at a distance r apart is $f = (mm')/(\mu r^2)$; μ depends upon the material in which the poles are immersed, and is called its permeability.

Pferdekraft. —See Cheval-vapeur.

Phot.—An illumination of one lumen per cm^2.

Photoelectric constant.—1. h/e. It is $1/\nu$ of the rise in potential required to impart to a negative electron the energy it has when emitted under the action of radiation of frequency ν. 2. hc/e. This is λ times the rise in potential mentioned in (1). λ = wave-length in vacuo.

Planck's constant of action.—(h). $[ml^2/t]$. A universal constant which fixes the amount of energy contained in the individual bundles, or quanta, of radiation emitted by a radiating body. Each such bundle contains an amount of energy = $h\nu$, ν = vibration frequency of the radiation. **h** is also called **Planck's quantum.**

Poise.—$[m/lt]$. The cgs unit of viscosity. If the tangential force, per unit area, which one layer of a fluid exerts upon an adjacent one is one dyne when the space rate of variation of the tangential velocity from layer to layer is unity, the viscosity of the fluid is one poise.

Poisson's ratio.—If a bar of uniform section be subjected to a pure tensile stress, the ratio of its transverse contraction per unit of transverse thickness to its elongation per unit of length is called the Poisson's ratio of the material.

Pole strength.—See Quantity of magnetism.

Poncelet.—Unit of power = 100 meter-kilograms per second.

Potential.—The excess of the potential at the point A over that at B, with reference to any quantity m, is the mechanical work per unit of m which must be done in carrying a very small positive amount of m from B to A. The difference in electrical potential is called **electromotive force, emf, potential difference;** in magnetic potential, is called **magnetomotive force, mmf.**

Potential gradient.—The space rate of increase in the potential. If the direction in which the rate to be measured is not stated, that corresponding to the maximum gradient is to be understood.

Pound weight.—See Weight.

Poundal.—The unit of force in the fps system. It is the force which, if acting continuously upon a mass of one pound, will impart to it a uniform acceleration of one foot per second^2 (cf. Dyne).

Power.—1. The time **rate of doing work.** 2. If when the two junctions of a bimetallic circuit differ in temperature by a small amount (dt), there is an open circuit emf (dE) around the circuit, then (dE)/(dt) is called the **thermoelectric power** of the circuit, corresponding to the average temperature of the two junctions. 3. The **ability** to do some specific thing; as in rotatory power.

Practical electric units.—A system of electrical units based upon 10^9 cm, 10^{-11} gram, sec, and the permeability of a vacuum, as fundamental units. The units of most interest are the ohm (= 10^9 cgsm), ampere (= 0.1 cgsm), and volt (= 10^8 cgsm). Frequently described as absolute (cf. Int. elec. units).

Precession of the equinoxes.—See Celestial sphere.

Pressure.—(p), (P). $[m/lt^2]$. Normal force per unit of area. A **hydrostatic pressure** is a pressure which is the same in all directions. For critical pressures, see Critical.

Quadrant.—1. Unit of **angle** = 90°. 2. Formerly used occasionally to denote the **henry.**

Quantity of electricity.—1. (es). The **electrostatic unit** is that quantity which when concentrated to a point and placed at unit distance from an equal point charge will exert upon it a unit force, the surrounding medium being a vacuum. 2. (em). The **electromagnetic unit** is that quantity which is transferred per unit of time across any section of an infinitely long, straight, linear conductor when the current is such that the intensity of the resulting magnetic field at unit distance from the conductor is unity. 3. For **other units**—coulomb, electronic charge, faraday—see corresponding names.

Quantity of magnetism.—Also called **pole strength.** 1. The **electromagnetic unit** is that quantity which when concentrated to a point pole and placed at a unit distance from an equal point pole will exert upon it a unit force, the surrounding medium being a vacuum. 2. The **electrostatic unit** is that quantity which when concentrated to a point pole and placed at a unit distance from an infinitely long, straight, linear conductor would experience a unit force as a result of a current in the conductor such that one electrostatic unit of electricity per second is transferred across each section of the conductor. 3. The Int. electric unit is not named, it is the same as the cgsm unit.

Quantum.—1. Certain processes are essentially discrete, and consequently parcel out into bundles the several quantities involved. If for a certain quantity and a particular process these bundles are all alike, it is now customary to call them quanta, without implying that the quantity so bundled has in itself any atomistic properties. 2. **Planck's quantum.** See Planck.

Radian.—An angle which encloses, of the circumference of a concentric circle, an arc = radius.

Radiance.—The radiance of a body, within the spectral range λ_1 to λ_2, is defined as the intensity of the radiant energy, having wave-lengths lying between λ_1 and λ_2, which the body emits in a direction perpendicular to its radiating surface. If the spectral range is not mentioned, all wave-lengths are to be included; this is frequently called the **total radiance.** The **spectral,** or **monochromatic, intensity** of the radiance of wave-length λ is defined as the ratio of the radiance within the range $(\lambda - \frac{1}{2}d\lambda)$ to $(\lambda + \frac{1}{2}d\lambda)$ to $d\lambda$, when the latter is indefinitely small (cf. Emissivity).

Radiation constants.—See Black body.

Rankine.—A name sometimes applied to the absolute Fahrenheit scale of temperature.

Réaumur.—(R). A thermometric system in which the freezing point of water is called 0°, and the boiling point, 80°.

Reflectivity.—The ratio of the intensity of the light specularly reflected from a surface to the intensity of the light incident upon it. It is a pure numeric.

Refraction.—1. The **index of refraction, refractive index,** or **refractive exponent** is n = sin i/sin r; i = angle of incidence from a vacuum upon the substance, and r = angle of refraction, each measured from the normal to the surface. 2. **Refractivity** is $(n - 1)$. 3. **Specific refractivity** (r_G) is $(n - 1)/d$. **Specific refraction** (r_L) is $(n^2 - 1)/d(n^2 + 2)$. d = mass per unit of volume. 4. **Molecular refractivity** = Mr_G. **Molecular refraction** = Mr_L. M = molecular weight. By replacing M by the atomic weight, the corresponding **atomic values** are obtained. 5. **Refractive constant** of a solute is its specific refractivity computed on the assumption that the refractivity of the solution is equal to the sum of the refractivities of its pure constituents each multiplied by the ratio of its mass per unit volume of the solution to its own density when pure.

Reluctance.—The magnetic reluctance of a body between two specified equipotential surfaces is the ratio of the difference in the two potentials divided by the flux of magnetic induction from [to] either surface to [from] the body. It has no significance unless these two fluxes are the same.

Resistance.—1. The **electrical resistance** of a body between two specified equipotential surfaces is E/I, where E is the unchanging difference in the potentials of the surfaces and I is the result-

ing current across any transverse section between them. 2. **Specific resistance.** See Resistivity.

Resistivity.—1. [resistance × length]. **Resistivity, or volume resistivity,** of a substance is the longitudinal resistance per unit of length of a uniform bar of the substance of unit sectional area. 2. [resistance × mass/(length)^2]. **Mass resistivity** of a substance is the longitudinal resistance per unit of length of a uniform bar of the substance of such a sectional area that it contains one unit of mass per unit of length. 3. [resistance] **Surface resistivity** is the resistance per unit of length of a strip of the surface of unit width. It has reference solely to the current which is restricted to the surface.

Rhe.—Name proposed for cgs unit of fluidity; = one reciprocal poise.

Right ascension.—See Celestial sphere.

Rigidity.—If to the four faces of a cube which are parallel to a given edge there be applied tangential stresses which are equal in absolute value, perpendicular to the given edge, and so directed as to produce a pure distortion, the other two faces will be deformed into diamond shaped figures if the material is isotropic. The modulus of rigidity is defined as the quotient of the stress on any one of the faces divided by the resulting change in any one of the angles of a distorted face. Also called **modulus of shear, Coulomb's modulus, modulus of torsion** (the last is undesirable).

Rotation.—See Rotatory power.

Rotatory power, Optical.—1. The **natural rotatory power** is θ/l, where θ is the rotation of the plane of polarization which occurs in a path of length l. The **specific rotatory power** ([α]) is θ/dl, d = density. The **molecular** [or **atomic**] **rotatory power** is $M\theta/dl$ [or $A\theta/dl$]; M = molecular, A = atomic weight. 2. The **magnetic rotatory power** is $\theta/(Hl \cos \alpha)$, where H = intensity of the magnetic field and α = angle between H and the path of the light. It is commonly called **Verdet's constant.** From the magnetic rotatory power, the **specific** ([ω]), **molecular,** and **atomic magnetic rotatory powers** are derived exactly as in the case of natural rotation. The ratio of any one of these quantities to the corresponding one for a chosen reference substance is called the **relative power.** Water is the reference substance commonly chosen, and [Ω] is used to denote the molecular magnetic rotatory power relative to water.

Rydberg's fundamental frequency, and series constant.—See Series, spectral.

Secohm.—A superseded name for the henry.

Second.—1. (sec). Time, $\frac{1}{86400}$ day. Mean solar day, unless contrary is indicated. 2. (″). Unit of angle, $\frac{1}{1600}$ degree. 3. (ˈ). **Centesimal second** = 0.0001 grade.

Seger cone.—One of a graded series of cones of refractory material which, by their softening and the resultant deformation, indicate the heat treatment to which they have been subjected.

Series, Spectral.—Spectral lines, or groups of lines, which occur in orderly sequence. Most of these sequences can be represented by an equation of the form $\frac{1}{\lambda} = A - \frac{BN}{(m + \alpha + \beta/m^2)^2}$; λ = wave-length in vacuo; m is an integer varying from one line (or group) to another; for any one series, A, B, N, α and β are constants; B is an integer; N is known as **Rydberg's constant,** its value is determined by the constitution of the radiating atom. On Bohr's theory, $N = N_\infty \frac{M}{M + m_0}$, where M = mass of the atom, m_0 = electronic mass, and $N_\infty = 2\pi^2 m_0 e^4/h^3 c \epsilon_0{}^2$; N_∞ is known as **Rydberg's universal series constant;** e = electronic charge; h = Planck's constant; ϵ_0 = dielectric constant of vacuum; c = velocity of light in vacuo. On this theory, B denotes the number of electrons displaced from their normal positions, m is the **principal quantum number,** α depends

upon the subordinate, or azimuthal, **quantum number, and** β = 0. For atoms of the type of hydrogen, $\alpha = 0$, $\beta = 0$; for others $(m + \alpha + \beta/m^2)$ is frequently called the **effective quantum number,** generally it is not an integer. **Rydberg's fundamental frequency** is $\nu_\infty = cN_\infty$.

Sidereal month.—The time required for the moon to complete one apparent circuit among the stars.

Siemens unit.—(S.E.). A superseded unit of electrical resistance proposed in 1860 by Werner von Siemens; defined as the resistance at 0°C of a column of mercury one meter long and of a uniform cross section = one mm^2.

Slug.—A unit of mass. 1. The mass which will acquire an acceleration of one foot per sec^2 when continuously acted upon by a force of one pound weight. Also called **geepound, and engineer's unit of mass.** 2. The **metric slug** is the mass which will acquire an acceleration of one meter per sec^2 when continuously acted upon by a force of one kilogram weight.

Solar month.—$\frac{1}{12}$ tropical year.

Solubility.—1. By solubility of the **non-gas** a in b is meant the mass of a per unit mass of b which is contained in the mixture which is in equilibrium with an excess of a. In this mixture b is said to be saturated with a. Data are frequently restricted to mass of a per unit mass of mixture, mass of a per unit volume of mixture, or moles of a per mole of mixture. 2. **Solubility of a gas** is C_s/C_g, C_s = concentration of gas in the solution, C_g = concentration of gas in overlying gas phase. 3. **Solubility product** of an ionized substance (A_nB_m) in a stated solvent = $[A]^n \cdot [B]^m$, where $[A]$ and $[B]$ denote the concentrations of the two ions when the solution is saturated with the substance.

Specific gravity.—($d_{t_1}^{t_2}$). The ratio of the mass of a certain volume of the substance at the temperature t_2 to that of the same volume of a reference substance (usually water) at temperature t_1. Frequently, but incorrectly, called density.

Specific heat.—1. **Heat capacity.** See Capacity. 2. **Specific heat of electricity.**—See Thomson effect. 3. **Einstein's specific heat constant** (β) = ratio of Planck's constant (h) to Boltzmann's molecular gas constant (k_0). 4. **Ratio of specific heats** = $\gamma = c_p/c_v$; c_p, c_v = specific heat at constant pressure and at constant volume, respectively.

Specific inductive capacity.—The ratio of the dielectric constant of the substance to that of a vacuum.

Specific refractive power.—Used indifferently to denote several of the refractive constants (cf. Refraction).

Sperm candle, English.—A superseded unit of luminous intensity = one Int. candle.

Spheradian.—See Steradian.

Spherical candlepower, Mean.—See Mean spherical candlepower.

Square.—(sq.), (²). Used in conjunction with the name of a unit of length to form the name of a related unit of area; e.g., square foot (sq. ft.), (ft.²) is the name of a unit of area equivalent to the area of a square with edges one foot long.

Square degree.—The solid angle enclosed by a cone of vanishingly small vertex angle 2θ is $k\pi\theta^2$. If θ is expressed in radians and the unit of solid angle is so chosen that $k = 1$, that unit is called a **steradian.** If θ is expressed in degrees, and $k = 1$, the corresponding unit of solid angle is called a **square degree.** One square degree = $(\pi/180)^2$ steradians. This procedure defines a definite unit of solid angle although the solid angles enclosed in cones of finite vertex angles are not proportional to the squares of those angles.

Stefan's constant.—See Black body.

Steradian.—The solid angle which encloses on the surface of a concentric sphere an area = (radius)^2.

Stoichiometric.—Pertaining to the ratio of the masses of the several elements contained in a pure chemical compound.

Strain.—1. For pure distortion the strain is measured by the change in a significant angle. 2. The ratio of change in size to original size.

Stress.—The force per unit of area over which it acts.

Surface tension.—(γ). [m/t^2]. Owing to molecular attraction, two fluids in contact adjust themselves so that the area of their interface is a minimum, consistent with other requirements. This adjustment may be pictured as arising from a tension residing in the surface itself; to this is given the name **surface tension.** Its value is defined as the normal, tensile force, per unit of length, across any line traced on the surface.

Susceptibility.—(κ). In the electromagnetic systems of units, $4\pi\kappa$ is the excess of the magnetic permeability of the substance over that of a vacuum.

Synodical.—In astronomy, the synodical period of a body is the interval between its successive returns to the same position with reference to the plane which is perpendicular to the plane of the ecliptic and which continuously passes through the centers of the earth and the sun.

Synonical month.—See Lunar month.

Temperature conductivity.—See Diffusivity.

Tension, Surface.—See Surface tension.

Tenth-meter.—10^{-10} meter; one Ångstrom unit.

Thermal.—See Heat.

Thermoelectric power.—See Power.

Thomson effect.—In a region in which the temperature of a homogeneous metallic conductor varies from section to section, there exists a potential gradient which is proportional to the product of the temperature and its gradient. This is the Thomson (or Kelvin) thermoelectric effect. The constant of proportionality is called the coefficient of the effect. If the coefficient is positive, a positive electric current flowing from hot section to cooler section tends to make the temperature more uniform; it is as if the current carried heat from hot portion to cooler portion, as if the electricity had a certain specific heat. This is what Thomson called the **specific heat of electricity.** It may be either positive or negative, depending upon the metal.

Time.—True noon, or local true noon, is the instant at which the sun is bisected by the meridional plane of the observer. **Mean noon,** or local mean noon, is the instant at which a fictitious mean sun is bisected by the meridional plane. This **mean sun** is one endowed with such a uniform, apparent angular velocity in the equatorial plane that in one tropical year it will make exactly the same number of apparent revolutions around the earth as are made by the true sun. Time measured from the true noon is called **true, or apparent, solar time;** that from mean noon is called **mean time.** The excess of mean time over true time is called **equation of time.** The earth has been divided into a series of time zones, each 15° of longitude in width, so that intercourse may be facilitated by all places in each zone using the mean time corresponding to the center of the zone; this is known as **standard time.** The first zone is centered on Greenwich, England.

Titer.—See Concentration.

Torque.—The moment of a force.

Tropical month.—The yearly average of the time required for the moon to traverse 360° of astronomical longitude.

Twist.—If a uniform bar of free length l be clamped rigidly at one end and the other end be twisted, about the axis of the bar, through an angle θ, the twist of the bar is defined as θ/l. Similarly for other cases.

Units, Systems of.—The fundamental units in most absolute systems are those of mass, length, time, thermometric degree, and the dielectric constant (or the magnetic permeability) of a vacuum. Other units are defined in terms of these by the use of established relations, arbitrary factors being made unity. The most common systems are the centimeter-gram-second-degree Centigrade (cgs), and the foot-pound-second-degree Fahrenheit (fps) systems. See also International electric units, practical electric units, and absolute.

Van der Waals.—See Waals.

Violle unit.—A superseded unit of luminous intensity based upon the brightness of fused platinum at the temperature of solidification.

Viscosity.—If a fluid is flowing in the plane yz with velocity v it exerts upon an adjacent plane a tangential drag $= \eta(dv)/(dx)$, per unit of area. η is called the viscosity, coefficient of viscosity, or coefficient of internal friction. Unit: poise.

Viscosity, Kinematic.—Viscosity divided by density.

Volt.—The electrical potential difference which, when steadily applied to a conductor having a resistance of one ohm, will produce in it a current of one ampere (cf. absolute and international units). The Int. Committee authorized by the London Conference, 1908, agreed to regard the emf of the Weston normal cell at 20°C as exactly 1.0183 Int. volts. This furnishes a subsidiary definition which is slightly discordant with the primary one. These tables distinguish between the two, and between units derived from them, by using (a) to denote those based on ampere and ohm, and (v) to denote those based on volt as defined by the Weston cell.

Volt-electronic charge.—Analogous to volt-faraday.

Volt-faraday.—The work which must be done in order to transfer one faraday of positive electricity from any point to another having a potential one volt higher than the former.

Volt-second.—Unit of flux of magnetic induction. The amount defined by the change per second, of the magnetic induction through an area, required to induce around the area an emf of one volt.

Volume, Specific.—Reciprocal of the density.

Waals, Van der.—In the equation $(p + a/v^2)(v - b) = 1 + at$, a and b are known as Van der Waals' constants; $a[b] = $ pressure [volume] constant.

Watt.—Unit of power; work done at rate of one joule per second.

Watt-hour.—Work expended by one watt in one hour (cf. kilowatt-hour).

Wave-length.—(λ). Distance between consecutive corresponding points in a monofrequent wave train. Occasionally applied to complex waves.

Wave number.—Reciprocal of wave-length.

Weight.—The force with which a body, left to itself, is urged towards the earth. In the absolute systems of units it is numerically equal to the mass of the body multiplied by the acceleration of gravity (g) at the position considered; hence varies with position. Such expressions as **gram weight** [pound weight] are to be interpreted as meaning the weight of a gram [a pound] at a place where g has the standard value, 980.665 cm/sec.²

Wien's displacement constant.—(w). See Black body.

Year.—(yr). Time required for earth to make one complete circuit of its orbit, as defined by its return to the same position as determined by the sun and some celestial point of reference. For the **tropical, equinoctial, or ordinary** year the reference point is the mean vernal equinox; for the **sidereal, or true,** year, it is a fixed star; for **anomalistic** year, it is perihelion of earth's orbit; for **eclipse** year, it is ascending node of moon's orbit.

Young's modulus.—If a bar of uniform section be subjected to a longitudinal tension, the ratio of this stress to the resulting elongation per unit of length is called its Young's modulus. Also called **modulus of elasticity, elastic modulus, longitudinal elasticity, coefficient of resistance to extension, modulus of traction.**

ELEMENTS AND ATOMS

ATOMIC WEIGHTS

The values given in column four were compiled for International Critical Tables (I. C. T.) by Prof. G. P. Baxter in 1923 and are those upon which all the data given in International Critical Tables are based.

Following these are shown the accepted atomic weights back to 1882. For the period since 1903 these are taken from the reports of the International Committee on Atomic Weights; for the period 1894 to 1903, from the reports of the American Chemical Society's Committee on Atomic Weights; for the year 1882, from F. W. Clarke's "A Recalculation of the Atomic Weights," reproduced in the first (1883) edition of "Landolt-Bornstein." These 1882 values (to two decimals) are given in parentheses. A date in parentheses indicates the first appearance of the element in the atomic weight table. All the values given are based upon O = 16.000.

Symbol	Atomic number	Name	I. C. T. at. wt.	Atomic weights (1925-1882)
A	18	Argon	39 91	'25, 39.91, '24-'19, 39 9; '18-'11, 39.88; '10-'03, 39.9; '02, 39.96 (1902)
Ac	89	Actinium	?	
Ag	47	Silver	107 880	'25, 107.880; '24-'09, 107.88; '08-'03, 107.93; '02-'94, 107.92 (107.92)
Al	13	Aluminium	26 96	'25, 26.97; '24 '22, 27 0; '21-'00, 27.1; '99-'96, 27.11; '95-'94, 27 (27.08)
As	33	Arsenic	74 96	'25-'10, 74.96; '09-'00, 75 0; '99-'97, 75.01; '96, 75.09; '95-'94, 75 0 (75.09)
Au	79	Gold	197 2	'25-'00, 197.2; '99 '97, 197.23; '96, 197.24; '95 '94, 197 3 (196.61)
B	5	Boron	10 82	'25, 10 82; '24-'19, 10 9; '18-'00, 11 0; '99 '96, 10.95; '95-'94, 11 (10.97)
Ba	56	Barium	137 37	'25-'09, 137 37; '08 '00, 137.10; '99-'94, 137.43 (137.01)
Be	4	Beryllium	9 02	'25, 9.02; '24-'00, 9.1; '99-'96, 9.08; '95-'94, 9 (9.11)
Bi	83	Bismuth	209 00	'25-'22, 209.0; '21-'07. 208.0; '06-'03, 208.5; '02-'00, 208.1; '99-'96, 208.11; '95, 208; '94, 208.9 (208.00)
Br	35	Bromine	79 916	'25, 79.916; '24-'09, 79.92; '08-'03, 79.96; '02-'94, 79.95 (79.95)
C	6	Carbon	12.000	'25, 12.000; '24-'16, 12.005; '15-'98, 12 00; '97-'96, 12.01; '95-'94, 12 (12.00)

Symbol	Atomic number	Name	I. C. T. at. wt.	Atomic weights (1925-1882)
Ca	20	Calcium	40 07	'25 '12, 40.07; '11-'09, 40.09; '08 '00, 40.1; '99-'97, 40.07; '96, 40.08; '95 '94, 40 (40.08)
Cb	41	Columbium	93 1	'25 '17, 93.1; '16-'09, 93.5; '08-'03, 94; '02-'00, 93.7; '99-'97, 93.73; '96 '94, 94.0 (94.03)
Cd	48	Cadmium	112 41	'25, 112.41; '24-'09, 112.40; '08-'00, 112.4; '99, 112.38; '98-'97, 111.95; '96, 111.93; '95 '94, 112 (112.09)
Ce	58	Cerium	140 25	'25-'04, 140.25; '03, 140; '02 '00, 139; '99-'98, 139.35; '97-'94, 140.25 (140.75)
Cl	17	Chlorine	35 458	'25, 35.457; '24-'09, 35.46; '08-'94, 35.45 (35.45)
Co	27	Cobalt	58 97	'25, 58.94; '24-'09, 58.97; '08-'00, 59.0; '99-'98, 58.99; '97, 58.93; '96, 58.95; '95, 59.5; '94, 59 (59 02)
Cp	71	Cassiopeium	175 0	See Lu
Cr	24	Chromium	52 01	'25, 52.01; '24-'10, 52.0; '09-'00, 52.1; '99-'96, 52.14; '95-'94, 52.1 (52.13)
Cs	55	Cesium	132 81	'25-'09, 132.81; '08-'04, 132.9; '03, 133.0; '02-'00, 132.9; '00-'96, 132.89; '95-'94, 132.9 (132.92)
Ct	72	Celtium		Same as Hf
Cu	29	Copper	63 57	'25 '09, 63.57; '08-'94, 63.6 (63.32)
Ds / Dy	66	Dysprosium	162 52	'25, 162.52; '24-'08, 162.5 (1908)
Em	86	Ra-emanation	222	See Rn
Er	68	Erbium	167 7	'25-'12, 167.7; '11-'09, 167.4; '08-'00, 166.0; '99 '97, 166.32; '96-'94, 166.3 (166.27)
Eu	63	Europium	152 0	'25-'07, 152.0 (1907)
F	9	Fluorine	19 00	'25 '03, 19.0; '02-'00, 19.05; '99-'97, 19.06; '96, 19.03; '95-'94, 19 (19.03)
Fe	26	Iron	55 84	'25-'12, 55.84; '11-'09, 55.85; '08-'01, 55.9; '00, 56.0; '99-'96, 56.02; '95-'94, 56 (56.04)
Ga	31	Gallium	69 72	'25, 69.72; '24-'19, 70.1; '18-'09, 69.9; '08-'00, 70.0; '99-'97, 69.91; '96-'94, 69.0 (68.96)
Gd	64	Gadolinium	157 26	'25, 157.26; '24-'09, 157.3; '08-'03, 156; '02, 156.4; '01-'00, 157.0; '99-'97, 156.76; '96-'94, 156.1

Symbol	Atomic number	Name	I. C. T. at. wt.	Atomic weights (1925–1882)
Ge	32	Germanium	72 38	'25, 72.60; '24 '00, 72.5; '99 '97, 72.48; '96–'94, 72 3
Gl	4	Glucinium	9 02	See Be
H	1	Hydrogen	1 0077	'25, 1.0077; '24 '94, 1.008 (1 00)
He	2	Helium	4 00	'25–'16, 4 00; '15–'11, 3.99; '10–'03, 4.0; '02, 3.96 (1902)
Hf	72	Hafnium	178 6	
Hg	80	Mercury	200 61	'25, 200.61; '23–'12, 200.6; '11–'94, 200.0 (200 17)
Ho	67	Holmium	163 4	'25, 163.4; '23–'13, 163.5 (1913)
I (J)	53	Iodine	126 932	'25, 126.932; '24–'09, 126.92; '08–'05, 126.97; '04–'94, 126.85 (126.85)
In	49	Indium	114 8	'25–'09, 114.8; '08–'05, 115; '04–'00, 114; '99–'97, 113.85; '96–'91, 113.7 (113.66)
Ir	77	Iridium	193 1	'25–'09, 193.1; '08–'03, 193.0; '02–'00, 193.1; '99–'96, 193.12; '95–'94, 193.1 (193.09)
K	19	Potassium	39 095	'25, 39.096; '24–'09, 39.10; '08–'03, 39.15; '02–'94, 39.11 (39.11)
Kr	36	Krypton	82 9	'25, 82.9; '24–'11, 82.92; '10, 83.0; '09–'03, 81.8; '02, 81.76 (1902)
La	57	Lanthanum	138 91	'25, 138.90; '24–'09, 139.0; '08–'03, 138.9; '02–'00, 138.6; '99–'97, 138.64; '96, 138.6; '95–'94, 138.2 (138.84)
Li	3	Lithium	6 939	'25, 6.940; '24–'11, 6.94; '10–'09, 7.00; '08–'96, 7.03; '95–'94, 7.02 (7.02)
Lu	71	Lutecium	175 0	'25 '16, 175.0; '15–'09, 174.0 (1909)
Ma	43	Masurium		
Mg	12	Magnesium	24 32	'25–'09, 24.32; '08–'03, 24.36; '02–'00, 24.3; '99–'97, 24.28; '96, 24 29; '95–'94, 24.3 (24.01)
Mn	25	Manganese	54 93	'25–'09, 54.93; '08–'00, 55.0; '99–'96, 54.99; '95–'94, 55 (54.03)
Mo	42	Molybdenum	96 0	'25–'00, 96.0; '99–'97, 95.99; '96, 95.98; '95–'94, 96 (95.75)
N	7	Nitrogen	14 008	'25–'19, 14.008; '18–'07, 14.01; '06–'96, 14.04; '95, 14.05; '94, 14.03 (14.03)
Na	11	Sodium	22 997	'25, 22.997; '24–'09, 23.00; '08–'94, 23.05 (23.05)
Nb	41	Niobium	93 1	See Cb
Nd	60	Neodymium	144.27	'25, 144.27; '24–'0 144.3; '08–'99, 143. '98–'97, 140.80; '96–'9 140.5
Ne	10	Neon	20.2	'25–'09, 20.2; '10–'0 20.0 (1904)
Ni	28	Nickel	58.69	'25, 58.69; '24–'09, 58.6 '08–'00, 58.7; '99–'9 58.69; '95–'94, 58. (58.06)
Nt	86	Niton	222.	See Rn
O	8	Oxygen	16 000	'25–'94, 16.000 (16.00)
Os	76	Osmium	190 8	'25, 190.8; '23–'09, 190.9 '08–'00, 191.0; '99–'96 190.99; '95–'94, 190. (198.95?)
P	15	Phosphorus	31 024	'25, 31.027; '24–'11 31.04; '10–'00, 31.0 '99–'94, 31.02; '95–'9 31 (31.03)
Pa	91	Protoactinium	?	
Pb	82	Lead	207.20	'25–'16, 207.20; '15–'09 207.10; '08–'03, 206.9 '02–'96, 206.92; '95–'94 206.95 (206.95)
Pd	46	Palladium	106.7	'25–'09, 106.7; '08–'03 106.5; '02–'00, 107.0 '99–'96, 106.36; '95 106.5; '94, 106.6 (105.98
Po	84	Polonium	(210)	
Pr	59	Praseodymium	140 92	'25, 140.92; '24–'16 140.9; '15–'09, 140.6 '08–'00, 140.5; '99–'97 143.60; '96–'94, 143.5
Pt	78	Platinum	195.23	'25, 195.23; '24–'11 195.2; '10–'09, 195.0 '08–'03, 194.8; '02–'00 194.9; '99–'96, 194.80 '95–'94, 195 (194.87)
Ra	88	Radium	225.95	'25, 225.95; '24–'16, 226 '15–'09, 226.4; '08–'03 225 (1903)
Rb	37	Rubidium	85 44	'25, 85.44; '24–'09, 85.45; '08–'05, 85.5; '04–'00, 85.4; '99–'96, 85.43 '95–'94, 85.5 (85.53)
Re	75	Rhenium		
Rh	45	Rhodium	102 91	'25, 102.91; '24–'09, 102.9; '08–'00, 103.0; '99–'96, 103.01; '95–'94, 103 (104.29)
Rn	86	Radon	222	'25, 222; '24–'12, 222.4 (1912)
Ru	44	Ruthenium	101 7	'25–'00, 101.7; '99–'96, 101.68; '95–'94, 101.6 (104.46?)
S	16	Sulfur	32.065	'25, 32.065; '24–'16, 32.06; '15–'09, 32.07; '08–'03, 32.06; '02–'96, 32.07; '95–'94, 32.06 (32.06)
Sa	62	Samarium	150.43	'25, 150.43; '24–'09, 150.4; '08–'05, 150.3;

Symbol	Atomic number	Name	I. C. T. at. wt.	Atomic weights (1925-1882)
Sa	62	Samarium	150.43	'04–'03, 150; '02–'00, 150.3; '99–'97, 150.26; '96–'94, 150.0
Sb	51	Antimony	121.77	'25, 121.77; '24–'03, 120.2; '02–'00, 120.4; '99–'96, 120.43; '95–'94, 120 (120.23)
Sc	21	Scandium	45 10	'25–'21, 45.10; '20–'00, 44.1; '99–'97, 44.12; '96–'94, 44.0 (44.08)
Se	34	Selenium	79 2	'25–'00, 79.2; '99, 79.17; '98–'97, 79.02; '96–'94, 79.0 (78.98)
Si	14	Silicon	28 06	'25, 28.06; '24–'22, 28 1; '21–'09, 28.3; '08–'94, 28.4 (28.26)
Sm	62	Samarium	150 43	See Sa
Sn	50	Tin	118 70	'25–'16, 118.70; '15–'00, 119.0; '99–'96, 119.05; '95–'94, 119 (117.97)
Sr	38	Strontium	87 62	'25–'11, 87.63; '10–'09, 87.62; '08–'00, 87.6; '99–'96, 87.61; '95, 87.66; '94, 87.6 (87.58)
Ta	73	Tantalum	181.5	'25–'10, 181.5; '11–'07, 181.0; '06–03, 183; '02–'00, 182.8; '99–'97, 182.84; '96–'94, 182.6 (182 56)
Tb	65	Terbium	159 2	'25–'07, 159.2; '06–'94, 160
Te	52	Tellurium	127 5	'25–'09, 127.5; '08–'03, 127.6; '02, 127.7; '01–'00, 127.5; '99–'97, 127.49; '96, 127; '95–'94, 125 (128.252)
Th	90	Thorium	232 15	'25–'19, 232.15; '18–'11, 232.4; '10–'09, 232.42; '08–'03, 232.5; '02–'00, 232.6; '99–'96, 232.63; '95–'94, 232.6 (233.95)
Ti	22	Titanium	47.9	'25–'03, 48.1; '02–'96, 48.15; '95–'94, 48 (49.96?)
Tl	81	Thallium	204.4	'25, 204.39; '24–'09, 204.0; '08–'03, 204.1; '02–'96, 204.15; '95–'94, 204.18 (204.18)
Tm / Tu	69	Thulium	169.4	'25, 169.4; '24–'22, 169.9; '21–'09, 168.5; '08–'03, 171; '02–'94, 170.7
U	92	Uranium	238.17	'25, 238.17; '24–'16, 238.2; '15–'03, 238.5; '02–'00, 239.6; '99–'96, 239.59; '95–'94, 239.6 (239.03)
X₂	91	Uranium-X₂	(234)	Isotope of Pa
V	23	Vanadium	50.96	'25, 50.96; '24–'12, 51.0; '11, 51.06; '10–'03, 51.2; '02–'00, 51.4; '99–'96, 51.38; '95–'94, 51.4 (51.37)

Symbol	Atomic number	Name	I. C. T. at. wt.	Atomic weights (1925-1882)
W	74	Tungsten	184 0	'25–'00, 184.0; '99–'97, 184.83; '96, 184.84; '95, 184.9; '94, 184 (184.03)
Xe	54	Xenon	130 2	'25–'11, 130.2; '10, 130.7; '09–'02, 128 (1902)
Y / Yt	39	Yttrium	89.0	'25, 88.9; '24–'19, 89.33; '18–'16, 88.7; '15–'00, 89.0; '99–'97, 89.02; '96, 88.95; '95–'94, 89.1 (90.027)
Yb	70	Ytterbium	173 6	'25, 173.6; '24–'16, 173.5; '15–'00, 172.0; '08–'03, 173; '02–'00, 173.2; '99–'97, 173.19; '96–'94, 173.0 (173.16)
Zn	30	Zinc	65 38	'25, 65.38; '24–'10, 65.37; '00, 65.7; '08–'00, 65.4; '99–'96, 65.41; '95–'94, 65.3 (65.05)
Zr	40	Zirconium	91	'25, 91; '24–'09, 90.6; '01–'97, 90.4; '96–'94, 90.6 (89 57)

TABLE OF ISOTOPES

F. W. ASTON

Element	Atomic number	I. C. T. atomic weight	Minimum number of isotopes	Mass numbers in order of the intensities of the mass-spectrum lines	Lit.
A	18	39.91	2	40, 36	(3, 5, 21)
Ag	47	107.880	2	107, 109	(15, 26)
Al	13	26.96	1	27	(10)
As	33	74.96	1	75	(4, 22)
B	5	10.82	2	11, 10	(4, 22)
Ba	56	137.37	1	138, 136	(17, 18)
Be	4	9 02	1	9	(33)
Bi	83	209.00	1	209	(19)
Br	35	79 916	2	79, 81	(4, 22)
C	6	12.000	1	12	(2, 21)
Ca	20	40.07	2	40, 44	(31, 32)
Cd	48	112.41	6	110, 111, 112, 113, 114, 116	(19)
Ce	58	140 25	2	140, 142	(18)
Cl	17	35 458	2	35, 37	(2, 21, 23)
Co	27	58 97	1	59	(15, 26)
Cr	24	52 01	1	52	(15, 26)
Cs	55	132 81	1	133	(6, 24)
Cu	29	63.57	2	63, 65	(14, 26)
F	9	19.00	1	19	(4, 22)
Fe	26	55.84	2	56, 54	(9, 17)
Ga	31	69.72	2	69, 71	(15, 26)
Ge	32	72 38	3	74, 72, 70	(13, 26)
Gl	4	9.02	1	9	(33)
H	1	1.0077	1	1	(3, 21)
He	2	4.00	1	4	(3, 21)
Hg	80	200.61	2,6	197–200, 202, 204	(2, 3, 31)
I	53	126.932	1	127	(5, 23)
In	49	114.8	1	115	(16)
K	19	39.095	2	39, 41	(6, 24)
Kr	36	82.9	6	84, 86, 82, 83, 80, 78	(3, 21)
La	57	138.91	1	139	(17)

PERIODIC CHART OF THE ELEMENTS WITH ATOMIC NUMBERS AND ATOMIC WEIGHTS

I	II	III	IV	V	VI	VII	VIII or 0
H (1) 1.0077							He (2) 4.00
Li (3) 6.939	Be (4) 9.02	B (5) 10.82	C (6) 12.00	N (7) 14.008	O (8) 16.000	F (9) 19.00	Ne (10) 20.2
Na (11) 22.997	Mg (12) 24.32	Al (13) 26.96	Si (14) 28.06	P (15) 31.024	S (16) 32.065	Cl (17) 35.458	A (18) 39.91
K (19) 39.095	Ca (20) 40.07	Sc (21) 45.10	Ti (22) 47.9	V (23) 50.96	Cr (24) 52.01	Mn (25) 54.93	(26)
Cu (29) 63.57	Zn (30) 65.38	Ga (31) 69.72	Ge (32) 72.38	As (33) 74.96	Se (34) 79.2	Br (35) 79.916	Kr (36) 82.9
Rb (37) 85.44	Sr (38) 87.62	Yt (39) 89.0	Zr (40) 91	Cb (41) 93.1	Mo (42) 96.0	Ma (43)	(44)
Ag (47) 107.880	Cd (48) 112.41	In (49) 114.8	Sn (50) 118.70	Sb (51) 121.77	Te (52) 127.5	I (53) 126.932	Xe (54) 130.2
Cs (55) 132.81	Ba (56) 137.37	La* (57) 138.91	Hf (72) (178.6)	Ta (73) 181.5	W (74) 184.0	Re (75)	(76)
Au (79) 197.2	Hg (80) 200.61	Tl (81) 204.4	Pb (82) 207.20	Bi (83) 209.00	Po (84) (210)	(85)	Rn (86) 222
	Ra (88) 225.95	Ac (89)	Th (90) 232.15	Pa (91)	U (92) 238.17		

Group VIII:
Fe (26) 55.84 — Co (27) 58.97 — (Ni) (28) 58.69
Ru (44) 101.7 — Rh (45) 102.91 — (46) 106.7
Os (76) 190.8 — Ir (77) 193.1 — (78) 195.23

Rare earths (* Indicates rare earths. See above):

*	La (57) 138.91	Ce (58) 140.25	(59) 140.92
	Nd (60) 144.27	Sa (61) 150.43	Eu (62) 152.0
	Gd (64) 157.26	Tb (65) 159.2 — Dy (66) 162.52	(67) 163.4
	Er (68) 167.7	Tu (69) 169.4 — Yb (70) 173.6	Lu (71) 175.0

THE RADIOACTIVE ELEMENTS
FREDERICK SODDY

α – ray ←   → β – ray (or rayless)

Group	III	IV	V	VI	VII	VIII or 0	I	II	III	IV	V	VI
Principal element	Tl	Pb	Bi	Po	—	Rn	—	Ra	Ac	Th	Pa	U
Atomic number	81	82	83	84	85	86	87	88	89	90	91	92

U-Ra Series: Ra-B, Ra-A, Rn = Ra-Em (or Radon), Ra, Ra-C'', Ra-C, Ra-Ω, Ra-C', Ra-D, Ra-E, Ra-F, Ra-Ω', UX$_1$, UX$_2$, U$_1$, U$_2$, Io, Ms-Th$_1$, Th, Ms-Th$_2$

Th Series: Th-C'', Th-B, Th-C, Th-A, Th-Em (Thoron), Th-X, Ra-Th, Th-Ω, Th-Ω', UY, Pa, U$_1$ or U$_2$, Ac, Ra-Ac

Ac Series: Ac-C'', Ac-B, Ac-C, Ac-A, Ac-Em (Actinon), Ac-X, Ac-C', Ac-Ω'', Ac-Ω'

TABLE OF ISOTOPES.—*Continued*

Element	Atomic number	I. C. T. atomic weight	Minimum number of isotopes	Mass numbers in order of the intensities of the mass-spectrum lines	Lit
Li	3	6.939	2	7, 6	(24, 27, 29, 30)
Mg	12	24.32	3	24, 25, 26	(28, 30)
Mn	25	54.93	1	55	(15, 26)
N	7	14.008	1	14	(3, 21)
Na	11	22.997	1	23	(6, 24)
Nd	60	144.27	3	142, 144, 146, 145	(17, 18)
Ne	10	20.2	2	20, 22	(1, 20, 21)
Ni	28	58.69	2	58, 60	(7)
O	8	16.000	1	16	(2, 21)
P	15	31.024	1	31	(4, 22)
Pr	59	140.92	1	141	(17)
Rb	37	85.44	2	85, 87	(6, 24)
S	16	32.065	1	32	(4, 22)
Sb	51	121.77	2	121, 123	(11, 25)
Sc	21	45.10	1	45	(15, 26)
Se	34	79.2	6	80, 78, 76, 82, 77, 74	(10)
Si	14	28.06	3	28, 29, 30	(4, 18, 22)
Sn	50	118.70	7,8	120, 118, 116, 124, 119, 117, 122, 121	(8)
Sr	38	87.62	2	88, 86	(15, 17, 26)
Te	52	127.5	3	128, 130, 126	(19)
Ti	22	47.9	1	48	(15, 26)
V	23	50.96	1	51	(15, 26)
Xe	54	130.2	7,9	129, 132, 131, 134, 136, 128, 130, 126, 124	(3, 5, 10, 21, 23)
Yt	39	89.0	1	89	(15, 26)
Zn	30	65.38	4	64, 66, 68, 70	(31)
Zr	40	91	3	90, 94, 92	(18)

LITERATURE

(For a key to the periodicals see end of volume)

¹) Aston, *58*, **104**: 334, 19 (²) *Ibid*, **104**: 393, 19. (³) *Ibid*, **105**: 8; 20 . (⁴) *Ibid*, **105**: 547; 20. (⁵) *Ibid*, **106**: 468, 20. (⁶) *Ibid.*, **107**: 72, 21 (⁷) *Ibid*, **107**: 520; 21. (⁸) *Ibid*, **109**: 843; 22. (⁹) *Ibid*, **110**: 312, 22. ⁸) Aston, *58*, **110**: 664; 22. (¹¹) *Ibid*, **110**: 732; 22. (¹²) *Ibid*, **111**: 739, 23 (¹³) *Ibid*, **111**: 771, 23. (¹⁴) *Ibid*, **112**: 162; 23. (¹⁵) *Ibid*, **112**: 449, 23 (¹⁶) *Ibid*, **113**: 192; 24. (¹⁷) *Ibid*, **113**: 856; 24. (¹⁸) *Ibid.*, **114**: 273, 24 (¹⁹) *Ibid*, **114**: 717, 24.
¹) Aston, *3*, **39**: 449, 20. (²¹) *Ibid.*, **39**: 611; 20. (²²) *Ibid*, **40**: 628, 20 (²³) *Ibid*, **42**: 140; 21. (²⁴) *Ibid*, **42**: 436; 21. (²⁵) *Ibid*, **45**: 924, 23 (²⁶) *Ibid.*, **47**: 385; 24. (²⁷) Aston and Thomson, *58*, **106**: 827; 21 (²⁸) Dempster, *166*, **52**: 559, 20 (²⁹) Dempster, *166*, **53**: 363, 21
¹⁰) Dempster, *2*, **18**: 415, 21 (³¹) *Ibid*, **19**: 431, 22. (³²) *Ibid.*, **20**: 631, 22. (³³) Thomson, *3*, **42**: 837, 21.

THE STRUCTURE OF THE ISOLATED ATOM

(Symbols, p. 50)

H. A. KRAMERS

According to the fundamental postulates of Bohr's atomic ~ory, a series of discrete "stationary states" has to be correlated ;h each atom. A definite "energy-content" can be assigned to *ery state, and an atom in a given state can change its energy only y performing a process of "transition" to another state. The nission of a spectral line of frequency ν is correlated with a
ontaneous transition from a stationary state of energy content ι to another of energy content E_2 by equation (1)

$$\nu = \frac{1}{h}(E_1 - E_2) \tag{1}$$

The stationary state with the smallest energy is termed the "normal state" of the atom. The properties of the stationary states can, to a considerable extent, be accounted for by assuming that the electrons surrounding the nucleus have definite motions, characterized by integral values of certain quantities. These integers are called the "quantum numbers" of the stationary state in question; by their values the energy of the state is completely fixed. For general treatment of the subject, see (¹, ³, ⁴, 10, 11, 18).

Of special interest are the recent attempts (²¹) to develop a rational "quantum mechanics" of the atom. This work clearly demonstrates the limited applicability of a picture of atomic structure, in which the behavior of the electrons inside the atom is visualized by orbits possessing definite kinematical properties.

Atoms Containing One Electron. — Only for atoms containing a single electron, can a fairly complete description of the electronic motion in the stationary state, and of the significance of the quantum numbers be given. The motion of the electron obeys quite approximately the laws of electrodynamics, and can be described as a Keplerian elliptic motion, with the centre of gravity of the nucleus and the electron in one focus. On this motion, a slow uniform precession in the plane of motion is superposed (effect of variability of mass or "relativity-effect"). Two quantum numbers (n, k) define the stationary states (n, k = 1, 2, 3 ; k ≤ n), k/n being the ratio of the minor to the major axis of the ellipse. The states are denoted by the symbol n_k.

In the **normal state**, $1_1(n = k = 1)$, the orbit is circular; and, omitting the correction due to the relativity effect, its constants are given by equations (2)

$$a_1 = \frac{1}{Z} \cdot \frac{h^2}{4\pi^2 e^2 m_0} = \frac{1}{Z} \cdot \frac{0.53}{Z} \times 10^{-8} \text{ cm}$$

$$\omega_1 = \frac{Z^2}{1 + \frac{m_0}{M}} \times \frac{4\pi^2 e^4 m_0}{h^3} = \frac{2\nu_\infty Z^2}{1 + \frac{m_0}{M}} = \frac{6.6 Z^2}{1 + \frac{m_0}{M}} \times 10^{15} \text{ sec}^{-1} \tag{2}$$

$$W_1 = \frac{Z^2}{1 + \frac{m_0}{M}} \times \frac{2\pi^2 e^4 m_0}{h^2} = \frac{Z^2 \nu_\infty h}{1 + \frac{m_0}{M}} = \frac{2.15 Z^2}{1 + \frac{m_0}{M}} \times 10^{-11} \text{ erg.}$$

In **higher quantum states**, the orbital constants are, with the same approximation, given by (3, 4):

$$a_n = n^2 a_1 = \frac{n^2}{Z} r_1$$

$$\omega_n = \frac{\omega_1}{n^3} = \frac{2Z^2 \nu_\infty}{n^3 \left(1 + \frac{m_0}{M}\right)} \tag{3}$$

$$W_n = \frac{W_1}{n^2} = \frac{Z^2 \nu_\infty h}{n^2 \left(1 + \frac{m_0}{M}\right)}$$

$$b_{n,k} = nka_1 = \frac{nkr_1}{Z}; \quad p_k = k^2 a_1 = \frac{k^2 r_1}{Z} \tag{4}$$

The number of revolutions corresponding to one rotation of the major axis, is, to a first approximation, given by (5):

$$\frac{\omega_n}{\sigma_{n,k}} = \frac{k^2}{Z^2} \times \frac{2}{\alpha^2} = \frac{k^2}{Z^2} \times 37,700 \tag{5}$$

$$\left(\alpha = \frac{2\pi e^2}{hc} = 7.30 \times 10^{-3} \cong \frac{1}{137}; \alpha^2 = 5.31 \times 10^{-5}\right)$$

The exact energy formula, neglecting terms containing m_0/M, is given by (6):

$$W_{n,k} = m_0 c^2 \left[\left\{1 + \left(\frac{\alpha Z}{n - k + \sqrt{k^2 - \alpha^2 Z^2}}\right)^2\right\}^{-\frac{1}{2}} - 1\right] \tag{6}$$

$$= \frac{Z^2}{n^2} \times \frac{2\pi^2 e^4 m_0}{h^2} \left\{1 + \alpha^2 Z^2 \left(\frac{1}{kn} - \frac{3}{4n^2}\right) + \cdots\right\}$$

(For general formula for W, including terms in m_0/M, see (⁹).) Figure 1 illustrates the stationary states in the hydrogen atom for which n = 1, 2, 3, 4. The arrows indicate the transitions giving

rise to the fine-structure components of the spectral lines, H_α and H_β. The numerical constants for these states are given in Table 1.

TABLE 1.—HYDROGEN ORBITS; $r_1 = 5.286 \times 10^{-9}$ cm [11]

n_k	a/r_1	b/r_1	p/r_1	$\omega \times 10^{-16}$	$\sigma \times 10^{8}$	ω/σ
1₁	1	1	1	65 78	1746	37 700
2₁	4	2	1	8 222	218 3	37 700
2₂	4	4	4	8 222	54 57	150 700
3₁	9	3	1	2 436	64 68	37 700
3₂	9	6	4	2 436	16 17	150 700
3₃	9	9	9	2 436	7.187	339 300
4₁	16	4	1	1 029	27 29	37 700
4₂	16	8	4	1 029	6 822	150 800
4₃	16	12	9	1 029	3 032	339 300
4₄	16	16	16	1.029	1.705	603 200

Fig 1 —Orbits in hydrogen to $n = 4$. (Reproduced by permission from The Journal of the Franklin Institute.)

Atoms Containing More than One Electron.—A complete theory of stationary states is lacking. Many properties of these states can be accounted for, however, on the basis of the principles applied to atoms containing one electron. As a first approximation, each electron may be considered as moving in a central field of force due to the nucleus and the other electrons, its motion being characterized by a "principal quantum number" n and a "subordinate quantum number" k. The electronic orbit can be described as a plane periodic orbit on which a uniform precession in the plane is superposed ("central orbit" cf. Fig. 2).

If the position of the electron in the orbital plane is defined by polar coordinate (r, φ), the quantum numbers are defined by Sommerfeld's quantum conditions (7)

$$k = \frac{2\pi m_0\beta r^2 \, d\phi}{h \quad dt} = \frac{2\pi P}{h} \qquad (n - k) = \frac{1}{h} \oint m_0\beta\left(\frac{dr}{dt}\right)^2 dt \quad (7)$$

where the factor β becomes equal to 1 if the relativity effect is neglected. P is equal to the angular momentum of the electron with respect to the nucleus; the integral has to be taken over a complete period of the radial motion, from A to B (Fig. 2).

In the normal state the electrons are distributed in groups, each of which is characterized by its quantum numbers (n, k). On passing from the nucleus to the surface of the atom, the successive groups correspond to successive integral values of the main quantum number n ("n-quantum group"), the innermost group being characterized by $n = 1$; each group is divided into subgroups corresponding to the different values which k may take. The possibility of reconciling such a picture with the dynamical properties of quantized central orbits is closely connected with the fact that in an orbit for which $k < n$ the electron will, in each revolution, dive into and leave again all regions occupied by

electronic orbits for which the principal quantum number is smaller than n but equal to or greater than k (conception of "penetrating orbits").

The maximum number of electrons which an n-quantum group can contain is equal to $2n^2$. If it contains this number, it contains sub-groups corresponding to all possible values for k ($k = 1$, $2 \ldots n$), and it is said to be a "**finally completed**" group. If a group, due to the dynamical properties of the atom under consideration, contains only sub-groups corresponding to $k = 1,$.

Fig. 2.—Central orbit.

$2 \ldots k_0$ ($k_0 < n$) it will be in a state which is termed "**provisionally completed**," if it contains $2k_0^2$ electrons. For example, the 4-quantum group has reached the state of a 2-group ($k_0 = 1$) in Ca (20), the state of an 8-group or 8-shell ($k_0 = 2$) in Kr (36), the state of an 18-group or 18-shell ($k_0 = 3$) in Ag (47), and its final state of a completed 32-group or 32-shell ($k_0 = 4$) in Lu (71). With the exception of the 2-groups it seems impossible to assign definite values to the number of electrons in the several sub-groups of a provisionally, or finally, completed group; in fact, the actual properties of the electronic groups seem to show that the simple conception of central orbits characterized by the symbol n_k is essentially insufficient for their description. (Originally Bohr assumed that a group of $2k_0^2$ electrons contained $2k_0$ electrons in each sub-group.) Closely connected herewith is the impossibility of assigning definite spatial arrangements to the orbits belonging to one and the same group. In Table 2 the number of electrons in each group is given as far as the theory allows of a definite statement; those in parentheses are uncertain.

From calculations based on Sommerfeld's quantum conditions and certain simplifying assumptions, a rough estimate of the dimensions of the different types of orbits may be made. Such estimates for neutral atoms and for positive ions containing only finally, or provisionally, completed groups are schematically represented in Fig. 3. The small vertical lines are so drawn that their distances from the dot at the left are proportional to the radius of the sphere inside which the electrons belonging to the respective groups are moving. The symbols g $(n_1, 2 \ldots k_0)$ means that the corresponding groups contain g electronic orbits of principal quantum number n, and of subordinate quantum numbers from 1 to k_0.

For the calculation of the dimensions of the outermost groups it has been necessary to consider also experimental data relative to the effective gas-kinetic radii of the atoms of the inert gases, the effective radii of ions in crystals, ionic refraction, etc. As a rule the effective radii are 1.5 to 2 5 times larger than the orbital dimensions. As regards the inner groups, the estimate is rather accurate; for the outer groups, errors of the order of 10% might be expected Special mention must be made of the uncertainty in the radius of the 5-quantum group for elements heavier than barium; the radii of this group as given in Fig 3 for the elements (72), 79, 80, 81, 82 are perhaps some 10% too high, as compared with radii of the homologous elements 47, 48, 49, 50.

For atoms containing only one electron in the outermost group, the dimensions of the orbit of this electron, and its frequency of revolution can with considerable accuracy be derived from the

TABLE 2

	1_1	$2_1\;2_2$	$3_1\;3_2\;3_3$	$4_1\;4_2\;4_3\;4_4$	$5_1\;5_2\;5_3\;5_4\;5_5$	$6_1\;6_2\;6_3\;6_4\;6_5\;6_6$	$7_1\;7_2$
1 H	1						
2 He	2						
3 Li	2	1					
4 Be	2	2					
5 B	2	2 1					
6 C	2	2 (2)					
10 Ne	2	8					
11 Na	2	8	1				
12 Mg	2	8	2				
13 Al	2	8	2 1				
14 Si	2	8	2 (2)				
18 A	2	8	8				
19 K	2	8	8	1			
20 Ca	2	8	8	2			
21 Sc	2	8	8 1	(2)			
22 Ti	2	8	8 2	(2)			
29 Cu	2	8	18	1			
30 Zn	2	8	18	2			
31 Ga	2	8	18	2 1			
36 Kr	2	8	18	8			
37 Rb	2	8	18	8	1		
38 Sr	2	8	18	8	2		
39 Y	2	8	18	8 1	(2)		
40 Zr	2	8	18	8 2	(2)		
47 Ag	2	8	18	18	1		
48 Cd	2	8	18	18	2		
49 In	2	8	18	18	2 1		
54 X	2	8	18	18	8		
55 Cs	2	8	18	18	8	1	
56 Ba	2	8	18	18	8	2	
57 La	2	8	18	18	8 1	(2)	
58 Ce	2	8	18	18 1	8 1	(2)	
59 Pr	2	8	18	18 2	8 1	(2)	
71 Lu	2	8	18	32	8 1	(2)	
72 Hf	2	8	18	32	8 2	(2)	
79 Au	2	8	18	32	18	1	
80 Hg	2	8	18	32	18	2	
81 Tl	2	8	18	32	18	2 1	
86 Rn	2	8	18	32	18	8	
87 —	2	8	18	32	18	8	1
88 Ra	2	8	18	32	18	8	2
89 Ac	2	8	18	32	18	8 1	(2)
90 Th	2	8	18	32	18	8 2	(2)
118 —	2	8	18	32	32	18	8

frequency of the lowest frequency term in the corresponding spectral series, provided we may adhere to the simple central orbit model. Figure 4 contains a schematic picture of the orbits of the outer electron in the normal state of neutral atoms of the alkali metals, and of Cu, Ag, Au. They are all penetrating orbits, since they correspond to $k = 1$. The regions inside which the electrons of the completed groups are moving are designated by circles. The atoms of the inert gases are added for the sake of comparison. The numbers at the left of the nucleus indicate the number of electrons contained in each group; the symbols $n_{1,2}$. at the right indicate the quantum numbers of the orbits contained in each group.

[For detailed calculations of electronic orbits, based on simplifying assumptions, see [12, 13, 20] (Cs and U); the work is semi-empirical. For detailed calculations on purely theoretical basis, see [15] (Ne, Na, Mg⁺, Al⁺⁺, Si⁺⁺⁺, P⁺⁺⁺⁺) and [16] (alkali metals); in Lindsay's work, the radii of outer groups in K⁺, Rb⁺, and Cs⁺ seem too large, probably on account of inadequacy of assumptions regarding numbers of electrons in sub-groups, as well as of the simplifying assumptions made. For critical review of work on effective atomic radii, see [14] and for recent work [8]. There is no simple direct connection between effective atomic radii and the magnitude of the space occupied by electronic orbits.]

In experiments on optical and X-ray spectra, we meet neutral atoms or atomic ions in **higher quantum states.** Several features of these states can be described on the simple central orbit model. In the case of "single excitation" all electronic orbits except one remain normal, and the other electron describes an orbit with quantum numbers which differ from those of the normal state. "Double excitation" corresponds to two electrons describing orbits different from those in the normal state, etc. We will here consider only singly-excited states.

In the stationary states (energy levels) involved in the emission of the ordinary X-ray spectra, one electron in the inner groups of the atom is lacking. In the states involved in the emission of the ordinary series-spectra, one electron belonging to the outermost group of the atom, the "series electron," moves in a central n_k orbit the dimensions of which are large as compared with those of the rest of the atom. It may move either quite outside the atomic residue or it may penetrate into it in each revolution.

As a first approximation, a non-penetrating orbit may be described as a Keplerian elliptical orbit performing a uniform precession in its plane, the shape of the ellipse being very nearly that of an n_k-orbit in an atom containing only one electron and having a nuclear charge Z^*e equal to the net-charge of the atomic residue. If the electron orbit is of the penetrating type, it may, as a first approximation, be described as a set of congruent outer Keplerian elliptical loops, connected by congruent inner loops, the angular distance between successive loops being the same. The semi-major axis, the semi-parameter p, and the semi-minor axis b of the outer loop can be found from the value of the corresponding spectral term (T) by means of the formulae

$$a = \frac{Z^* N r_1}{T} \qquad p = \frac{k^2}{Z^*} r_1 \qquad b = \sqrt{ap} \qquad (8)$$

where $N \left(= \frac{\nu_\infty}{c} \times \frac{1}{1 + m_0/M} \right)$ is the Rydberg constant for the element in question, and Z^*e is the net-charge of the atomic residue. If we introduce the effective quantum number n^* $(n^{*2} = Z^{*2}N/T)$, these formulae may be written:

$$a = \frac{n^{*2}}{Z^*} r_1 \qquad p = \frac{k^2}{Z^*} r_1 \qquad b = \frac{n^* k}{Z^*} r_1 \qquad (9)$$

The greater the ratio n^*/k (or a/b) the closer the approximation to which this description of the outer loops may be considered to hold. The maximum distance of the electron from the nucleus is equal to $a + \sqrt{a^2 - b^2}$, or very nearly equal to $2a - \frac{1}{2}p$.

Fig. 3 — Maximum elongations of electrons of several groups.

The values to be assigned to the precessional frequency characterizing the penetrating central orbits are very uncertain. For the alkali elements, the ratio ω/σ for the n_1 orbits probably lies between 0.3 and 0.5, for the n_2 orbits (except lithium) between 0.5 and 1.0. Based on the above formulae, an illustration of the shapes of the orbits of the series electron corresponding to the stationary states of the K-atom, is given in Fig. 5. [For connection between spectra and the group structure of atoms, see (6, 8); for spectra and central field of force, see (12, 13); for series spectra and electronic orbits, see (2, 7); for recent development of formal theory of electronic groups, see (17, 19)].

SYMBOLS

The symbols c, e, h, m_0, λ have their usual significance (see p. 16); others which occur more than once are:

a_n Semi-major axis of electronic orbit, state n.
$b_{n,k}$ Semi-minor axis of electronic orbit, state n, k.
k Subordinate, or azimuthal, quantum number defining a stationary state.
M Nuclear mass
n Principal quantum number defining a stationary state.

Fig. 5. — Orbits of the series electron of potassium. (*Reproduced by permission from The Journal of the Franklin Institute.*)

Fig. 4.—Normal orbit of outer electron.

n^* Effective quantum number $= Z^*N/T$.

n_k Designation of the state characterized by the numbers
             n, k.

N_a Rydberg constant.

p Semi-parameter of the electronic orbit (semi-latus rectum).

r_l Radius of first Bohr ring for hydrogen.

T Spectral term $= $ a wave number $(1/\lambda)$ of a spectral series.

v Speed of electron in its orbit

W_a Energy expenditure required to remove the electron to
             infinity.

Z Atomic number: $Ze = $ nuclear charge.

Z^*e Charge of atomic residue

α $2\pi e^2/hc$.

β $(1 - v^2/c^2)^{-1/2}$

ν Frequency of emitted radiation.

ν_∞ Rydberg fundamental frequency.

$\sigma_{n,k}$ Frequency of precession of electronic orbit.

ω_n Frequency of revolution of electron; for penetrating orbits,
             the radial frequency, one revolution being from A to
             B, Fig. 2.

LITERATURE

(For a key to the periodicals, see end of volume)

(1) Andrade, The Structure of the Atom, 1923. (2) Birge and Blackett, 48, 8: 213; 24 (3) Bohr, The Theory of Spectra and Atomic Constitution, 2nd ed., 1924 (4) Bohr, 58, 112: 29; 23. 218, 11: 606; 23. (5) Bohr, 8, 71: 228, 23 (6) Bohr and Coster, 96, 12: 342; 23. (6a) Born, Vorlesungen über Atommechanik. (7) Born and Heisenberg, 96, 23: 388; 24. (8) Davey, 2, 22: 211; 23. (9) Darwin, 5, 39: 537; 23. (10) Foote and Mohler, Origin of Spectra, 1923. (11) Foote, 145, 198: 344, 517; 24. (12) Fuess, 96, 11: 364; 22. 12: 1; 22. 21: 265; 24. 8, 76: 299; 25. (13) Hartree, 201, 21: 630; 23. 22: 409, 464; 24. 5, 106: 552; 24. (14) Herzfeld, 200, 19: 259; 23. (15) Kramers and Urey, O. (16) Lindsay, 2, 23: 552, 24. 285, 3: 191; 24. 2, 25: 239; 25. (17) Pauli, Jr., 96, 31: 765; 25. (18) Sommerfeld, Atombau und Spektrallinien, 4th ed., 1925. (19) Stoner, 3, 48: 719, 24

(20) Urey, O. (21) Heisenberg, 96, 33: 879, 25.

THERMOMETRY

E. F. MUELLER, L. H. ADAMS, F. O. FAIRCHILD AND H. T. WENSEL

1. THERMOMETRIC SCALES

E. F. MUELLER

Centigrade or Celsius scale, °C

Fahrenheit scale, °F

Réamur scale, °R

Centigrade absolute or Kelvin scale, °K

Fahrenheit absolute or Rankine scale, °R'

By definition or as basic values adopted for I. C. T., the ice
and steam points under a pressure of $1A_n$ have the following
values:

Ice point: $0°C = 32°F = 0°R = 273.1°K = 491.58°R'$.

Steam point: $100°C = 212°F = 80°R = 373.1°K = 703.58°R'$.

$°C = \frac{5}{9} (°F - 32) = \frac{5}{4}°R = °K - 273.1$.

$°F = \frac{9}{5}°C + 32 = °R' - 459.58$.

2. THE STANDARD THERMODYNAMIC SCALE

E. F. MUELLER

The thermodynamic scale, which is based solely on the laws
of thermodynamics and is independent of the properties of any
material substance, is accepted as the standard scale of tem-
perature. Temperatures on the thermodynamic scale are pro-
portional to the pressures (or to the volumes) of an ideal gas in a
perfect constant volume (or constant pressure) gas thermometer.
The standard scale is realized in practice by use of gas thermom-
eters, the indications of which can be reduced to the standard
scale, or for higher temperatures, by use of the relations between
the intensity of radiation from a black body and its temperature.

The experimental difficulties in the use of gas thermometers
and the relatively low precision attainable in a single measure-
ment have led to the introduction of a standard practical or work-
ing scale. This working scale is defined by certain base points,
the temperatures of which have been determined by gas ther-
mometer measurements, and by the indications of suitable
instruments used for interpolation between the base points or for
extrapolation to higher temperatures. It is possible in this way,
without actually using a gas thermometer, to establish a working
scale which does not differ to a demonstrable extent from the
standard scale at any temperature within the range of the working
scale. The practice of the various national standardizing labo-
ratories in defining the working scale is substantially uniform
at present, and it requires only minor adjustments and formal
agreement to give the working scales of these laboratories the
status of an international temperature scale. Such a scale would
bear essentially the same relation to the standard scale, as do the
international electric units to the absolute units.

The standard working scale may be defined by assigning nu-
merical values to the temperatures defined by the boiling point of
oxygen, the melting point of ice, the boiling point of water, the
boiling point of sulfur, and the freezing points of antimony,
silver and gold. The platinum resistance thermometer is the
standard for interpolation in the range $-195°$ to $0°C$ and from
$0°$ to $650°C$; the platinum-platinum rhodium thermocouple for
the range from $650°$ to $1063°$; and the luminous filament pyrom-
eter above $1063°C$.

Wien's law is accepted as expressing the brightness-tempera-
ture relation for a black body. For the purpose of defining the
temperature scale above $1063°C$ the present practice of the
national laboratories tends to favor the use of the value 1.430 cm
degrees for the constant C_2 in this equation but the value 1.433
cm degrees has been adopted for I. C. T.

LITERATURE

(For a key to the periodicals, see end of volume)

(1) Reichsanstalt, 8, 48: 1034; 15. (2) Griffiths and Schofield, 83, 12: 222; 18. (3) Waidner, Mueller and Foote, Pyrometry, p 46 (pub. by Am Soc. Min. and Met. Engrs., 1920). (4) Day and Sosman, Dictionary of Applied Physics, 1: 836; 22. (5) Henning, 245, 44: 349; 24. (6) Reichsanstalt, 245, 44: 517; 24.

Reduction of Gas Thermometer Indications to the Thermodynamic Scale

The temperature t_g on the scale of a constant volume or constant pressure gas thermometer filled with any real gas, is proportional to the pressure the gas would exert or the volume it would occupy, respectively, if all of the gas were at the uniform temperature to be measured, and if the volume or the pressure, respectively, were the same at all temperatures. At 0° and 100°C, the temperature t_g is by definition identical with the thermodynamic temperature t, while at other temperatures t_g departs from t by amounts which are proportional to the pressure at 0°, called the initial pressure. The tabular values are accordingly given only for an initial pressure equivalent to 1 m of mercury.

The values of $t - t_g$ obtained by various methods cover a wide range, so that only the order of magnitude of the values can be considered as known with any certainty. The tendency in modern work in gas thermometry has been to employ hydrogen or helium as the thermometric gas, and for these gases the magnitude of $t - t_g$ is comparable with the experimental error of the gas thermometer itself, so that the importance of an exact knowledge of the departure of the scales of these gas thermometers from the thermodynamic scale is correspondingly reduced.

REDUCTION OF GAS THERMOMETER INDICATIONS, t_g, TO THE THERMODYNAMIC CENTIGRADE SCALE, t

Values of $t - t_g$ for an initial pressure of 1 meter of mercury

t °C	Helium Const. vol.	Helium Const. press.	Hydrogen Const. vol.	Hydrogen Const. press.	Nitrogen Const. vol.	Nitrogen Const. press.
- 250	+0.04	..	+0 12		+0 5	
- 200	+ .02	+0 04	+ 06	+0 3	+ 2	
- 150	+ .01	+ 02	+ 03	+ .1	+ 06	+1 3
- 100	+ .005	+ .005	+ 015	+ .04	+ 03	+ 4
- 50	+ .002	+ .002	+ 005	+ 02	00	+ 12
0	000	.000	000	000	− .008	00
+ 25	− 001	− 001	− 001	− 003	− .010	− 02
50	− .001	− .000	− 002	− 004	− .005	− 03
75	− 001	+ 000	− 001	− 003	000	− 02
100	.000	000	000	000	+ .01	00
150	+ .002	+ 001	+ 01	+ 01	+ 02	+ 05
200	+ 006	+ .001	+ .02	+ 02	+ 03	+ 12
250	+ .01	+ .002	.	+ 03	+ .04	+ 2
300	+ .02	+ .003	..	+ 04	+ .07	+ 3
350	+ 03	+ .005			+ 10	+ 1
400	+ 04	+ 006	.		+ .14	+ 5
450	+ .05	+ .008			+ .17	+ 6
500	..				+ 2	+ 7
600					+ 3	+ 9
800	.	.			+ 5	+1 3
1000	. .			.	+ 7	+1 8
1200					+1 0	+2 3

LITERATURE

(For a key to the periodicals see end of volume)

[1]Rose-Innes, 5, 2: 131; 01. 16: 301, 08. (2)Callendar, 5, 5: 48; 03. (3)Berthelot, 238, 13B: 113p; 07. (4)Buckingham, 314, 3: 237, 07. (5)Cath and Onnes,168, No. 156a; 22. 18, 6: 1, 22. (6)Holborn and Otto, 96, 23: 77; 24. 30: 320, 24. (7)Keesom and Onnes, B60: 15; 24.

3. FIXED POINTS

E. F. MUELLER

t = Temperature on standard scale.

p = Pressure in millimeters of Hg (1 mm Hg = $\frac{1}{760}$ A$_B$) where p is between 680 and 780 mm.

BASE POINTS USED IN DEFINING THE STANDARD WORKING SCALE

(I. C. T. temperature scale)

Substance	Phenomenon	Temperature, °C
Liquid O$_2$	Vapor pressure	$t = \begin{bmatrix} -183.00 + 0.245\ (t + \\ 273\ 1)\log_{10} p/760 \text{ or} \\ -183\ 00 + 0.0126\ (p - 760) \\ -0\ 0000065\ (p - 760)^2 \end{bmatrix}$
Solid CO$_2$*	Vapor pressure	$t = \begin{bmatrix} -78\ 51 + 0\ 1443\ (t + \\ 273.1)\log_{10} p/760 \text{ or} \\ -78\ 51 + 0.01595\ (p - 760) \\ -0\ 000011\ (p - 760)^2 \end{bmatrix}$
Mercury*	Freezing	$t = -38\ 87°$
Ice	Melting	$t = 0\ 000°$
Steam	Condensing	$t = \begin{bmatrix} 100.000 + 0.1727\ (t + \\ 273\ 1)\log_{10} p/760 \text{ or} \\ 100.000 + 0.0367\ (p - \\ 760) -0\ 000023\ (p - 760)^2 \end{bmatrix}$
Sulfur	Condensing	$t = \begin{bmatrix} 444\ 60 + 0\ 2215\ (t + \\ 273\ 1)\log_{10} p/760 \text{ or} \\ 444\ 60 + 0\ 0909\ (p - 760) \\ -0\ 000048\ (p - 760)^2 \end{bmatrix}$
Antimony	Freezing	To be determined with resistance thermometer t = approx. 630.5°
Silver	Freezing	$t = 960\ 5°$ (reducing atmosphere).
Gold	Freezing	$t = 1063°$

* Not needed according to one suggested definition of the scale.

SECONDARY FIXED POINTS USEFUL IN CALIBRATING TEMPERATURE MEASURING INSTRUMENTS

(I. C. T. temperature scale)

Substance	Phenomenon	Temperature °C
Hydrogen	Boiling	$t = -252.7s + 0.0044\ (p - 760)$
Nitrogen	Vapor pressure	$t = -195\ 80 + 0\ 0109\ (p - 760)$
Naphthalene	Condensing	$t = 217\ 96 + 0.207s\ (t + 273.1)\log_{10} (p/760)$
Tin	Freezing	$t = 231\ 8s$
Benzophenone	Condensing	$t = 305.9 + 0\ 194\ (t + 273.1)\log_{10} (p/760)$
Cadmium	Freezing	$t = 320\ 9$
Lead	Freezing	$t = 327.4$
Zinc	Freezing	$t = 419.4s$
Aluminum (99.85 %)	Freezing	$t = 658.9$
Copper	Freezing	$t = 1083$ (reducing atmosphere)
Palladium	Freezing	$t = 1555 \pm 2$
Platinum	Melting	$t = 1755 \pm 6$
Tungsten	Melting	$t = 337s \pm 30$

The above values are in accord with the temperature scale used throughout I C T For the last three points the following slightly different values have been suggested for future adoption as secondary points on an international practical scale.

Palladium	Freezing	$t = \begin{bmatrix} 1555 \text{ for } C_2 = 1\ 430 \\ 1554 \text{ for } C_2 = 1\ 433 \end{bmatrix}$
Platinum	Melting	$t = \begin{bmatrix} 1765 \text{ for } C_2 = 1\ 430 \\ 1763 \text{ for } C_2 = 1\ 433 \end{bmatrix}$
Tungsten	Melting	$t = \begin{bmatrix} 3400 \text{ for } C_2 = 1\ 430 \\ 3386 \text{ for } C_2 = 1\ 433 \end{bmatrix}$

ADDITIONAL USEFUL SECONDARY POINTS

Substance	Formula	Phenomenon	Temperature, °C
Isopentane	C_5H_{12}	Freezing	− 159 6
Methylcyclohexane	$C_6H_{11}CH_3$	Freezing	− 126 3
Ether	$(C_2H_5)_2O$	Slow freezing (unstable)	− 123 3
Ether	$(C_2H_5)_2O$	Rapid freezing or slow melting	− 116 3
Carbon disulfide	CS_2	Freezing	− 111 6
Toluene	C_7H_8	Freezing	− 95 1
Ethyl acetate	$CH_3CO_2C_2H_5$	Freezing	− 83 6
Chloroform	$CHCl_3$	Freezing	− 63 5
Chlorobenzene	C_6H_5Cl	Freezing	− 45 2
Carbon tetrachloride	CCl_4	Freezing	− 22 9
Sodium sulfate	$Na_2SO_4.10H_2O$	Transition	32 384
Potassium dichromate	$K_2Cr_2O_7$	Melting	397 5
30 5 NaCl + 69 5 Na_2SO_4		Melting	637 0
Potassium chloride	KCl	Melting	770 3
Sodium chloride	$NaCl$	Melting	800 4
Sodium sulfate	Na_2SO_4	Melting	884 7
Potassium sulfate	K_2SO_4	Inversion	583 0
Potassium sulfate	K_2SO_4	Melting	1069 1
Nickel	Ni	Melting or freezing	1452
Cobalt	Co	Melting or freezing	1490
Lithium metasilicate	Li_2SiO_3	Melting	1202
Diopside	$CaMgSi_2O_6$	Melting	1395
Anorthite	$CaAl_2Si_2O_8$	Melting	1555

LITERATURE

(For a key to the periodicals see end of volume)

(1)Holborn and Day, 8, 2: 505, 00   12, 10: 171, 00 (Sb, Ag, Au, Cu)   (2)Buckingham, 31A, 3: 281, 07 (Review of values for S boiling point). (3)Wadner and Burgess, 31A, 7: 1, 11 (Naphthalene, benzophenone, Sn, Cd, Zn)   (4)Holborn and Henning, 8, 35: 761, 11 (Naphthalene, benzophenone, S, Sn, Cd, Zn) (5)Day and Sosman, 152, No 187; 11 (Zn, Sb, Ag, Au, Cu, Pd, Pt)   (6)Day and Sosman, 12, 33: 517, 12   8, 38: 849, 12 (Benzophenone, Zn, Sb, S). (7)Henning, 8, 43: 282, 14 (O, CO2, Hg) (8)Eumorfopoulous, 5, 90A: 189; 14 (8) (9)Wilhelm, 31A, 12: 655, 16 (Hg)

(10)Chappuis, 258, 16: 17 (8) (11)Bureau of Standards, Cir No 66; 17 (Sn, Zn, Al, Cu)   (12)Cath, 168, No 152d; 18 64P, 21: 656, 19 (O, N). (13)Martines and Onnes, 168, No 156b; 22   18, 6 31, 22 (H) (14)Worthing, 90, 22: 0, 24 (W) (15)Henning and Heuse, 8, 23: 104, 24 (O, N, H). (16)Finck and Wilhelm, 1, 47: 25 (Naphthalene, benzophenone) See also References under Standard Scale of Temperature.

Additional Fixed Points Timmermans, Van der Horst and Onnes, 168, No 187; 22 (Organic liquids below 0°). Dickinson and Mueller, 31A, 3: 641; 07 (Na2SO4 transition) Roberts, 2, 23: 386, 24 (Salts) Day and Sosman, Dictionary of Applied Physics, 1: 836, 22 (Metals and silicates)   Richards, et al, 1, 36: 485, 14 (Na2CO3 hydrates transitions) 40: 89, 18 (SrCl2 and SrBr2 transitions) 41: 2019, 19 (C4H8)

THE LEIDEN TEMPERATURE SCALE

In certain sections of International Critical Tables (where so indicated) the Leiden temperature scale will be employed. (Onnes and Holst, 168, No 141a. 64V, 23: 175; 14. Cath and Onnes, 168, No. 152a. 64V, 26: 137, 190; 17. Cath, 168, No. 152d. 64V, 27: 553; 18.) The relation between the Leiden and the I. C. T scales is shown by the following table:

Point	I. C. T	Leiden	Leiden − I. C. T.
H₂ (B. P.)	−252 8°	−252 74°	+0 06°
O₂ (B. P.)	−183 0°	−182 95°	+0 05°
ca. −40°			+0 04

4. RESISTANCE THERMOMETERS

E. F. MUELLER

Standard methods of calibration have been developed only for platinum resistance thermometers. Data on the resistance-temperature relation for particular thermometers of other metals, such as gold and lead, are available, and formulae to represent the relation have been published, but standardized methods for the calibration of such thermometers have not been developed.

The standard working scale, in the interval 0° to 650°C, is defined by means of a resistance thermometer of pure platinum, for which the relation between resistance R and temperature t is given by the equation:

$$R = R_0(1 + at + bt^2). \tag{1}$$

This may be transformed into the Callendar equations:

$$(pt) = \left(\frac{R - R_0}{R_{100} - R_0}\right)100; \quad t - (pt) = \delta\left[\left(\frac{t}{100} - 1\right)\frac{t}{100}\right]. \tag{2}$$

The three constants in these equations, namely R_0, a, and b or R_0, R_{100} and δ respectively, are determined by calibration at the ice point, the steam point, and the sulfur boiling point.

The purity of the platinum must be such that $R_{100}/R_0 > 1.390$ and $R_{444 6}/R_0 > 2.645$, the latter requirement being equivalent to $\delta < 1.50$.

The Callendar equations were devised to facilitate computations by the method of successive approximations. The platinum temperature, symbol (pt), is proportional to the resistance above R_0 and the amount by which it differs from the true temperature is given by the correction term,

$$\delta\left(\frac{t}{100} - 1\right)\frac{t}{100}.$$

Consequently, a value of t sufficiently exact for use in computing the value of the correction term is readily obtained, if not by the first, then certainly by a second or third approximation.

In the interval −195° to 0°C the standard reference scale is defined by means of the platinum resistance thermometer, using the equation

$$t - (pt) = \delta\left[\left(\frac{t}{100} - 1\right)\frac{t}{100}\right] + \beta\left[\left(\frac{t}{100} - 1\right)\frac{t^3}{100^3}\right]. \tag{3}$$

The constants R_0, R_{100} and δ are determined just as for the range above 0° and the additional constant β is determined by a calibration at the boiling point of oxygen. A criterion for the purity of the platinum is that $R_{-183}/R_0 < 0.250$.

Thermometers which are not to be heated above ordinary temperatures may be calibrated at the freezing point of mercury, the CO_2 point and the oxygen point, using the interpolation formula:

$$R = R_0(1 + at + bt^2 + ct^4). \tag{4}$$

The constant c in the equation is approximately equal to 5×10^{-12} and when this value is assumed, calibration at the CO_2 point may be omitted.

Equations (3) and (4) will yield substantially equivalent results, but they are not algebraically interconvertible.

Equation (1) or equation (2) may be used for temperatures up to 1000° or even 1100°C and the temperatures so determined will not depart appreciably from the standard scale.

LITERATURE

(For a key to the periodicals see end of volume)

(1) Callendar, 62, 178: 160; 87. (2) Wadner and Burgess, 31A, 6: 149; 09. (3) Holborn and Henning, 8, 35: 761, 11. (4) Henning, 8, 40: 635; 13 (Pt and Pb at low temperatures). (5) Henning, 8, 43: 282; 14. (6) Cath, Onnes and Burgers, 168, No 152c; 17 64P, 20, 1163, 18 (Pt and Au at low temperatures) (7) Henning and Heuse, 96, 23: 95; 24. (8) Van Dusen, 1, 47: 326, 25

5. TEMPERATURE SCALES DEFINED BY LIQUID-IN-GLASS THERMOMETERS

E. F. MUELLER

The readings of any particular thermometer, taken when all of the liquid in the thermometer is at a uniform temperature, may be reduced to those which would have been obtained if the thermometer had been perfect and used under ideal conditions, by applying corrections for non-uniformity of the capillary bore, corrections for the change of reading due to departure of the external and internal pressures from arbitrary constant values, a correction for the departure of the ice-point reading, taken immediately after the temperature measurement, from the 0° mark, and

a correction to allow for the value of the mean scale degree, in case the difference between the readings of the thermometer taken first at 100°C and then at 0°C, does not correspond to 100 scale degrees. The reading of a thermometer, when so corrected, may be defined as the temperature on the liquid-in-glass scale for the particular liquid and the particular kind of glass of which the thermometer is made.

The temperature scales of mercury thermometers made of French hard glass (verre dur), Jena 16III, Jena 59III, Jena 1565III and Jena combustion tubing are defined as above. For Kew glass, the temperature scale is defined in a somewhat different way, in that the point of reference is the (single) ice point reading taken after the thermometer has been held for a sufficiently long period at ordinary temperature (about 10°C) instead of the (variable) ice point reading taken immediately after each temperature measurement. It is apparent that temperatures on the mercury-in-glass scale are not proportional to the relative increase of volume of mercury-in-glass.

Constants characteristic of the several glasses are the ice-point depression, the softening point, and the average coefficient of expansion of mercury-in-glass, between 0° and 100°C.

The ice point depression is the difference between the ice point reading of the thermometer taken after it has been kept a sufficiently long time (a few days or weeks) at 0° and the ice point reading taken immediately after the thermometer has been kept a sufficiently long time (a few minutes or hours) at 100°C. Good thermometric glasses are characterized by small ice point depression (less than 0.1°C) and rapid recovery. Some glasses have an ice point depression of nearly 1°C.

The softening point determines the upper limit of temperature at which thermometers made of the glass can be used

The expansion coefficient is useful in calculating corrections for emergent stem.

Values of these characteristic constants are.

Glass	Ice point depression °C	Softening point °C	Coefficient of cubical exp. of mercury-in-glass 0° to 100°C
Verre dur	0 07-0 11	500	0 000158
"Kew" glass	0 20		
Jena 16III	0 04-0 08	505	0 000158
Jena 59III	0 03-0 04	510	0 000164
Jena 1565III	0 01	660	0 000172
Jena combustion	0 03	560	

Thermometers containing alcohol, toluene or pentane are not adapted for observation at 100°C, and for such thermometers the mean scale degree is conveniently referred to the interval 0° to −78.5°, the sublimation temperature of carbon dioxide serving to fix the latter temperature.

The tabular values are the result of comparisons of mercury-in-glass thermometers with gas thermometers or platinum resistance thermometers which served to establish the standard scale of temperature. The data for Jena 16III glass and Jena 59III glass may be used for Corning normal and Corning borosilicate thermometer glasses respectively.

Data of this kind were of great importance during the latter part of the 19th and even during the early part of this century, when calibrated mercury-in-glass thermometers were used to distribute the standard scale of temperature. At present the data are useful principally for minor purposes, such as calculation of factors for determining emergent stem correction, calculation of setting factors for metastatic thermometers, such as the Beckmann thermometer, graduation of thermometers by mercury thread calibration in the absence of standards and thermally controlled baths, etc.

In the tables, t represents the temperature on the standard working scale (platinum resistance thermometer) except for verre dur, where t represents temperatures on the former International hydrogen scale, which in practice is not distinguishable from the standard reference scale, while t_{gl} represents corresponding temperatures on the several liquid-in-glass scales.

VALUES OF $t - t_{gl}$ FOR MERCURY-IN-GLASS THERMOMETERS

t = temperature on standard scale, t_{gl} = temperature on mercury-in-glass scale.

t°C	French hard (verre dur)	Kew glass	Jena 16III	Jena 59III	Jena 1565III	Jena combustion
− 39	+0 120					
− 30	+0 290		+0 28	+0.13		
− 20	+.172		+.16	+.07		
− 10	+.073		+.07	+.03		
0	.000	0 00	00	00	0.00	0.00
+10	− .052	00	− .06	− .02	− .03	
20	− .085	00	− .09	− .04	− .05	
30	− .102	+.005	− 11	− .04	− .06	
40	− .107	+.01	− 12	− .03	− .06	
50	− .103	+.01	− 12	− .03	− .05	
60	− .090	+.01	− 10	− .02	− .04	
70	− .072	+.015	− .08	− .01	− .03	
80	− .050	+.02	− .06	00	− .02	
90	− .026	+.025	03	+.02	− .01	
100	.000	.00	00	00	.00	0.00
120	+.06		+.03	05	+.06	
140	+.07		+.02	− .16	+.03	
160	+.03		− .02	− .31	− .13	
180	− .04		− 12	− 52	− 38	
200	− 12		− 29	− 84	− .90	− 1.13
220			− 5	− 1 3	− 1.3	− 1.6
240			− 9	− 1 9	− 1 8	− 2.2
260			−1 4	− 2 6	− 2.4	− 3.0
280			−2 0	3 4	3.1	− 4.0
300			−2 7	4 4	− 3.9	− 5.1
320				5 8	− 4.8	− 6.4
340				7 2	− 5.9	− 7.8
360				− 8 8	− 7 3	− 9.5
380				10 6	− 8.9	−11.4
400				−12.6	−10 5	−13.5
420				14 9	−12.4	−15.9
440				− 17 4	−14.7	−18.6
460				−20 2	−17 2	−21.5
480				−23 3	−20 0	−24.8
500				−26 9	−23 1	−28.4
550					−32	−39.
600					−44	
650					−58	

VALUES OF $t - t_g$ FOR LIQUID-IN-GLASS THERMOMETERS

t	Pentane in 16III glass	Toluene in verre dur	Alcohol in verre dur
−190	−23 4		
−180	−21 0		
−170	−18 6		
−160	−16 2		
−150	−13 9		
−140	−11 6		
−130	− 9 4		
−120	− 7 3		
−110	− 5 3		

VALUES OF t − t₁ FOR LIQUID-IN-GLASS THERMOMETERS.—Continued

t	Pentane in 16^III glass	Toluene in verre dur	Alcohol in verre dur
−100	− 3 4		
− 90	− 1 7		
− 80	− 0 2	0.0	
− 78 5	0 0	0 0	0 0
− 70	+ 1 0	+ 4	+0 3
− 60	+ 2 0	+ 8	+ 6
− 50	+ 2 6	+ 1 1	+ 7
− 40	+ 3 0	+ 1 2	+ .9
− 30	+ 2 9	+ 1 2	+ 9
− 20	+ 2 1	+ 1 0	+ .8
− 10	+ 1 5	+ 0 6	+ 5
0	0 0	0 0	.0
+ 10	2 0		
20	− 4 4		
30	− 7 6		− 3 6
100		− 24 4	

LITERATURE

(For a key to the periodicals see end of volume)

Guillaume, Traité pratique de la thermometerie Gauthier-Villars, Paris, 1889 (General). Chappuis, 238, 6: 1, 88 (Verre dur −25° to 100°). Harker, 5, 78A: 225; 06 (Kew glass) Scheel, Deut Mechan Ztg, 1916: 170 and Holborn, Scheel and Henning, 863 (Jena glasses and organic liquids in glass).

Emergent Stem Correction for Liquid-in-glass Thermometers

If a liquid-in-glass thermometer standardized for total immersion is used with a portion of the liquid column at a temperature below that of the bulb, the reading will be too low for this reason, and an emergent stem correction should be applied to the observed reading.

The emergent stem correction is calculated by the formula,

$$\text{Correction} = Kn(t - t_s)$$

in which

K = coefficient of cubical expansion of mercury-in-glass, per °C,

t = temperature of bulb, °C,

t_s = average temperature °C of the mercury column n°C degrees in length.

The value of t is to be determined by means of an auxiliary thermometer or thermometers, preferably with a capillary thermometer. The sign as well as the magnitude of the correction is given by the formula.

For many purposes, in using mercury-in-glass thermometers K may be treated as a constant of the glass, using the values given above for the apparent coefficient of expansion of mercury-in-glass. The value of K does, however, change with temperature. For purposes of computing the emergent stem correction, it may be considered as depending on the average of t and t_s, that is $\frac{t + t_s}{2}$ and is here so tabulated.

If the coefficients of expansion of mercury and of glass were both constant, K would also be constant Most of the change in K is the result of the varying coefficient of the mercury, so that the change in K with temperature for one glass may with some certainty be inferred from the change for some other glass.

The use of the formula requires that t, the temperature of the bulb, be known. In case t is not known, but is to be determined from the indication of the thermometer, the reading of the thermometer may be substituted in the formula in place of t, as a first approximation and the true magnitude of the correction then calculated by means of a second, or if necessary, a third approximation.

In many cases, in calculating the emergent stem correction for thermometers containing organic liquids, it is sufficient to use the approximate value, K = 0.001. The tables show to what extent this is justified for pentane, toluene, and alcohol. In such thermometers, K is practically independent of the kind of glass used.

With the abandonment of the mercury-in-glass thermometer as an instrument of high precision there has been an increasing tendency to use partial immersion thermometers, graduated and standardized for a particular depth of immersion, thus avoiding the necessity of determining and applying the correction for emergent stem.

TABLE OF EMERGENT STEM CORRECTION FACTORS
Mercury-in-glass Thermometers

$\frac{t + t_s}{2}$ °C	Verre dur	Jena 16^III	Jena 59^III	Jena 1565^III	Jena combustion
50	0 000158	0 000158	0.000164	0 000172	0.000164
100	158	158	164	172	164
150	158	158	165	173	165
200	159	159	167	175	167
250		161	170	177	171
300		164	174	180	174
350			177	184	178
400			182	188	182
450			187	194	188
500			195	200	195

Liquid-in-glass Thermometers

$\frac{t + t_s}{2}$	Pentane	Toluene	Alcohol
−180	0.0009		
−160	09		
−140	09		
−120	10		
−100	10		
− 80	10	0 0009	0 0010
− 60	11	09	10
− 40	12	10	10
− 20	13	10	10
0	14	10	10
+ 20	15	11	10

LITERATURE

(For a key to the periodicals see end of volume)
Buckingham, 31a, 8: 239, 12

Example: A thermometer of Jena 59^III (or Corning borosilicate glass) indicated a temperature, t, of 470° after application of corrections peculiar to the instrument. The thermometer was immersed to the 150° mark, and the average temperature t_s of the 320° (n°) of exposed mercury column was found to be 190°. The average of t and t_s is 330° and the value of the factor K for this temperature is 0 000176. Accordingly

Correction = 0.000176(320)(470 − 190) = 15.8°

The corrected temperature is therefore 470° + 15.8° = 485.8°. Since the bulb temperature was considerably higher than 470° a second approximation may be tried:

Correction = 0.000176(320)(486 − 190) = 16.7°

The second approximation yields a corrected temperature of 470° + 16.7° = 486.7° which in view of the rather large emergent stem correction, may properly be reported as 487°.

Possible short cuts in making the second approximation will be readily apparent.

The example given is purposely somewhat exaggerated by assuming an unusually high temperature (190°) for the emergent

stem, in order to show that the factor K may differ appreciably from the conventional value of 0.00016.

For computations in Fahrenheit temperatures, the proper value of K is $5/9$ of the tabulated value.

6. THERMOCOUPLES

L. H. Adams

"Standard" Calibration Tables (for Use with Deviation Curve)

Standard tables such as these do not necessarily have any absolute significance; primarily, they are arbitrary reference curves which, although representing fairly well the temperature-emf functions for certain thermocouples, are intended for use with an appropriate deviation-curve. This correction-curve is determined for each couple by calibration at several—preferably three or more—fixed points within the "applicability range of the couple." This curve is constructed by plotting ΔE as ordinate ($\Delta E = E_{obs.} - E_{stand}$) against E_{stand} as abscissa. In order to obtain the temperature corresponding to the emf indicated by the couple, the appropriate value of ΔE (as obtained from its deviation curve) is subtracted algebraically from the observed value of E before the latter is converted into degrees by means of the table. Example: At a certain temperature a copper-constantan couple gave an emf of 8720 microvolts. From the previously determined deviation curve of the particular couple the value of ΔE at 8720 microvolts is found to be 12 microvolts. The "standard" emf is therefore 8720 − 12 or 8708 microvolts and from the copper-constantan table this may be seen to correspond to 189.08°, which is the required temperature.

The fixed (i.e., cold) junction is supposed to be maintained at 0°C.

TEMPERATURES AND TEMPERATURE DIFFERENCES FOR EVERY 100 MICROVOLTS

Platinum: Platinrhodium (90-10). Standard range, 630°-1083°C. Applicability range, 0-1754°C

E µV	0	1000	2000	3000	4000	5000	6000	7000	8000	9000	E µV
0	0	147.1	265.4	374.3	478.1	578.3	675.3	769.5	861.1	950.4	0
	17.8	12.6	11.2	10.6	10.2	9.8	9.5	9.3	9.0	8.8	
100	17.8	159.7	276.6	384.9	488.3	588.1	684.8	778.8	870.1	959.2	100
	16.7	12.4	11.1	10.5	10.1	9.8	9.5	9.2	9.0	8.8	
200	34.5	172.1	287.7	395.4	498.4	597.9	694.3	788.0	879.1	968.0	200
	15.8	12.3	11.0	10.5	10.1	9.8	9.5	9.2	8.7		
300	50.3	184.3	298.7	405.9	508.5	607.7	703.8	797.2	888.1	976.7	300
	15.1	12.0	11.0	10.4	10.1	0.7	9.5	9.2	9.0	8.7	
400	65.4	196.3	309.7	416.3	518.6	617.4	713.3	806.4	897.1	985.4	400
	14.6	11.8	10.9	10.4	10.0	9.7	9.4	9.2	9.0	8.7	
500	80.0	208.1	320.6	426.7	528.6	627.1	722.7	815.6	906.1	994.1	500
	14.1	11.6	10.9	10.4	10.0	9.7	9.4	9.1	8.9	8.7	
600	94.1	219.7	331.5	437.1	538.6	636.8	732.1	824.7	915.0	1002.8	600
	13.7	11.5	10.8	10.3	10.0	9.7	9.4	9.1	8.9	8.7	
700	107.8	231.2	342.3	447.4	548.6	646.5	741.5	833.8	923.9	1011.5	700
	13.4	11.5	10.7	10.3	9.9	9.6	9.4	9.1	8.9	8.0	
800	121.2	242.7	353.0	457.7	558.5	656.1	750.9	842.9	932.8	1020.1	800
	13.1	11.4	10.7	10.2	9.9	9.6	9.3	9.1	8.8	8.0	
900	134.3	254.1	363.7	467.9	568.4	665.7	760.2	852.0	941.6	1028.7	900
	12.8	11.3	10.6	10.2	9.9	9.6	9.3	9.1	8.8	8.0	
1000	147.1	265.4	374.3	478.1	578.3	675.3	769.5	861.1	950.4	1037.3	1000

E µV	10,000	11,000	12,000	13,000	14,000	15,000	16,000	17,000	18,000		E µV
0	1037.3	1122.2	1205.9	1289.3	1372.4	1454.8	1537.5	1620.9	1704.3		0
	8.6	8.4	8.3	8.4	8.4	8.2	8.3	8.3	8.3		
100	1045.9	1130.6	1214.2	1297.7	1380.7	1463.0	1545.8	1629.2	1712.6	....	100
	8.5	8.4	8.4	8.3	8.3	8.2	8.5	8.4	8.4		
200	1054.4	1139.0	1222.6	1306.0	1389.0	1471.2	1554.1	1637.6	1721.0	....	200
	8.5	8.4	8.3	8.3	8.3	8.2	8.3	8.3	8.3		
300	1062.9	1147.4	1230.9	1314.3	1397.3	1479.4	1562.4	1645.9	1729.3		300
	8.5	8.4	8.4	8.3	8.3	8.3	8.4	8.4	8.4		
400	1071.4	1155.8	1239.3	1322.6	1405.6	1487.7	1570.8	1654.3	1737.7	....	400
	8.5	8.4	8.3	8.3	8.2	8.3	8.3	8.3	8.3		
500	1079.9	1164.2	1247.6	1330.9	1413.8	1496.0	1579.1	1662.6	1746.0	....	500
	8.5	8.3	8.3	8.3	8.2	8.3	8.4	8.3	8.3		
600	1088.4	1172.5	1255.9	1339.2	1422.0	1504.3	1587.5	1670.9	1754.3		600
	8.5	8.4	8.4	8.3	8.2	8.3	8.3	8.4			
700	1096.9	1180.9	1264.3	1347.5	1430.2	1512.6	1595.8	1679.3	....	....	700
	8.5	8.3	8.3	8.3	8.2	8.3	8.4	8.3			
800	1105.4	1189.2	1272.6	1355.8	1438.4	1520.9	1604.2	1687.6	....		800
	8.4	8.4	8.4	8.3	8.2	8.3	8.3	8.4			
900	1113.8	1197.6	1281.0	1364.1	1446.6	1529.2	1612.5	1696.0	....		900
	8.4	8.3	8.3	8.3	8.2	8.3	8.4	8.3			
1000	1122.2	1205.9	1289.3	1372.4	1454.8	1537.5	1620.9	1704.3	....		1000

TEMPERATURES AND TEMPERATURE DIFFERENCES FOR EVERY 100 MICROVOLTS
Copper: Constantan

E µV	6000	5000	4000	3000	2000	1000	0	-0	-1000	-2000	-3000	-4000	-5000
0	135.91 (2.03)	115.31 (2.09)	94.07 (2.16)	72.08 (2.23)	49.20 (2.33)	25.27 (2.45)	0 (2.59)	0 (2.60)	-26.82 (2.79)	-55.81 (3.05)	-87.86 (3.42)	-124.46 (4.01)	-169.14 (5.20)
100	137.94 (2.02)	117.40 (2.08)	96.23 (2.15)	74.31 (2.23)	51.53 (2.32)	27.72 (2.43)	2.59 (2.57)	-2.60 (2.62)	-29.61 (2.81)	-58.86 (3.08)	-91.28 (3.46)	-128.47 (4.09)	-174.34 (5.40)
200	139.96 (2.02)	119.43 (2.08)	98.38 (2.14)	76.54 (2.22)	53.85 (2.31)	30.15 (2.43)	5.16 (2.56)	-5.22 (2.63)	-32.42 (2.84)	-61.94 (3.11)	-94.74 (3.51)	-132.56 (4.18)	-179.74 (5.64)
300	141.98 (2.01)	121.56 (2.07)	100.52 (2.14)	78.76 (2.21)	56.16 (2.30)	32.57 (2.42)	7.72 (2.55)	-7.85 (2.65)	-35.26 (2.86)	-65.05 (3.15)	-98.25 (3.57)	-136.74 (4.28)	-185.38 (5.89)
400	143.99 (2.01)	123.63 (2.06)	102.66 (2.13)	80.97 (2.20)	58.46 (2.30)	34.98 (2.41)	10.27 (2.53)	-10.50 (2.67)	-38.12 (2.89)	-68.20 (3.19)	-101.82 (3.63)	-141.02 (4.39)	-191.27 (6.17)
500	146.00 (2.00)	125.69 (2.06)	104.79 (2.12)	83.17 (2.20)	60.76 (2.28)	37.38 (2.40)	12.80 (2.52)	-13.17 (2.69)	-41.01 (2.90)	-71.39 (3.22)	-105.45 (3.68)	-145.41 (4.50)	-197.44 (6.51)
600	148.00 (2.00)	127.75 (2.05)	106.91 (2.11)	85.37 (2.19)	63.04 (2.27)	39.77 (2.38)	15.32 (2.51)	-15.88 (2.71)	-43.91 (2.93)	-74.61 (3.26)	-109.13 (3.74)	-149.91 (4.61)	-203.95 (6.97)
700	150.00 (1.99)	129.80 (2.04)	109.02 (2.10)	87.56 (2.18)	65.31 (2.27)	42.15 (2.36)	17.83 (2.49)	-18.57 (2.73)	-46.84 (2.96)	-77.57 (3.29)	-112.87 (3.80)	-154.52 (4.73)	-210.92 (7.55)
800	151.99 (1.99)	131.84 (2.04)	111.12 (2.10)	89.74 (2.17)	67.58 (2.25)	44.51 (2.35)	20.32 (2.48)	-21.30 (2.75)	-49.80 (2.99)	-81.16 (3.33)	-116.67 (3.86)	-159.25 (4.87)	-218.47
900	153.97 (1.98)	133.88 (2.03)	113.22 (2.09)	91.91 (2.16)	69.83 (2.25)	46.86 (2.34)	22.80 (2.47)	-24.05 (2.77)	-52.79 (3.02)	-84.49 (3.37)	-120.53 (3.93)	-164.12 (5.02)	
1000	155.95	135.91	115.31	94.07	72.08	49.20	25.27	-26.82	-55.81	-87.86	-124.46	-169.14	

E µV	19,000	18,000	17,000	16,000	15,000	14,000	13,000	12,000	11,000	10,000	9000	8000	7000
0	369.61 (1.64)	353.08 (1.66)	336.36 (1.68)	319.49 (1.70)	302.42 (1.72)	285.13 (1.74)	267.60 (1.76)	249.82 (1.79)	231.74 (1.82)	213.36 (1.85)	194.62 (1.89)	175.50 (1.93)	155.95 (1.97)
100	371.25 (1.64)	354.74 (1.66)	338.04 (1.68)	321.19 (1.69)	304.14 (1.71)	286.87 (1.74)	269.36 (1.76)	251.61 (1.79)	233.56 (1.82)	215.21 (1.85)	196.51 (1.89)	177.43 (1.93)	157.92 (1.97)
200	372.89 (1.64)	356.40 (1.66)	339.72 (1.68)	322.88 (1.69)	305.85 (1.71)	288.61 (1.74)	271.12 (1.76)	253.40 (1.78)	235.38 (1.82)	217.06 (1.85)	198.40 (1.88)	179.36 (1.92)	159.89 (1.97)
300	374.53 (1.64)	358.06 (1.66)	341.40 (1.67)	324.57 (1.69)	307.56 (1.71)	290.35 (1.73)	272.88 (1.76)	255.18 (1.78)	237.20 (1.81)	218.91 (1.84)	200.28 (1.88)	181.28 (1.92)	161.86 (1.96)
400	376.17 (1.63)	359.72 (1.65)	343.07 (1.67)	326.26 (1.69)	309.27 (1.71)	292.08 (1.73)	274.64 (1.76)	256.96 (1.78)	239.01 (1.81)	220.75 (1.84)	202.16 (1.88)	183.20 (1.91)	163.82 (1.96)
500	377.80 (1.63)	361.37 (1.65)	344.74 (1.67)	327.95 (1.69)	310.98 (1.71)	293.81 (1.73)	276.40 (1.75)	258.74 (1.78)	240.82 (1.81)	222.59 (1.84)	204.04 (1.87)	185.11 (1.91)	165.78 (1.96)
600	379.43 (1.63)	363.02 (1.65)	346.41 (1.67)	329.64 (1.68)	312.69 (1.70)	295.54 (1.72)	278.15 (1.75)	260.52 (1.77)	242.63 (1.80)	224.43 (1.83)	205.91 (1.87)	187.02 (1.91)	167.73 (1.95)
700	381.06 (1.63)	364.67 (1.65)	348.08 (1.67)	331.32 (1.68)	314.39 (1.70)	297.26 (1.72)	279.90 (1.75)	262.29 (1.77)	244.43 (1.80)	226.26 (1.83)	207.78 (1.86)	188.93 (1.90)	169.68 (1.94)
800	382.69 (1.63)	366.32 (1.65)	349.75 (1.67)	333.00 (1.68)	316.09 (1.70)	298.98 (1.72)	281.65 (1.74)	264.06 (1.77)	246.23 (1.80)	228.09 (1.83)	209.64 (1.86)	190.83 (1.90)	171.62 (1.94)
900	384.32 (1.63)	367.97 (1.64)	351.42 (1.66)	334.68 (1.68)	317.79 (1.70)	300.70 (1.72)	283.39 (1.74)	265.83 (1.77)	248.03 (1.79)	229.92 (1.82)	211.50 (1.86)	192.73 (1.89)	173.56 (1.94)
1000	385.95	369.61	353.08	336.36	319.49	302.42	285.13	267.60	249.82	231.74	213.36	194.62	175.50

TEMPERATURES AND TEMPERATURE DIFFERENCES FOR EVERY 0.5 MILLIVOLT

Chromel-alumel

E mv	0	10	20	30	40
0	0.0	244.5	482 8	719.2	970 4
	12.3	*12 2*	*11 7*	*12 2*	*13 0*
0.5	12 3	256.7	494 5	731 4	983.4
	12 1	*12.2*	*11 7*	*12 3*	*13 1*
1.0	24.4	268.9	506 2	743 7	996 5
	12.0	*12.1*	*11 7*	*12 3*	*13 2*
1.5	36.4	281 0	517 9	756 0	1009 7
	12 0	*12 1*	*11 7*	*12 3*	*13 3*
2.0	48 4	293 1	529 6	768 3	1023 0
	12 0	*12 0*	*11 7*	*12 4*	*13 3*
2.5	60 4	305 1	541 3	780 7	1036 3
	12.0	*12.0*	*11 7*	*12 4*	*13 4*
3.0	72 4	317.1	553 0	793 1	1049 7
	12 0	*12 0*	*11 7*	*12 5*	*13 5*
3.5	84 4	329 1	564.7	805 6	1063 2
	12 0	*11 9*	*11 7*	*12 5*	*13 6*
4.0	96 4	341 0	576 4	818 1	1076 8
	12 1	*11 9*	*11 8*	*12 5*	*13 7*
4.5	108.5	352 9	588 2	830 6	1090 5
	12 1	*11 9*	*11 8*	*12 6*	*13 7*
5.0	120 6	364 9	600 0	843 2	1104 2
	12 2	*11 9*	*11 8*	*12 6*	*13 8*
5.5	132 8	376 8	611 8	855 8	1118 0
	12 4	*11 9*	*11 8*	*12 6*	*13 8*
6.0	145 2	388.6	623 6	868 4	1131 8
	12.5	*11 8*	*11 8*	*12 6*	*13 9*
6.5	157.7	400 4	635 4	881 0	1145 7
	12 6	*11 8*	*11 8*	*12 7*	*13 9*
7.0	170 2	412.2	647 2	893 7	1159 6
	12 5	*11 8*	*11 9*	*12 7*	*14*
7.5	182.7	424.0	659 1	906 4	(1174)
	12 5	*11 8*	*11 9*	*12 7*	*14*
8.0	195 2	435 8	671 0	919 1	(1188)
	12 4	*11 8*	*12 0*	*12 8*	*14*
8.5	207 7	447 6	683 0	931 9	(1202)
	12 3	*11 8*	*12 0*	*12 8*	
9.0	220'0	459 4	695 0	944 7	
	12 3	*11.7*	*12 1*	*12 8*	
9.5	232 3	471 1	707 1	957 5	
	12 2	*11 7*	*12 1*	*12 9*	
10.0	244.5	482 8	719 2	970 4	

Fixed-junction Corrections

If the fixed or "cold" junction be not maintained at 0°C, a correction must be applied. This may be done by any one of several methods, of which the following are suggested.

A. Let the temperature of the fixed junction be t_c and that of the variable or "hot" junction be t. Then to the emf as read E_{t-t_c} add the emf corresponding to t_c. This gives E_t which may at once be converted into degrees by means of the proper table.

B. Multiply the fixed-junction temperature by the factor, $f = (dE/dt)_0/(dE/dt)$, which is the ratio of the mean emf-temperature gradient between 0° and t_c to the gradient at t, and add the product to t', the uncorrected temperature. That is, $t = t' + ft_c$. These emf-temperature gradients may be obtained by taking the reciprocals of the numbers appearing in the difference columns of the calibration tables.

COMPARISON OF THE MORE COMMON THERMOCOUPLES

E mv	Temperature, °C				E mv	Temperature, °C		
	Iron: constantan	Chromel (X): copel	Chromel (P): alumel	Platinrhodium: gold-palladium		Platinum: platinrhodium (Heraeus)	Platinum: Platinrhodium (Johnston-Matthey)	Copper: constantan
0	0	0	0	0	0	0	0	0
5	95	105	121	131	1	147	146	25
10	186	195	244	237	2	265	260	49
15	277	277	365	335	3	374	364	72
20	367	353	483	429	4	478	461	94
25	457	425	600	513	5	578	553	115
30	546	495	719	607	6	675	641	136
35	632		843	694	7	769	725	156
40	713		970	779	8	861	806	176
45	792		1104	866	9	950	884	195
50	871			954	10	1037	959	213
55	950			1044	11	1122	1032	232
60				1136	12	1206	1103	250
					13	1289	1173	268
					14	1372	1242	285
					15	1455	1311	302
					16	1537	1379	320
					17	1620	1447	336
					18	1704	1515	353

* 10 % Rh, 40 % Pd.

LITERATURE

(For a key to the periodicals see end of volume)

[1] Adams, *128*, **3**: 469; 13. *1*, **36**: 65; 14. *255*, **1919**: 2111. [2] Adams, O. [3] Adams and Johnston, *12*, **33**: 534; 12. [4] Foote, Fairchild and Harrison, *52*, No *170*, 21. [5] Hoskins Mfg Co., Catalog D; 24. [6] Roberts, O. [7] Sosman, *12*, **30**: 7, 10.

OPTICAL PYROMETRY

C. O. FAIRCHILD AND H. T. WENSEL

The temperature scale above the melting point of gold is based upon Wien's Law, $J_\lambda = c_1\lambda^{-5} e^{-\frac{C_2}{\lambda T}}$, in which the constant C_2 (1.433 cm deg) and the value 1336°K for the melting point of gold determine the scale. In optical pyrometry temperatures are usually measured by comparing the brightness of a glowing object with that of the filament of a lamp mounted in the image plane of a simple telescope. For highest accuracy the current through the lamp is kept at or near the value corresponding to 1336°K and higher temperatures are measured by reducing the brightness of the image of the object to match that of the filament by means of a suitable screen such as a rotating sector or an absorption glass of known transmission. The temperature is then found from the following formula derived from Wien's Law:

$$\frac{1}{T} = \frac{1}{1336} + \lambda_e \cdot \frac{\log_{10} R}{6222},$$

in which R is the transmission of the absorption device and λ_e is the "mean effective wave-length" of a color filter in the pyrometer for the temperature interval 1336° to T. Values of λ_e can be obtained in some cases by the use of Table 2.

For practical purposes the pyrometer is ordinarily calibrated in the range 700° to 1400°C (occasionally to 1550°C) in terms of filament current. A satisfactory empirical relation between the current I through the lamp filament and temperature $t°C$ is:

$I = a + bt + ct^2 + dt^3$. For tungsten lamps with short 3 mil filaments dI/dt varies from about 0.00015 ampere per degree at 700°C ($I = 0.3$) to 0.0003 ampere per degree at 1400° ($I = 0.5$). For measurements above 1400° an absorption glass of such type is employed that $A (= \lambda_0 \log_{10} R/6223)$ is a constant or varies slightly with temperature. If the spectral transmission, Tr, of the absorption device is of the form $Tr_\lambda = e^{\frac{K}{\lambda}}$, A will be a constant and equal to K/c_2. For sector discs $A = $ constant $\cdot \lambda e$.

Table I

Temperatures extrapolated from 1336°K, using Wien's Law, compared with those obtained using Planck's Law. The values in this table were computed from the relation:

$$T_p = \frac{c_2}{\lambda \log_e\left[1 + e^{\frac{c_2}{\lambda T_w}}\right]}$$

taking $\lambda = 0.65\mu$.

T_v	T_p	$T_w - T_p$	T_u	T_p	$T_w - T_p$
1336	1336 000		4500	4493	7
2000	1999 997	0 003	5000	4986	14
2500	2499 958	.042	6000	5959	41
3000	2999 74	.26	8000	7825	175
3500	3499 0	1 0	10 000	9550	450
4000	3997	3	∞	31 800	∞

Table 2

Effective wave-length and mean effective wave-length of optical pyrometer red glass filters. The effective wave-length λ_T is found from the formula

$$\frac{1}{\lambda_T} = a - \frac{b}{T}$$

Equation*	Corning H. T. red glasses				Visibility
	A	B	C	D	
a	1 5509	1 5415	1 5369	1 5319	
b	29.6	28 2	28 0	26 8	
Wave-length microns	Transmission				
0.615	0.000	0.000	0.000	0.000	0 442
.625	.085	.007	.000	.000	.323
.635	.520	.270	.141	.080	.220
.645	.730	.533	.389	.350	.141
.655	.798	.637	.508	.520	.084
.665	.815	.664	.541	.580	.046
.675	823	.677	.557	.605	.024
.685	828	686	567	605	0126
.695	830	.689	572	.603	0061
.705	830	689	572	598	0031
.715	826	.682	564	590	00158
.725	824	.679	.559	.580	00078
.735	822	.676	.555	.572	00038
.745	820	.672	551	567	00018
.755	818	.669	.517	.550	00009
.765	815	.664	541	535	00003
.775	813	661	537	510	00000

* The constants a and b are given for four typical red glasses of the transmissions indicated. The change in effective wave-length with temperature of glass filter itself is closely 0 00009μ per deg C at ordinary room temperatures.

Angular apertures required in the telescope of the disappearing filament type of optical pyrometer for a balance between reflection and diffraction at the filament. Under such conditions disappearance of the filament is obtained without resorting to low magnification or very low resolving power.

Table 3.—Tungsten Filament of Circular Cross-section

Exit aperture radians	Entrance aperture, radians	
	Filament diameter 0.04 to 0.06 mm	Filament diameter 0.1 mm
0.005	very low resolving power	
.01	0.04 and larger	0.04 and larger
02	.06 to .16	.055 to .07
.04	.08 to .13	
.06	non-disappearance	

Table 4.—Brightness Temperature versus True Temperature for Red Light($\gamma = 0.65\mu$)

Observed brightness temperature	True temperature						
	Platinum[1]	Iron[2]	Iron oxide[3]	Nickel oxide[4]	Copper[5]	Copper oxide[5]	Nichrome or chromel[6]
700	745		700	701			702
800	857		801	802			804
900	972		902	904		903	906
950					1083	958	
975					1181		
1000	1090		1004	1007	1156	1020	1010
1025					1193		
1050					1231	1087	
1100	1210	1183	1106	1110		1159	1116
1150						1233	
1200	1332	1296	1210	1215			1224
1300	1455	1410		1320			
1400		1525					
1500		1641					
1600		1758					
1700		1877					
1750		1936					

LITERATURE

(For a key to periodical see end of volume)

(1) Waidner and Burgess, *31a*, 3: 163, 07.   (2) Computed for an emissivity of 0.4, *cf.* Burgess, *32*, No. 91: 17.   (3) Burgess and Foote, *31a*, 12: 83; 15   (4) Burgess and Foote, *31a*, 11: 41; 15.   (5) Burgess, *31a*, 6: 111; 09.   (6) Foote, Bureau of Standards, *O* For data on C, Ta, W and other substances see sections on emissivity, color temperature, etc.

General References

Burgess and Le Chatelier, Measurement of High Temperature, 1912. Pyrometry: Symposium of American Institute of Mining and Metallurgical Engineers, 1919. Foote, Fairchild and Harrison, *52*, No. 170: 21. Foote, Mohler and Fairchild, *128*, 7: 18; 17. Foote, *85*, 13: 3; 18. Forsythe, *85*, 15: 3, 20. Fairchild and Hoover, *48*, 7: 7; 23.

From −80° to −50°.—35. *Ethyl ether 80% + ethyl alcohol 20%:* Fl. Used down to −78°. η < alcohol Less turbid from moisture than is ether *(25)*. 36. *H₂SO₄ 38% in H₂O*, E.: S. at −75°. η+ at low temps. Cor. *(23)*. 37. *Chloroform:* S. at −63.5°. Non-Fl. η− at −63° *(21)*. Cor−. SS. *(37)*. A small quantity of alcohol prevents decomposition. 38. *CaCl₂ 29.8% in H₂O.* E.: S. at −55°. η+ at −55° *(38)*. Cor.+ *(32, 41)*. Cor. diminished by addition of K₂CrO₄ *(27)*.

From −50° to −25°. 39. *Gasolene + CCl₄:* Depending upon the density of the gasolene the following %'s of CCl₄ should be used to reduce Fl. 0.765, 30%; 0.725, 45%; 0.700, 60%; 0.680, 70% *(2, 28)*. The 65% CCl₄ may be used at −50°. Flash pt. ca. 50°. Cor− *(8)*. 40. *Chlorobenzene:* S. at −45.2°. Fl. SS. *(37)*. 41. *NaCNS 500 g per 1 H₂O*, E.: S. at ca. −33°. Cor. < NaCl or CaCl₂ *(36)*. 42. *Ethyl alcohol 25% + glycerine 25% + water 50%:* Used to −30° *(40)*.

From −25° to 0°. 43. *Carbon tetrachloride:* S. at −22.9°. Non-Fl. η− at −23° *(21)* Cor−. SS. *(37)*. 44. *NaCl 22.4% in water*, E.: S. at −21.2°. η−. Cor.

DISTILLATES FROM GALICIAN PETROLEUM[11]

Fractionation temp	24°–40°	40°–60°	60°–80°	80°–100°	100°–120°
d₄¹⁵ ...	0.6324	0.6593	0.7005	0.7351	0.7495
S. at	−203°	−198°	−185°	−170°	−151°

Fractionation temp.	120°–110°	140°–160°	160°–180°	180°–200°	200°–220°
d₄¹⁵	0.7625	0.7738	0.7872	0.7962	0.8072
S. at	−139°	−127°	−112°	−104°	−93°

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Archibald and Ure, *4*, **125**: 726, 24   (²) Associated Factory Mutual Fire Insurance Cos, Quart. Nat Fire Protect Assoc **11**: 173, 17.   (³) Baudin, *34*, **132**: 1207, 01   (⁴) Baume, *42*, **12**: 216, 14   (⁵) Beckmann and Waentig, *85*, **67**: 17, 10   (⁶) Cabot, *54*, **26**: 813, 07   (⁷) Cardoso, *42*, **13**: 312; 16.   (⁸) Cragoe, McKelvy and O'Connor, *31.4*, **18**: 707; 23.   (⁹) Fabre, *42*, **21**: 268, 20.

(¹⁰) Fischer, Die neueren Arzneimittel, 6th ed, p. 74.   (¹¹) Formánek, Knop and Korber, *136*, **41**: 731, 17   (¹²) Hammerl, *75*, **78**: 59; 78.   (¹³) Henning, U S Pat. 1,393,124, Brit Pat. 158,494, 20.   (¹⁴) Henning, U. S. Pat 1,386,497; Canadian Pat 213,825.   (¹⁵) Henning, *243*, **33**: 33; 13   (¹⁶) Henning, *Bd2*, p 261   (¹⁷) Hoffmann and Rothe, *243*, **27**: 265; 07.   (¹⁸) Holborn and Wien, *8*, **59**: 213, 96   (¹⁹) Jenkin, *85*, **16**: 197; 22.

(²⁰) Jenkin and Shorthose, *115*, **116**: 761, 23   *240*, **66**: 317, 24   (²¹) Kanolt, Bur. Stands, O   (²²) Kohlrausch, *8*, **60**: 463; 97.   (²³) Pickering, *4*, **57**: 331; 90   (²⁴) Loomis and Walters, Bur Mines, O.   (²⁵) Maass and Wright, *1*, **43**: 1098, 21   (²⁶) Meyerhoffer and Saunders, *7*, **31**: 381; 99   (²⁷) Pedersen, U S Pat. 1,105,320   (²⁸) Remington and Wood, U. S. Dispensatory, 20th ed, 18   (²⁹) Roozeboom, *7*, **4**: 42; 99

(³⁰) Rothe, *245*, **22**: 14, 33; 02   (³¹) Rothe, *243*, **22**: 192,02.   (³²) Rudnick, *48*, **11**: 608, 19   (³³) Ruff and Fischer, *25*, **36**: 421; 03.   (³⁴) Saposhnikov, *245*, **6**: 384   (³⁵) Smits and Berckmans, *64P*, **21**: 401; 19   (³⁶) Sperr, U S Pat 1,173,327   (³⁷) Timmermans, Van der Horst and Onnes, *34*, **174**: 365, 22   Timmermans, *28*, **32**: 95, 23.   (³⁸) Tucker, *67*, **25**: 111, 13   (³⁹) Wahl, *5*, **87**: 371, 12

(⁴⁰) Walton and Judd, *50*, **18**: 717, 14   (⁴¹) Zimmerman, *244*, **9**: 307, 21.   (⁴²) Thiele and Schulte, *7*, **96**: 312, 20.

LABORATORY METHODS FOR THE PRODUCTION OF COLD

C. W. KANOLT

(a) Liquids for Cooling by Vaporization into the Atmosphere

The liquid may be sprayed onto the object to be cooled *(2, 3,* it may be vaporized by a current of air passed through it, forming a bath in which the object to be cooled is immersed *(5)*; it may b vaporized from a porous vessel *(1)*; or in other ways. The temperatures obtainable from the liquids are approximately in the order of their boiling points given below, but are much lower. Gases with critical temperatures above 20° are not included.

The data given below are, in the order given; boiling point, name of liquid, remarks, and literature.

Remarks: 1. Harmless. 2. Harmful. 3. Flammable. 4. Non-flammable. 5. Anaesthetic.

100°, *Water* (1, 4). 61.2°, *Chloroform* (4, 5). 46.2°, *Carbon disulphide* (2, 3). 40°, *Methylene chloride* (4, 5). 38.4°, *Ethyl bromide* (4, 5). 35°–39°, *Amylene, techn.* (3, 5). 34.6°, *Ethyl ether* (3, 5) produces −15° to −20° *(2, 5)*. 13.1°, *Ethyl chloride* (3, 5) produces −35°*(2)*. 0°—70°, *Volatile petroleum distillates* (1, 3). −10.0°, *Sulfur dioxide* (2, 4). −24.1°, *Methyl chloride* (3, 5) produces −55° to −60° *(1, 2)*. −33.4°, *Ammonia* (2, 3). *Carbon dioxide* (1, 4). (The liquid can not exist at atmospheric pressure. Solid can be obtained by the release of liquid from pressure. Sublimation temperature −78.5°. Used mixed with a liquid *(6)*, produces −112° to −115° *(1)*. −89.8°, *Nitrous oxide* (4, 5).

LITERATURE

(For a key to the periodicals see end of volume)

(¹) d'Arsonval, *34*, **133**: 980; 01   (²) Braun, Die Lokalanaesthesie, Chapt. 4.   (³) Kanolt, *48*, **9**: 416; 24.   (⁴) Krause, *Bd1*, **6**: 635; 19.   (⁵) Lawrence, *247*, No. **18**: 10; 16.   (⁶) Thiele and Schulte, *7*, **96**: 312; 20.

(b) Freezing Mixtures

To absorb the largest amount of heat, an aqueous freezing mixture should be made with ice, rather than with water, and the other substance used should be cooled to 0°, or as low as possible, before mixing with the ice. To absorb at a given temperature the maximum amount of heat per unit mass of mixture, the proportions of ice and the other cooling agent should be those of a solution, the freezing point of which is the required temperature *(8)*. The eutectic (cryohydric) temperature is the lowest attainable, if the ingredients are precooled sufficiently. Most, if not all, salts when mixed at room temperature with ice, produce sufficient cooling to reach this temperature.

For more extensive information than given here relative to the freezing points of solutions, together with the literature references, see the separate tables of freezing points.

The following mixtures are among the most useful:

(a) Sodium chloride with ice for temperatures down to −21.2°.

(b) Hydrated calcium chloride, CaCl₂·6H₂O, with ice, for temperatures down to −55°.

Aqueous solutions of sulfuric acid or hydrochloric acid with ice have an advantage over salts with ice in avoiding the delay incident to the solution of the salt.

Substances	Composition of mixture (% anhydrous salt, unless otherwise stated). E = eutectic composition	Freezing point of solution	Initial condition of freezing mixture	Lowest attained temperature recorded	Heat absorbed at temperature of mixing, cal per g of mixture	Heat absorbed (at freezing or saturation point of solution) from objects to be cooled, cal per g of mixture. The * values are heats of fusion of the eutectic, r. [8]
NaCl—H₂O (4, 12)	22 4 (E for NaCl 2H₂O)	−21 2°				56 4*
	23 1 (E for NaCl)	22 4°				
	24 8		salt and ice at −1°	−21 3°		
			with ice	−21°		
NaNO₃—H₂O (12, 13)	33 3		salt and ice at −1°	−17 75°		
	37 E	−18 5°				57 5*
	42 9		water and salt 13 2°	−5 3°		
Na₂CO₃.10H₂O—H₂O (12)	5 93E	2 1°				77 2*
	16 7		salt and ice at −1°	2 0°		
Na₂SO₄ 10H₂O—H₂O	3.8E	−1 2°				80 1*
Na₂S₂O₃.5H₂O—H₂O (13)	30 0E	−11°				
	52 4		water and salt 10 7°	8 0°		
NaOOCCH₃.H₂O—H₂O (13)	45 9		water and salt 10 7°	4 7°		
KCl—H₂O (12)	19 3	−9 0				71 2*
	23 1		salt −1° ice 0°	10 9°		
KNO₃—H₂O (12)	11 2E	3 0°				80 7*
	11 5		salt and ice at −1°	2 85°		
K₂SO₄—H₂O (12)	6 54E	−1.55°				
	9 1		salt and ice at −1°	1 9°		
KSCN—H₂O (13)	60 0		water and salt 10 8	23 7°		
NH₄Cl—H₂O (12)	18 7E	15 8°				75 0*
	20.0		salt and ice at −1°	15 4°		
	16.6	−6°	water and salt 0°	14 0°	12 2	2 6
			ice and salt 0°		78 8	73 6
	31 0		ice and salt at −1°	−16 75°		
	31 2	12°	water and salt 0°	−26 0°	19 7	6 8
			ice and salt 0°		74 6	65.6
	37 5		water and salt 13 6°	13 6°		
	41 2	17.4°				68 4*
NH₄NO₃—H₂O (12, 13, 15)	43 3E	−17.5°	water and salt 0°	33 9°	21 3	8 2
			ice and salt 0°		69 5	57.1
	46 8	−12°	water and salt 0°	36 4°	25 5	13 6
			water and salt 20°			3 1
			ice and salt 0°		68 1	59 8
	50 3	−6°	water and salt 0°	39 3°	26 5	19 0
			water and salt 20°			8 9
			ice and salt 0°		66 2	62 1
	54 1	0°	water and salt 0°	42 2°	27 6	24 3
			water and salt 20°			14.5
			ice and salt 0°		64 4	64.4
	57 1	5°	water and salt 0°	−44 7°	28 1	28.4
			water and salt 20°			18 8
NH₄SCN—H₂O (13)	57.1		water and salt at 13 2°	18 0°		
CaₐClₐ6H₂O—H₂O (6)	% of hydrated salt 16 9	−4 0°	ice and salt 0°		69 9	66 2

Substances	Composition of mixture (% anhydrous salt, unless otherwise stated) E = eutectic composition	Freezing point of solution	Initial condition of freezing mixture	Lowest attained temperature recorded	Heat absorbed at temperature of mixing, cal. per g of mixture	Heat absorbed (at freezing or saturation point of solution) from objects to be cooled, cal. per g of mixture. The * values are heats of fusion of the eutectic, v. (*)
	26 8	−8 1°	ice and salt 0°		63 8	57.3
	34 6	−12 4°	ice and salt 0°		59.3	50.2
	45 7	−22 7°	ice and salt 0°		53 0	38.4
	54 9	−39 0°	ice and salt 0°		48 0	26.0
	58 8L	−54 9?	ice and salt 0°		45 8	17.7
	63 7	−33 3?	ice and salt 0°		43.7	27.9
			water and salt 0°		14 4	none
CaCl₂6H₂O—H₂O (*) — Continued	67 1	19 7°	ice and salt 0°		41 9	33.2
			water and salt 0°		15 4	6.7
	69 0	−14 1°	ice and salt 0°		41 0	35.0
			water and salt 0°		16 0	10.1
			water and salt 20°		none	1.5
	74 1	0°	ice and salt 0°		38 7	38 7
			water and salt 0°		17.7	17.7
			water and salt 20°		none	10 2
	77 5	7.6°	water and salt 0°		19 0	21 6
			water and salt 20°		none	14.7
MgSO₄ 12H₂O—H₂O(⁵)	% anhyd salt 19 0	−3 9°			58 2	
CuSO₄ 5H₂O—H₂O(¹⁸)	11 9	−1 6°			69 0	
ZnSO₄ 7H₂O—H₂O(⁵)	27 2	0 55°			50 9	
FeSO₄ 7H₂O—H₂O(⁵)	13 0	−1.8°			67 2	
	% of 66 19% H₂SO₄ 7 1		ice and acid at 0°	−16°	−2.1°†	68.6
	11 2		ice and acid at 0°	−20°	−3.1°†	62.0
	17 2		ice and acid at 0°	−24°	−5.5°†	52.9
66 19% H₂SO₄ H₂O (¹¹)	23 9		ice and acid at 0°	−28°	−9.5°†	43.0
	33 6		ice and acid at 0°	−32°	−16 5°†	24.5
	44 2		ice and acid at 0°	−36°	−30 2°†	7 5
	47 7		ice and acid at 0°	−37°	−37°†	0
HCl—H₂O	% HCl 24 8E	−86°				
	% of Na₂SO₄ 10H₂O 21 05		0°		6 09	
	30 33		0°		9 17	
	36 59		0°		11 15	
	37 69		21 2°	−8 1°		
	42 37		0°		13 15	
	50 22		21.6°	−12 2°		
	62 67		15°			21.2 at 0° / 12.0 at −15°
Na₂SO₄ 10H₂O—36 69% HCl (¹⁴)	62 96		21 6°	−15 3°		
	63 88		0°		28 89	
	74 64		15°			30.6 at 0° / 19.1 at −15°
	74 68		0°		30 85	
	75 30		21 5°	−14 8°		
	78 90		0°		27 43	
	86 63		15°			24.5 at 0° / 13 4 at −15°
	86 72		0°		19.44	
	88 53		20.1°	−15 6°		

(Salt separates — noted vertically in the CaCl₂6H₂O—H₂O section)

† Temperature when all ice is melted

Substances	Composition of mixture (% anhydrous salt, unless otherwise stated). E = eutectic composition	Freezing point of solution	Initial condition of freezing mixture	Lowest attained temperature recorded	Heat absorbed at temperature of mixing, cal per g of mixture	Heat absorbed (at freezing or saturation point of solution) from objects to be cooled, cal. per g of mixture. The ° values are heats of fusion of the eutectic, e (⁵)
Na₂SO₄.10H₂O—30 13% HCl (14)	% of Na₂SO₄ 10H₂O 46.04		19 7°	−11 8°		
	49.74		19 7°	−11 8°		
	63.46		19 7°	−14.4°		
	65.23		20 4°	−15 0°		
	75 43		20.0°	−14 8°		
	82 54		19 9°	−17.2°		
	86 31		20 0°	−12.6°		
	89 88		20 1°	ca 0°		
Na₂SO₄ 10H₂O—24 47% HCl (14)	% of Na₂SO₄ 10H₂O 35.54		0°		12 07	
	38.16		19.9°	8 2°		
	50 42		19 8°	10 0°		
	62 22		0°		26.84	
	63 86		20 5°	−12 0°		
	67 57		0°		27.18	
	71.46		0°		25 72	
	75 36		21.0°	−11 8°		
	78 40		0°		20 21	
C₂H₅OH—H₂O (16)	% alc 50	−37°	alc at 2° ice at 0°	−24 2°		
			alc at 1 5° ice at −1°	−29 4°		
	51 3	−38°	alc at 4° ice at 0°	ca −30°		
CS₂—(CH₃)₂CO	A temperature of −43 5° in a volume of 20 cc was maintained by mixing 100 cc of carbon disulfide and 70 cc of acetone per hour, using a heat interchanger (³).					

Salts	Temperature produced by mixing salts with water	Lit	Reduction of temperature produced by water with an equal weight of a salt or of a mixture of salts in equal parts (⁷)	Salts	Temperature produced by mixing salts with water	Lit.	Reduction of temperature produced by water with an equal weight of a salt or of a mixture of salts in equal parts (⁷)
NH₄Cl			14°	NaNO₃—KCNS	−37 4°	(1)	
NaCl			4°	KNO₃—NH₄CNS	−28.2°	(1)	
KCl			12°	NH₄Cl—NH₄NO₃—KNO₃	−22 6°	(9)	
NH₄NO₃			25°	NH₄Cl—NH₄NO₃—NaNO₃	−30.1°	(9)	
NaNO₃			9 5°	NH₄Cl—Na₂SO₄.10H₂O—KNO₃			17°-23°
KNO₃			10°	NH₄Cl—(NH₄)₂SO₄—K₂SO₄	−15.2°	(9)	
NH₄SO₄			8°	NH₄Cl—(NH₄)₂SO₄—Na₂SO₄.10H₂O	−19.9°	(9)	
Na₂SO₄.10H₂O			7 5°	NaCl 2H₂O—NaNO₃—KNO₃	−24 6°	(9)	
K₂SO₄			4 5°	KCl—KNO₃—K₂SO₄	−11.55°	(2)	
NH₄Cl—KNO₃	−18 2°	(9)	20°	NH₄NO₃—KNO₃—NaNO₃			16°-27°
NH₄Cl—NaNO₃	−31 5°	(9)	17°	NH₄NO₃—KNO₃—Na₂SO₄.10H₂O			17°-26°
NH₄Cl—NH₄NO₃			22°	NH₄NO₃—(NH₄)₂SO₄—Na₂SO₄.10H₂O	−19.5°	(9)	
NH₄Cl—Na₂SO₄.10H₂O	−17 6°	(9)	19°				
NH₄Cl—K₂SO₄	−18 0°	(9)					
NaCl—KNO₃			10°				
NaCl.2H₂O—KNO₃	−24 9°	(9)					
KCl—NaNO₃			11°				
KCl—NH₄NO₃			20°				
NH₄NO₃—KNO₃			22°				
NH₄NO₃—Na₂SO₄.10H₂O	−19.5°	(9)	26°				
Na₂NO₃—NaSO₄.10H₂O			10°				

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Brendel, Diss , Charlottenburg; 92. (²) Bruni, 56, 27, I: 537; 97. (³) Duclaux, 54, 151: 715, 10. (⁴) Gortner, 106, 39: 584; 14. (⁵) Gröber, Diss , Techn. Hochschüle, München; 08. (⁶) Hammerl, 75, 78: 59; 78. (⁷) Hanamann, 112, 173: 314; 64. (⁸) Kanolt, 48, 9: 416; 24. (⁹) Mazzotto, 72, 23: 545, 633; 90. (¹⁰) Morits, 156, 6: 1374; 82. (¹¹) Pfaundler, 75, 71: 509; 75. (¹²) Rüdorff, 8, 122: 337; 64. (¹³) Rüdorff, 8, 136: 276; 69. 26, 2: 68; 69. (¹⁴) Szydlowski, 75, 116: 855; 07. (¹⁵) Tollinger, 75, 72: 535; 75.

2. TEMPERATURES ABOVE 0°C

Olaf A. Hougen and Roland A. Ragatz

(a) *Bath Liquids or Vapor Baths with Boiling under Constant External Pressure.* For heterogeneous systems and solutions v. (13). For fire hazards on certain of these liquids v. p. 61. For a more extensive series of liquids arranged in order of boiling points v. p. 310.

Substance	Boiling point At 760 mm	At 100 mm	Actual range used	Lit
Ethyl chloride	12 2'	31 3°	13' to 30°	(23)
Ethyl ether	34 5°	-12 1°		(2, 11, 13)
Carbon disulfide	46 3'	4 8°	16° to -26°	(3, 11, 13, 26, 27, 31, 41)
Acetone	56 1'	7 5°		(13, 21)
Chloroform	61 2°	9 7°		(11, 21)
Methyl alcohol	64.5'	20 62'	65° to 49°	(2, 10, 11, 13, 21, 30)
Ethyl alcohol	78 5°	34 1°	78° to 40°	(2, 10, 11, 13, 21, 31)
Benzene	79 8'	25 8°	81° to 40°	(10, 11, 13, 39)
Water	100'	51 7'	145° to 25°	(2, 3, 9, 11, 13, 14, 18, 26, 27, 29, 30, 32, 43)
Toluene	110 5'	51 8'	130° to 70°	(10, 13, 21, 29, 32, 39, 45)
Chlorobenzene	132 1'	70 3°	132° to 70°	(31, 39)
m-Xylene	139 0°	77 8°	140° to 70°	(10, 21, 28, 32, 39, 45)
Isoamyl acetate	142 5'		141° to 119°	(30, 45)
Bromobenzene	156.2°	90 7°	160° to 120°	(28, 31)
Aniline	184.1°	119 1°	184° to 150°	(27, 31, 32, 39, 42, 44)
Ethyl benzoate	213 2°	142°		(21, 27, 45)
Naphthalene	217 9'	144 3°		(28, 39)
Methyl salicylate	223 3'	151°	223° to 175°	(31)
Quinoline	237 7°	166 7'	208° to 170°	(18, 21, 39, 45)
Isoamyl benzoate	262°			(21, 28, 45)
α-Bromonaphthalene	281 1°	198 8°	281° to 215°	(28, 31)
Diphenylamine	302 0°	221°		(5, 18, 28, 39, 45)
Benzophenone	305 4°	224°	306° to 257°	(28, 39)
Mercury	356 9°	261 5'	Various ranges	(2, 8, 31, 39)
Sulfur	444 6°	330 7'	Various ranges	(2, 8, 8, 39)
Phosphorus pentasulfide	52°			(5)
Zinc	907°	758°		(2)

(b) *Solid-liquid Non-variant Points* 1. Ice-water, v (11, 24, 29, 46). 2. Transformation temperatures of crystalline hydrates.

Salt	Hydration temperature °C	Lit
Sodium chromate	19 71	(12, 33)
Sodium sulfate	32 383	(11, 12, 32, 33, 34, 35)
Sodium carbonate	35 3	(12, 33)
Sodium thiosulfate	48 0	(12, 33)
Sodium bromide	50 8	(12, 33)
Manganese chloride	57 8	(12, 33)
Trisodium phosphate	73 1	(12, 33)
Barium hydroxide	78 0	(12, 33)

(c) *Bath Liquids with Thermostatic Control*

Liquid	Useful range	Lit
Water	0° to 90°	(17, 18, 21, 40)
Mineral oils	To 20° below the flash point	(5, 19, 22, 37, 38, 40)
Paraffin	M.P. to 300°	(5, 27, 29, 40)
10 parts cottonseed oil, 1 part beeswax	M.P. to 300°	(7)
Hydrogenated sesame oil	60° to 300°	(36)
Hydrogenated cottonseed oil	60° to 285°	(36)

Fused salts	Melting point	Lit.
NaNO₃ (45%), KNO₃ (55%).	218°	(8, 14, 21, 32, 44)
NaNO₃ (55%), NaNO₂ (45%)	221°	(44)
KNO₃	337°	(1)
NaCl (28%), CaCl₂ (72%)	500°	(44)
NaCl (50%), K₂CO₃ (50%)	560°	(44)
Na₂CO₃ (50%), KCl (50%)	560°	(44)
CaCl₂ (50%), BaCl₂ (50%)	600°	(44)
NaCl (35%), Na₂CO₃ (65%)	620°	(44)
NaCl (22%), BaCl₂ (78%)	654°	(44)
NaCl (44%), KCl (56%)	663°	(44)

Molten metals	Useful range	Lit.
Lead	327° to 700°	(4, 5, 6, 29)
Lead (30%), Tin (70%)	Above 183°	(14)
Lead (50%), Tin (50%)		(5)

Other liquids	Useful range	Lit.
Naphthalene	80° to 217°	(20, 21, 25)
Benzophenone	49° to 305°	(20, 21, 25)
Sulfur	113° to 444°	(20, 25)

(d) *Metal Blocks*—Aluminum and copper blocks have been used up to 600°, with a uniformity of temperature of 1° (39).

(e) *Gas Baths and Furnaces.*—For temperatures above 900°, an electrically heated gas bath is usually employed, although for the higher temperatures a bath material is not essential since heat transfer takes place primarily by radiation. For lower temperatures, heat transfer and temperature uniformity are promoted by packing with a granular non-oxidizing metal.

The following references (compiled by the Geophysical Laboratory) deal with the construction and temperature regulation of high temperature furnaces: Kolovrat, 51, 8: 495; 09. Haughton and Hanson, 47, 14: 145; 15. 18: 173; 17. White and Adams, 2, 14: 44; 19. Haugn, 101, 40: 670; 19. Roberts, 128, 11: 409; 21. 48, 6: 965; 22 Bunting, 38, 6: 1209; 23. Adams, 48, 9: 599; 24. Roberts, 48, 10: 723; 25.

LITERATURE

(For key to the periodicals see end of volume)

(1) Aten, 7, 78: 13, 12 (2) Barus and Hallock, 156, No 84; 89. (3) Brown, 5, 7: 411, 79 (4) Bodenstein, 7, 29: 665, 99 (5) Bodenstein, 7, 30: 113; 99 (6) Bodenstein, 7, 30: 125, 99 (7) Bosart, 1, 31: 724; 09. (8) Day and Sosman, 8, 38: 849, 853, 12. (9) Dupre, 175, 38: 308; 13. (10) Forster, 155, 106: 80, 12. (11) Freas, Thesis, Chicago; 11. (12) Geer, 50, 6: 85; 02. (13) Golodetz, 136, 38: 1253, 14. (14) Goodwin and Mailey, 2, 25: 469, 07 (15) Gordon, 7, 28: 305; 99 (16) Grutzmacher, Deutsch. Mech.-Ztg 1902: 193 (17) Grutzmacher, Deutsch Mech.-Ztg 1902: 184 (18) Grutzmacher, 89, 3: 248, 260, 00 (19) Holborn and Henning, 8, 23: 810, 07 (20) Holborn and Henning, 8, 26: 860, 08. (21) Holborn, Scheel and Henning, B63 (22) Holborn and Schultze, 8, 47: 1101; 15. (23) Jenkin, 85, 18: 197, 22. (24) Marshall, 83, 7: 249; 11 (25) Meisner, 8, 39: 1230; 12. (26) Meyer, 15, 165: 303, 73 (27) Moser, 92, 34: 625, 21. (28) Noyes, 152, No 63: 12, 73, 194, 210, 07. (29) Ostwald-Luther, B64, p 100. (30) Pomplun, 245, 11: 1, 91. (31) Ramsay and Young, 4, 47: 640, 85. (32) Richards, 45, 4: 910; 12 (33) Richards and Churchill, 7, 28: 313; 99. (34) Richards and Mark, 65, 38: 417; 02. (35) Richards and Wells, 65, 38: 431, 02 (36) Robertson, 45, 18: 701, 23. (37) Rothe, 245, 19: 144; 99. (38) Shaw, 69, 11, III: 129, 17 (39) Stähler, B65, 1: 501. (40) Stähler, B65, 1: 498. (41) Stock, Henning and Kuss, 25, 54: 1119; 21. (42) Sudborough, 54, 18: 16, 99 (43) Thiesen, Scheel and Sell, 89, 2: 140; 95. (44) Tour, 218, 6: 171, 24, (45) Wiebe and Böttcher, 245, 10: 16; 90. (46) Washburn and Williams, 1, 35: 741, 13.

MAXIMUM TEMPERATURES THAT CAN BE REACHED AND MAINTAINED FOR OBSERVATIONAL PURPOSES BY VARIOUS MEANS

W. E. FORSYTHE

	Maximum temperature, °C
Electric furnaces operating in open air	
Iron tube or iron wire wound furnace	500
Nicrome wound refractory tube	800
Platinum wound refractory tube—double winding (2)	1530
Iridium tube	1900
Carbon resistor furnace	2200
Carbon arc furnace	3200
Electric furnaces operating in vacuo or inert gas	
Tungsten wound refractory tube limited by refractory tube	2000
Carbon tube furnace	2700
Tungsten tube furnace (in vacuo)	2200
Tungsten tube furnace (in inert gas)	2800
Gas-fired furnaces	
Special makes of furnaces(5) with flames entering the furnace in tangential direction so as to give a good distribution of the heat, if gas and air are well mixed, can be raised up to about	1700

	Maximum temperature °C
The regenerative furnaces, such as are used in open hearth steel furnaces, can be heated up to about the same temperature of	1700
Special furnaces and methods	
High-frequency induction furnace. Limited only by melting point of refractory or metal used	
Filament in vacuum or inert gas limited only by rate of vaporization or melting point of filament used	
Arc under pressure	
Carbon (4)	5790
Tungsten (3)	4785
Exploding fine wires by discharging a condenser charged to high voltage through them gives a temperature up to about (1)	19700

LITERATURE

(For a key to the periodicals see end of volume)

(1) Anderson, *21*, **51**: 37; 20. (2) Hyde and Forsythe, *21*, **51**: 247; 20. (3) Luckey, *2*, **9**: 134, 17. (4) Lummer, *254*, **21**: 8, 17. (5) McCauley, Corning Glass Works, Corning, N Y, O Spencer, National Lamp Works, Nela Park, Cleveland, Ohio, O.

LABORATORY METHODS FOR MAINTAINING CONSTANT HUMIDITY

HUGH M. SPENCER

A saturated aqueous solution in contact with an excess of a definite solid phase at a given temperature will maintain a constant humidity within any enclosed space around it. By properly selecting the salt to be used almost any desired degree of humidity can be secured and controlled in this way. A number of salts suitable for this purpose are displayed in the accompanying chart and tables, together with the % humidity prevailing above their saturated solutions at different temperatures. To convert "% humidity" into "aqueous tension" multiply it by the vapor pressure of pure water at the same temperature.

SOLID PHASE

1. CaCl₂ 2H₂O (19)
2. CoCl₂.6H₂O (8).
3. CoSO₄ 6H₂O (7).
4. CuCl₂.2H₂O (8, 13, 22).
5. CuSO₄.5H₂O (11, 16).
6. K₂C₄H₄O₆.½H₂O (4).
7. KCl (4, 5, 9, 18, 21).
8. KClO₃ (5, 11, 16).
9. KNO₃ (4, 5, 9, 16).
10. K₂SO₄ (4, 5, 15, 20).
11. MgCl₂.6H₂O (8, 13)
12. MgSO₄.6H₂O (7).
13. MnSO₄ H₂O (7).
14. NH₄NO₃ (9, 18).
15. NaCl (4, 5, 18, 21).
16. Na₂CO₃.H₂O (10, 22)
17. Na₂C₄H₄O₆.2H₂O (14)
18. NaKC₄H₄O₆.4H₂O (14).
19. NaNO₃ (4, 5, 9, 18, 21).
20. Na₂SO₄ (4, 16, 24, 26)

Solid phases	t, °C	% humidity	Lit
BaCl₂.2H₂O	24 5	88	(15)
CaCl₂.6H₂O	5	39 8	(20)
	10	38	(19)
	18 5	35	(15)
	20 0	32 3	(19)
	24 5	31	(15)
Ca(NO₃)₂.4H₂O	18 5	56	(15)
	24 5	51	(15)

Solid phases	t, °C	% humidity	Lit.
CaSO₄.5H₂O	20	98	(15)
CrO₃...	20	35	(15)
H₂C₂O₄.2H₂O	20	76	(15)
H₃PO₄.½H₂O	24 5	9	(15)
KC₂H₃O₂	20	20	(15)
	168	13	(11)
KBr	20	84	(15)
	100	69.2	(5)

Solid phases	t, °C	% humidity	Lit.
K₂CO₃.2H₂O	18.5	44	(15)
	24.5	43	(15)
KCNS	20	47	(15)
K₂CrO₄	20	88	(15)
KF	100.0	22.9	(5)
K₂HPO₄	20	92	(15)
KHSO₄	20	86	(15)
KI	100.0	56.2	(5)
KNO₂	20	45	(15)
LiCl.H₂O	20	15	(15)
Mg(C₂H₃O₂)₂.4H₂O	20	65	(15)
Mg(NO₃)₂.6H₂O	18.5	56	(15)
NH₄Cl	24.5	52	(15)
	20.0	79.2	(9)
	25.0	79.3	(9)
	30.0	79.5	(9)
NH₄Cl and KNO₃	20.0	72.6	(9)
	25.0	71.2	(9)
	30.0	68.6	(9)
NH₄H₂PO₄	20.0	93.1	(9)
	25.0	93.0	(9)
	30.0	92.9	(9)
(NH₄)₂SO₄	20.0	81.0	(9)
	25.0	81.1	(9)
	30.0	81.1	(9)
	108.2	75	(11)
NaBr	100.0	22.9	(5)
NaBr.2H₂O	20	58	(15)
NaBrO₃	20	92	(15)
NaCl and KClO₃	16.39	36.58	(6)
NaCl and KNO₃	16.39	32.57	(6)
NaCl, KNO₃ and NaNO₃	16.39	30.49	(6)
NaC₂H₃O₂.3H₂O	20	76	(15)
Na₂CO₃.10H₂O	18.5	92	(15)
	24.5	87	(15)
NaClO₃	20	75	(15)
	100.0	54	(5)

Solid phases	t, °C	% humidity	Lit.
Na₂Cr₂O₇.2H₂O	20	52	(15)
NaF	100.0	96.6	(5)
Na₂HPO₄.12H₂O	20	95	(15)
NaHSO₄.H₂O	20	52	(15)
NaI	100.0	50.4	(5)
NaNO₂	20	66	(15)
Na₂SO₃.7H₂O	20	95	(15)
Na₂S₂O₃.5H₂O	20	78	(15)
Na₂SO₄.10H₂O	20	93	(15)
Pb(NO₃)₂	20	98	(15)
	103.5	88.4	(11)
TlCl	100.097	99.7	(4)
TlNO₃	100.317	98.7	(4)
Tl₂SO₄	104.7	84.8	(4)
ZnCl₂.1½H₂O*	20	10	(15)
Zn(NO₃)₂.6H₂O	20	42	(15)
ZnSO₄.7H₂O	5	94.7	(20)
	20	90	(15)

* Unstable at this temperature.

LITERATURE

(For a key to the periodicals see end of volume)

(1) Badger and Baker, 53, 23: 569; 20. (2) Baker and Waite, 53, 25: 1137; 21. (3) Baker and Waite, 53, 25: 1174, 21. (4) Berkeley and Appleby, 5, 85A: 489, 11. (5) Bronsted, 7, 82: 633; 03. (6) Bronsted, 137, 1: 5; 18. (7) Carpenter and Jette, 1, 45: 578, 23. (8) Derby and Ingve, 1, 38: 1439; 16. (9) Edgar and Swan, 1, 44: 570; 22.
(10) Gerasimov, 53, 45: 1666; 13. (11) Gerlach, 91, 26: 413; 87. (12) Lescoeur, 34, 103: 1260, 86. (13) Lescoeur, 6, 2: 78; 94. (14) Lowry and Morgan, 1, 46: 2192; 24. (15) Obermuller, 7, 109: 145; 24. (16) Pawlowitsch, 7, 84: 169, 13. (17) Prideaux, 54, 39: 182; 20. (18) Rodebush, 1, 40: 1204; 18. (19) Roozeboom, 7, 4: 31, 89.
(20) Sidgwick and Ewbank, 4, 125: 2268, 24. (21) Speranski, 7, 70: 519; 10. (22) Speranski, 7, 78: 86; 12. (23) Speranski, 7, 84: 166; 13. (24) Tammann, 8, 24: 530, 85. (25) van't Hoff, 7, 45: 288, 03. (26) Wuite, 7, 86: 369; 14.

BAROMETRY AND MANOMETRY

H. H. KIMBALL

1. *Gravity Correction.*—The equivalent barometric, or other manometric, height (B_s) corresponding to standard gravity ($g_s = 980.665$ cm sec⁻²) is related to the height (B_l) corresponding to local gravity (g_l) as shown by equation (1):

$$B_s = B_l \frac{g_l}{g_s} = B_l + C_g; \quad C_g = B_l \frac{g_l - g_s}{g_s} \quad (1)$$

When g_l and g_s are expressed in cm sec⁻²,

$$C_g = B_l \left[\frac{(g_l - g_s)(1.0197)}{1000} \right]$$

Any desired unit may be used for B_l; C_g and B_s are in the same unit as B_l. [For most barometric purposes, a sufficiently accurate correction (within ±0.01% of B_l) is obtained by the use of the approximate correction $C_g' = B_n \frac{g_l - g_s}{g_s}$, in which B_n is the usual barometric pressure at the station]

Example: $B_l = 29.851$, $g_l = 978.053$ cm sec⁻². Then $(g_l - g_s) = -2.612$ cm sec⁻²; $0.0197(g_l - g_s) = -0.0515$ cm sec⁻²; $1000\, C_g = -2.663 B_l = -79.49$. ∴$B_s = 29.851 - 0.079 = 29.772$.

2. *Temperature Correction.*—The equation by which the equivalent barometric, or other manometric, height (B) at the standard temperature (t_m) can be computed from the nominal height (B') at the temperature t, is generally written in the form

$$B = B' + C_t; \quad C_t = B' \frac{l(t - t_s) - m(t - t_m)}{1 + m(t - t_m)} \quad (2)$$

where t_m = standard temperature of the manometric liquid, t_s = temperature at which the scale, after correction for errors of graduation, reads correctly, m = coefficient of cubical expansion of the manometric liquid, l = coefficient of linear expansion of the material on which the scale is engraved.

The value of m which is generally used for mercury, and which has been adopted by the International Meteorological Tables, is $m = 181.8 \times 10^{-6}$ per °C. For temperatures between 0°C and 30°C this value appears (5, 6, 8, 15, 17) to be correct within ±0.1 × 10⁻⁶ per °C. The value of l, for brass, which has been adopted by the International Meteorological Tables, is $l = 18.4 \times 10^{-6}$ per °C. The best determinations (1, 2, 11) of this coefficient for temperatures between 0° and 30° yield values varying from

17.5 × 10⁻⁴ per °C to 19.3 × 10⁻⁴ per °C, or by ± 5%. For glass scales the approximate value l = 8.5 × 10⁻⁴ per °C is usually satisfactory. (For silicate flint glasses [13] l varies from 7.88 × 10⁻⁴ per °C to 9.35 × 10⁻⁴ per °C; for crown glasses [13] it varies from 6.75 × 19⁻⁴ to 9.54 × 10⁻⁴ per °C.

For barometers with metric scales, the combined effect of an error of ±0.1 × 10⁻⁴ per °C in m and of ±0.9 × 10⁻⁴ per °C in l

will cause an error in C_t of $\pm \dfrac{B't \times 10^{-4}}{1 + mt}$. For t = 30°C and B' = 760 mm, the error would be ±0.023 mm; while for t = 10°C, B' = 100 mm, it would be only ±0.001 mm. At ordinary room temperatures, the error so produced in C_t will be less for barometers graduated in inches than for one graduated in millimeters. (For barometers graduated in inches t_m = 62°F, t_m = 32°F).

TABLE 1.—TEMPERATURE CORRECTION (C_t) FOR MERCURIAL MANOMETERS AND BAROMETERS

$B = B' + C_t$; (B' = nominal height at $t°$; B = equivalent height for mercury at 0°C; B, B', and C_t are all in the same unit, which may be anything desired)

A. Brass scale correct at 62°F, inches, °F; t_m = 32°F, t_s = 62°F, m = 181.8 × 10⁻⁴ per °C, l = 18.4 × 10⁻⁴ per °C
(Applies directly to commercial barometers graduated in inches)

t(°F) \ B'	10	20	30	40	50	60	70	80	90
+12	+0 015	+0 030	+0 045	+0 061	+0 076	+0 091	+0 106	+0 121	+0 136
22	+0.006	+0.012	+0 018	+0 024	+0 030	+0 036	+0 042	+0 048	+0 054
32	−0 003	−0 006	−0 009	−0 012	−0 015	−0 018	−0 021	−0 024	−0.028
42	−0 012	−0 024	−0 036	−0 049	−0 061	−0 073	−0 085	−0 097	−0.109
52	−0.021	−0 042	−0 064	−0 085	−0 106	−0 127	−0 148	−0 169	−0 191
62	−0 030	−0 060	−0 091	−0 121	−0 151	−0 181	−0 211	−0 242	−0.272
72	−0 039	−0 078	−0 118	−0 157	−0 196	−0 235	−0 275	−0 314	−0.353
82	−0.048	−0 096	−0 145	−0 193	−0 241	−0 289	−0 338	−0 386	−0.434
92	−0.057	−0 114	−0 172	−0 229	−0 286	−0 343	−0 400	−0 458	−0.515

B. Brass scale correct at 0°C, millimeters, °C; t_m = t_s = 0°C, m = 181.8 × 10⁻⁶ per °C, l = 18.4 × 10⁻⁶ per °C

t(°C) \ B'	100	200	300	400	500	600	700	800	900
−10	+0 16	+0 33	+0 49	+0 65	+0 82	+0 98	+1 15	+1 31	+1 47
− 5	+0 08	+0 16	+0 25	+0 33	+0.41	+0 49	+0 57	+0 65	+0.74
0	0 00								
+ 5	−0 08	−0 16	−0 24	−0 33	−0 41	−0 49	−0.57	−0 65	−0.73
10	−0 16	−0 33	−0 49	−0 65	−0 82	−0 98	−1 14	−1 30	−1.47
15	−0 24	−0 49	−0 73	−0 98	−1 22	−1 47	−1 71	−1 96	−2 20
20	−0 33	−0 65	−0 98	−1 30	−1 63	−1 95	−2 28	−2 60	−2 93
25	−0 41	−0 81	−1 22	−1 63	−2.03	−2 44	−2 85	−3 25	−3.66
30	−0 49	−0 98	−1 46	−1 95	−2 44	−2 93	−3 41	−3 90	−4.39
35	−0 57	−1 14	−1 70	−2 27	−2 84	−3 41	−3 98	−4 55	−5.11
40	−0 65	−1 30	−1 95	−2 60	−3 24	−3 89	−4 54	−5 19	−5 84

C. Glass scale correct at 0°C, t_m = t_s = 0°C, m = 181.8 × 10⁻⁶ per °C, l = 8.5 × 10⁻⁶ per °C

t(°C) \ B'	100	200	300	400	500	600	700	800	900
−10	+0 17	+0 35	+0 52	+0 69	+0 87	+1 04	+1 22	+1 39	+1.56
− 5	+0.09	+0 17	+0 26	+0 35	+0 43	+0 52	+0 61	+0 69	+0.78
0	0 00								
+ 5	−0 09	−0 17	−0 26	−0 35	−0 43	−0 52	−0 61	−0 69	−0 78
10	−0 17	−0 35	−0 52	−0 69	−0 86	−1 04	−1 21	−1 38	−1.56
15	−0 26	−0 52	−0 78	−1 04	−1 30	−1 56	−1 81	−2 07	−2.33
20	−0 34	−0 69	−1 04	−1 38	−1 73	−2 07	−2 42	−2.76	−3.11
25	−0 43	−0 86	−1 29	−1 73	−2 16	−2 59	−3 02	−3 45	−3.88
30	−0 52	−1 03	−1 55	−2 07	−2 59	−3 10	−3.62	−4 14	−4.65
35	−0 60	−1 21	−1 81	−2 41	−3.01	−3.62	−4 22	−4.82	−5.42
40	−0 69	−1 38	−2 06	−2 75	−3 44	−4 13	−4 82	−5 51	−6.19

Example: Barometer graduated in inches, brass scale correct at 62°F; B' = 29.564 in., t = 76.8°F. From section A it is found that at 72°, C_t for B' = 29.564 is −0.1155, at 82° it is −0.1421; hence at 76.8°, C_t = $-0.1155 + \dfrac{4.8}{10}(-0.0266)$ = −0.1155 − 0.0128 = −0.128. Hence B = 29.564 − 0.128 = 29.436 in.

3. Capillary Corrections.—The curvature of the surfaces of the manometric liquid introduces pressures directed towards the centers of curvature of the surfaces. For each surface, this pressure is

$\gamma\left(\dfrac{1}{r_1} + \dfrac{1}{r_2}\right)$ dynes cm⁻² = $\dfrac{\gamma}{dg}\left(\dfrac{1}{r_1} + \dfrac{1}{r_2}\right)$ cm of the manometric liquid. [γ = surface tension (in dynes cm⁻¹), d = density of the liquid (in g cm⁻³), g is the acceleration of gravity (in cm sec⁻²), and r_1 and r_2 are the principal radii of curvature (in cm) of the surface at the point considered.] At the vertex of the meniscus in a tube of circular section, r_1 = r_2 = r, and if the angle of contact of the liquid with the tube is either 0° or 180°, and if the tube is not too large, r is practically equal to the internal radius of the tube. If

the liquid surface is in an annular space between coaxial, circular cylinders (as in the reservoir of a Fortin barometer), if the angle of contact is 0°, and if neither r_1 nor $(r_3 - r_2)$ is very great as compared with the capillary constant, (18), then $h' = \dfrac{2dhr_1}{(r_3 - r_2)^2}$, approximately; h' and h are the respective capillary pressures (in terms of unit column of the liquid) at the vertices of the surfaces in the annular space of width $(r_3 - r_2)$, and in a tube of radius r_1; and d is the depth of the annular meniscus

Laplace (12) has shown that, except for sign, the equations for a convex meniscus are the same as those for a concave one. Hence, this expression can probably be accepted as a first approximation to the value for h' for any liquid, provided that the angle of contact of the liquid with the solid is the same at all three surfaces, and that r_1 and $(r_3 - r_2)$ are not too great. In the case of the ordinary mercurial cistern barometers, $(r_3 - r_2)$ is quite large as compared with the capillary constant of mercury, and the angles of contact may not be the same at all three surfaces; for these reasons, no great confidence can be placed in the actual value of h', as so computed, for such barometers, but its order of magnitude will probably be correct.

TABLE 2.—CAPILLARY DEPRESSION OF THE APEX OF A MERCURIAL COLUMN IN A GLASS TUBE OF CIRCULAR SECTION*

Depression in millimeters

Radius of the tube, mm	Height of the meniscus, mm								
	0 2	0 4	0 6	0 8	1 0	1 2	1 4	1 6	1 8
1 0	2 46	4 40							
1 4	1 26	2 36	3 22						
1 8	0 75	1 44	2 02	2 48					
2 2	0 49	0 95	1 36	1 70	1 98				
2 6	0 34	0 66	0 96	1 22	1 44	1 61			
3 0	0 24	0 48	0 70	0 90	1 07	1 21	1 32		
3 5	0 17	0 34	0 49	0 64	0 76	0 87	0 96	1 04	
4 0	0 12	0 24	0 35	0 46	0 56	0 61	0 71	0 77	0 82
4 5	0 09	0 18	0 26	0 34	0 41	0 47	0 53	0 58	0 62
5 0	0 07	0 13	0 19	0 25	0 30	0 35	0 40	0 44	0 47
5 5	0 05	0 10	0 14	0 19	0 23	0 27	0 30	0 33	0 36
6 0	0 04	0 07	0 11	0 14	0 18	0 20	0 23	0 25	0 27
6 5	0 03	0 06	0 09	0 11	0 14	0 16	0 18	0 20	0 21
7 0	0 02	0 04	0 06	0 08	0 10	0 12	0 14	0 15	0 16

* From the Schleiermacher-Delcros (4, 9, 10) table, as revised by Süring (14). The values are about 5% larger than those obtained from Bravais's (3) table, in which the arguments are the diameter of the tube, and the angle of incidence of the meniscus of the mercurial column with the walls of the tube.

Example In a barometer cistern for which $r_2 = 6$ mm, $r_3 = 16$ mm, d was found to be 0.5 mm ; the radius of the barometer tube was $r_1 = 5$ mm, and the height of the meniscus in it was 1.0 mm. From Table 2 it is found that the depression h, due to the meniscus in the 5 mm tube, is 0.30 mm; hence $h' = 0.015$ mm. That is, the pressure due to the annular surface is of the order of 0.02 mm; and the total depression of the column is $H = 0.30 - 0.02 = 0.28$ mm, subject to the uncertainty regarding the actual value of h'.

4. *Possible Residual-gas Error in Good Barometers.*—Under ordinary laboratory conditions, errors amounting to as much as 4.1 mm (0.163 in.) have been observed, and errors of 1.1 mm (0.043 in.) are not uncommon; but in most barometers, this error

does not exceed 0.25 mm (0.010 in.) when the instrument is shipped by the manufacturer. Air may be introduced during shipment and by handling. The smaller the tube of the barometer, the more likely is the error to be large. The magnitude of the error varies with the temperature and with the volume of the space above the mercury column, as indicated by equation (3):

$$x = x_0 \frac{V_0}{V}[1 + 0.00367(t - t_0)] \qquad (3)$$

where x_0 and x are, respectively, the errors corresponding to the volume V_0 temperature t_0, and to the volume V temperature t; temperatures being expressed in °C.

5. *Conversion of Water Column at $t°C$ to the Equivalent Water Column at 4°C.*—If h_t and h_4 are the equivalent true heights (corrected for scale errors of graduation and expansion, and for capillary pressures), and if d_t and d_4 are the respective densities (7, 16) then, if $\delta = (d_4 - d_t)/d_4$, $h_4 = h_t(1 - \delta)$.

TABLE 3.—VALUES OF 100δ

t (°C)	Units of t				
	0	2	4	6	8
tens					
0	0 013	0 003	0 000	0 003	0 012
1	0.027	0 048	0 073	0.103	0.138
2	0.177	0.221	0 268	0 320	0.375
3	0.435	0 497	0.563	0 633	0.706

Example.—$h_{25} = 67.53$ cm. At 25°, $100\delta = 0.294$. $\therefore \delta h_{25} = 0.199$, $h_4 = h_{25}(1 - \delta) = 67.53 - 0.20 = 67.33$ cm.

6. *Conversion of Water Column at 4°C to Equivalent Mercury Column at Standard Density* (13.5951 $g\ cm^{-3}$); *and the Reverse.*— If h_w and h_m are the equivalent true heights (corrected for the scale errors of graduation and expansion, and for capillary effects) of the water and the mercury, respectively, $h_m = 0.073554h_w$.

TABLE 4.—EQUIVALENT COLUMNS OF WATER (h_w) AND OF MERCURY (h_m)

(Density of water = 0.999973 $g\ cm^{-3}$; of mercury = 13.5951 $g\ cm^{-3}$)

h_w	h_m	h_w	h_m	h_m	h_w	h_m	h_w
100	7.3554	600	44 132	1	13 5955	6	81.573
200	14 7108	700	51 488	2	27 1909	7	95 168
300	22 0662	800	58.843	3	40 7864	8	108.764
400	29 4216	900	66 199	4	54 3818	9	122.359
500	36 7770	1000	73 554	5	67 9773	10	135 955

LITERATURE

(For a key to the periodicals see end of volume)

(1) Bein, *88*, 14: 1113; 12   (2) Benoit, *238*, 6: 190, 88.   (3) Bravais, *6*, 5: 492, 42   (4) Bravais and Martins, *239*, 14: 47, 41   (5) Broch, *238*, 2: 21, 83   (6) Chappuis, *238*, 13: 28, 07   (7) Chappuis, *238*, 13D: 39; 07. (8) Chappuis, *238*, 16: 31, 17   (9) Delcros, Annaire Météorologique de la France, 169–170; 49. (10) Delcros, *240*, 6: 3; 18   (11) Dittenberger, *98*, 46: 1535; 02.   (12) Laplace, Mécanique Céleste (Bowditch translation) 4: 737   (13) Pulfrich, *8*, 45: 661; 92   (14) Süring, Ber u d Tätigk d Kgl. Pruss Meteor. Inst., 24–42; 16   (15) Thiessen, *89*, 4: 4, 04   (16) Thiessen, Scheel and Dusselhorst, *89*, 3: 68; 00   (17) Thiessen, Scheel and Sell, *89*, 2: 180; 95.   (18) Verschaffelt, *168*, No 32. *64V*, 24: 175; 88.

PSYCHROMETRY; DENSITY OF MOIST AIR; CHANGE IN BAROMETRIC PRESSURE WITH ALTITUDE

F. W. J. Whipple

$B; B_h$ — Barometric pressure, in general; at h

C — Instrumental constant

$d; d_h; d_o$ — Density of air, in general; at h; at T_o and A_n

$e; e'$ — Pressure of water vapor, present; when in equilibrium with water (or ice) at temperature t'

$g; g_*$ — Acceleration of gravity, actual; standard value

$h; H$ — Altitude above sea level, cm; meters

$t; t'$ — Readings of dry bulb; of wet bulb

$T; T_o; T'$ — Absolute temperatures in °C, general; of ice point, "virtual"

x — Ratio (mass of vapor)/(mass of dry air)

1. Psychrometry.—The pressure of the water vapor contained in the air is commonly deduced from the simultaneous readings of wet bulb and of dry bulb thermometers. The difference in these two readings depends upon the heat received by radiation as well as upon that furnished directly by the air. When the air flow is slow, the radiation is an important factor. In the Assmann psychrometer the bulb is surrounded by a double metal sheath; this largely eliminates radiation effects. It is important to secure adequate ventilation by the use of a thermometer with a bulb much smaller than the sheath. The standard bulb is 12 mm long and 4 mm in diameter. Alternatively, the thermometers may be "slung," *i.e.*, whirled on a suitable holder. In this case, direct radiation from sun or sky should be avoided as it affects the dry-bulb readings and therefore the psychometric difference.

The general formula for the computation of vapor pressure is

$$e' - e = CB(t - t') \times 10^{-4}$$

B, e, and e' are expressed in the same units, which may be anything desired. Within the order of accuracy of psychrometer observations, C is constant for a given velocity of the air-flow past the wet bulb. The relation of C to the air velocity has not been determined very precisely. The variation of C with temperature is negligible. If temperatures are expressed in °C, the value of C for thermometers with adequate ventilation (a relative velocity of 3 m per second or more) is 6.6 when the cover of the wet-bulb is saturated with water. On theoretical grounds, a lower factor, 5.8, is appropriate for an ice-covered bulb, but in the tables in general use 6.6 is adopted in this case as well. (Aspirations Psychrometer Tafeln, Braunschweig. 1908. Ferrel, Report of Chief Signal Officer, p. 218. Washington, 1886.) For the reduction of the readings of thermometers exposed in a Stevenson screen, Regnault's values of C, 8 for water and 7 for ice, are generally recommended (Études sur l'Hygrométrie, p. 102. Paris, 1845.) As, however, the ventilation is indeterminate, the accuracy obtainable is of a lower order.

Relative Humidity is computed by expressing e, determined by the psychrometric formula, as a percentage of the pressure of vapor in equilibrium with water (not ice) at the temperature of the dry bulb.

2. Density of Moist Air*

$$T, T_* = \text{absolute temperature in °C}$$

*If d_w/d_a = density of vapor and of dry air at same pressure and temperature, d_w/d_a = 0.6217 and $(d_a - d_w)/d_a$ = 0.3783.

Pressure unit	d
Any unit	$\dfrac{d.T.}{T}\left(\dfrac{B - 0.3783e}{A_n}\right);$ $\dfrac{d.T.B}{TB_*}\left(\dfrac{0.6217(1 + x)}{0.6217 + x}\right)$
Mm Hg	$\dfrac{464.6}{10^5}\left(\dfrac{B - 0.3783e}{T}\right)$ g/cm³; $\dfrac{288.9}{10^6}\left(\dfrac{B(1 + x)}{(0.6217 + x)T}\right)$ g/cm³
Kilodynes per cm²	$\dfrac{348.5}{10^6}\left(\dfrac{B - 0.3783e}{T}\right)$ g/cm³; $\dfrac{216.7}{10^6}\left(\dfrac{B(1 + x)}{(0.6217 + x)T}\right)$ g/cm³

$$x = \frac{\text{mass of vapor}}{\text{mass of dry air}} = \frac{0.6217\,e}{B - e}$$

Tables in Dictionary of Applied Physics 3: 76, and in paper by Shaw and Fahmy in Quart. J. Roy. Meteorological Soc., 1925, 210

$$\text{Specific humidity} = \frac{\text{mass of vapor}}{\text{total mass}} = \frac{0.6217\,e}{B - 0.3783\,e}$$

3. Relations Connecting Pressure and Altitude. −V. Bjerknes defines "virtual" temperature (T') as $T' = TB/(B - 0.3783e)$.

$$\frac{dB}{B} = d(\log_e B) = -\frac{gd}{B}dh = -0.03416\frac{g}{g_*}\cdot\frac{dH}{T'} = -\frac{g}{29.26\,g_*}\cdot\frac{dH}{T'} \quad (1)$$

$$d(\log_{10} B) = -\frac{0.014842}{g_*}\frac{g\,dH}{T'} = -\frac{g}{67.38\,g_*}\cdot\frac{dH}{T'} \quad (2)$$

If suffix $_1$ refers to the lower station and $_2$ to the upper, then

$$\log_{10}\frac{B_1}{B_2} = 0.014842\frac{g}{g_*}\cdot\frac{2(H_2 - H_1)}{T'_1 + T'_2}, \text{ approx.} \quad (3)$$

$$B_1 = B_2\left[1 + 0.03416\frac{g}{g_*}\cdot\frac{2(H_2 - H_1)}{T'_1 + T'_2 - 0.03416(H_2 - H_1)\frac{g}{g_*}}\right], \text{ approx.} \quad (4)$$

$$H_2 - H_1 = \frac{29.26\,g_*}{g}\cdot\frac{B_1 - B_2}{B_1 + B_2}(T'_1 + T'_2), \text{ approximately.} \quad (5)$$

For $(H_2 - H_1)$ not exceeding 1000 m, equations (4) and (5) are equivalent to the logarithmic formula. The factor $g/g_* = (1 - 0.002640 \cos 2\phi)(1 - 3.14H \times 10^{-7})$ may generally be taken as unity. The distinction between virtual and actual temperature may be ignored except when high temperatures are involved.

In the determination of heights in an extended barometric survey of a country, allowance must be made for the horizontal pressure gradient. When daily weather maps are available, B_1 may be taken from them as the pressure at sea-level in the neighborhood. If T_1 is not known, the conventional value (adopted by Intern. Meteorological Conference, Innsbruck, 1905) $T_1 = T_2 + 0.005 (H_2 - H_1)$ may be used, but in hot weather $T_1 = T_2 + 0.01 (H_2 - H_1)$ is a better approximation. Value of T_2 observed at a mountain station may differ considerably from the temperature of free atmosphere at same level; this is especially true in calm weather, at night, and in the early morning. (*cf.* Hesselberg, Int. Meteorol. Conference, Utrecht, 1923, App. L.) Tables of

virtual temperatures: V. Bjerknes, Dynamic Meteorology, etc., Washington, 1911. Values of $0.01484/T$: Computer's Handbook of Meteorological Office, London, **2**: 45.

Graduation of Aneroids. —The height scales on aneroids designed for the use of travellers, are graduated on the assumption that the temperature of the atmosphere is constant and independent of the altitude. Various standard temperatures, such as 50°F and 0°C have been used. For such scales, especially when applied to aircraft use, the difference between the indicated and the true height may be excessive.

The International Commission for Aerial Navigation adopted in 1925 a scale based on the following conventions (cf. Dict. Applied Physics **3**: 182): (a) Pressure at sea-level is $A_0 = 1.0132 \times 10^6$ dynes/cm²; (b) temperature at sea-level is 15°C; (c) temperature decreases by 6.5°C per km, up to 11 km; and above 11 km is constant at -56.5 C; (d) humidity may be ignored; (e) value of g is same at all heights and $= g_{45}$ (essentially $g.$). Whence, denoting the pressure and density at sea-level by B_1, and d_1; those at 11 000 m by $B_{11\ 000}$ and $d_{11\ 000}$:

$$\frac{B}{B_1} = \left(\frac{288 - 0.0065\ H}{288}\right)^{4.256}; \frac{d}{d_1} = \left(\frac{288 - 0.0065\ H}{288}\right)^{4.256};$$

if $H \not> 11\ 000$ m.

$$\log_{10}\frac{B_{11\ 000}}{B} = \log_{10}\frac{d_{11\ 000}}{d} = \frac{H - 11\ 000}{14\ 600}, \text{ if } H > 11\ 000\ \text{m}$$

	Unit	Value	Log₁₀
B_1	mm	760	2.88081
B_1	kilodyne/cm²	1013.2	3.00570
d_1	g/m³	1226	3.08849
$B_{11\ 000}$	mm	169.6	2.22943
$B_{11\ 000}$	kilodyne/cm²	226.1	2.35432
$d_{11\ 000}$	g/m³	364	2.56104

As the regulations drawn up by the I. C. A. N. are ambiguous, attention must be drawn to the fact that whilst the altimeter reading, H, gives the pressure uniquely, it cannot give the temperature and density of the air. Hence the formulae for d are on quite a different footing from those for B. (Cf. Section on Aerodynamics, Ed.)

VOLUMES OF LIQUID MENISCI

F. A. GOULD

As used in this section, the volume (V_m) of the liquid meniscus in a vertical, circular tube = volume of the liquid which lies below the capillary surface and between two horizontal planes, one tangent to the meniscus, and the other passing through the line in which the meniscus meets the wall of the tube. The value of V_m depends upon the surface tension (γ), the acceleration of gravity (g), the difference (ρ) in the densities of the fluids separated by the surface, the radius (r) of the cylinder, and the angle (θ) at which the capillary surface meets the wall of the cylinder. If θ is variable and not too small, it is more convenient to use the height (h_m) of the meniscus (= distance between the planes mentioned), than θ, as one of the variables. This has been done in Tables 1 and 2, which give the volume of the mercury meniscus for $\gamma = 400$ mg wt./cm ($= 392.27$ dynes/cm, $g = 980.665$), $\rho =$

13.55g/cm³. This value of γ is close to the mean of the values corresponding to the experimental determinations of V_m by Scheel and Heuse (*8*, **33**: 295; 10) (425 mg/cm), and by Palacios (*139*, **17**: 295; 19. *63*, **24**: 152; 23) (406 to 326 mg/cm); an idea of the error which is associated with a departure of the actual value of γ from that assumed may be obtained by comparing their values with those here given. (*See also* Schalkwijk, *168*, No. **67**, and *64* V, **8**: 462; 00. **9**: 512; 01.)

If $\theta = 0$, it is convenient to tabulate the dimensionless quantities V_m/r^3 and $h_c/r = V_m/\pi r^3$ as functions of $g\rho r^2/\gamma$, as is done in Table 3. [$g\rho r^2/\gamma = r^2/a_1^2$, where a_1 is capillary constant (British usage), *see* section Technical Terms (p. 34); h_c = length of circular cylinder of radius r and volume V_m].

TABLE 1.—VOLUME (V_m) OF MERCURY MENISCUS

h_m = height of meniscus, d = internal diameter of tube. Accuracy for the larger menisci = 0.3 %, for the smaller = 1 %. Unit of V_m = 0.001 cm³; of h_m and d = 1 mm. Assumes γ = 400 mg wt./cm

h_m	d 1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	d h_m
0 1	0 040	0 159	0 360	0 616	1 02	1 50	2 08	2 75	3 55	4 46	5 49	6.67	7 97	9 42	11 1	12 8	14 8	16 9	19 2	21 6	24 2	27 0	30 0	33 1	0 1
0 2	0 083	0 321	0 723	1 30	2 05	3.00	4.16	5 53	7.12	8 95	11 0	13.4	16.0	18 9	22 2	25 7	29 6	33 9	38 5	43 4	48.6	54 1	60 0	66.3	0.2
0 3	0 134	0 490	1.09	1 95	3 09	4 52	6.26	8 32	10.7	13.5	16.6	20 2	24.1	28 5	33.4	38.7	44 6	51 0	57 8	65 2	73 0	81 3	90 2	99 6	0.3
0 4	0 195	0 669	1 47	2 63	1 14	6 04	8 37	11 1	14 3	18 0	22.3	27.0	32 3	38.1	44 7	51 8	59 6	68 1	77 3	87 1	97 5	109	120	133	0.4
0 5		0 801	1 87	3 31	5 21	7 59	10.5	14 0	18.0	22.7	28 0	33 9	40.6	47.9	56 1	65.0	74 7	85 4	96 9	109	122	136	151	167	0.5
0 6		1 07	2 29	4 01	6 30	9 16	12 7	16.8	21.7	27.3	33 7	40.9	48 9	57.8	67 6	78.3	90 0	103	117	131	147	164	181	200	0 6
0 7		1 31	2 72	4 74	7 43	10 8	14 9	19 7	25 4	32 0	39 5	47.9	57.4	67 8	79.2	91 7	105	120	136	154	172	191	212	234	0.7
0 8		1 56	3 17	5 50	8 58	12 4	17.1	22 6	29 2	36 8	45 4	55 1	65 9	77.8	91 0	105	121	138	156	176	197	219	243	268	0.8
0 9		1 85	3 67	6 29	9 77	14 1	19 4	25 6	33 0	41 6	51 4	62 3	74 5	88 0	103	119	137	160	177	199	222	248	274	303	0.9
1.0			4 19	7 12	11 0	15.8	21 7	28 6	36 9	46 5	57 3	69.6	83 2	98.3	115	133	153	174	197	222	248	276	306	337	1.0
1 1			4 76	7 99	12 3	17.6	24.1	31.8	40 0	51.4	63.5	77 0	92 1	109	127	147	169	192	218	245	274	305	338	372	1.1
1 2			5 39	8 90	13 6	19.5	26 6	35 0	44.9	56 5	69 7	84 5	101	119	139	161	185	211	238	268	300	334	369	407	1.2
1 3			6 07	9 88	15 0	21 4	29 1	38 2	49 1	61.7	76 0	92.1	110	130	152	176	201	229	260	292	326	363	402	443	1.3
1 4				10 9	16 4	23 4	31 7	41 7	53 3	66 9	82.4	99 9	119	141	164	190	218	248	281	316	353	392	434	478	1.4
1 5				12 1	18 0	25 4	34 5	45 2	57 8	72 3	89 0	108	129	152	177	205	235	268	303	340	380	422	468	515	1.5
1 6				13 3	19 6	27 6	37 3	48 8	62 3	77 8	95 7	116	138	163	190	220	253	287	325	365	407	453	501	552	1 6
1.7					21 4	29 8	40 2	52 6	67 0	83 6	103	124	148	175	204	236	270	307	347	390	436	484	535	589	1.7
1 8					23 2	32 2	43 3	56 4	71.7	89 5	110	133	158	186	218	251	288	328	370	415	464	515	570	628	1.8
1 9						34 8	46 5	60 4	76 7	95 5	117	141	168	199	232	268	306	349	393	441	493	547	605	666	1.9
2.0						37.4	49.8	64 6	81 9	102	124	150	179	211	246	284	325	370	417	468	522	580	641	706	2.0
2.1							53.4	69 1	87 3	108	132	159	190	224	261	301	344	391	441	495	552	614	678	746	2.1
2 2								73.7	93 0	115	140	169	201	237	276	318	364	413	466	523	583	648	716	787	2.2
2.3									98.9	122	149	179	213	250	291	336	384	436	492	552	615	683	754	829	2.3
2.4										130	158	189	225	264	307	354	405	459	518	581	648	719	794	872	2.4
2.5											167	200	237	279	324	373	427	484	546	612	681	755	833	915	2.5

TABLE 2.—HEIGHT (h_c) OF CYLINDER EQUIVALENT TO VOLUME (V_m) OF MERCURY MENISCUS

$h_c = V_m/\pi r^2$ = length of tube of radius r and volume V_m; h_m = height of meniscus; $d = 2r$ = diameter of tube. Accuracy and basis are same as for Table 1

Unit of h_c, h_m, and $d = 1$ mm. Assumes $\gamma = 400$ mg wt./cm

h_m \ d	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	d \ h_m
0.1	0.051	0.051	0.051	0.051	0.052	0.053	0.054	0.055	0.056	0.057	0.058	0.059	0.060	0.061	0.063	0.064	0.065	0.066	0.068	0.069	0.070	0.071	0.072	0.073	0.1
0.2	0.106	0.102	0.102	0.103	0.104	0.106	0.108	0.110	0.112	0.114	0.116	0.118	0.121	0.123	0.126	0.128	0.131	0.133	0.136	0.138	0.140	0.142	0.144	0.147	0.2
0.3	0.171	0.156	0.155	0.155	0.157	0.160	0.163	0.165	0.168	0.172	0.175	0.178	0.182	0.185	0.189	0.192	0.196	0.200	0.204	0.208	0.211	0.214	0.217	0.220	0.3
0.4	0.248	0.213	0.209	0.209	0.211	0.214	0.218	0.221	0.225	0.230	0.234	0.239	0.243	0.248	0.253	0.257	0.263	0.268	0.273	0.277	0.281	0.286	0.290	0.294	0.4
0.5		0.274	0.265	0.263	0.265	0.268	0.273	0.278	0.283	0.288	0.294	0.300	0.306	0.311	0.318	0.323	0.329	0.336	0.342	0.348	0.353	0.358	0.363	0.368	0.5
0.6		0.341	0.323	0.319	0.321	0.324	0.329	0.335	0.341	0.348	0.355	0.362	0.368	0.375	0.383	0.389	0.397	0.404	0.411	0.418	0.421	0.430	0.437	0.443	0.6
0.7		0.416	0.385	0.377	0.378	0.381	0.386	0.392	0.400	0.408	0.416	0.424	0.432	0.440	0.448	0.456	0.465	0.473	0.481	0.489	0.496	0.504	0.511	0.518	0.7
0.8		0.497	0.449	0.438	0.437	0.440	0.444	0.450	0.459	0.468	0.478	0.487	0.496	0.506	0.515	0.524	0.533	0.542	0.551	0.561	0.569	0.577	0.585	0.593	0.8
0.9		0.588	0.519	0.501	0.497	0.499	0.503	0.510	0.519	0.530	0.540	0.551	0.561	0.572	0.582	0.592	0.602	0.613	0.623	0.633	0.642	0.651	0.660	0.669	0.9
1.0			0.593	0.567	0.559	0.560	0.563	0.570	0.580	0.592	0.604	0.615	0.627	0.638	0.650	0.661	0.673	0.684	0.695	0.706	0.716	0.726	0.736	0.746	1.0
1.1				0.674	0.636	0.624	0.623	0.626	0.632	0.643	0.655	0.668	0.681	0.694	0.706	0.719	0.731	0.743	0.756	0.767	0.779	0.790	0.802	0.812	0.823	1.1
1.2				0.762	0.708	0.692	0.689	0.691	0.696	0.707	0.720	0.733	0.747	0.761	0.775	0.788	0.802	0.815	0.828	0.841	0.854	0.866	0.878	0.890	0.900	1.2
1.3				0.859	0.786	0.763	0.756	0.757	0.761	0.772	0.785	0.799	0.815	0.829	0.844	0.859	0.873	0.887	0.902	0.915	0.929	0.942	0.955	0.967	0.979	1.3
1.4					0.896	0.837	0.826	0.825	0.829	0.839	0.852	0.867	0.883	0.899	0.915	0.930	0.946	0.961	0.976	0.991	1.01	1.02	1.03	1.05	1.06	1.4
1.5					0.961	0.915	0.899	0.896	0.900	0.908	0.921	0.936	0.953	0.969	0.986	1.00	1.02	1.04	1.05	1.07	1.08	1.10	1.11	1.13	1.14	1.5
1.6						1.06	1.00	0.976	0.969	0.972	0.980	0.9911	01	02	04	06	08	10	11	13	15	16	18	19	21	1.6
1.7							1.06	1.04	1.05	1.05	1.06	1.08	10	12	13	15	17	19	21	22	24	26	27	29	30	1.7
1.8							1.18	1.14	1.12	1.12	1.13	1.14	15	17	19	21	23	25	27	29	31	32	34	36	37	1.8
1.9								1.23	1.21	1.20	1.21	1.22	23	25	27	29	31	33	35	37	39	40	42	44	46	1.9
2.0								1.32	1.30	1.29	1.29	1.30	31	33	35	37	39	41	43	45	47	49	51	53	54	2.0
2.1									1.39	1.37	1.37	1.38	39	41	43	45	47	50	52	54	56	58	59	61	63	2.1
2.2										1.47	1.46	1.47	48	50	52	54	56	58	60	62	64	66	68	70	72	2.2
2.3										1.55	1.56	1.57	58	60	63	65	67	69	71	73	76	78	80	82	83	2.3
2.4										1.65	1.66	1.68	69	72	74	76	78	80	83	85	87	89	91	93	2.4	
2.5										1.76	1.77	1.79	81	83	86	88	90	92	95	97	99	01	02	2.5		

TABLE 3.—VOLUME (V_m) OF LIQUID MENISCUS, $\theta = 0$

(Meniscus concave upwards)

As quantities tabulated are dimensionless, any consistent system of units may be used. g = acceleration of gravity, r = radius of tube, h_c = length of tube of radius r and volume V_m. (Computed from tables of Bashforth and Adams as given in their "Capillary Action.")

$g\rho r^2/\gamma$	V_m/r^3	h_c/r	$g\rho r^2/\gamma$	V_m/r^3	h_c/r
0	1.048	0.333	4.0	0.649	0.206
0.1	1.029	0.327	4.5	0.623	0.198
0.2	1.010	0.321	5.0	0.599	0.190
0.4	0.978	0.311	5.5	0.578	0.184
0.6	0.947	0.301	6.0	0.557	0.177
0.8	0.919	0.292	6.5	0.537	0.171
1.0	0.894	0.284	7.0	0.518	0.165
1.5	0.837	0.266	7.5	0.501	0.159
2.0	0.789	0.251	8.0	0.484	0.1540
2.5	0.747	0.238	8.5	0.470	0.1493
3.0	0.711	0.226	9.0	0.456	0.1449
3.5	0.678	0.216	9.5	0.442	0.1406
			10.0	0.429	0.1365

Example 1. A gas is collected in a eudiometer over mercury. The volume to the plane through the line of contact of the mercury with the wall of the tube $= V_o$. If this portion of the eudiometer is a vertical, circular cylinder of diameter $d = 10$ mm, and if height of meniscus is $h_m = 1.5$ mm, then $V_m = 0.0723$ cm^3 (Table 1), and the actual volume of the gas is $V = V_o - 0.072$ cm^3.

If volumes are expressed in terms of a linear scale engraved upon the cylindrical portion of the eudiometer, and if the scale reading at the line of contact is h_o, and if $d = 10$ mm, $h_m = 1.5$ mm, then $h_c = 0.921$ mm (Table 2), and the actual volume of the gas corresponds to $h_o - h_c = h_o - 0.921$ mm.

Example 2. A gas is collected in a eudiometer over water. The volume to the plane tangent to the bottom of the meniscus = V_o. If this portion of the eudiometer is a vertical, circular cylinder of radius $r = 0.5$ cm, if $\gamma = 73$ dynes/cm, $g = 980.7$ cm/sec^2, $\rho = 1.000$, and $\theta = 0$ (the tube is perfectly wetted by the water), then $g\rho/\gamma = 13.43$ cm^{-1}, $g\rho r^2/\gamma = 3.36$. Hence $V_m/r^3 = 0.689$ (Table 3), and $V_m = 0.086$ cm^3. Hence the actual volume of the gas is $V_o - V_m = V_o - 0.086$ cm^3.

If volumes are expressed in terms of a linear scale engraved upon the cylindrical portion of the eudiometer, and if the scale reading corresponding to the bottom of the meniscus is h_o, then for $g\rho r^2/\gamma = 3.36$, $h_c/r = 0.219$ (Table 3), and if $r = 5$ mm, $h_c = 1.10$ mm, and the actual volume of the gas corresponds to $h_o - h_c = h_o - 1.10$ mm.

WEIGHTS AND WEIGHING

A. T. PIENKOWSKY

In this section are considered:—(A) Weights—the basis upon which they are adjusted or tested, and their constancy; (B) the correcting of weighings for the buoyant effect of the air, including the weighing of substances in containers; and (C) the correcting of density determinations for the buoyant effect of the air.

WEIGHTS

Basis of Adjustment.—Most weights are adjusted by the maker according to their apparent weight in air against brass standards. This is equivalent to adjusting brass weights according to their real mass (or "weight in vacuo"), but the true mass values of other

weights (*e.g.*, those of platinum, aluminum, or quartz) may be much different from their nominal values. When a set of weights is calibrated, however, the values found may be either true mass or apparent values, depending on the standard used and the method of conducting the test. Certificates from different standardizing laboratories may give values on either basis, or on both.

"Weight in Air against Brass."—Commercial weighing is all based on apparent weight in air against brass standards, this basis being more or less accurately defined in some countries. Precise scientific weighing is based on true mass values (*i.e.*, on "weight in vacuo"), but weights below one gram may be tested and used as if they were of brass, even for work of rather high precision. In so testing these weights, their apparent "values" are computed on the assumption that their density is Δ_b – density of brass (generally Δ_b is taken as 8.4 g per cm³); and in using them the apparent values so found are used as though they were the true masses of the weights, Δ_b being at the same time used just as though it were the true density of the weights. In such cases the error ($m_f - m$) so introduced, arises solely from the fact that the density (σ_1) of the air at the time the values of the weights were determined differs from that (σ) at the time they were used in weighing the object. This error is given approximately by equation (1) in which m is the correct, and m_f is the false mass, s is the nominal value of the weight, Δ_b is the density assumed for brass weights and Δ the actual density of the weights used.

$$m_f - m = s\left(\frac{1}{\Delta_b} - \frac{1}{\Delta}\right)(\sigma_1 - \sigma) \tag{1}$$

Example: If the value of a platinum 500 mg weight ($\Delta = 21.5$ g/cm³) is determined according to "weight in air against brass" ($\Delta_b = 8.4$ g/cm³) at sea level ($\sigma_1 = 0.0012$ g/cm³), and this value is used at an altitude of 5000 ft. ($\sigma = 0.0010$ g/cm³) the error in the mass of a body as so weighed will be $m_f - m = 0.007$ mg.

"Apparent" densities or specific gravities determined according to apparent "weight in air against brass" are subject not merely to variations in the density of the air, but also to differences in experimental technique (see p. 78 to 80).

Constancy. Data on changes in weights can indicate only the order of magnitude of such changes, and as a rule can show only what *may* happen, since such changes are extremely irregular.

Ordinary brass weights with knobs screwed in (whether gold plated, platinum plated, or lacquered) may continue to gain in weight for many years, and may do so without developing any visible signs of such change. The following examples are typical of extreme changes that sometimes occur. Larger changes have been recorded.

Denomination	g	100	50	20	10	5	2	1
Gain in 6 yr	mg	1 7	1 2	0 8	0 7	0 6	0.8	0 3
Gain in 14 yr	mg	3 3	3 9	1 8	2 5	0 8	0 3	1 1

The following is typical of what has often happened when new weights were not used and were carefully protected.

Denomination	g	100	50	20	10	5	2	1
Gain in 5 mo	mg	0 1	0 1	0 0	0 1	0 1	0 0	0 0
Gain in 1 yr	mg	0 2	0 1	0 0	0 0	0 1	0 0	0 0

Lacquered weights of good quality are less subject to spotting and general surface tarnishing than are the gold or platinum plated weights often sold. Lacquered weights, however, are subject to rapid variations caused by changes in the relative humidity of the air. Lacquered weights of about 20 to 100 g may be expected to vary 0.1 or 0.2 mg with large variations in humidity. Changes of over 0.5 mg have been recorded.

Sets of weights of the ordinary type may, however, be very constant. For example, one set was used for over a year with changes less than 0.02 mg and few changes over half that amount; and two sets were used occasionally for 17 and 18 yr, respectively, with no changes over 0.2 mg.

For reference standards, one-piece weights are very much more reliable than the common screw-knob type. The following changes in a high grade, gold plated, bronze set of this type are typical for weights used little and with great care. Positive changes are gains, negative changes losses.

Denomination.	g	50	20	20	10	5	2	2	1
Changes in 15 yr	mg	-0 12	0 00	0.02	-0 01	-0 00	0 001	0 008	-0.007

Solid platinum or platinum-iridium weights of moderate size may be expected to remain constant within about 0.01 mg if handled with sufficient care and protected from dust and other deposits. The sheet metal weights below one g are not much more constant than this; very good weights kept with extreme care as reference standards may stay within 0.001 mg for some years, but this cannot safely be assumed. If these small weights are much used, even with good care, losses of 0.01 mg may soon be expected in the larger ones.

CORRECTING OF WEIGHINGS FOR BUOYANT EFFECT OF THE AIR

("Reduction of Weighings to Vacuo")

In addition to a sufficiently sensitive balance, accurate weighing requires (1) that the balance itself maintain a sufficiently constant zero point and ratio of arms of the beam; (2) that the effect of inequality of the arms of the beam be eliminated by the method of weighing, since it cannot as a rule be corrected for with sufficient accuracy; (3) that the object and the weights have definite constant values, free from such effects as variable surface loss by evaporation, magnetic attractions, etc.; (4) that surrounding conditions be maintained free from sources of disturbance and error, such as electrostatic attractions, convection currents, variable or unsymmetrical heat radiations, etc.; and (5) that proper correction be made for the buoyant effect of the air.

The first four types of requirements are matters of technique, and no general methods of correction can be used for errors arising from them. They are therefore outside the scope of these tables.

The fifth requirement demands definite formulae and facts, some of the most fundamental or general of which are given below.

The phrase "apparent weight" is commonly used for the result of a weighing in which no correction has been made for the buoyant effect of the air. The phrase is ambiguous[1] and often leads to a confusion of ideas. Therefore this term is not used in the equations of this section, but reference is made directly to the weights that would be used on an equal-arm balance to make the weighings. The phrase "weights needed" must be understood to include the proper fraction of the rider or other small weights needed to make up the total amount; and it refers to *actual* values of the weights, which may or may not equal the nominal values marked on them.

Symbols —

a mass of the contents of the "empty" portions of the container. (In weighing gases a is zero. In weighing solids or liquids it may be the mass of air or of vapor of the solid or liquid. In weighing a pyknometer with the liquid which fills it at a temperature different from that at which it is weighed, the volume occupied by a results from the unequal expansion of pyknometer and liquid)

b $(v_a - v_c)/r_a$. Relative size of the container and its counterpoise

c mass of counterpoise

k buoyancy reduction factor

l mass of liquid that fills the pyknometer at the established filling temperature

m mass of object; in general or where its volume is not fixed by the volume of a pyknometer

p mass of pyknometer or other container

r error resulting from use of approximate buoyancy formula

[1] Compare equations (8) and (9); in each case $s'' - s'$ would be called the apparent weight, but its value in (9) is r_{ac} greater than in (8).

s   mass of weights needed on an equal arm balance, whether with or without special counterpoise, to balance the objects being weighed. (Regarding use of other than true mass values, *see* p. 73)

s_i $s - v_i\sigma = s(1 - \sigma/\Delta)$. This is *not* "weight in vacuo" as that phrase is often used

t   temperature. If accented it is the temperature at the time of the indicated weighing; if unaccented, it is the temperature at which the pyknometer is filled. In so far as their temperatures have any effect upon the operation considered, all objects (*e.g.*, the balance, its loads, and the surrounding air) are assumed to be at the same temperature

v   volume or capacity; when without subscript it is capacity of the container at time of weighing; with one of the subscripts *a, c, l, m, p, s,* or *w,* it is volume of the object whose mass is indicated by the subscript (*e.g.*, v_m = volume of the object whose mass is *m*)

v_t capacity of the pyknometer at the temperature of filling

v_p volume of the pyknometer itself, excluding the space that would be filled by liquid at the temperature of filling. (Ordinarily v_p = volume of the material of which the pyknometer is constructed)

v_e "exterior volume" of the pyknometer or other container. With pyknometers, at temperature of filling, $v_e = v_p + v_t$; at another temperature, t'', $v_e''-v_p'' + v'' = v_p'' + v_w'' + v_a''$

w   mass of the calibrating liquid (*e.g.*, water) which is used to determine a volume or to serve as a standard of density

β cubical coefficient of thermal expansion

Δ density of the weights at the time of weighing

σ density of the air at the time of weighing

ρ density of object being studied or of calibrating liquid. If accented it is density at time of weighing; if unaccented it is density at temperature (*t*) at which the pyknometer was filled

Density is true mass per unit of volume.

Accents denote the weighing to which the quantity applies. In general ′ denotes the weighing of the object alone or of the container; ″ denotes the weighing of the combined container and object studied, or of the container filled with the calibrating liquid or of the object suspended in the calibrating liquid; ‴ denotes the weighing of the pyknometer "filled" with liquid to be studied, or "filled" with object studied plus calibrating liquid.

Subscripts.—*i* denotes false or erroneous values. For *i* see above (s_i and v_e). Other subscripts indicate the object to which the quantity applies; *e.g.*, ρ_a = density of material whose mass is *a*.

Fundamental Exact Equation.—The use of the direct, fundamental, exact equation (2) avoids many complications and approximations introduced by most formulae based on densities.

$$m = s + (v_m - v_s)\sigma \qquad (2)$$

The equation using densities, in one of the exact forms (3) given below, is useful chiefly for computing exact tables, or the effect of errors, approximations, etc. As a rule, either the densities are not known well enough to warrant its use, or the volumes involved will have been measured, thus going back to equation (2).

$$m = s\left(\frac{1 - \dfrac{\sigma}{\Delta}}{1 - \dfrac{\sigma}{\rho_m}}\right) = s\frac{\rho_m(\Delta - \sigma)}{\Delta(\rho_m - \sigma)} = s\left\{1 + \frac{\sigma(\Delta - \rho_m)}{\Delta(\rho_m - \sigma)}\right\} =$$
$$s + s\frac{\sigma(\Delta - \rho_m)}{\Delta(\rho_m - \sigma)} \qquad (3)$$

In the last form of (3), the second term is the exact "buoyancy correction term," and in this correction term the factor (fraction) by which *s* is multiplied is the exact "buoyancy reduction factor" (*k*). See Tables 2 and 3.

Common Equation Using Densities.—Some form of equation (4) is commonly used for reducing weighings. This equation is not exact. It is entirely inapplicable to weighing gases, but is amply accurate for much work with solids and liquids.

$$m = s + s\sigma\left(\frac{1}{\rho_m} - \frac{1}{\Delta}\right) \qquad (4)$$

The factor $\sigma\left(\dfrac{1}{\rho_m} - \dfrac{1}{\Delta}\right)$ is the "buoyancy reduction factor" commonly given. When the densities lie between 0.5 and 21.5 g per cm³, and are known with sufficient accuracy, the error (*r*) introduced by the use of this formula does not exceed one part in 100 000 of the mass of the object weighed. Its value, and that of the proportional error ($r' = r/s$) may be calculated by formula (5); their orders of magnitude may readily be determined from Table 1, which is based on $\sigma = 0.0012$ g/cm³.

$$r' = \frac{r}{s} = \frac{\sigma^2(\Delta - \rho_m)}{\Delta\rho_m(\rho_m - \sigma)} \qquad (5)$$

TABLE 1

Unit of Density is g/cm³

ρ_m	100 r'		
	$\Delta = 21.5$	$\Delta = 8.4$	$\Delta = 2.65$
1.00	0.0001	0.0001	0.0001
0.5	0.0006	0.0005	0.0005
0.05	0.06	0.06	0.06
0.005	8.	8.	7.

Density of the Air.—Variations in the density of the air under standard conditions,[1] as well as the uncertainties of its experimental determination, limit the precision with which very large or extremely precise buoyancy corrections can be calculated from tables of air density. The former seems at present to be the larger, and therefore sets a fixed limit which can be exceeded only by eliminating or reducing the size of the correction, or by making an experimental determination of the density of the air at the time of the weighing. These limiting uncertainties are of the order of 5 in 10⁴ and affect the total buoyancy correction in the same ratio. Since they affect only the fourth significant figure in the buoyancy reduction factor they are negligible in the use of Tables 2 and 3.

In weighing gases, the density of the air must be found from precise tables (consult index). When the volume of the gas is not compensated by a counterpoise of the same size, the density of the air must be known with approximately the same precision as is desired for that of the gas; when it is so compensated, the buoyancy correction is generally the total buoyancy on the weights, and therefore is still relatively large.

For most work with solids and liquids an approximate value of the density of the air is sufficient. The precision to which it must be known can be found from an examination of Table 2. It should be noted that a precision of 1 in 10ⁿ in the mass to be determined requires a precision of 1 in the n'th decimal place of the buoyancy reduction factor (i.e., in the actual factor *k*, not in the printed value of 1000*k*). In getting the buoyancy reduction factor from Table 2, and in similar work, to a precision not greater than one in about 10⁵, the density of the air may be found from the "Air Density Chart," Fig. 1.

The precision to which temperature, pressure, and humidity must be known in order to find the density of the air to the necessary precision, may be inferred from Fig. 1, except in the case of very large corrections, or of corrections to be determined with extreme precision. In the latter cases this information must be sought in other places.

Density of the Weights.—If the density of the air in which the weights are used is the same as that in which their values were determined, errors in the density assumed for the weights will have

[1] Treuthart, *34*, **172**: 1598, 21. Moles, *34*, **172**: 1600, 21.

no effect on the accuracy with which the mass of the object may be determined, provided the same density that was assumed for them in determining their values is assumed for them when they are used. It is not necessary, therefore, to know the density of the weights as accurately as that of the object weighed.

If weights are used in air whose density differs by not more than 20% from that of the air in which their values were determined, the amount by which the density of ordinary weights is likely to differ from the values used in Tables 2 and 3 will not cause errors greater than one part in about 100,000 in the determination of the mass of the object weighed; provided that the density used in determining the value of the weight is the same as that used in the computation of the mass.

For a precision above one part in a million, it is frequently necessary to measure the volume or density of each weight.

Fig. 1.—Air density chart   (For use with Tables 2 and 3.)

Ordinary two-piece weights are not used for such work because they cannot safely be put into liquids for hydrostatic weighing.

Aluminum is not used for weights above 0.02 g in high quality weights, nor above 0.5 g in second quality sets. When the values of such weights have been determined on the assumption of a density of 2.7 g per cm³ at 0°C, the use of the buoyancy reduction factors given for quartz in Table 2 introduces an error in the mass of the object weighed, of less than 0.0002 mg for amounts up to 0.02 g, and of less than 0.005 mg for amounts up to 0.5 g.

The densities of most gold alloys used for weights lie between 16 and 18 g per cm³. For gold within this range, the use of the factors given in Tables 2 and 3 will not introduce errors greater than one part in 200,000, or not over 0.005 mg in weighing amounts under one g.

In Tables 2 and 3, the densities used for weights of platinum or platinum-iridium, for those of brass or bronze, and for those of aluminum, are those which were adopted many years ago for certifying weights at the National Bureau of Standards of the United States of America, and were assumed as the densities at 0°C. The following coefficients of cubical expansion are assumed in reducing the volumes of such weights to the volumes at 20°C.

Platinum and Platinum-iridium .... 0 000 026 per deg. C
Brass or bronze   .... 0 000 054 per deg. C
Aluminum. ... . 0.000 069 per deg. C

The densities of gold and of crystal quartz are assumed as the densities at 20°C. All buoyancy reduction factors are based on differences in volume at 20°C.

Density of Object Weighed.—A change of one in 10^n of the mass of the object corresponds to a change of one in the n'th decimal place of the buoyancy reduction factor. Therefore, to the precision obtainable by the use of Table 2, the precision required in the density of the object may be found by noting in that table what change in density (at approximately the density under consideration) corresponds to the allowable variation in the buoyancy reduction factor.

The use of "standard" or "adopted" densities for the object weighed may give an accuracy which is entirely fictitious. There is no compensation as in the case of weights, and the actual error or uncertainty in the density of the particular object weighed has its full effect in the error or uncertainty of the calculated mass

A fictitious "apparent" density derived from weighings uncorrected for buoyancy of the air must be corrected to true density before being inserted in the formulae given in this section unless only an approximate value of density is needed (see p. 78).

Temperature of Objects and Weights.—In weighing gases, and to secure the highest precision in many other cases, it is necessary to compute all volumes or densities at the actual temperature of the observations, unless the coefficient of expansion of the object happens to be nearly the same as that of the weights. If the temperature is entirely neglected, and weighings are made at room temperatures, the extreme error likely to be introduced in the mass calculated for solids and liquids is less than three in 10^4. (This would be the error for material having a density of 0.2 g per cm³ at 0°C, and a coefficient of cubical expansion of 1.6×10^3, when compared with weights whose actual volumes or densities are those used in the calculation.)

Example 1: The actual mass of the weights used was $s = 10\ 0105$ g; the corrected barometric height was 758 mm; air temperature, 19.6°C; relative humidity 25%; density of object 3.5 g/cm³; weights were of brass.

Referring to Fig. 1, the air density corresponding to these conditions is seen to be close to 0.0012 g/cm³. Entering Table 2 with $\rho_m = 3.5$ and the column for brass weights, under $1000\sigma = 1.2$, it is found that $1000\ k$ is 0 20; hence the mass of the object is $m = s + ks = 10.0105 + 0.000\ 20 \times 10.0105 = 10.0105 + 0.0020 = 10.0125$ g.

Example 2: The factor for $\rho_m = 3.0$ differs by 6 in the fifth decimal place from that for $\rho_m = 3.5$. The error in mass produced by using 3.0 in place of 3.5 as the density of the object is therefore 6 parts in 10^5. For the object in Example 1 this would be an error of 0.000 6 g. Similarly the use of 7.0 instead of 7.5 for ρ_m would produce an error of about one part in 10^4 in the mass of the object.

Example 3: In Fig. 1 the point corresponding to barometric height 720 mm, air temperature 21°C, and relative humidity 50%, lies to the right of the line for 0.0011 g/cm³, 50%, by $^{19}/_{32}$ of the distance between the 0.0011 and the 0.0012 lines. Hence, $\sigma = 0.0011 + 0.0001 \times {}^{19}/_{32} = 0.001131$ g/cm³. (For most work for which Table 2 is suited the density can be estimated by eye with sufficient accuracy; as in this case, 0.00113 g/cm³.) The factor from Table 2 may then be found either by multiplying the factor for $1000\sigma = 1.0$ by 1.13 or by interpolating between the factor for $1000\sigma = 1.1$ and that for $1000\sigma = 1.2$. For brass weights and $\rho_m = 3.5$ the former gives $0.17 \times 1.13 = 0.19_2$ as the value of $1000k$. A calculated interpolation between 0.18 and 0.20 gives 0.18₆, which agrees with the other value within the accuracy of such tabular interpolations.

Weighing Objects in Containers.—Two weighings are required; one of the container alone and the other with the object in the

TABLE 2.—BUOYANCY REDUCTION FACTOR (k)

$$m = s + ks, \text{ where } k = \frac{\sigma(\Delta - \rho_m)}{\Delta(\rho_m - \sigma)}$$

(*Cf* equation (3). Symbols, p. 74.) Unit of density is g/cm³ or, to precision of this table, g/ml

1000 k

Density of object weighed ρm	Δ = 21.5 Pt or Pt-Ir 1000σ =			Δ = 17 Gold 1000σ =			Δ = 8.4 Brass or bronze 1000σ =			Δ = 2.65 Crystal quartz or aluminum* 1000σ =		
	1.0	1.1	1.2	1.0	1.1	1.2	1.0	1.1	1.2	1.0	1.1	1.2
0.2	4 98	5 48	5 98	4 97	5 47	5 97	4 91	5 40	5 89	4 65	5 11	5.58
0.3	3 30	3 63	3 96	3 29	3 62	3 95	3 22	3 55	3 87	2 97	3 26	3 56
0.4	2 46	2 71	2 95	2 45	2 69	2 94	2 39	2 63	2 87	2 13	2.34	2.55
0.5	1 96	2 15	2 35	1.95	2 14	2 34	1 88	2 07	2 26	1 63	1.79	1 95
0.6	1 62	1 79	1.95	1 61	1 77	1 93	1 55	1 71	1 86	1 29	1 42	1.55
0.7	1 38	1 52	1 66	1 37	1 51	1 65	1 31	1 44	1 57	1 05	1.16	1.26
0.75	1.29	1.42	1.55	1.28	1 40	1 53	1 22	1 34	1 46	0 96	1 05	1 15
0.80	1 20	1.33	1 45	1 19	1 31	1 43	1 13	1 25	1 36	0 87	0 96	1.05
0.82	1 17	1 29	1 41	1 16	1 28	1 39	1 10	1 21	1 32	0 84	0 93	1 01
0.84	1 15	1 26	1 37	1 13	1 25	1 36	1 07	1 18	1 29	0 81	0 90	0.98
0.86	1 12	1 23	1 34	1 11	1 22	1 33	1 04	1 15	1 25	0 79	0 86	0 94
0.88	1 09	1 20	1 31	1 08	1 19	1 29	1 02	1 12	1 22	0.76	0.88	0 91
0 90	1.07	1 17	1 28	1 05	1 16	1 26	0 99	1 09	1 19	0 73	0 81	0 88
0.91	1 05	1 16	1 26	1 04	1 15	1 25	0 98	1 08	1 18	0 72	0.79	0.87
0.92	1 04	1 15	1 25	1 03	1 13	1 24	0 97	1 06	1 16	0 71	0 78	0.85
0 93	1.03	1 13	1 24	1 02	1 12	1 22	0 96	1 05	1 15	0 70	0 77	0.84
0.94	1 02	1.12	1 22	1 01	1 11	1 21	0 95	1 04	1 13	0 69	0 76	0.82
0.95	1 01	1 11	1 21	0 99	1 09	1 19	0 93	1 03	1 12	0 68	0 74	0 81
0.96	1 00	1 10	1 20	0 98	1 08	1 18	0 92	1 02	1 11	0 67	0 73	0 80
0 97	0 99	1 08	1 18	0 97	1 07	1 17	0 91	1 00	1 09	0 65	0 72	0 79
0.98	0 97	1 07	1 17	0 96	1 06	1 16	0 90	0 99	1 08	0 64	0 71	0.77
0.99	0 96	1 06	1 16	0.95	1 05	1 14	0 89	0 98	1 07	0 63	0 70	0.76
1.00	0 95	1 05	1 15	0.94	1 04	1 13	0 88	0 97	1 06	0 62	0 69	0.75
1 01	0 94	1 04	1 13	0 93	1 03	1 12	0 87	0 96	1 05	0 61	0 67	0 74
1.02	0 93	1 03	1 12	0 92	1 01	1 11	0 86	0 95	1 03	0 60	0 66	0.72
1.03	0 93	1.02	1 11	0 91	1 00	1 10	0 85	0 94	1 02	0 59	0 65	0.71
1.04	0.92	1.01	1.10	0 90	0 99	1 08	0 84	0 93	1 01	0 58	0 64	0.70
1.05	0 91	1 00	1 09	0 89	0 98	1 07	0 83	0 92	1 00	0 58	0 63	0 69
1 06	0 90	0 99	1 08	0 89	0 97	1 06	0 82	0 91	0 99	0.57	0 62	0 68
1 07	0 89	0 98	1 07	0 88	0 96	1 05	0 82	0 90	0 98	0 56	0 61	0 67
1.08	0 88	0 97	1 06	0 87	0 95	1 04	0 81	0 89	0 97	0 55	0 60	0 66
1.09	0 87	0 96	1 05	0 86	0 94	1 03	0 80	0 88	0 96	0 54	0 59	0 65
1.10	0 86	0 95	1 04	0 85	0 94	1 02	0 79	0.87	0 95	0 53	0 59	0.64
1.12	0 85	0 93	1 02	0 83	0 92	1 00	0 77	0 85	0 93	0 52	0 57	0.62
1.14	0 83	0 91	1 00	0 82	0 90	0 98	0 76	0 83	0 91	0 50	0 55	0.60
1.16	0 82	0 90	0 98	0 80	0 88	0 96	0 74	0.82	0 89	0 49	0 53	0.58
1.18	0 80	0 88	0 96	0.79	0 87	0 95	0 73	0.80	0 87	0 47	0 52	0.56
1.20	0 79	0 87	0 95	0 78	0 85	0 93	0 71	0 79	0 86	0 46	0 50	0.55
1.25	0 75	0 83	0 91	0 74	0 82	0 89	0 68	0.75	0.82	0 42	0 47	0.51
1.30	0 72	0 80	0 87	0 71	0 78	0 85	0 65	0.72	0 78	0 39	0 43	0.47
1.35	0 69	0 76	0 83	0 68	0 75	0 82	0 62	0.68	0 75	0 36	0.40	0.44
1.40	0.67	0.74	0 80	0 66	0 72	0 79	0 60	0 66	0 71	0 34	0 37	0 40
1.50	0 62	0 68	0 74	0 61	0 67	0 73	0 55	0 60	0 66	0 29	0 32	0 35
1 6	0 58	0.64	0 69	0 57	0 62	0 68	0 51	0 56	0 61	0 25	0 27	0 30
1 7	0.54	0 60	0.65	0 53	0 58	0 64	0 47	0 52	0 56	0 21	0 23	0.25
1 8	0 51	0 56	0 61	0 50	0 55	0 60	0 44	0 48	0 52	0 18	0 20	0.21
1.9	0.48	0.53	0 58	0 47	0 51	0 56	0 41	0 45	0.49	0.15	0 16	0.18
2 0	0 45	0 50	0 54	0 44	0 49	0 53	0 38	0 42	0 46	0 12	0 14	0 15
2.2	0 41	0 45	0 49	0 40	0 44	0 48	0 34	0 37	0 40	0 08	0.08	0.09
2.4	0 37	0 41	0 44	0 36	0 39	0 43	0 30	0 33	0 36	0 04	0 04	0.05
2.6	0 34	0 37	0 41	0 33	0 36	0 39	0 27	0 29	0 32	0 01	0 01	0.01
2.8	0.31	0 34	0 37	0 30	0 33	0 36	0 24	0 26	0.29	-0 02	-0 02	-0 02
3.0	0.29	0.32	.0 34	0 27	0 30	0 33	0 21	0 24	0 26	-0 04	-0 05	-0 05
3.5	0 24	0 26	0 29	0 23	0 25	0 27	0 17	0 18	0 20	-0 09	-0 10	-0.11
4	0 20	0 22	0 24	0 19	0 21	0 23	0 13	0 14	0 16	-0 13	-0 14	-0.15
5	0.15	0.17	0 18	0 14	0 16	0 17	0 08	0 09	0 10	-0 18	-0 20	-0 21
6	0 12	0 13	0 14	0 11	0 12	0 13	0 05	0 05	0 06	-0 21	-0 23	-0.25
7	0 10	0 11	0 12	0 08	0 09	0 10	0 02	0 03	0.01	-0 23	-0 26	-0.28
8	0 08	0 09	0 09	0 07	0 07	0 08	0 01	0 01	0.01	-0 25	-0 28	-0 30
9	0 06	0 07	0 08	0 05	0 06	0 06	-0 01	-0 01	-0 01	-0 27	-0 29	-0 32
10	0 05	0 06	0 06	0 04	0 05	0 05	-0 02	-0 02	-0 02	-0 28	-0 31	-0.33
12	0 04	0.04	0 04	0 02	0 03	0 03	-0 04	-0 04	-0 04	-0 29	-0 32	-0 35
14	0 02	0 03	0 03	0 01	0 01	0 02	-0 05	-0 05	-0 06	-0 31	-0 34	-0 37
16	0.02	0 02	0 02	0 00	0 00	0 00	-0 06	-0 06	-0 07	-0 31	-0 35	-0 38
18	0 01	0 01	0 01	0 00	0 00	0 00	-0 06	-0 07	-0 08	-0 32	-0 35	-0.39
20	0 00	0.00	0 00	-0 01	-0 01	-0 01	-0 07	-0 08	-0 08	-0 33	-0.36	-0.39
22	0 00	0 00	0 00	-0 01	-0 01	-0 01	-0 07	-0 08	-0 09	-0 33	-0 37	-0 40

* See Density of Weights, p. 75

container. The exact equations connecting the masses and corresponding to equation (2) are:

$$(p' + a') = (s' + c') + [v_s' - (v_s' + v_c')]\sigma'$$

and

$$(p'' + m + a'') = (s'' + c'') + [v_s'' - (v_s'' + v_c'')]\sigma''$$

Assuming p and c to be constant, as must generally be done, and subtracting, gives the general equation (6).

$$m = (s'' - s') - (a'' - a') + [v_s'' - (v_s'' + v_c'')]\sigma'' -$$
$$[v_s' - (v_s' + v_c')]\sigma' \quad (6)$$

If also v_s, v_c, Δ and σ are the same for both weighings, which requires the same temperature and equivalent atmospheric conditions,

$$m = (s'' - s') - (a'' - a') - (v_s'' - v_s')\sigma \quad (7)$$

Table 3.—Buoyancy Reduction Factor (k) for Use in Intercomparison of Weights

(For other factors and for symbols, see Table 2 and p. 74)

$m = s + ks$ Unity of density = g/cm³

Density of weight tested ρ_m	$\Delta^* = 21.5$ Pt or Pt-Ir 1000σ =			$\Delta\dagger = 17$ Gold 1000σ =			$\Delta^* = 8.4$ Brass or bronze 1000σ =			$\Delta^* = 2.7$ Aluminum 1000σ =			$\Delta\dagger = 2.65$ Crystal quartz 1000σ =		
	1 0	1 1	1 2	1 0	1 1	1 2	1.0	1 1	1.2	1 0	1.1	1 2	1 0	1.1	1.2
21 5*	0.000	0.000	0 000	−0 012	−0.014	−0 015	−0.073	−0.080	−0.087	−0.324	−0.357	−0 389	−0 331	−0.364	−0 397
17†	0.012	0 014	0.015	0 000	0 000	0 000	−0 060	−0.066	−0 072	−0.312	−0 343	−0 374	−0.319	−0.350	−0 382
8.4*	0 073	0 080	0.087	+0 060	+0.066	+0 072	0 000	0 000	0.000	−0 252	−0.277	−0.302	−0.258	−0.284	−0.310
2.7*	0.324	0.357	0 389	0 312	0 343	0 375	+0 252	+0 277	+0.302	0.000	0.000	0.000	−0.006	−0.007	−0.008
2 65†	0 331	0 364	0 397	0 319	0 351	0 382	0 258	0 284	0.310	+0.006	+0.007	+0.008	0.000	0.000	0.000

* Density at 0°C, see "Density of Weights," p. 75
† Density at 20°C, see "Density of Weights," p 75

If also $\rho_a{}'' = \rho_a{}' = \sigma$, as when the "empty" portion of the container is filled with air of the same density as the surrounding atmosphere, and the vapor of the "object" weighed is negligible or should be included in m,

$$m = (s'' - s') + (v_m - v_{s''-s'})\sigma \qquad (8)$$

or

$$m = (s'' - s')\left(1 - \frac{\sigma}{\Delta}\right) + v_m\sigma = (s'' - s')\left(\frac{1 - \frac{\sigma}{\Delta}}{1 - \frac{\sigma}{\rho_m}}\right) \qquad (8')$$

In equations (8) and (8') the effect of the container has been eliminated; the equation is of the form of equation (2), and the buoyancy reduction factor from Table 2 may be used.

If the container is exhausted[1] when weighed alone; and if, when the object is being weighed there is in the container only material whose mass should be part of m, then $a' = a'' = 0$ and instead of equations (8) and (8') we have

$$m = (s'' - s') - v_{s''-s'}\,\sigma = (s'' - s')\left(1 - \frac{\sigma}{\Delta}\right) \qquad (9)$$

In this case the buoyant effect of the air on the object weighed has been eliminated, and the ordinary buoyancy reduction factors or equations do not apply (cf. (2) and (3)); Table 2 can not be used.

CORRECTING DENSITY DETERMINATIONS FOR THE BUOYANT EFFECT OF THE AIR

Correcting "Apparent" Values.—Radical differences in the constancy of temperatures or air densities, or such differences as that between equations (8) and (9) above, make it impossible to develop any single correction formula for correcting what are often called "apparent" values of specific gravity, or of density—values which have been determined without proper correction for the buoyant effect of the air. Such values can, however, be corrected in so far as the method and conditions of their determination are known.

Limitations. In general: (1) It is impossible to correct each weighing on which the determination depends, because some unknown mass, volume, or density will generally be needed in order to find the volume of the air displaced. In some cases, however, approximate values may be known with sufficient accuracy for this purpose.

(2) Some special experimental requirements are always involved. Among these may be equal temperatures for two operations, constant volumes (e.g., of pyknometer), negligible changes in the density of the air, etc., or a combination of several of them. A variety of combinations of such requirements may be used, each

[1] As v, is assumed to remain constant, pressure effects must be suitably eliminated.

having its peculiar advantages, and each leading to a different equation.

(3) If the number of experimental requirements is made very small, the resulting equation for true density is very complex. Simplification of the final solution can be accomplished only by increasing the experimental requirements or by introducing approximations into the solution.

No method can be selected as "best."[1] Hence, the material given here is limited to the general fundamental equations, and to the exact solutions for certain cases that are of wide applicability in work of moderate precision. From these it is possible to arrange procedures suited to many different conditions, and to determine the accuracy of the corresponding solutions, and the effects of different errors under various circumstances.

In every case, ρ_m is obtained in the same units as those in which ρ_w is expressed. For the purposes of the following equations, σ may, in general, be expressed either as g/cm³ or as g/ml.

Density of Gases.—The general equations for weighing gases are the same as those for pyknometer determinations of liquids, particularly those for cases in which the pyknometer is exhausted when weighed alone, as in equation (17).

Experimental Requirements.—All the following equations involve two general requirements: (1) That in any one weighing or other operation all objects involved are at the same temperature (in weighing, the temperature of the atmosphere is involved); and (2) that changes in pressure produce no change in any of the volumes; e.g., the volume of the pyknometer or other container must not change when it is exhausted. In addition, each equation involves one or more of the following special requirements:

A. Mass of pyknometer and its counterpoise remains constant: $p' = p'' = p'''$ and $c' = c'' = c'''$.

B. Coefficient of expansion of counterpoise is the same as that of the pyknometer: $\beta_p = \beta_c$. This makes b the same for all weighings.

C. Temperature at which pyknometer is filled is the same for the material being studied as for the calibrating liquid. Therefore $w'' = \rho_w v_t$ and $l''' = \rho_l v_t$.

D. Temperature for all three weighings is the same as that at which the pyknometer is filled. This results in all volumes being constant, in $v_w'' = v_l''' = v'' = v'''$, in $a'' = a''' = 0$, and in the density of each material being constant.

E. Density of the atmosphere the same for all three weighings: $\sigma' = \sigma'' = \sigma'''$.

F. Density of the weights the same in all weighings. This demands that the temperature be the same for all three weighings. See also p. 75.

[1] The advantages and disadvantages of different experimental arrangements, such as the size and mass of the counterpoise used, or the temperature control, do not depend on the form of solution of the equations so much as on the effect of variations and errors that are not shown in the fundamental equations.

G. **Density of air or other material in the "empty" portion of the pyknometer equal to that of the surrounding atmosphere:** $\rho_a' = \sigma', \rho_a'' = \sigma'', \rho_a''' = \sigma'''$.

H. **Pyknometer evacuated when weighed empty.**

I. **Volume of counterpoise equal to "exterior" volume of pyknometer.** $v_c = v_e$.

J. **Volume of counterpoise equals that of the pyknometer itself, excluding the space that would be filled by liquid at the temperature of filling:** $v_c = v_p$.

Pyknometer Determinations.—(1) *Liquids.*—Three weighings are required, from which, under experimental requirement *A*, w'' and l''' are obtained directly by equation (6). Under requirement *C*, $\rho_l = \dfrac{l'''}{w''}\rho_w$.

Therefore under requirements *A* and *C*:

$$\rho_l = \frac{(s'''-s')-(a'''-a')+[v_e'''-(v_e'''+v_e''')]\sigma'''-[v_e'-(v_e'+v_e')]\sigma'}{(s''-s')-(a''-a')+[v_e''-(v_e''+v_e'')]\sigma''-[v_e'-(v_e'+v_e')]\sigma'}\rho_w \quad (10)$$

and

$$v_l = \frac{(s''-s')-(a''-a')+[v_e''-(v_e''+v_e'')]\sigma''-[v_e'-(v_e'+v_e')]\sigma'}{\rho_w} \quad (11)$$

Under requirement *B*, *b* may be introduced for $\dfrac{v_e - v_c}{v_e}$. If also a part of the buoyancy correction for each weighing is made by calculating s_e', s_e'', and s_e'', then the remaining buoyancy reduction terms can be combined and simplified. Then under requirements *A*, *B*, and *C* the equations may be put in the form

$$\rho_l = \frac{s_e'''-s_e'}{s_e''-s_e'}\left[\rho_w + \frac{a''-a'}{v_l} - \frac{b}{v_l}(v_e''\sigma''-v_e'\sigma')\right] - \frac{a'''-a'}{v_l}$$
$$+ \frac{b}{v_l}(v_e'''\sigma'''-v_e'\sigma') \quad (12)$$

and

$$v_l = \frac{(s_e''-s_e')-(a''-a')+b(v_e''\sigma''-v_e'\sigma')}{\rho_w} \quad (13)$$

Under the conditions noted, these equations are perfectly general. They do not involve any mathematical approximations in their derivation and therefore show the proper effect of each quantity. However, in using them, approximate data must, in general, be used, because v_c which is needed in computing v_l cannot be accurately known until after v_l has been computed. If a first approximation is not sufficiently accurate the accuracy may be increased by successive approximations.

[The values of v_e', v_e'' and v_e''' may be computed from the relation $v_e = v_p + v_l = \dfrac{p}{\rho_p} + \dfrac{w}{\rho_w}$, and if the capacity depends solely on temperature (and not on pressure or other factors),
$v_e' = v_e[1 + \beta_p(t' - t)]; \; v_e'' = v_e[1 + \beta_p(t'' - t)];$
$v_e''' = v_e[1 + \beta_p(t''' - t)] \quad (14)$

The values of a', a'', and a''' may be computed from known values of ρ_a and the equations

$$\left.\begin{array}{l} v_a' = v' = v_l[1 + \beta_p(t'-t)] \\ v_a'' = v'' - v_w'' = v_l(\beta_p - \beta_w)(t''-t) \\ v_a''' = v''' - v_l''' = v_l(\beta_p - \beta_l)(t'''-t) \end{array}\right\} \quad (15)$$

Under requirements *D*, *E*, *F*, and *G*, in addition to *A*, *B*, and *C*, (12) becomes

$$\rho_l = \frac{s'''-s'}{s''-s'}(\rho_w - \sigma) + \sigma \quad (16)$$

And under requirement *H* in addition to *A*, *B*, *C*, *D*, *E*, *F*, and *G*

$$\rho_l = \frac{s'''-s'}{s''-s'}\rho_w \quad (17)$$

As shown in equations (16) and (17), experimental requirements *A* to *G* inclusive render the results independent of the size or nature of the counterpoise and of the value of the density of the weights used, though these quantities must be the same for all observations. Including requirement *H* renders the results independent of the

actual value of the density of the air also, but still requires that this value shall be the same for all three weighings.

Under requirement *I*, with *A*, *B*, and *C*, (10) becomes

$$\rho_l = \frac{(s_e'''-s_e')-(a'''-a')}{(s_e''-s_e')-(a''-a')}\rho_w \quad (18)$$

and its equivalent (12), and (13) become

$$\rho_l = \frac{s_e'''-s_e'}{s_e''-s_e'}\left[\rho_w + \frac{a''-a'}{v_l}\right] - \frac{a'''-a'}{v_l} \quad (19)$$

and

$$v_l = \frac{(s_e''-s_e')-(a''-a')}{\rho_w} \quad (20)$$

Under requirement *J*, with *A*, *B*, and *C*, (10) becomes

$$= \frac{(s'''-s')-(a'''-a')+[v'''-v_e''']\sigma'''-[v'-v_e']\sigma'}{(s''-s')-(a''-a')+[v''-v_e'']\sigma''-[v'-v_e']\sigma'}\rho_w \quad (21)$$

and its equivalent (12), and (13) become

$$\rho_l = \frac{s_e'''-s_e'}{s_e''-s_e'}\left[\rho_w + \frac{a''-a'}{v_l} - \frac{1}{v_l}(v''\sigma''-v'\sigma')\right] - \frac{a'''-a'}{v_l} +$$
$$\frac{1}{v_l}(v'''\sigma'''-v'\sigma') \quad (22)$$

and

$$v_l = \frac{(s_e''-s_e')-(a''-a')+v''\sigma''-v'\sigma'}{\rho_w} \quad (23)$$

Pyknometer Determinations.—(2) *Solids.*—The following equations are based on two pyknometer weighings and a separate determination of the mass of the object. If the pyknometer is used as a container for weighing the object this requires two weighings. (*See* p. 76 to 78.)

The symbol " refers to the weighing with the calibrating liquid alone; ''' to the weighing with both this liquid and the object being studied.

Under requirements *A* and *C* only,

$$\rho_m'' = \frac{m\rho_w''}{m - (s'''-s'') + (a'''-a'') - [v_e'''-v_e''-v_e''']\sigma''' + [v_e''-v_e''-v_e'']\sigma''} \quad (24)$$

Under requirement *B*, in addition to *A* and *C*, equation (24) may be put into the form (25) by combining the terms in *s* with those in $v_e\sigma$.

$$\rho_m'' = \frac{m\rho_w''}{m - (s_e'''-s_e'') + (a''' - a'') - b(v_e'''\sigma'''-v_e''\sigma'')} \quad (25)$$

Under requirements *D* and *E*, in addition to *A*, *B*, and *C*,

$$\rho_m = \frac{m\rho_w}{m - (s_e'''-s_e'')} \quad (26)$$

This equation is independent of the magnitudes of σ, c, and v_c, merely requiring their constancy.

Hydrostatic Weighings for Density of Solids.—These equations are based on two weighings; one with the object in air and one with it suspended in a liquid (*e.g.*, water) of known density. The equilibrium equations for these weighings are

$$m' - v_m'\sigma' = s' - v_s'\sigma'$$

and

$$m'' - v_m''\rho_w'' = s'' - v_s''\sigma''$$

the notation being similar to that used for pyknometer weighings. If the mass of the object remains constant (*i.e.*, $m' = m''$), (27) is an exact solution of these equations.

$$\rho_m' = \frac{s_e'}{s_e'-s_e''}(\rho_w''[1+\beta_m(t''-t')] - \sigma') + \sigma' \quad (27)$$

If also all temperatures, the air density, and the density of the weights are the same in the two weighings,

$$\rho_m = \frac{s'}{s'-s''}(\rho_w - \sigma) + \sigma \quad (28)$$

Correction Formula.—When the result of a density determination is calculated without any correction for the buoyant effect

of the air, a false value (ρ_f) is obtained except for pyknometer determinations in which the conditions of the work are those specified for equation (17).

If for pyknometer determinations, these false values were computed by means of the equation $\rho_f = \dfrac{s''' - s'}{s'' - s'}\rho_w$ and for hydrostatic

weighings of solids by means of the equation $\rho_f = \dfrac{s'}{s' - s''}\rho_w$, then to the precision attainable by assuming that the conditions were those specified for equations (16) or (28) the values may be corrected by the equation

$$\rho = \rho_f\left(1 - \frac{\sigma}{\rho_w}\right) + \sigma \tag{29}$$

VOLUME OF A MASS OF LIQUID OF KNOWN WEIGHT IN AIR
(See also p. 73)

Verney Stott and Philip H. Bigg

Symbols. $F = \dfrac{1 - \dfrac{\sigma}{\Delta}}{\rho - \sigma}$; t = temperature of the liquid when its volume is V; t_o = temperature of the liquid when weighed; V = volume of the liquid at temperature t; W = weight of the liquid in air against weights of density Δ; ρ, ρ_o = density of the liquid at t and at t_o, respectively; σ = density of air at time of weighing.

If densities are expressed in g/cm³, and W in g, V is in cm³; if densities are in g/ml and W in g, V is in ml; if densities are in lb./gal., and W in lb., V is in gal.; etc.

The exact relations connecting these quantities are given by the equation

$$V = \frac{W\left(\dfrac{1 - \dfrac{\sigma}{\Delta}}{1 - \dfrac{\sigma}{\rho_o}}\right)}{\rho} = \frac{W\left(\dfrac{1 - \dfrac{\sigma}{\Delta}}{1 - \dfrac{\sigma}{\rho}}\right)}{\rho}\left(\dfrac{1 - \dfrac{\sigma}{\rho}}{1 - \dfrac{\sigma}{\rho_o}}\right) = FW\left(\dfrac{1 - \dfrac{\sigma}{\rho}}{1 - \dfrac{\sigma}{\rho_o}}\right)$$

VALUES OF F FOR WATER AND MERCURY
(Liquids are air-free)

$$V = FW\cdot\frac{1 - \dfrac{\sigma}{\rho}}{1 - \dfrac{\sigma}{\rho_o}}$$

In many cases the factor $\left(\dfrac{1 - \dfrac{\sigma}{\rho}}{1 - \dfrac{\sigma}{\rho_o}}\right)$ does not differ significantly from unity. If $t_o = 20°C$, the greatest value of this factor for the temperature range covered by the following table differs from unity by only 7.3×10^{-6} for water and by 0.48×10^{-6} for mercury.

If $t_o = t$, $V = FW$. For water, $F = 1 + 0.001\,K_{H_2O}$; for mercury, $F = 0.07 + 0.001\,K_{Hg}$.

Unit of F = milliliter per g of W; of t = °C. Assumes[*] $\sigma = 0.0012$ g/ml; $\Delta = 8.3$ g/ml.

t	K_{H_2O}	K_{Hg}	t	K_{H_2O}	K_{Hg}	t	K_{H_2O}	K_{Hg}	t	K_{H_2O}	K_{Hg}	t	K_{H_2O}	K_{Hg}
0	1 18$_9$	3 55$_0$	10	1 33$_0$	3 68$_3$	20	2.83$_2$	3.817	30	5.41$_0$	3.951	40	8 89$_0$	4.08$_5$
1	1 13$_0$	3 56$_3$	11	1 42$_5$	3 697	21	3 04$_4$	3 83$_0$	31	5.72$_0$	3 964	41		4.09$_8$
2	1 08$_9$	3 57$_6$	12	1 53$_3$	3 71$_0$	22	3 267	3 84$_4$	32	6 03$_8$	3.977	42		4.111
3	1 06$_5$	3 59$_0$	13	1 65$_4$	3 72$_3$	23	3 50$_1$	3 857	33	6 36$_6$	3.991	43		4.12$_5$
4	1 057	3 60$_3$	14	1 78$_8$	3 737	24	3 74$_4$	3 87$_0$	34	6.70$_2$	4 00$_4$	44		4.13$_8$
5	1 06$_5$	3 61$_6$	15	1 93$_3$	3 75$_0$	25	3 99$_8$	3 88$_4$	35	7 04$_6$	4.01$_8$	45		4.15$_2$
6	1.08$_9$	3 63$_0$	16	2 09$_0$	3 76$_3$	26	4 26$_1$	3 897	36	7 39$_9$	4 03$_1$	46		4.16$_5$
7	1 127	3 64$_3$	17	2 25$_9$	3 777	27	4 53$_4$	3 91$_0$	37	7.76$_0$	4.044	47		4 17$_8$
8	1 18$_1$	3 65$_6$	18	2 43$_8$	3 79$_0$	28	4.817	3 92$_4$	38	8 12$_9$	4 05$_8$	48		4 19$_2$
9	1 24$_8$	3.67$_0$	19	2.63$_0$	3 80$_3$	29	5.10$_9$	3.937	39	8.50$_5$	4.071	49		4 20$_5$
												50		4.21$_9$

[*] The increase (dK) produced in K by changing Δ to $\Delta(1 + \delta)$ and σ to $\sigma(1 + s)$ is closely given ($\pm ca$ 1 %) for the range of this table by the equations:

$$dK_{H_2O} = 0.145(7.3s + 0.997\delta + 8.3s\delta)\frac{1}{1 + \delta}$$

$$dK_{Hg} = 0.00078(-5.3s + 13.6\delta + 8.3s\delta)\frac{1}{1 + \delta},$$

units being those of this table. For uncertainties in σ, and for the variation of σ with pressure, temperature, and humidity, see p. 78. When brass weights are not used, δ will, in general, be large, in such cases it is desirable to transform the equations once for all by inserting the proper value for δ; they will take the convenient form $dK = a + bs$. If $\delta = 0$, $dK_{H_2O} = 1.06\,s$, $dK_{Hg} = 0.00414\,s$. If $s = 0$, $dK_{H_2O} = 0.145\dfrac{\delta}{1 + \delta}$; $dK_{Hg} = 0.0105\dfrac{\delta}{1 + \delta}$.

Example.— (1) If $\sigma = 0.00132$ and $\Delta = 8.383$, $s = 0.1$, $\delta = 0.01$ and $dK_{H_2O} = 0.145(0.73 + 0.01 + 0.008)\dfrac{1}{1.01} = 0.144(0.75) = 0.108$. Hence, if $t = 19°C$, $K_{H_2O} = 2.63 + 0.108 = 2.74$.

(2) If $\sigma = 0.00132$ and $\Delta = 2.65$ (quartz), $s = 0.1$, $(1 + \delta) = \dfrac{2.65}{8.3}$, $\delta = -\dfrac{5.65}{8.3}$, and $dK_{Hg} = 0.00078(-0.53 - 9.26 - 0.565)(3.13) = -0.025_3$. Hence, if $t = 25°C$, $K_{Hg} = 3.884 - 0.025 = 3.85_9$.

STANDARD BUFFER SOLUTIONS AND ACID-BASE INDICATORS

Mansfield Clark

In the following tables pH represents (formalistically) $\log_{10}$ $\frac{1}{[H^+]}$ where $[H^+]$ is the symbol for grams of hydrogen ions per liter. Since there is a disagreement concerning the precise interpretation of experimental values, the experimental meaning of pH is defined by the set of conditions described below (8, 57).

The normal hydrogen-electrode is regarded as a properly coated, noble metal, under one atmosphere partial-pressure of hydrogen, immersed in a solution normal with respect to hydrogen ions. The difference of potential between electrode and solution is regarded as zero at all temperatures.

The following values are regarded as *standard* differences of potential (E_c) (liquid-junction potential-difference being eliminated) between the tenth-normal KCl—Hg_2Cl_2—Hg half-cell and the hypothetical, normal hydrogen-electrode.

$T°$	18	20	25	30	37.5	40	50	60
E_c	0 3380	0 3379	0 3376	0 3372	0 3364	0 3360	0 3341	0 3317

For present purposes it is assumed that the liquid-junction potential-difference between an Hg_2Cl_2 half-cell solution and the solution the pH of which is under measurement has been eliminated when there has been interposed a saturated solution of KCl, or when there has been employed the Bjerrum extrapolation (4) from measurements made with $3.5N$ KCl and $1.75N$ KCl as interposed solutions.

When the electromotive force, e m f, of the "chain",

H_2Pt	$[H^+]$	KCl saturated	KCl 0.1N	Hg_2Cl_2Hg

is measured under the above conditions, and the Hg is positive to the Pt, pH is calculated from the equation

$$\frac{E.M.F. - E_c}{0.000\ 198\ 37(273.09 + t)} = pH.$$

(See (8, 37, 45, 64) and references therein on potentiometric measurement of pH.)

The chief modes of employing indicators for the determination of pH may be illustrated by the following examples.

I. A solution having been found to induce a blue color with brom thymol blue (see No. 139, Table 3A), a yellow color with thymol blue (No. 129), and a color intermediate between yellow and red with phenol red (No. 142) is judged to have a pH value between 7.0 and 7.8. Then to 10 ± 0.05 cc of solution are added 5 drops 0.04 % phenol red solution (made by dissolving 0.1 g phenol red in 28.5 cc 0.01N NaOH solution and diluting to 250 cc). The resulting mixture is then compared with standards made by adding 5 drops of the same phenol red solution to each of 10 ± 0.05 cc portions of buffers having pH values of 7.0, 7.2, 7.4, 7.6, etc. (See Table 1A.)

The comparison is made in containers of identical dimensions and under uniform illumination. It is found that the tested solution has a color intermediate and half-way between those of buffers 7.4 and 7.6, and since the total salt contents of the tested solution and of the buffers are of the same order of magnitude, and since the solution contains no protein or substance known to affect the indicator, 7.5 is judged to be the true pH value of the tested solution (8, 11, 31, 37, 45, 53, 54, 56).

II. A solution is found to induce a partial color transformation of phenol red. Using uniform containers (e g., test tubes) there are prepared:

(1) A mixture of 10 ± 0.05 cc solution under test and 10 drops standard phenol red solution (see I).

(2) A mixture of x drops of indicator and sufficient buffer solution of the value shown in column B of Table 3A to equal the total volume of solution 1.

(3) A mixture of $10 - x$ drops of indicator and sufficient buffer of the value shown in column C of Table 3A to equal the total volume of solution 1.

x is varied and there is found at $x = 4$ a match in color between solution 1 and superposed solutions 2 and 3. From the relation:

$$pH = pK + \log_{10} \frac{r}{x}, \text{ and the value 7 8 for pK given in Table}$$

3A it is calculated that the value of the tested solution is 7.6 (see in addition to the general references under I (2, 19, 20, 22, 34, 63).

III A solution is found to induce a partial color-transformation in m-nitrophenol (No 15, Table 3C) It is found that 10 cc of the tested solution plus 1 cc of 0.3% m-nitrophenol matches in color 11 cc of an alkalinized solution containing 0.2 cc of 0.3% m-nitrophenol. It is thus shown that the tested solution has induced a 20% transformation If a is the percentage transformation of the indicator, pH is calculated from

$$pH = pK + \log \frac{a}{100 - a}$$

In the case at hand $a = 20$, the temperature of the measurement was 25° and the total salt content of the solution was of the order of magnitude of 0.15M. Hence from Table 3C, pK is taken as 8 16. By the above equation pH = 7.56

The equation $pH = pK + \log \frac{a}{100 - a}$ cannot be used with picric acid, phenolphthalein or Alizarine yellow GG listed in Table 3C, since these indicators do not behave as monoacidic within the range of pH specified. Empirical data (38) for phenolphthalein and Alizarine yellow GG are shown in Table 4. It is best to vary the amounts of indicator used till the most favorable color-differences are found. (In addition to the material found in the general references under I see (30, 31, 38, 39) for method III.)

pK in the tables represents the pH at which there is an apparent half-transformation of the indicator. For indicators behaving as monoacidic or monobasic, within the zone of pH designated, pK is log $1/Ka$ when Ka is the "apparent dissociation constant" (43). When an indicator, such as phenolphthalein, is known not to behave as monoacidic within the range of pH designated, pK is bracketed

pK values listed in Tables 3A and 3C are uniform with respect to the bases of reference Those of the indicators in the general list (Table 2) are referred to such a variety of bases that tabulation is impracticable. The reader is therefore referred to original articles (8, 31, 37, 43, 45, 51, 58, 59, 60, 61, 67).)

The values assigned to useful pH ranges are somewhat arbitrary, depending upon concentration of indicator, the spectral distribution of illumination, and psychological preferences.

Indicator solutions are affected to various degrees by

   a. Total salt content.

   b. Specific ions: e g., alizarine red S is affected by borates differently than by phosphates (67).

   c. Colloidal suspensions, protein solutions, etc.: e.g., congo red in a gelatine solution of pH 3.6 behaved as if the pH were 5.6 (53). Neutral red in soap solutions forms a fatty acid complex (27).

   d. Presence of immiscible solvents: e.g., chloroform used for disinfection removes benzene-azo-benzyl-aniline from the aqueous phase (53).

e. Mixed solvents and change of solvent (3, 31, 32, 40, 62).

f. Temperature. See Table 3A, 3C.

g. Time: *e.g*, water blue changes color slowly and propyl red precipitates.

h. Destructive agents: *e.g.*, methyl red is irreversibly reduced in some bacterial cultures.

Since it is impracticable to tabulate all available data, only representative "salt" and temperature effects are given in Tables 3A, 3B and 4.

The indicators of Table 3 include the better of those which may be used in acidimetric and alkalimetric titration. (For principles see (5, 31, 43, 45).)

Table 1.—Standard Buffer Solutions

The following tables give the compositions of solutions which furnish, at the temperatures indicated, values of pH which conform in essential respects to the specifications listed in the general notes above. Recalculation to make the conformity rigid would involve changes in the original data which would be less than the uncertainties of the working standards used in the experiments. The solutions listed may serve as standards for the colorimetric measurements of pH. The solutions suffer relatively slight displacement of pH with addition or subtraction of small proportions of acid or alkali. This property is referred to as that of a *buffer* (*puffer, tampon*). (For buffer solutions see (8, 37, 45, 64).)

A. Standard Buffer Solutions of Clark and Lubs (10) at 20° 50 cc A + x cc B diluted to 200 cc

A = 0.2M KCl* B = 0.2M HCl		A = 0.2M KH o-phthal- ate B = 0.2M HCl		A = 0.2M KH o-phthal- ate B = 0.2M NaOH		A = 0.2M KH₂PO₄ B = 0.2M NaOH		A = 0.2M H₃BO₃† + 0.2M KCl B = 0.2M NaOH	
pH	cc B	pH	cc B	pH	cc B	pH	cc B	pH	cc B
1.2	64.5	2.2	46.70	4.0	0.40	5.8	3.72	7.8	2.61
1.4	41.5	2.4	39.60	4.2	3.70	6.0	5.70	8.0	3.97
1.6	26.3	2.6	32.95	4.4	7.50	6.2	8.60	8.2	5.90
1.8	16.6	2.8	26.42	4.6	12.15	6.4	12.60	8.4	8.50
2.0	10.6	3.0	20.32	4.8	17.70	6.6	17.80	8.6	12.00
2.2	6.7	3.2	14.70	5.0	23.85	6.8	23.65	8.8	16.30
		3.4	9.90	5.2	29.95	7.0	29.63	9.0	21.30
		3.6	5.97	5.4	35.45	7.2	35.00	9.2	26.70
		3.8	2.63	5.6	39.85	7.4	39.50	9.4	32.00
				5.8	43.00	7.6	42.80	9.6	36.85
				6.0	45.45	7.8	45.20	9.8	40.80
				6.2	47.00	8.0	46.80	10.0	43.90

B. Sørensen's Glycocoll-NaCl-HCl Mixtures(56)

Glycocoll solution: 0.1M Glycocoll + 0.1M NaCl per l; HCl 0.1N. Values hold between 10°-70° (66)

Glycocoll (cc)	0.0	1.0	2.0	3.0	4.0	5.0
HCl (cc)	10.0	9.0	8.0	7.0	6.0	5.0
pH	1.04	1.15	1.25	1.42	1.65	1.93

Glycocoll (cc)	6.0	7.0	8.0	9.0	9.5
HCl (cc)	4.0	3.0	2.0	1.0	0.5
pH	2.28	2.61	2.92	3.34	3.68

C. Sørensen's Citrate-HCl Mixtures (56)

Citrate solution: 21.008 g crystn. citric acid + 200 cc N NaOH per l; HCl: 0.1N. Values hold between 10°-70° (66)

Citrate (cc)	0.0	1.0	2.0	3.0	3.33	4.0	4.5	4.75
HCl (cc)	10.0	9.0	8.0	7.0	6.67	6.0	5.5	5.25
pH	1.04	1.17	1.42	1.93	2.27	2.97	3.36	3.53

* The pH values of these mixtures are given by Clark and Lubs as *preliminary* measurements
† The old atomic weight (11.0) of boron is used throughout these tables

Citrate (cc)	5.0	5.5	6.0	7.0	8.0	9.0	9.5	10.0
HCl (cc)	5.0	4.5	4.0	3.0	2.0	1.0	0.5	0.0
pH	3.69	3.95	4.16	4.45	4.65	4.83	4.89	4.96

D. Sørensen's Phosphate Mixtures (55, 56)

9.078 g KH₂PO₄, 11.876 g Na₂HPO₄.2H₂O each per l. Values hold between 10°-70° (66).

Na₂HPO₄ (cc)	0.25	0.5	1.0	2.0	3.0	4.0
KH₂PO₄ (cc)	9.75	9.5	9.0	8.0	7.0	6.0
pH	5.29	5.59	5.91	6.24	6.47	6.64

Na₂HPO₄ (cc)	5.0	6.0	7.0	8.0	9.0	9.5
KH₂PO₄ (cc)	5.0	4.0	3.0	2.0	1.0	0.5
pH	6.81	6.98	7.17	7.38	7.73	8.04

E. Sørensen's Citrate-NaOH Mixtures (56); Walbum's Values (66)

Citrate solution; 21.008 g crystn. citric acid + 200 cc N NaOH per l; NaOH: 0.1N

Volume parts		Temperature						
Citrate	NaOH	10°	20°	30°	40°	50°	60°	70°
10.0	0.0	4.93	4.96	5.00	5.04	5.07	5.10	5.14
9.5	0.5	4.99	5.02	5.06	5.10	5.13	5.16	5.20
9.0	1.0	5.08	5.11	5.15	5.19	5.22	5.25	5.29
8.0	2.0	5.27	5.31	5.35	5.39	5.42	5.45	5.49
7.0	3.0	5.53	5.57	5.60	5.64	5.67	5.71	5.75
6.0	4.0	5.94	5.98	6.01	6.04	6.08	6.12	6.15
5.5	4.5	6.30	6.34	6.37	6.41	6.44	6.47	6.51
5.25	4.75	6.65	6.69	6.72	6.76	6.79	6.83	6.86

F. Sørensen's Borate-HCl Mixtures (56); Walbum's Values (66)

Borate: 12.404 g H₃BO₃ + 100 cc N NaOH per l; HCl: 0.1N

Volume parts		Temperature						
Borate	HCl	10°	20°	30°	40°	50°	60°	70°
10.0	0.0	9.30	9.23	9.15	9.08	9.00	8.93	8.86
9.5	0.5	9.22	9.15	9.08	9.01	8.94	8.87	8.80
9.0	1.0	9.14	9.07	9.01	8.94	8.87	8.80	8.74
8.5	1.5	9.06	8.98	8.92	8.86	8.80	8.73	8.67
8.0	2.0	8.96	8.89	8.83	8.77	8.71	8.65	8.59
7.5	2.5	8.84	8.79	8.72	8.67	8.61	8.55	8.50
7.0	3.0	8.72	8.67	8.61	8.56	8.50	8.45	8.40
6.5	3.5	8.54	8.49	8.44	8.40	8.35	8.30	8.26
6.0	4.0	8.32	8.27	8.23	8.19	8.15	8.11	8.08
5.75	4.25	8.17	8.13	8.09	8.06	8.02	7.98	7.95
5.5	4.5	7.96	7.93	7.89	7.86	7.82	7.79	7.76
5.25	4.75	7.64	7.61	7.58	7.55	7.52	7.49	7.47

H. Sørensen's Borate-NaOH Mixtures (56); Walbum's Values (66)

Borate: 12.404 g H₃BO₃ + 100 cc N NaOH per l; NaOH: 0.1N

Volume parts		Temperature						
Borate	NaOH	10°	14°	18°	22°	26°	30°	34°	37°
10	0.0	9.30	9.27	9.24	9.21	9.18	9.15	9.13	9.11
9	1	9.42	9.39	9.36	9.33	9.29	9.26	9.23	9.20
8	2	9.57	9.54	9.50	9.46	9.43	9.39	9.35	9.32
7	3	9.76	9.72	9.68	9.63	9.59	9.55	9.50	9.47
6	4	10.06	10.02	9.97	9.91	9.86	9.80	9.75	9.71
5	5	11.24	11.16	11.08	10.99	10.91	10.82	10.74	10.68
4	6	12.64	12.51	12.38	12.25	12.13	12.00	11.87	11.77

G. Sørensen's Glycocoll-NaCl-NaOH Mixtures (**); Walbum's Values (**)

Glycocoll: 7.505 g glycocoll + 5.85 g NaCl per l; NaOH: 0.1N

Volume parts		Temperature														
Glycocoll	NaOH	10°	12°	14°	16°	18°	20°	22°	24°	26°	28°	30°	32°	34°	37°	40°
9 5	0.5	8.75	8.70	8 66	8 62	8 58	8 53	8 49	8 45	8 40	8 37	8 32	8 28	8 24	8.18	8.12
9.0	1.0	9.10	9.06	9.02	8 97	8 93	8 88	8 84	8.79	8.75	8.71	8 67	8.62	8.58	8.52	8.45
8.0	2.0	9.54	9 50	9.45	9 40	9 36	9 31	9 26	9 22	9 17	9.13	9 08	9 04	9 00	8.92	8.85
7.0	3.0	9 90	9.85	9 80	9 75	9.71	9 66	9 61	9 56	9.51	9 46	9 42	9 37	9 32	9.25	9.18
6 0	4.0	10 34	10.29	10.24	10.18	10 14	10 09	10 03	9 98	9.93	9 88	9 83	9 78	9.73	9.66	9.58
5.5	4.5	10.68	10.63	10.58	10 53	10.48	10 42	10 37	10 32	10.27	10 22	10 17	10 12	10 07	9.99	9.91
5.1	4 9	11 29	11.24	11 18	11.12	11 07	11 01	10 96	10 90	10.85	10 79	10.74	10 68	10.62	10.54	10.46
5 0	5.0	11.53	11.48	11 42	11 36	11.31	11 25	11 20	11.14	11 09	11 03	10 97	10 92	10 86	10.78	10.70
4 9	5.1	11.80	11 74	11.68	11 62	11 57	11 51	11 45	11 39	11 33	11.27	11 22	11 16	11.10	11.02	10.93
4 5	5.5	12 34	12 28	12 22	12 16	12 10	12 04	11 98	11 92	11 86	11 80	11 74	11 68	11.62	11.53	11.44
4 0	6.0	12.65	12.59	12 52	12.46	12 40	12 33	12 27	12 21	12 15	12 09	12 03	11.96	11 90	11.81	11.72
3.0	7.0	12.92	12.86	12 80	12.73	12.67	12 60	12 54	12.48	12 42	12 35	12 29	12 23	12.17	12.07	11.98
2 0	8 0	13 12	13 06	12 99	12 92	12.86	12 79	12 73	12.66	12.60	12.53	12.47	12 41	12 34	12.25	12.15
1 0	9 0	13.23	13.16	13 09	13 03	12 97	12 90	12 83	12 77	12 70	12 64	12 57	12 51	12 45	12.35	12.25

Volume parts		Temperature														
Glycocoll	NaOH	42°	44°	46°	48°	50°	52°	54°	56°	58°	60°	62°	64°	66°	68°	70°
9.5	0 5	8.07	8.03	7 99	7 95	7 91	7 86	7 82	7.78	7 74	7 69	7 65	7.61	7 56	7.52	7.48
9 0	1 0	8.41	8 37	8 32	8 28	8 24	8 19	8 14	8 10	8 06	8 02	7 97	7 93	7 88	7.84	7.79
8.0	2.0	8.81	8 76	8 72	8 67	8 63	8 58	8 53	8 49	8 44	8.40	8 35	8 30	8 26	8.21	8.16
7 0	3.0	9.13	9 08	9 03	8 99	8 94	8 89	8 84	8 79	8 74	8 70	8.65	8 60	8.55	8.50	8.45
6.0	4 0	9.53	9 48	9 43	9 38	9 33	9 28	9 23	9 18	9 13	9 08	9 03	8 98	8.93	8.88	8.82
5.5	4 5	9 86	9 81	9 76	9 71	9 66	9 61	9.56	9 51	9 46	9 41	9 35	9 30	9.25	9.20	9.15
5.1	4.9	10.40	10.35	10 29	10 24	10 18	10.13	10 07	10 02	9 96	9 90	9 85	9 79	9.74	9.68	9.62
5.0	5 0	10 64	10 59	10 54	10.48	10 43	10 37	10 32	10 26	10 20	10.14	10 09	10 04	9 98	9.93	9.87
4.9	5 1	10.87	10.81	10 75	10 69	10 64	10 58	10 52	10.46	10 40	10 35	10 29	10 23	10 17	10.11	10.05
4.5	5.5	11.38	11.32	11 26	11 20	11.14	11 08	11 02	10 96	10 90	10 84	10 78	10 72	10.66	10.60	10.54
4.0	6.0	11 65	11 59	11 53	11 47	11.41	11 34	11.28	11.22	11 16	11 10	11 03	10 97	10.91	10.84	10.78
3 0	7.0	11 91	11 85	11 79	11.73	11.66	11 60	11 54	11 47	11 41	11 35	11.28	11.22	11.16	11.09	11.03
2 0	8 0	12.08	12.02	11 96	11 89	11.83	11.77	11.70	11.64	11.57	11.51	11.44	11.38	11.31	11.25	11.18
1 0	9 0	12 19	12.13	12 06	12 00	11 94	11 87	11 80	11 74	11 67	11 61	11 54	11 48	11 41	11.35	11.28

J. pH Values of Borax-borate Mixtures at 18°C and "Salt-effects" for Phenolphthalein and α-Naphtholphthalein Palitzsch (44)

Borax solution: 19 108 g Na₂B₄O₇.10H₂O in 1 l.   Boric acid solution: 12.404 g H₃BO₃ + 2.925 g NaCl in 1 l

Standard solutions			True pH values of sea water containing S parts per 1000 salinity at color-match with standard										
Borax cc	Boric acid cc	pH	S = 36	S = 30	S = 26	S = 22	S = 18	S = 14	S = 10	S = 6	S = 4	S = 2	S = 1	
6.0	4 0	8 69	8 48	8 49	8 50	8 52	8 54	8 57	8 59	8 63	8 66	8 69	8.72	Phenolphthalein
5.5	4.5	8 60	8 39	8 40	8 41	8 43	8 45	8 48	8 50	8 54	8 57	8 60	8.63	
5 0	5 0	8 51	8 30	8 31	8 32	8 34	8 36	8 39	8 41	8 45	8 48	8 51	8.54	
4.5	5 5	8 41	8.20	8 21	8 22	8 24	8 26	8 29	8 31	8 35	8 38	8.41	8 44	
4.0	6 0	8 31	8.10	8.11	8 12	8 14	8 16	8 19	8 21	8 25	8 28	8.31	8.34	
3.5	6 5	8 20	7 99	8 00	8 01	8 03	8 05	8 08	8 10	8.14	8 17	8 20	8.23	
4.5	5.5	8 41	8 19	8 20	8 21	8 23	8 25	8 28	8.32	8 37	8 40	8.45	8.48	α-Naphtholphthalein
4.0	6.0	8 31	8 09	8.10	8 11	8 13	8 15	8 18	8 22	8 27	8 30	8.35	8.38	
3 5	6 5	8 20	7.98	7.99	8 00	8 02	8 04	8.07	8 11	8 16	8 19	8.24	8.27	
3.0	7.0	8.08	7 86	7 87	7 88	7 90	7 92	7 95	7 99	8.04	8 07	8 12	8.15	
2.5	7.5	7.94	7 72	7.73	7.74	7.76	7 78	7 81	7.85	7 90	7 93	7.98	8.01	
2 3	7.7	7.88	7 66	7.67	7 68	7 70	7 72	7.75	7 79	7.84	7.87	7.92	7.95	
2.0	8.0	7 78	7.56	7 57	7 58	7.60	7 62	7 65	7.69	7.74	7 77	7.82	7.85	
1.5	8.5	7.60	7.38	7 39	7.40	7.42	7 44	7.47	7.51	7.56	7.59	7.64	7.67	
1.0	9.0	7 36	7 14	7.15	7.16	7 18	7 20	7 23	7.27	7.32	7.35	7.40	7.43	
0 6	9.4	7.09	6 87	6 88	6.89	6 91	6 93	6.96	7.00	7.05	7.08	7.13	7.16	
0.3	9 7	6.77	6.55	6 56	6 57	6 59	6 61	6 64	6 68	6.73	6.76	6.81	6.84	

H. Sørensen's Borate-NaOH Mixtures.—(Continued)

Volume parts		Temperature						
Borate	NaOH	40°	44°	48°	52°	56°	60°	64°	70°
10	0 0	9 08	9 05	9 02	9 00	8 97	8.93	8 90	8.86
9	1	9 18	9 15	9 11	9 08	9 05	9 01	8 98	8 94
8	2	9 30	9 26	9 22	9 18	9 15	9.11	9 08	9 02
7	3	9 44	9 40	9 35	9 31	9 27	9 22	9 18	9 12
6	4	9 07	9 62	9 56	9 51	9 46	9 40	9 35	9 28
5	5	10 61	10 53	10 44	10 36	10 27	10 19	10.10	9 98
4	6	11 68	11 55	11 42	11 29	11 17	11 04	10 91	10 72

I. Acetic Acid-acetate Mixtures; Walpole's Values (Recalculated) [68]

CH_3CO_2H M.	...	0 185	0 176	0 164	0.147	0.126	0.102
CH_3CO_2Na M	...	0 015	0 024	0 036	0 053	0 074	0.098
pH		3 6	3 8	4 0	4 2	4 4	4 6

CH_3CO_2H M	..		0 080	0.059	0.042	0.029	0.019
CH_3CO_2Na M			0 120	0 141	0 158	0 171	0 181
pH			4 8	5 0	5 2	5 4	5 6

TABLE 2.—GENERAL LIST OF INDICATORS

The following list of indicators includes all those for which data on the pH-ranges have been found. Many of the data of this table are to be regarded with caution, because in some cases the names proposed are inadequate for complete identification, and in others names have been given to materials of uncertain composition (*8, 11, 31, 37, 45, 53, 54, 56, 64*).

The Schultz (S......) and Rowe (R......) numbers are taken from the 1923 (*52*) and 1924 (*48*) editions, respectively, of these works. Delicate shades of meaning in the color nomenclature have been avoided, as data regarding the purity of the compounds have often been lacking. The abbreviations used are as follows: b, blue; br, brown; c, colorless; f, fades; fl, fluorescent; g, green; o, orange; p, pink; pu, purple; r, red; v, violet; y, yellow. pK is the pH at which there is an apparent half-transformation of the indicator. * indicates that the indicator has been studied in sufficient detail to be used in supplementing the lists of Table 3.

Nitro Compounds

Index No.	Indicator	Color and useful range pH	Lit.
1	2, 4, 6-Trinitrophenol; Picric acid [S 5; R. 7]	c 0 0– 1.3 y	(31, 39)
2	2, 6-Dinitrophenol [Michaelis' β]	c 2.0– 4.0 y	(31, 38, 39)
3	2, 4-Dinitro-α-naphthol; Manchester yellow [S. 6; R 9]	y 2.0– 4.0 y	(9)
4	2, 4-Dinitrophenol [Michaelis' α]	c 2.6– 4.4 y	(31, 38, 39)
5	Dinitrohydroquinol	3–10	(23, 46)
6	Nitrohydroquinol	3–11	(46)
7	2, 3-Dinitrophenol [Michaelis' ε]	c 3.0– 5 9 y	(31, 38, 39)
8	2, 5-Dinitrophenol [Michaelis' γ]	c 4.0– 5 8 y	(31, 38, 39)
9	2, 6-Dinitro-4-aminophenol; Isopicramic acid.	p 4 1– 5 6 y	(67)
10	3, 4-Dinitrophenol [Michaelis' δ]	c 4 3– 6 3 y	(38, 39)
11	4-Nitro-6-aminoguaiacol	y 4 5– 8 0 r	(35)
12	p-Nitrophenol	c 5.6– 7 6 y	(31, 38, 39, 56)
13	o-Nitrophenol	c 5 0– 7 0 y	(46)
14	*Dinitrobenzoylene urea	c 6 0– 8 0 y	(6)
15	m-Nitrophenol	c 6 8– 8 6 y	(31, 38, 39)
16	2, 4, 6-Trinitrophenyl-methyl-nitroamine; Nitramine	c 10 8–13 0 br	(31, 33)
17	sym.-Trinitrobenzene.	c 12 0 14 0 o; f	(50)
18	2, 4, 6-Trinitrotoluene	p 11 5–14 0 o	(9)

Mono-azo Compounds

19	p-Toluene-azo-phenyl-aniline	1 0– 2 0	(53, 54, 56)
20	p-Carboxybenzene-azo-dimethylaniline; Para methyl red	r 1 0– 3 0 y	(9, 60)
21	p-Toluene-azo-phenyl-α-naphthylamine	1 1– 1 9	(53, 54, 56)
22	Benzene-azo-diphenylamine	p 1 2– 2 1 y	(56)
23	m-Benzenesulfonic acid-azo-diphenylamine; Metanil yellow [S. 134; R. 138]	r 1.2– 2 3 y	(56)
24	Benzene-azo-phenyl-α-naphthylamine	v 1 4– 2 6 o	(53, 54, 56)
25	p-Benzenesulfonic acid-azo-diphenylamine; Tropaeolin OO [S. 139; R. 143]	r 1 4– 2 6 y	(56, 60)
26	o-Toluene-azo-o-toluidine; Spirit yellow R [S. 68; R. 17]	1 4– 2 9	(53, 54, 56)
27	p-Toluene-azo-benzyl-α-naphthylamine.	1 6– 2 6	(53, 54, 56)
28	p-Toluene-azo-benzyl-aniline	1 6– 2 8	(53, 54, 56)
29	Benzene-azo-benzyl-α-naphthylamine. ...	1 9– 2 9	(53, 54, 56)
30	Benzene-azo-aniline; Amino-azo-benzene [S. 31; R. 15]	y 1 9– 3 3 y	(53, 54, 56, 60)
31	p-Benzenesulfonic acid-azo-aniline	r 1 9– 3 3 y	(52, 53, 54, 60)
32	p-Benzenesulfonic acid-azo-benzylaniline	r 1 9– 3 3 y	(56, 60)
33	m-Carboxybenzene-azo-dimethylaniline	p 2 3– 3 3 y	(56)
34	Benzene-azo-benzylaniline . . .	p 2 3– 3 3 y	(56)
35	p-Benzenesulfonic acid-azo-m-chlorodiethylaniline	r 2 6– 4 0 y	(56, 60)
36	m-Nitrobenzene-azo-β-naphthol-3, 6-disulfonic acid; Orange III [S. 47; R. 39]	r 2 6 4 6 y	(9)
37	Benzene-azo-dimethylaniline; Topfer's indicator [S. 32; R. 19] ...	r 2 9– 4 0 y	(56, 60)
38	o-Carboxybenzene-azo-α-naphthylamine	r 2 9– 5 8 y	(61)
39	p-Benzenesulfonic acid-azo-o-toluidine	mid-point 2 9	(60)

MONO-AZO COMPOUNDS.—(Continued)

Index No.	Indicator	Color and useful range pH	Lit.
40	p-Benzenesulfonic acid-azo-m-xylidine.	mid-point 2 9	(60)
41	o-Carboxybenzene-azo-diphenylamine	p 3 0- 4 6 y	(11)
42	p-Benzenesulfonic acid-azo-methylaniline	r 3.1- 4 2 y	(53, 54, 56, 60)
43	p-Benzenesulfonic acid-azo-ethyl aniline	r 3.1- 4 4 y	(53, 54, 56, 60)
44	p-Benzenesulfonic acid-azo-dimethylaniline; Methyl orange [S. 138; R. 142]	r 3 1- 4.4 y	(56, 60)
45	p-Benzenesulfonic acid-azo-diethylaniline; Ethyl orange.	r 3 5- 4.5 y	(53, 54, 56, 60)
46	o-Benzenesulfonic acid-azo-dimethylaniline	mid-point 3 5	(60)
47	p-Benzenesulfonic acid-azo-m-toluidine.	mid-point 3 5	(60)
48	p-Benzenesulfonic acid-azo-p-xylidine	mid-point 3 6	(60)
49	*p-Sulfo-o-methoxybenzene-azo-dimethyl-α-naphthylamine	b 3 5- 4 9 o	(42)
50	p-Benzenesulfonic acid-azo-α-naphthylamine	r 3 5- 5 7 y	(56, 61)
51	p-Benzenesulfonic acid-azo-phenyl-α-naphthylamine	v 3 5 6 5 o	(61)
52	o-Carboxybenzene-azo-phenyl-α-naphthylamine.	v 3 5 6 5 o	(61)
53	Benzene-azo-α-naphthylamine.	r 3 7 5 0 y	(56, 61)
54	p-Toluene-azo-α-naphthylamine	3 7 5 0	(53, 54, 56)
55	o-Carboxybenzene-azo-methylaniline	r 4 0 6 0 y	(11)
56	Benzene-azo-m-phenylenediamine; Chrysoidine [S. 33; R. 20]	o 4 0- 7 0 y	(9)
57	o-Carboxybenzene-azo-ethylaniline	r 4 2- 6 2 y	(11)
58	o-Carboxybenzene-azo-n-propylaniline	r 4 2 6 2 y	(11)
59	o-Carboxybenzene-azo-dimethylaniline; Methyl red [R. 211]	r 4 2- 6 3 y	(11, 14, 56, 60)
60	o-Carboxybenzene-azo-diethylaniline; Ethyl red	r 4 4- 6 2 y	(11, 60)
61	*o-Carboxybenzene-azo-di-n-propylaniline; Propyl red	r 4 6 6 6 y	(11)
62	o-Carboxybenzene-azo-m-phenylenediamine	o 4 6- 7 6 y	(9)
63	Benzene-azo-dimethyl-α-naphthylamine	4 8- 5 5	(53, 54, 56)
64	p-Benzenesulfonic acid-azo-dimethyl-α-naphthylamine	r 5 0- 5 7 o	(53, 54, 56, 61)
65	o-Carboxybenzene-azo-α-naphthylamine	p 5 6- 7 0 y	(11)
66	o-Carboxybenzene-azo-(di or mono?)-amyl aniline	o 5 6 7 6 y	(11)
67	o-Carboxybenzene-azo-dimethyl-α-naphthylamine	r 5 6 7 6 o	(11, 61)
68	4-Sulfo-α-naphthalene-azo-α-naphthol; Naphthylamine brown [S. 160; R. 175]	o 6 0- 8 4 p	(9)
69	Tropaeolin?.	y 7 0 9 0 r	(50)
70	6-Sulfo-α-naphthol-1-azo-m-hydroxybenzoic acid	o 7 0- 8 0 b / v 12 13 r	(67)
71	Curcumine?.	y 7 4 8 6 b	(31)
72	p-Benzenesulfonicacid-azo-α-naphthol; Tropaeolin OOO No. 1 [S. 144; R. 150]	y 7 6 8 9 p	(56)
73	p-Benzenesulfonicacid-azo-β-naphthol; Tropaeolin OOO No. 2 [S. 145; R. 151]	7 6 8 9(?)	(45)
74	m-Nitrobenzene-azo-salicylic acid; Alizarine yellow GG [S. 48; R. 36]	c(?)10 0- 12 0 y	(38, 39)
75	p-Nitrobenzene-azo-salicylic acid; Alizarine yellow R [S. 58; R. 40]	y 10.0- 12 4 y	(56)
76	α-Naphthylaminosulfonic acid-azo-β-naphthol; Red I [S. 161; R. 176]	10 5 12 1	(53, 54, 56)
77	α-Naphthalene-azo-β-naphthol-3, 6-disulfonic acid; Bordeaux B [S. 112; R. 88]	p 10 5 12 5 o	(9)
78	p-Benzenesulfonic acid-azo-resorcinol; Tropaeolin O [S. 143; R. 148]	y 11 1- 12 7 o	(56)
79	Benzene-azo-β-naphthol-6, 8-disulfonic acid; Orange GG [S. 38; R 27]	y 11 5- 14 0 p	(9)
80	Crocein?.	p 12 0- 14 0 v	(50)
81	Helianthin (Grübler)?	o 11 0-12 0 r	(9)
82	Helianthin I?.	o 11 0-13 0 r	(50)
83	Helianthin II?	y 13 0- 14 0 v	(50)
84	Curcumein?.	o 0 0 1 0 y / y 13 0 15 0 g	(50)

DIS-AZO COMPOUNDS

Index No.	Indicator	Color and useful range pH	Lit.
85	Ditolyl-disazo-bis-β-naphthylamine-6-sulfonic acid; Benzopurpurin B [S. 365; R. 450]	b 0 3- 1 0 v / v 1 0- 5 0 y / y 12 0-14 0 r	(50)
86	Ditolyl-disazo-bis-α-naphthylamine-4-sulfonic acid; Benzopurpurin 4B [S. 363; R. 448].	v 1 3- 4 0 r	(31)
87	Diphenyl-disazo-bis-α-naphthylamine-4-sulfonic acid; Congo red [S. 307; R. 370]	b 3 0- 5 0 r	(50)
88	Ditolyl-disazo-bis-α-naphthol-4-sulfonic acid; Azo blue [S. 377; R. 463]	v 10 5-11 5 p	(9)
89	Curcumin W [Probably Rowe, 364 (21)]	mid-point 7.3 / mid-point 7 6	(49) (18)

TRIPHENYLMETHANE DERIVATIVES

Index No.	Indicator	Color and useful range pH	Lit.
90	Methylated pararosaniline; Crystal violet [S. 516; R. 681]	g 0.0– 2.0 b	(9)
91	p, p'-Tetramethyldiamino-triphenylcarbinol; Malachite green [S. 495; R. 657]	y 0.0– 2.0 g / b 11.5–14.0 f	(50)
92	Hofmann's violet; Methylated rosanilines and pararosanilines [S. 514; R. 679]	g 0.0– 2.0 b	(9)
93	Tetraethyl-diamino-triphenyl-carbinol; Brilliant green [S. 499; R. 662]	y 0.0– 2.6 g	(9)
94	Heptamethylrosaniline; Iodine green [R. 686]	y 0.0– 2.6 b	(9)
95	Hexaethylpararosaniline; Ethyl violet [S. 518; R. 682]	y 0.0– 3.6 b	(9)
96	Ethyl-hexamethyl-pararosaniline; Ethyl green [R. 685]	y 0.3– 2.0 b	(31)
97	Methyl violet 6B; Benzylated tetra- and pentamethyl-pararosaniline [S. 517; R. 683]	y 0.15– 3.2 v	(56)
98	Gentian violet; mixture	0.4– 2.7	(53, 54, 56)
99	Aniline red; Rosaniline and pararosaniline [S. 512; R. 677]	pu 1.2– 3 0 f	(9)
100	Red violet 5RS; Di- and tri-sulfonate of ethylrosaniline [S. 525; R. 693]	p 3.6– 6.0 c	(9)
101	Resazurin [R. 727 note]	o 3.8– 6.5 v	(31)
102	China blue [S. 539; R. 707]; Mixture	b 4.7– 7.0 c	(9)
103	Rosolic acid [S. 555; R. 724]; Mixture	br 6.9– 8 0 r	(56)
104	Alkali blue 4B [S.,536; R. 704]; Mixture	v 9.4–14 0 p	(9)
105	XL Soluble blue [S. 538; R. 706]; Mixture	b 10 0–13 0 p	(9)
106	Poirrier's blue	b 11.0–13.0 r	(8)
107	Acid fuchsin; Di- and tri-sulfonic acids of rosaniline and pararosaniline [S. 524; R. 692]	r 12 0–14 0 f	(50)

PHTHALEINS AND RELATED COMPOUNDS

Index No.	Indicator	Color and useful range pH	Lit.
108	Diethyl-m-amino-phenolphthalein; Rhodamine B [S. 573; R. 749]	o 0.1– 1 2 p	(9)
109	Pyrogallol-phthalein; Gallein [S. 599; R. 781]	variable 0–14	(50)
110	Tetrabromofluorescein; Eosine Y S [S. 587; R. 768].	y 0 – 3.0 fl	(9)
111	Erythrosin (iodeosin); Di- or tetra iodated fluorescein [S. 591, 592?; R. 772, 773?]	o 0.0– 3.6 fl	(9)
112	Phloxin red B.H. (Grübler)?	p 1.4– 3.6 r	(9)
113	Dihydroxyfluoran; Uranin (fluorescein) [S. 585; R. 766]	y 3.6– 5.6 fl	(9)
114	Dichlorofluorescein	y 4.0– 6.6 fl	(9)
115	o-α-Naphthol phthalein	y 8.9– 9 5g(f)	(17)
116	p-α-Naphthol phthalein	y 7 0– 9 0 b	(56)
117	Tetrabromophenol phthalein	c 8 0– 9 0 v	(45)
118	o-Cresoltetrachlorophthalein	c 8.5– 9 0 pu	(1)
119	o-Cresolphthalein	c 8 2– 9 8 r	(11, 14)
120	Phenolphthalein [R. 764]	c 8.3–10.0 r	(38, 39, 56)
121	*1, 2, 3-Xylenolphthalein	c 8.9–10.2 b	(17)
122	Thymolphthalein	c 9 3–10.5b(f)	(56)
123	Dibromo-dinitrofluorescein; Eosin BN [S. 590; R. 771]	p 10.5–14.0 y	(9)
124	R = SCH₃	c 8.4–10.0 v	(25)
125	R = SC₄H₉	c 8 6– 9 8 v	(25)
126	R = SC₂H₅	c 9 0–10 0 v	(25)

SULFONPHTHALEINS

Index No.	Indicator	Color and useful range pH	Lit.
127	Catecholsulfonphthalein	p 0 2– 0 8 o / y 4 0– 7.0 g / v 8 5–10 2 b	(41)
128	m-Cresolsulfonphthalein; Metacresol purple	r 0 8– 2 4 y / y 7.6– 9 2 pu	(11, 14)
129	Thymolsulfonphthalein; Thymol blue	r 1 2– 2 8 y / y 8.0– 9 6 b	(11, 14)
130	Tetranitrophenolsulfonphthalein	2 8– 3 8?	(11)
131	Tetrabromophenolsulfonphthalein; Bromphenol blue	y 3.0– 4 6 b	(11, 14)
132	*Tetrachlorophenolsulfonphthalein	y 3 0– 4 6 b	(11)
133	*Dichloro-dibromo-phenol-sulfonphthalein; Brom-chlorphenol blue	y 3 2– 4 8 b	(14)
134	Tetrabromo-m-cresolsulfonphthalein; Bromcresol green	y 3 8– 5 4 b	(11, 14)
135	Dichlorophenolsulfonphthalein; Chlorphenol red	y 5 0– 6 6 r	(11, 14)
136	Dibromo-o-cresolsulfonphthalein; Bromcresol purple	y 5 2– 6 8 pu	(11, 14)
137	Dibromophenolsulfonphthalein; Bromphenol red.	y 5 4– 7 0 r	(11, 14)
138	*Diiodophenolsulfonphthalein	y 5.7– 7 3 pu	(9)
139	Dibromothymolsulfonphthalein; Bromthymol blue	y 6 0– 7 6 b	(11, 14)
140	*Brom Xylenol Blue, dibrominated No. 145	y 6 0– 7 6 b	(11, 14)
141	Phenol-nitrosulfonphthalein	y 6.6– 8.4 pu	(11)

SULFONPHTHALEINS.—(Continued)

Index No.	Indicator	Color and useful range pH	Lit.
142	Phenolsulfonphthalein; Phenol red..		
143	o-Cresolsulfonphthalein; Cresol red	y 6 8- 8 4 r	(11, 14)
144	Salicylsulfonphthalein	y 7 2 8 8 r	(11, 14)
145	*1.4-Dimethyl-5-hydroxybenzenesulfonphthalein; Xylenol blue	y 7 2 - 9 2 p	(9)
146	α-Naphtholsulfonphthalein	y 8 0- 9 6 b	(12)
147	Carvacrolsulfonphthalein .	y 7 5- 9 0 b	(11)
148	Orcinsulfonphthalein..	y 7 8- 9 6 b	(11)
149	Nitro-thymolsulfonphthalein..........	y 8 6-10 0 fl	(11)
		v 9 2-11 5 y	(11)

QUINOLINE COMPOUNDS

150	α-(p-Dimethylaminophenylethylene)-quinoline ethiodide; Quinaldine red. Eastman Kodak Co. No. 1361.....		
151	Quinoline blue (cyanin); 1, 1' Disoamyl-4, 4'-quinocyanine iodide [S. 611; R. 806]	1 0 2 0	(36)
		r 7 0 8 0 v	(52, 54, 56)

Index No. 152 INDOPHENOLS (15)

Color changes: from brownish or clear red in acid to deep blue in alkali. All indophenols are somewhat unstable

Indophenol

Substituents	pK
2, 6, 3' Tribromo-.	5 1
2, 6-Dibromo-3'-chloro-.	5 4
2, 6-Dibromo-3'-methyl-.	5.4
2, 6-Dichloro-3'-chloro-..	5.8
2, 6-Dichloro-3'-methyl-.	5.5
2, 6-Dibromo-3'-methoxy-..	5 6
2, 6-Dichloro-. .	5.7
2, 6-Dibromo- ..	5 7
2, 6-Dibromo-2'-methyl-	5.9
2, 6-Dibromo-2'-bromo-.	6 3
2-Chloro- .	7 0
2-Bromo-	7 1
3-Bromo-. . .	7 8
Indophenol	8.1
2-Methyl- ,	8.4
3-Methyl- .	8.6
2-Methoxy-	8.7
2-Isopropyl-5-methyl-	8 8
2-Methyl-5-isopropyl	8 9

Orthoindophenol

Substituents	pK
3' Bromo-	7 1
Orthoindophenol	8 4
2'-Methyl-	8 8

Indonaphthol-2'-sulfonic acid

Substituents	pK
2, 6 Dichloro-.	6.1
Indonaphthol-2'-sulfonic acid...... ..	8.7
2-Methyl-.	9.0

AZINES

Index No.	Indicator	Color and useful range pH	Lit.
153	Safranine (Which?) .	b —0 3 - 1 0 r	(50)
154	Amino-dimethylamino-phenyl-diphenazonium chloride; Methylene violet B.N. [S. 680; R. 842]	pu 0 0 - 1 2 v	(9)
155	Amino-phenylamino-p-tolyl-ditolazonium sulphate; Mauve [S 688; R. 846] . .	0 1 2 9	(56)
156	Magdala red; Mixture amino- and diamino-naphthyl-dinaphthazonium chlorides [S. 694; R. 857] . ..	p 3 0- 4 0 fl	(50)
157	Induline, spirit soluble [S. 697; R. 860]; Mixture . . .	b 5 6- 7 0 v	(9)
158	Amino-dimethylamino-toluphenazonium chloride; Neutral red [S. 670; R. 825] ...	r 6 8- 8.0 y	(56)
159	Dimethylamino-phenyl-naphtho-phenazonium chloride; Neutral blue [S. 676; R. 832]	9 3-10 2	(52, 54, 56)

OXAZINE COMPOUNDS

160	Dihydroxy-dinaphthazoxonium sulfonate; Alizarin green B [S. 657; R. 918]	v –0 3- 1.0 p / y 12 0-14.0 br	(60)
161	Diethylamino-benzylamino-naphtho-phenazoxonium chloride; Nile blue 2B [S. 654; R. 914]	b 7 2- 8.6 p	(9)
162	Diethylamino-aminonaphtho-phenazoxonium sulfate; Nile blue A [S. 653; R. 913]....... ..	b 10.2-13.0 p	(9)

ANTHRAQUINONE COMPOUNDS

Index No.	Indicator	Color and useful range pH	Lit.
163	1, 2-Dihydroxy-anthraquinone-β-quinoline; Alizarin blue ABI [S. 803; R. 1066]	p 0.0– 1 6 y y 6 0– 7 6 g	(9)
164	1, 2, 4-Trihydroxy-anthraquinone; Purpurin [S. 783; R. 1037]	y 0 0– 4 0 o o 4.0– 8 0 p	(9)
165	Alizarin sulfonic acid; Alizarin red S [S. 780; R. 1034]	y 3 7– 4 2 p	(67)
166	1, 2-Dihydroxy-anthraquinone; Alizarin [S. 778; R. 1027]	y 5 5– 6 8 r v 10 1–12 1 pu	(53, 54, 56)
167	Alizarin blue S	various 6–14	(45)

INDIGOS

| 168 | Indigo disulfonate; Indigo carmine [S. 877; R. 1180] | b 11 6–14 0 y | (9) |

MISCELLANEOUS AND NATURAL INDICATORS

169	Echtrot?	y 0 – 1 0 r	(50)
170	Logwood [S 938; R. 1246]	various 0–14	(45)
171	*Red cabbage extract	r 2 4– 4 5 g	(65)
172	1-Oxynaphtho-quinomethane; Nierenstein's indicator	c 2 7– 3 7 pu	(67)
173	Tröger and Hille's Indicator, C₁₄H₁₅N₄SO₃H	o 2 8– 3 9 y	(67)
174	Phenacetolin	y 3 0– 6 0 r r 10 0 13 0 c	(45)
175	Lacmosol	r 4 4– 5 5 b	(26)
176	Lacmoid [R. 908 note]	r 4 4– 6 2 b	(53, 54, 56)
177	Azolitmin (litmus) [R. 1242]	r 4 5– 8 3 b	(53, 54, 56)
178	Cochineal [S. 932; R. 1239]	y 4 8 6 2 v	(53, 54, 56)
179	Archil (orchil) [S. 934; R. 1242]	p 5 6 7 6 v	(9)
180	Brazilein [S. 935; R. 1243]	c 6 0– 8 0 p	(9)
181	Di-o-hydroxy-styryl ketone; Lygosine	y 7 3– 8 7 g	(67)
182	Mimosa flower extract	7 7 9 6	(67)
183	Turmeric (curcuma) [S. 927; R. 1238]	y 7 8– 9 2 br	(31)
184	Alkannin [R. 1240, note] cf. alizarin	8 3–10 0	(53, 54, 56)
185	α-Naphtholbenzein	v 8.5– 9 8 g	(53, 54, 56)

COMMON SYNONYMS OF INDICATORS

Among synonyms given in this table are several which apply to dyes which are not listed in preceding table or which have been applied to two or more of the indicators listed. Such cases are indicated by *.

Acid bordeaux, 77
Acid brown R,* 68
Acid fuchsin,* 107
Acid magenta H, 107
Acid roseine, 107
Alizarin, 166
Alizarin blue ABI, 163
Alizarin blue S, 167
Alizarin blue X, 163
Alizarin carmine, 165
Alizarin green B, 160
Alizarin red S, 165
Alizarin sulfonate or S, 165
Alizarin yellow GG, 74
Alizarin yellow R, 75
Alkali blue 4B, 104
Alkanet, 184
Alkanin, Alkannin, 184
Alphanaphtholbenzein, 185
Alphanaphtholphthalein,* 116
Amido-azo-benzol, 30
Amido-azo-toluol, 26
Amino-azo-bensene, 30
Amino-azo-toluene, 26
Amyl red, 66
Anchusin, 184
Aniline orange,* 31
Aniline red, 99
Aniline yellow,* 3, 25, 30
Archil, 179
Aurin, 103
Azo blue, 88

Azolitmin, 177
Azoresorcin, 101
Benzopurpurin B, 85
Benzopurpurin 4B, 86
Benzyl violet, 97
Beta naphthol orange, 73
Bitter almond oil green, 91
Blauholz, 170
Boettger's indicator, 184
Bordeaux B, 77
Brasilein, brasilin, brazilin, 180
Brazil wood, 180
Brilliant green, 93
Brilliant yellow,* 89
Brom-chlor-phenol blue, 133
Brom cresol green, 134
Brom cresol purple, 136
Brom phenol blue, 131
Brom phenol red, 137
Brom thymol blue, 139
Brom xylenol blue, 140
Butter yellow,* 26, 37
Cabbage red, 171
Campeachy wood, 170
Carmine, 178
Carminic acid, 178
Catechol sulphonphthalein, 127
China blue, 102
Chlor phenol red, 135
Chrome printing orange R, 75
Chrome printing yellow G, 74
Chrysoidine,* 56

Chrysoine, 78
Coccus, 178
Cochenille, cochineal, 178
Congo, 87
Congo red, 87
Corallin, 103
Cresol red, 143
Cresolphthalein,* 119
Cresolsulphonphthalein,* 143
Crismer's indicator, 101
Crocein,* 80
Crystal violet, 90
Curcuma, 183
Curcumein,* 84
Curcumin,* 183
Curcumin W, 89
Curcumin,* 183
Cyanin, 151
Dechan's indicator, 109
Degener's indicator, 174
Dianil red,* 87
Dichlorofluorescein, 114
Diethylamine orange, 45
Dihydroxyanthraquinone, 166
Dimethylamine orange, 44
Dimethyl orange, 44
Dimethyl yellow, 37
Dinitroaminophenol, 9
Dinitrohydroquinone, 5
Echtrot,* 169
Echtrot A, 76
Echtrot B, 77
Eosine, 110
Eosine BN, 123

Eosine YS, 110
Erythrosine,* 111
Ethyl green,* 96
Ethyl orange, 45
Ethyl red,* 60
Ethyl violet, 95
Fast red A, 76
Fast red B,* 77
Fluorescein, 113
Formanek's indicator, 160
Fuchsin, 154
Fuchsin,* 99
Fuchsin S, 107
Galeine, 109
Gallein, 109
Gentian violet, 98
Golden orange, 44
Haematein,* 170
Haematoxylin,* haematoxylon,* 170
Helianthine,* 44, 81, 82, 83
Hematein,* hematine,* 170
Hematoxylin, * 170
Henderson & Forbes' indicator, 5
Herzberg's indicator, 87
Hofmann's violet, 92
Holt & Reid's indicators, 124–126
Indigo carmine, 168
Indigo disulphonate, 168
Indophenols, 152
Induline spirit-soluble, 157
Iodeosine,* 111
Iodine green, 94
Kosmos red, 87

¹ Haematoxylin is the leuco-compound of Haematein or Hematine as obtained from logwood although the name is sometimes given to the oxidised form. Haematein or Hematine should not be confused with Hematin of the blood pigment.

TABLE 3

A CLARK AND LUBS' SELECTION OF INDICATORS SUPPLEMENTED BY COHEN (11, 14)

A = Cubic centimeters of $0.01N$ NaOH required per 0.1 g acid indicator to form sodium salt. Dilute to 250 cc for 0.04 % reagent. Use alcoholic solutions of methyl red (59) and cresolphthalein (119).

B = Approximate pH value of solution required for full "acid color" appertaining to range indicated

C = Approximate pH value of solution required for full "alkaline color" appertaining to range indicated

Index No.	A	B	C	Useful range pH	pK†
129	see below	conc. HCl	6	1 2 2 8	1.5
131	15 0	0	7	3 0-4 6	4.0
134	14 5	1	8	4 0-5 6	4.7*
59		?	9	4 4-6 0	[5.0]
135	23 5	3	10	5 0 6.6	6.2*
136	18 5	3	10	5 2-6 8	6.3
139	16 0	4	10	6 0 7 6	7.1
142	28 5	5	11	6 8-8 4	7.8
143	26 3	5	11	7 2-8 8	8.2
128	26 5	5	11	7 6 9 2	8.4*
129	21 5	6	12	8 0 9 6	8.9
119	6	6	12	8 2 9 8	[9.4]

* No salt and protein errors determined
† pK values are weighted means of values found in (2, 7, 11, 14, 19, 20, 24, 24).

Representative Corrections of Colorimetric Readings with Indicators of Table 3A to Bring Readings to Electrometric pH

	Peptone-beef infusion	10% gelatine sol.	2% egg-white	Urine
131 Brom phenol blue	0 05			
59 Methyl red	−0 10		0 24	0.05
136 Brom cresol purple	0 01	0 04		0.01
139 Brom thymol blue	0 10	0 04		0.02
142 Phenol red	0 04	0 20		0.00
143 Cresol red	0 03	0 20		
129 Thymol blue	0 04	0 20		
119 Cresolphthalein	−0 03	0 20		

Corrections at different salt content [after Kolthoff (29)]

Thymol blue (acid range)	0 1N KCl	. . . . . .	−0.06
	1 0N KCl	. . . . . .	+0.05
Brom phenol blue	0.1N KCl	. . . . . .	−0.05
	1 0N KCl	. . . . . .	−0.35
Methyl red	0 5N NaCl	. . . .	+0.10
Brom cresol purple	0 5N NaCl	. . .	−0.25
Phenol red	0 5N NaCl	. .	−0.15
Thymol blue	0 5N NaCl	. . . . . .	−0.17

With color match between a solution at 70° and a standard buffer at 20° the solution at 70° will have the pH of the standard corrected by the following values according to Kolthoff (26).

Thymol blue (acid range)	. .	0.0
Brom phenol blue	. .	0.0
Methyl red	. .	−0.2
Brom cresol purple	. . . .	0.0 to +0.2
Phenol red	. .	−0.3
Thymol blue (alk.)	.	−0 4

Corrections in sea water of salinity S [parts per 1000] after Ramage and Miller 1925 (unpublished).

S	5	10	15	20	25	30	35
Cresol red	− 11	− 17	− 21	− 24	− 25	− 26	− 27

B. Sørensen's Selection of Indicators (56)

Index No.	Composition of test solution	Useful range pH	Sensitivity to neutral salts	Usefulness in presence of			Stability on standing
				True proteins	High conc. of products of proteolysis	Chloroform and toluene	
97	0.01%–0.05% aqueous	0 1–3 2	high	fair	good	with chloroform not, with toluene useful	acid solutions fade
155	0.01%–0.05% aqueous	0 1–2 9	high	fair	good	as above	as above
22	0.01 g in 1 cc N HCl + 50 cc alcohol + 49 cc water	1 2 2 1	low	not	fair	not	moderate
25	0.01% aqueous	1 4–2 6	low	not	fair	good	good
23	0.01% aqueous	1 2–2 3	low	not	fair	good	good
34	0.02 g in 1 cc N/10 HCl + 50 cc alcohol + 49 cc water	2 3 3 3	low	not	good	not	moderate
32	0.01% aqueous	1 9 3 3	low	not	fair	good	good
35	0.01% aqueous	2 6–4 0	low	not	fair	good	good
37	0.01 g 0.1 cc N/10 HCl + 80 cc alcohol + 20 cc water	2 9–4 0	low	not	good	not	moderate
44	0.01% aqueous	3 1–4 4*	low	not	fair	good	good
53	0.01 g in 0.4 cc N/10 HCl + 30 cc alcohol + 70 cc water	3 7–5 0	low	not	good	not	moderate
50	0.01 g in 60 cc alcohol + 40 cc water	3 5 5 7	low	not	good	good	good
59	0.02 g in 60 cc alcohol + 40 cc water	4 2 6 3*	low	S.C.	good	good	moderate
12	0.04 g in 6 cc alcohol + 94 cc water	5 0–7 0*	moderate	good	good	good	good
158	0.01 g in 50 cc alcohol + 50 cc water	6 8–8 0*	low	S.C.	good	S.C.	good
103	0.04 g in 40 cc alcohol + 60 cc water	6 9 8 0	low	fair	good	fair	good
72	0.01% aqueous	7 6–8 9	low	good	good	good	good
116	0.1 g in 150 cc alcohol + 100 cc water	7 3 8 7	moderate	S.C.	good	good	fair
120	0.05 g in 50 cc alcohol + 50 cc water	8 3–10 0*	moderate	S.C.	good	good	good—fades in strong alkali
122	0.04 g in 50 cc alcohol + 50 cc water	9 3–10 5	moderate	S.C.	good	good	fades in moderate alkali
75	0.01% aqueous	10 1 12.1			good		good
78	0.01% aqueous	11 1–12 7			fair		good

S.C. = useful in special cases

* Apparent pK values referred to standard buffers: Methyl orange (44) 3 7 (34 cf 40). Methyl red (59) see Table 3A (59, 40). Paranitrophenol (12) see Table 3C. Neutral red (158) 6 85 (34). Phenolphthalein see Table 3C.

Representative average corrections of colorimetric readings with indicators of Table 3B to bring readings to electrometric pH (see also Table 2).

Index No. of indicator	Corrections (after Sørensen (53))		Corrections in solutions containing salts
	In 2% peptone 0.01–0.3N salt	In 2% egg-white 0.07–0.3N salt	
97	−0 02	−0 19	
155	−0 04	−0 19	
22	−0 06	> −0 90	
25	−0 27	> −1.40	
23	−0 30	> −1 40	
34	+0 01	> −0 80	
32	−0 22	> −0 80	
35	−0 41		
37	−0 08	−0 53	
44	−0 18		0.1N KCl, −0.08; 1.0N KCl, +0.23 Kolthoff
53	−0 02		
50	−0 03	+0 15	0.5N NaCl, + 0.10 Sørensen
12	−0 06	−0 04	0.5N NaCl, − 0.15 Sørensen (−0.05 Kolthoff)
158	+0 13	+0 68	0.5N NaCl, + 0.09 Sørensen

Index No. of indicator	Corrections (after Sørensen (53))		Corrections in solutions containing salts
	In 2% peptone 0.01–0.3N salt	In 2% egg-white 0.07–0.3N salt	
103	+0.08	+0.44	0.5N NaCl, − 0.06 Sørensen
72	−0.12	+0.10	0.5N NaCl, − 0.12 Sørensen
120	−0.01	+0 18	0.5N NaCl, − 0.12 Sørensen (−0.17 Kolthoff)
122	+0.01	+0 40	
75		+0.29	
78		−0 30	0.1N KCl, + 0.38; 1.0N KCl, + 0.62 Kolthoff

C. Michaelis' Selection of One-color Indicators

Index No.	Useful range pH	Conc. % in H₂O	pK (Michaelis and coworkers (38, 39))			pK (Kolthoff (31) at 15° and 0.05M salt)
			In low salt content	In 0 15M salt	In 0.5M salt	
1	0.03–1.3		[0.26]			
2	2 0–4.0	sat.	3 71 +0.006 (15 − t°)	3.59	3.41	3.58

C. Michaelis' Selection of One-color Indicators.—(*Continued*)

Index No.	Useful range pH	Conc. % in H_2O	pK (Michaelis and coworkers (38, 39))			pK(Kolthoff (31) at 15° and 0.05M salt)
			In low salt content	In 0 15M salt	In 0 5M salt	
4	2.6–4.4	0.05	4 08 + 0 006 (15 − *t*°)	3.98	3 88	3 95
7			4 87	4 76	4 71	
8	4.0–5.8	0 025	5 16 + 0 005 (15 − *t*°)	5.08	5 01	5 15
10			5 35	5.30	5 25	
12	5 6–7 6	0 10	7 22 + 0 011 (15 − *t*°)	7 22	7.17	7 03
15	6 8–8 .6	0.30	8 35 + 0 008 (15 − *t*°)	8 24	8 19	8 30
120	8 0–10.0	0.04	[9 76] + 0.011 (18 − *t*°)	9 6	9 5	
74	10 0–12 0		[11 2] + 0 013 (20 − *t*°)			

TABLE 4

RELATION BETWEEN PERCENTAGE, a, OF AVAILABLE COLOR AND pH (AFTER MICHAELIS AND GYEMANT (38))

Phenolphthalein.. 18° a 1 0 1 4 3 0 4 7 6 9 9 0
pH 8 45 8 5 8 6 8 7 8 8 8 9

Phenolphthalein 18° a 12 0 16 0 21 0 27 0 34 0 40 0
pH 9 0 9 1 9 2 9 3 9 4 9 5

Phenolphthalein 18° a 45 0 50 0 55 0 60 0 65 0
pH 9 6 9 7 9 8 9 9 10 0

Phenolphthalein 18° a 70 0 75 0 80 0 84 5 87 3
pH 10 1 10.2 10 3 10 4 10 5

Alizarine yellow GG. 20° a 13 16 22 29 36 46
pH 10 0 10 2 10.4 10.6 10.8 11 0

Alizarine yellow GG.. 20° a 56 66 75 83 88
pH 11 2 11 4 11 6 11.8 12.0

LITERATURE

(For a key to the periodicals see end of volume)

(1) Arnold, *1*, **46**: 489, 24. (2) Barnett and Barnett, **127**, **18**: 127; 21. (3) Bishop, Kittredge and Hildebrand, *1*, **46**: 135, 22. (4) Bjerrum, *9*, **17**: 58, 380, 11. (5) Bjerrum, *196*, **21**: 1, 14. (6) Bogert and Scatchard, *1*, **33**: 1606, 16. (7) Brode, *1*, **46**: 581; 24. (8) Clark, *The Determination of Hydrogen Ions*, 2nd. ed , 1922. (9) Clark, Cohen and Elvove, *O*; *cf*. (8). (10) Clark and Lubs, *141*, **26**: 479; 16. (11) Clark and Lubs, *149*, **2**: 1, 109, 191. 17 (12) Cohen, A., **230**, **16**: 31; 22. (13) Cohen, A , **230**, **17**: 335; 23. (14) Cohen, B , **231**, **18**: 199; 23. (15) Cohen, B , Gibbs and Clark, **231**, **29**: 381, 804; 24. (16) Cohn, *Indicators and Test Papers*, 1914. (17) Csanyi, *9*, **27**: 64; 21. (18) Fels, *9*, **10**: 208, 04. (19) Gillespie, **233**, **9**: 115; 20

(20) Gillespie, *1*, **42**: 742; 20. (21) Glaser, *Indikatoren der Acidimetrie und Alkalimetrie*, 1901. (22) Hatfield, *1*, **46**: 940; 23. (23) Henderson and Forbes, *1*, **32**: 687; 10. (24) Holmes and Snyder, *1*, **47**: 221, 226; 25. (25) Holt and Reid, *1*, **46**: 2333, 24. (26) Hottinger, **205**, **65**: 177; 14. (27) Jarisch, **205**, **184**: 177; 22. (28) Kolthoff, *70*, **40**: 775; 21. (29) Kolthoff, *70*, **41**: 54, 22.

(30) Kolthoff, **233**, **69**: 104; 22. **60**: 949; 23. (31) Kolthoff, *Der Gebrauch von Farbenindicatoren*, 1923. (32) Kolthoff, *70*, **42**: 251; 23. (33) Kolthoff, *70*, **43**: 964; 23. (34) Kolthoff, *70*, **43**: 144; 24. (35) McClendon, **247**, **21**: 348, 24. (36) McClendon, *141*, **59**: 437; 24. (37) Michaelis, *Die Wasserstoffionenkonzentration*, 1914. (38) Michaelis and Gyemant, **205**, **109**: 165; 20 (39) Michaelis and Krüger, **205**, **119**: 307; 21.

(40) Michaelis and Mizutani, **205**, **147**: 7; 24 (41) Moir, **234**, **3**: no. 2; 20. (42) Moir, **234**, **6**: no 2; 23. (43) Noyes, *1*, **32**: 815, 10. (44) Palitasch, *Bull Inst Oceanographie*, No **409**: 22. (45) Prideaux, *The Theory and Use of Indicators*, 1917. (46) Prideaux and Nunn, *4*, **126**: 2110; 24. (47) Ramage and Miller, *O*. (48) Rowe, *Color Index*, 1924. (49) Salessky, *9*, **10**: 204; 04.

(50) Salm, *7*, **57**: 471, 06. (51) Scudder, *Electrical Conductivity and Ionisation Constants of Organic Compounds*, 1914. (52) Schultz, *Farbstofftabellen*, 1923. (53) Sørensen, **235**, **8**: 1; 09. (54) Sørensen, **205**, **21**: 131, 201; 09. (55) Sørensen, **205**, **22**: 352; 09. (56) Sørensen, **236**, **12**: 393; 12. (57) Sørensen and Linderstrøm-Lang, **235**, **18**: no. 6; 24. (58) Thiel, *196*, **16**: 307, 11. (59) Thiel and Dassler, **25**, **86**: 1667; 23.

(60) Thiel, Dassler and Wülfken, **257**, **18**: 1; 24. (61) Thiel and Wülfken **93**, **136**: 393, 24. (62) Thiel, Wülfken and Dassler, **93**, **136**: 406; 24. (63) Van Alstine, **252**, **10**: 467; 20. (64) Vincent, *La concentration en ions hydrogène et sa mesure par la méthode électrometrique*, 1924. (65) Walbum, **205**, **46**: 291; 13. (66) Walbum, **205**, **107**: 209; 20 (67) Walpole, **230**, **8**: 628; 14. (68) Walpole, *4*, **105**: 2501, 2521; 14.

HIGH VACUUM TECHNIQUE

SAUL DUSHMAN

SELECTED FORMULAE

1. *Amount of Gas Striking* 1 Cm^2 *per Sec*—

$$m = \frac{1}{4}\rho\Omega = p\sqrt{\frac{M}{2\pi RT}}$$

where ρ = density and Ω = average velocity

= $43.74 \times 10^{-6} \times p\sqrt{M/T}$ g cm^{-2} sec^{-1} (*p* in baryes)

= $58.32 \times 10^{-3} \times p\sqrt{M/T}$ g cm^{-2} sec^{-1} (*p* in mm of Hg)

n = number of molecules

= $6.062 \times 10^{23} \frac{m}{M} = 2.653 \times 10^{19} \frac{p}{\sqrt{MT}} cm^{-2} sec^{-1}$ (*p* in baryes)

= $3.535 \times 10^{22} p/\sqrt{MT}$ cm^{-2} sec^{-1} (*p* in mm of Hg)

2. *Laws of Molecular Flow (Flow of Gases at Very Low Pressures).—Q* = amount of gas flowing through any tube or opening in cm^3 per sec

$$= \frac{p_2 - p_1}{W\sqrt{\rho_1}}$$

where $p_2 - p_1$ = difference of pressure

ρ_1 = density at 1 barye pressure

$$= \frac{M}{83.15 \times 10^6 T}$$

W = "resistance" of tube or opening

For a circular opening (diam., *d* cm) in a thin plate

$$W = \frac{3.184}{d^2}$$

For a tube of diameter *d* and length *l*

$$W = \frac{2.394 l}{d^3} + \frac{3.184}{d^2}$$

3. *Speed of Exhaust* (*S*) *of Given Volume* (*v*).—

$$S = \frac{v}{t} \log_e \frac{p_2}{p_1}$$

For p_2/p_1 = 10, *t* in sec and *v* in cm^3

$$S = \frac{2.303 v}{t} cm^3 sec^{-1}$$

For pump exhausting through resistance

$$\frac{1}{S_o} = \frac{1}{S_p} + \frac{1}{F}$$

where S_o = observed speed of exhaust,

S_p = speed of pump through negligible resistance, and

F = rate of flow through resistance (cm^3/sec)

$$S = \frac{Q}{p_2 - p_1} = \frac{1}{W\sqrt{\rho_1}}$$

TABLE OF MOLECULAR DATA

	H$_2$	He	N$_2$	O$_2$	A	Hg	CO	CO$_2$	H$_2$O
Mean Free path (cm) at 25°C and 1 barye.	19 2	29 6	10.0	10.7	10.6	[3.24]*	9.92	6.68	[6.03]*
(1/d^2) × 10^{-16} (Number of molecules per cm^2)	1 74	2 74	1 01	1.11	1.19	1.11	0.98	0.92	1.19
Micrograms (10^{-6} g) of gas striking 1 cm^2 per sec at 25°C and 1 barye.	3 597	5.062	13 42	14.33	16.01	35.89	13.42	16.81	10.76
Number of molecules striking 1 cm^2 per sec at 25°C and 1 barve. Unit = 10^{15}	1082	769 3	283 7	271 7	243 3	10.85	283 7	231.7	362.0

* Values in square brackets refer to 0°C   Note: 1 barye = 0 75 × 10⁻³ mm mercury. Values of mean free path calculated from viscosity coefficients

RATE OF FLOW OF AIR AND HYDROGEN AT LOW PRESSURES AND 20°C

l	d	W	F (air)	F (H$_2$)
1 cm	1 cm	5 58	5 204	197 10
10	1	27 12	1 070	40 53
1	0 1	2 712 4	10 70	40 53
10	0 1	24 258	1 196	3 60

(Note.—These relations are valid only for pressures so low that the mean free path is equal to or greater than d.)

DATA ON VARIOUS TYPES OF PUMPS

	S_p cm^3 sec^{-1}	Fore pump pressure	Min. pressure attainable
Gaede rotary mercury	100 (max.)	ca. 1 cm	10^{-4} mm
Gaede molecular	1 400	0 01 mm	<10^{-6} mm
Gaede diffusion	80	0 01 mm	<10^{-6} mm
Langmuir condensation (metal)	4 000	0 01 mm	<10^{-6} mm
Gaede two stage metal	60 000	20 mm	<10 ^{6} mm

Evolution of Gas from Glass.—For rate at which gas is evolved at different temperatures, v. R. G. Sherwood (1, 40: 1645; 18) and J. E. Shrader (2, 13: 434; 19).

Chemical Clean-up Reagents for Producing Low Pressures.—1. Charcoal in liquid air. 2. Ca or Mg volatilized in sealed-off device, cleans up all gases except those of group 0. 3. P$_2$O$_5$, efficient for water vapor. 4. Palladium black at low temperatures, very good for hydrogen.

SOME VAPOR PRESSURES AT LOW TEMPERATURES

Substance	t°C	p, mm	p, baryes
Hg	− 78	3 × 10^{-9}	4 × 10^{-6}
H$_2$O	−111	0 75 × 10^{-6}	1 × 10^{-3}
CO$_2$	−182	0 75 × 10^{-5}	1 × 10^{-2}
CO$_2$	−193	0 75 × 10^{-6}	1 × 10^{-3}
CO	−190	863	
CH$_4$	−185 8	79 8	
C$_2$H$_4$	−188	0 076	
C$_2$H$_6$	−180	0.076	
Vaseline (Stopcock grease)	−190 (fresh liquid air)		<10^{-6}

PSYCHOLOGICAL DATA PERTAINING TO ERRORS OF OBSERVATION

R. S. WOODWORTH

(Additional data pertaining to sight and hearing are given in other sections of International Critical Tables treating of the mechanical equivalent of light, colorimetry, and the physical aspects of audition. Consult index. Editor.)

SIGHT

Much of the available data pertaining to the sensitivity of the eye have been obtained under such conditions that the exact value of the stimulus cannot satisfactorily be determined. Some are expressed in terms of the illumination, others in terms of the brightness, of a screen; the latter procedure is to be preferred. If the illuminated screen were a perfect diffuser of the light, and also a perfect reflector, if illuminated from the front, or a perfect transmitter, if illuminated from the rear, then its brightness (B) expressed in millilamberts would be numerically equal to 0.1 of its illumination (I) expressed in meter-candles. In the following data, this relation has been used to reduce to the basis of B, data which have been given in terms of I. Although in many cases the screens surely did not possess the properties thus assumed, it seems probable that the error so introduced is of less importance than those arising from other sources. Data for reaction times will be found near the end of this report.

Spectral range (41) for daylight vision is λ = 397mμ to 760 mμ; for twilight vision (illumination too low for color perception), λ = 440 mμ to 670 mμ.

Threshold value = minimum stimulus which can be visually perceived as light; the perception of form is not involved. For white light and a thoroughly light-adapted eye, luminous area subtending an angle of 10°, it is that corresponding to a brightness of 0.1 millilambert (37). For white light and a dark-adapted eye, it varies with the area of the luminous area and with the duration of stimulus as shown in Table 1.

TABLE 1.—THRESHOLD OF VISION FOR DARK-ADAPTED EYE (45)

D = distance; θ = visual angle subtended by shortest dimension of area; B = brightness required for perception; P = power entering eye; t = duration of exposure. Diameter of pupil = 8.3 mm.

Unit of: Area = 1 cm^2; D = 1 cm; B = 1 microlambert; P = 1 milliwatt = 10^{-10} erg sec^{-1}; t = 1 sec.

Form	Area	D	θ	B	P	t	B‡
Star*	0 00785	300	1 2'	7.20	17.1	0.002	0.362
Star*	0 00785	150	2.30	2.60	24 8	0 006	0 098
Star*	0 00785	35	9 8	0 24	42.1	0.011	0.0446
Square	0 04	35	19.6	0 028 3	25 3	0 020	0.0239
Square	0 25	35	50	0.006 62	37	0.034	0.0123
Square	1.00	35	1° 30'	0 002 41	54	0 160	0.0071
Square	4 00	35	3 16	0 001 02	91	0 250	0 0051
Square	9 00	35	4 54	0 000 45	91	0.500	0.003 54
Square	36.0	35	9 44	0 000 258	208	1.000	0.002 62
Square	144 0	35	18 56	0 000 175	564	2.000†	0.000 77

* Circle, Diameter = 1 mm
† If t = ∞, B = 0 000 45, t = 4, B = 0 000 63
‡ For square, area = 9 cm^2, D = 35 cm, θ = 4 9°.

TABLE 2.—CHANGE IN THRESHOLD DURING ADAPTATION

Threshold = brightness (B) of a surface which can just be seen   Sensitivity $(S) = 1/B$. In light adaptation, I = illumination to which dark adapted eye was subjected for the time t; S was measured 10 sec after this exposure. Unit of t = 1 min; B = 1 microlambert; S = 0 1 millilambert[-1], I = meter-candle

*Dark adaptation [18]				†Light adaptation [34, 39]			
t	B	S	$\frac{I}{t}$	5 S	25 S	60 S	Day ‡ S
0	100	1		23 000	9950	5800	435
0 5	50 0	20		17 500	7440	3700	230
4	1 33	75	1	10 400	5200	3250	200
9	0 054	1850	2	8130	3360	2600	115
14	0 0096	10 400	3	5200	2740	2038	87
19	0 0038	26 000	6	3470	2040	1600	48
23	0 001 43	69 500	10	3000	1450	1130	40
26	0.001 56	94 700	15		1000	312	
31	0 000 57	174 000	60		95	36	
39	0 000 51	195 000	80		54	28	
51	0 000 48	208 000	110		54	24	
61	0.000 46	215 000					

* Following nearly complete light adaptation Luminous surface was 10 cm n diameter and 57 cm from eye (θ = 10°)
† Following nearly complete dark adaptation Luminous surface was 1 m square and 1 m from eye (θ = 45°); initial S, just before exposure to I, was 10 000 millilambert[-1].
‡ Moderate diffused day-light.

The rates of adaptation to darkness and to light are indicated in Table 2 in which are given the threshold values at various intervals (1) after removal from daylight, and (2) immediately (10 seconds) after removal from a specified exposure, the eye before exposure having been kept in darkness for 45 min. The visibility of monochromatic light varies with the wave-length, and the *relative visibility* of lights of different wave-lengths depends upon their intensities. (Figs. 1, 2.) For a large surface with a brightness of

Fig. 1.—Relative visibility (V) [28, 40].
B = brightness, unit = 1 millilambert;   abscissae = wavelengths.

Fig. 2.—Position (λ_{max}) of maximum visibility [28, 40].
Unit of brightness = 1 millilambert.

5 to 80 millilamberts, the maximum visibility for the average observer, is near [9] λ = 557.6 mμ, but even normal subjects exhibit individual differences; out of 125 subjects, the percentage finding the maximum at each of the several wave-lengths was as follows [9]:

λ	%	λ	%	λ	%	λ	%	λ	%	λ	%
549	1	553	2	557	12	561	2	565	2	569	0
550	2	554	7	558	13	562	3	566	2	570	2
551	5	555	9	559	12	563	2	567	0		
552	3	556	8	560	7	564	1	568	2		

All of the preceding refer to direct vision. The sensitivity of other portions of the retina is greater.

Complementary colors are those pairs of colors which, when superposed upon the retina in suitable proportions, produce the sensation of white. Grunberg states that if their wave-lengths are $\lambda m\mu$, $\lambda' m\mu$, then $(\lambda - 559)(498 - \lambda') = 424$, $\lambda > 559$, $\lambda' < 498$ [47]; there are no complementaries to the colors in the range 498mμ to 559mμ.

Stable, or invariable, colors are those which do not change in hue, except to become gray, as they are moved from the fovea to the periphery of the retina. They are: yellow of λ = 570mμ; bluish green of λ = 490mμ: blue of λ = 460mμ; and a non-spectral bluish red [21].

Discrimination of Brightnesses.—For large adjacent fields, differences of 1 % or even of 0.8 % in the brightness can be detected [31] if the brightness is of the order of 100 millilamberts. Under such

conditions the color of the light has no effect upon the discrimination. At lower brightnesses, the sensitiveness to change in brightness depends upon both the color and the brightness (Fig. 4).

Resolving power of the eye is the smallest angular separation at which two points, under the best illumination, can be seen as distinct. For different observers, it varies from 50″ to 93″ [20]; the generally accepted normal value is 1′. It varies with the color of the light. In day-light and on a bright background, a dark line a few minutes long can be seen if it is 1.2″ wide; but, on a dark background, a bright line is not visible unless it is at least 3.5″ wide [48].

Aligning power, the ability to detect a lack of alignment of two similar, adjacent lines of the same width, as in setting a vernier, exceeds the resolving power. The average error [48] of skilled observers under best conditions corresponds to a visual error of not over 3″; in coincidence range-finders, the images can be aligned with an error not greater than 12″ and sometimes as small as 2″.

Fig. 3.—Discrimination of Hue [49].
$\Delta\lambda$ = Change in wave-length (λ) corresponding to the least noticeable difference in color

Fig. 4.—Discrimination of brightnesses [39, 40].
ΔB = least noticeable increase in the brightness (B). Unit of B is 1 millilambert; of wave-length (Δ) is 1mμ.

Acuity, or discrimination of form, is closely related to the resolving power, but differs from that in dealing, in general, with extended, interpenetrating, bright and dark areas, and frequently with low brightnesses. The *absolute acuity* (A) is the reciprocal of the smallest visual angle for which neighboring contrasted portions of the field can be seen as separated. Its variation with the brightness (B) of the brighter portions of the field is given by the equation [25] $A = c + k \log B$; the values of the constants c and k are determined by the units, the character of the field, and the eye; some values are given in Table 3. The unit commonly employed for A is 1 reciprocal minute.

TABLE 3.—ABSOLUTE ACUITY (A) AND BRIGHTNESS (B)

$A = c + k \log_{10} B$ (cf. Fig. 5)
Unit of: A = 1 minute[-1]; B = 1 millilambert

Limits of B	c	k	Field	Lit.
0 01 to 43.5	1 05	0 415	Snellen and similar charts	[27]
40 to 1000	1 69	0.000	Snellen and similar charts	[27]
0 1 to 18	1 44	0 573	Snellen and similar charts	[12]
0 02 to 21	1 23	0 282	Crossed gratings	[8]
0 06 to 26	1 33	0 262	Crossed gratings	[7]

When the test field is a Snellen test chart, the acuity is commonly expressed as the ratio of the maximum distance (d_m), at which the characters can be distinguished, to the standard distance (d_s). This ratio (d_m/d_s) may be called the Snellen acuity; it is numerically equal to the reciprocal of the visual angle (in minutes) subtended by the sides of the elementary squares of the chart. As expressed in these units, the acuity of the average good eye exceeds 1.00; for the E-hooks, the mean of 100 subjects was 1.74, ranging from 1.00 to 2.45 [54].

Fig. 5.—Acuity in white and in chromatic illumination [44]
Unit of acuity = 1 Snellen unit; of illumination = 1 meter-candle

Fig. 6.—Relative acuity in indirect vision [20].
Abscissa indicates angular position of image upon the retina.

The effect of dark adaptation upon acuity may be obtained by determining, at various intervals (t) after the light adapted eye had been placed in darkness, the minimum illumination (I) in which it can distinguish Snellen test characters placed at a known distance. For a distance corresponding to a Snellen acuity of $\frac{2}{10}$ (= 0.2), the median[1] values of I for 6 observers having in daylight a Snellen acuity of $\frac{6}{4}$ (= 1.5) were found to be as follows [13]:

t	0	5	10	15	25	35	45 minutes
I	1.09	0 70	0 56	0 40	0 34	0 42	0 42 meter-candles

The acuity depends also upon the color of the light, and upon the position of the image upon the retina. See Figs. 5, 6.

Detection of Differences in Length.—About 1% of the length is the least noticeable difference for simultaneously presented parallel lines which are relatively displaced (result of several old investigations). More recent work shows that a variable line, 1 to 5 cm long, can, by eye, be set to equality with a standard line with a probable error, for a single setting, of only 0.4%; for shorter lines the error is greater, attaining 0.5% for lines 1 mm long [36]. When the time allowed for observation and judgment is short, the differences which can be detected with certainty are considerably greater. If the sign of the difference is to be judged correctly in 75% of the trials, then, for a 10 cm line, the difference must be 3.5 mm if the time is 4 seconds, and over 5 mm if the time is only 0.5 second [18].

Decimal Subdivision of a Small Distance.—When a fine line is set on a millimeter scale to successive positions in random order, and the subject is required to estimate its position to the nearest 0.1 mm, the average actual setting, for each tenth as estimated by 10 subjects (total of 6000 readings), for horizontal and for vertical scales was as follows [3, 52]:

Estimate	0 1	0 2	0 3	0 4	0 5	0 6	0 7	0 8	0 9	1 0
Horizontal	0 126	0 231	0 336	0 423	0 509	0 591	0 676	0 773	0 886	1 001
Vertical	0 106	0 202	0 308	0 393	0 486	0 576	0 652	0 757	0 875	0 992

The lines of the scale were presumably of the same width as the "fine line" of variable position. Settings were distributed over a length of 30 mm, the illumination was good, and the distance was that for best reading.

SENSES OTHER THAN SIGHT

Range of audible tones is from 18 to 18 600 double vibrations per second [44, 53]; at high intensities the lower limit may be reduced

[1] For each value of t, the 6 observed values of I are arranged in order of magnitude, the mean of the third and the fourth of the values is by definition the *median* of the set.

to 12. At the upper limit, individuals varied from 15 000 to 22 000 d.v. per sec. As the age increases, the upper limit becomes lower (Fig. 7).

Fig. 7.—Dependence of highest audible tone upon age of subject [4].
 * It is probable that these frequencies should be divided by two.

Fig. 8.—Aural sensitivity.

J = minimum audible power, unit = 1 erg cm^{-2} sec^{-1}. Data in terms of effective, or r.m.s., pressure (P) in dynes cm^{-2} have been reduced to erg cm^{-2} sec^{-1} (E) by means of the relation $P = \sqrt{dvE} = 6.5\sqrt{E}$; d = density of air, v = velocity of sound in air, both in cgs units.

REACTION TIMES

The *simple reaction time*, or, briefly, the *reaction time*, is the interval which elapses between the application of a definite,

Fig. 10.—Discrimination of differences in pressure [50].
P = initial pressure applied to ball of the little finger by means of a force F exerted on a circular disk 4 mm in diameter. ΔP = smallest noticeable change in P.

Fig. 9.—Discrimination of pitch.

N = number of double vibrations per sec; ΔN = smallest noticeable change in N. o = Knudsen [26], x = Stücker [51], Δ = Vance & Schaefer [53].

expected stimulus and the performance of a prescribed movement (usually a finger movement) indicating that it has been perceived.

Light.—For foveal stimulation of medium intensity, reaction time is 0.190 (±0.008) sec; individuals range from 0.150 to 0.225 sec. It is the same for withdrawal as for initiation of stimulus [22]. For faint stimulation, near threshold, interval is increased by 0.04 to 0.05 sec [16]; reaction to withdrawal is 0.005 to 0.025 sec quicker than to initiation of stimulus [22]. For photo-

Fig. 11.—Discrimination of differences in lifted weights.

ΔW = smallest noticeable change in the weight W.

* Weights had horizontal handles, were lifted successively with same hand

** Cylindrical boxes lifted successively with same hand, ΔW is change for which 50 % of the estimates were of proper sign.

§ Cylindrical boxes lifted successively with same hand; ΔW is change for which 75 % of the estimates were of proper sign.

† Weights lifted by downward pressure of finger on a lever, several series of observations; curves represent the extremes.

Fig. 12.—Discrimination of differences in temperature [1].

Both hands were adapted by immersion in water of temperature T_a, they were then separately placed simultaneously in water at temperatures T and T_1. ΔT = least value of $(T_1 - T)$ which could be detected

metrically equal stimuli of different colors, reaction time is independent of the color [22]. Reaction time for eye to turn towards a stimulus in indirect vision is 0.151 sec (or 1 181 sec) if stimulus lies 1° (or 5°) from fixation point [10]. For medium intensity, reaction time to monocular stimulation is about 0.015 sec greater than for binocular [43].

TABLE 4.—DISCRIMINATION REACTION TIME

Unit of: T = 0 001 sec; L_1, L_2 = 1 cm, λ = 1mμ = 10Å

Position of squares* or circles†			Contrast [21]			Lengths‡ [21]		
Contrast [21]								
	λ	II		λ	II	L_1	L_2	T
Black and White		205	Red (640) and Orange red	627	270	1	3	312
Red	640	222	Orange	614	257	1	25	313
Orange	614	218	Yellow	585	237	1	2	318
Yellow	585	211	Green	521	222	1	15	326
Green	521	218	Blue	452	231	1	1	335
Blue	453	226	Yellow and Green	521	232	1	05	351
†Circles [24]		296	Blue	452	222			

* Two colored squares each 3 by 3 cm, placed side by side; observer was to react with corresponding hand to indicate on which side the previously specified square was placed This type of discrimination reaction is the quickest The same procedure was used in the discrimination of lengths.

† On a background of approximately 2 6 millilamberts and at a visual angle of 45' to each side of fixation point was a circle of angular diameter = 21', brightness = 3 5% greater than that of background Either circle could be made to disappear, and the subject, by a reaction with the corresponding hand, indicated which disappeared.

Sound.—For finger reaction to sound of medium intensity, reaction time = 0.136 (±0.002) sec; individuals range from 0.082 to 0.195 sec. For very faint sound, the interval is increased by 0.06 to 0.07 sec [16].

Touch.—For finger reaction to tactile stimulus of medium intensity, reaction time is 0.148 sec [23].

The *discrimination reaction time* is the interval which elapses between the application of one of two possible, definite, expected stimuli and the performance of the prescribed movement indicating which of the two stimuli has been applied. For printed letters, 10-point type, average for the alphabet, the reaction time for Roman capitals is 0.327 sec, Roman lower case 0.325, for short words 0 353, for long words 0.355, for small (1 cm square) pictures of familiar objects 0 336 sec [6]. For other data, see Table 4.

Number Limitation and *Span of Apprehension.*—For college students, the greatest number of digits which an individual can repeat correctly immediately after a single auditory presentation averages 7.6 [5, 19], individuals range from 5 to 11 [8]; for visual presentation the average is 8.0 [19].

Fig. 13.—Reaction time for non-foveal stimulation [48].

ΔT = excess of reaction time over that required for foveal excitation. Abscissa indicates angular position of image upon the retina. Finger reaction.

Fig. 14.—Span of apprehension [41].

N = number of dots exposed; ordinates = % of judgments which were correct.

When a number of black dots irregularly arranged upon a well illuminated white background were exposed to view for a very short interval (0.038 sec) and the subject was required to determine the number of dots presented, the average number of correct judgments made after considerable, but not extreme, practice was as shown in Fig. 14. The visual angle subtended by the dots was well above the threshold value.

LITERATURE

(For a key to the periodicals see end of volume)

[1] Abbot, *550*, No **68**; 14 [2] Baird, *Color Sensitivity of the Peripheral Retina*, Washington, 1905 [3] Bauch, *Fortschritte d Psychol.*, **1**: 169; 13. [4] Bruner, *551*, **11**: 46, 08 [5] Carothers, *551*, **46** 29, 36; 21. [6] Cattell, *552*, **3**: 463, 86 [7] Cobb, *553*, **21**: 23, 14 [8] Cobb and Geissler, *553*, **20**: 425, 13. [9] Coblents and Emerson, *51A*, **14**: 167, 18. [10] Dodge, *Psych Rev*, *Monograph Suppl*, **36**: 19; 07 [11] Fechner, *Psychophysik*, **1**: 193, 89. [12] Ferree and Rand, *Trans Illum Eng Soc*, **16**: 769, 20 [13] Ferree, Rand and Buckley, *554*, **3**: 352, 20. [14] Fernberger, *555*, **21**: 346, 14. *555*, **27**: 269; 16 *554*, **1**: 515, 16; **3**: 141; **20.** **4:** 71, 21 [15] Fletcher and Wegel, *8*, **19**: 533, 22 [16] Froeberg, *551*, **6**: 00, 07 [17] Fullerton and Cattell, *On the Perception of Small Differences*, 1892 [18] Garrett, *551*, **86**: 56, 22. [19] Gates, *U of Cal Publ in Psych*, **1**: 327, 16

[20] Helmholtz, *B69*, **2**: 36, 24 [21] Henmon, *Time of Perception, etc*, New York, 1906 [22] Holmes, *Diss*, Columbia Univ, 1923 [23] Jastrow, *Time Relations of Mental Phenomena*, 1890 [24] Johnson, *554*, **7**: 34; 24. [25] Johnson, *554*, **7**: 5, 24 [26] Knudsen, *8*, **31**: 84; 23 [27] König, *76*, **1897**: 559 [28] König, in Helmholtz Festgruss, 1891. [29] König and Brodhun, *76*, **1888**: 000

[30] Koester, *Centralbl. f Physiol*, **10**: 433; 96 [31] Kraepelin, *552*, **2**: 306; 85 [32] Kranz, *8*, **22**: 66, 23 [33] Lane, *8*, **19**: 492; 22 [34] Lohmann, *Z Sinnesphysiologie*, **41**: 307; 06 [35] Merkel, *552*, **5**: 253; 89. [36] Merkel, *552*, **9**: 409, 94. [37] Nagel, *B69*, **2**: 318; 24. [38] Nagel, *B69*, **2**: 320, 24. [39] Nagel, *B69*, **2**: 325, 24

[40] Nutting, *51A*, **5**: 261. 08 09 [41] Oberly, *555*, **35**: 336-338; 24 [42] Parsons, *B70*, 28, 54, 59, 15 [43] Poffenberger, *551*, **23**: 48; 12. [44] Rice, *551*, **31**: 404; 20 [45] Reeves, *21*, **47**; 143, 145; 18. [46] Rice, *551*, **20**: 30; 12. [47] Southall, *B69*, **2**: 128 (footnote); 24. [48] Southall, *B69*, **2**: 33 (footnote), 24 [49] Steindler, *75*, **115**: 115; 06. [50] Stratton, *552*, **12**: 538, 96 [51] Stücker, *75*, **96**: 367; 07. [52] Urban, *Arch. ges. Psychol.*, **21**: 1; 14 [53] Vance and Schaefer, *550*, **69**: 114, 115, 14. [54] Woodworth and Bruner, *O*

ARRANGEMENT OF CHEMICAL SUBSTANCES

ARRANGEMENT OF CHEMICAL SUB-

ARRANGEMENT DES SUBSTANCES CHIMIQUES

Throughout I. C. T., except when otherwise indicated, the tabular arrangement of all chemical substances and of all systems capable of representation by formula is in accordance with a system called the "Standard Arrangement," which will now be explained and which should be learned by every user of I. C. T.

L'arrangement tabulaire de toutes les substances chimiques et de tous les systèmes susceptibles d'une représentation par formule est, dans les T. C. I, excepté lorsqu'il y a une autre indication, en accord avec un système appelé "arrangement type," (standard arrangement) expliqué ci-dessous, qui devra être appris par chaque personne qui veut utiliser les T. C. I.

Elementary Substances

Substances Élémentaires

All tables containing *only* elementary substances (𝔄-Tables) are arranged in alphabetical order of the symbols of the elements. In tables containing both elements and compounds (𝔄𝔅-Tables) the elements follow the "standard arrangement," *v. infra*

Toutes les tables ne contenant que les substances élémentaires (Tables 𝔄) sont arrangées dans l'ordre alphabétique des symboles des éléments. Dans les tables contenant les éléments et les corps composés (Tables 𝔄𝔅) les éléments se trouvent suivant l' "arrangement type" voir *infra*.

Chemical Compounds and Other Systems Represented by Formula

Composés Chimiques et Autres Systèmes Représentés Par Formule

The arrangement is based upon the following table of "Key-numbers" of the elements:

L'arrangement est basé sur la table suivante des "nombres clés" des éléments

KEY-NUMBERS OF THE ELEMENTS / NOMBRES CLÉS DES ÉLÉMENTS																									
-6	5	-4	-3	2	1		1	2	3	4	5		6	7	8	9	10		11	12	13	14	15		16	17	18	19	20
(He	Ne	A	Kr	Xe	Rn)		O	H	F	Cl	Br		I	(85)	S	Se	Te		N	P	As	Sb	Bi		C	Po	Si	Ti	Ge

| | | | | | | | 46 | 47 | 48 | 49 | 50 | | 51 | 52 | 53 | 54 | 55 | | 56 | 57 | 58 | 59 | 60 | | 61 | 62 | 63 | 64 | 65 |
| | | | | | | | Cr | Mo | W | U | V | | Cb(Nb) | Ta | Pa | B | Al | | Sc | Y | La | Ce | Pr | | Nd | (61) | Sa | Eu | Gd |

| Ac | Ag | Al | As | Au | | | B | Ba | Be | Bi | Br | | C | Ca | Cb | Cd | Ce | | Cl | Co | Cr | Cs | Cu | | Dy | Er | Eu | F | Fe |
| 74 | 32 | 55 | 13 | 33 | | | 54 | 79 | 75 | 15 | 5 | | 16 | 77 | 51 | 29 | 59 | | 4 | 44 | 46 | 85 | 31 | | 67 | 69 | 64 | 3 | 43 |

| | | | | | | | Os | P | Pa | Pb | Pd | | Po | Pr | Pt | Ra | Rb | | Re | Rh | Ru | S | Sa | | Sb | Sc | Se | Si | Sn |
| | | | | | | | 35 | 12 | 53 | 23 | 41 | | 17 | 60 | 37 | 80 | 84 | | 34 | 40 | 39 | 8 | 63 | | 14 | 56 | 9 | 18 | 22 |

To locate a given compound, first write its "key-formula," neglecting water of crystallization, thus:

Afin de situer un composé donné, il faut d'abord écrire sa "formule-clé," en négligeant l'eau de cristallisation, ainsi:

Compound / Composé	Na_2SO_4	$HClO_4.3H_2O$	$Hg(C_{18}H_{33}O_2)_2$	$2Fe_2O_3.P_2O_6.12H_2O$	$Na_2Pr_2(NO_3)_{12}.24H_2O$	$I_2C_6H_3SO_3H$	$(NH_4)_2CO_4$
Key formula / Formule-clé	82 8 1	4 2 1	30 16 2 1	43 12 1	60 45 11 1	16 8 6 2 1	16 11 2 1

In writing a key-formula the key-numbers must be written *in descending order*.

All chemical compounds (𝔅-Tables) are arranged in the inverse numerical order of their key-formulae. *Example:* to find the compound $Hg(C_{18}H_{33}O_2)_2 = 30 - 16 - 2 - 1$; First, turn to section 30 of the table. Then follow down the column of chemical formulae until element 16 (C) is first encountered. From this point continue until element 2 (H) is found, and then on until element 1 (O) is reached. At this point will be found all the compounds composed of the four elements Hg, C, H, and O and these compounds are arranged in an obvious manner according to the subscripts in the chemical formula. To facilitate the use of the tables, key-numbers are inserted at frequent intervals either along the top of the page or down the left hand column or both

In looking for a chemical compound *always consult the 𝔅-Table*, the scope of which provides for *all* chemical compounds except those of the radioactive elements, of which only compounds of U, Th and Ra are given in the 𝔅-Table. For the others see p. 364. In certain of the 𝔅-Tables, at the point where key-formulae beginning with 16 occur, there will be found frequently only a few of the simpler compounds, and the reader will be referred to a

Lorsqu'on écrit une formule-clé, les nombres clés doivent être écrits *dans l'ordre des valeurs décroissantes*.

Tous les composés chimiques dans toutes les tables (Tables 𝔅.) sont arrangés d'après l'ordre numérique inverse de leurs formules-clés. *Exemple:* pour trouver le composé Hg $(C_{18}H_{33}O_2)_2 = 30$ 16-2-1; il s'agit premièrement de chercher la section 30 de la table; ensuite de suivre en descendant la colonne des formules chimiques jusqu'à ce qu'on trouve l'élément 16 (C). De ce point, on continue jusqu'à ce qu'on rencontre l'élément 2 (H), et ensuite jusqu'à ce que l'élément 1 (O) soit atteint. On trouvera alors à ce point tous les composés renfermant les quatre éléments Hg, C, H et O et ces composés sont arrangés d'une manière apparente en relation avec les indices de leurs formules chimiques. Afin de faciliter l'usage des tables, les nombres-clés sont inscrits, à de fréquents intervalles, ou au haut de la page ou le long de la colonne gauche, ou aux deux places.

Pour la recherche d'un composé chimique, il s'agit de *consulter toujours la Table 𝔅* dont le but est de renseigner sur *tous* les composés chimiques, à l'exception des éléments radio-actifs, dont seuls ceux de U, Th et Ra sont donnés dans la Table 𝔅. Pour les autres, voir p. 364. Dans certaines des Tables 𝔅, au point où les

STANCES AND SYSTEMS IN I. C. T.

DIE ANORDNUNG DER CHEMISCHEN VERBINDUNGEN

Durch die ganzen I. C. T., ausgenommen es ist etwas anderes angegeben, ist die tabellarische Anordnung aller chemischen Verbindungen und aller durch chemische Zeichen oder Formeln darstellbarer Systeme, nach der "Normal-Anordnung" (standard arrangement), durchgeführt. Sie ist im folgenden dargelegt und soll von jedem Leser der I. C. T. erlernt werden.

Elementare Stoffe

Alle Tafeln, welche *nur* elementare Stoffe (A-Tabellen) enthalten, sind in alphabetischer Reihenfolge nach den Symbolen der Elemente angeordnet. In den Tafeln, welche beides, Elemente und Verbindungen (AB-Tabellen), enthalten, folgen die Elemente der "Normal-Anordnung." Siehe weiter unten.

Die chemischen Verbindungen und andere durch Formeln darstellbare Systeme

Die Anordnung ist auf der folgenden Tafel begründet, welche die "Schlüsselnummern" der Elemente enthält:

ORDINE DI ELENCAZIONE DELLE SOSTANZE

In tutti i volumi delle T. C. I. l'ordine in cui le sostanze ed i sistemi rappresentabili con formule sono disposti nelle tabelle è (tranne che non sia diversamente indicato) quello "standard" illustrato più avanti. Chiunque voglia servirsi delle T. C. I. deve anzitutto apprendere in che consiste questo sistema "standard."

Sostanze Elementari

Tutte le Tabelle contenenti soltanto sostanze elementari (tabelle A) sono disposte secondo l'ordine alfabetico dei simboli degli elementi. Nelle tabelle che comprendono elementi e composti (tabelle A-B) gli elementi sono ordinati secondo la disposizione "Standard" e infra.

Composti Chimici ed Altri Sistemi Rappresentati da Formule

La disposizione è basata sul quadro seguente di "numeri chiave" degli elementi.

SCHLÜSSELNUMMERN DER ELEMENTE / NUMERI CHIAVE DEGLI ELEMENTI

21	22	23	24	25		26	27	28	29	30		31	32	33	34	35		36	37	38	39	40		41	42	43	44	45
Zr	Sn	Pb	Th	Ga		In	Tl	Zn	Cd	Hg		Cu	Ag	Au	Re	Os		Ir	Pt	Ma	Ru	Rh		Pd	Mn	Fe	Co	Ni

66	67	68	69	70		71	72	73	74	75		76	77	78	79	80		81	82	83	84	85	86
Tb	Dy	Ho	Er	Tm		Yb	Lu	Hf	Ac	Be(Gl)		Mg	Ca	Sr	Ba	Ra		Li	Na	K	Rb	Cs	(87)

Ga	Gd	Ge	Gl	H		Hf	Hg	Ho	I	In		Ir	K	La	Li	Lu		Ma	Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O
25	65	20	75	2		73	30	68	6	26		36	83	58	81	72		38	76	42	47	11		82	51	61	45	1

Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr		(61)	(75)	(85)	(87)
78	52	66	10	24		19	27	70	49	50		48	57	71	28	21		62	34	7	86

Um eine gegebene Verbindung aufzufinden, hat man zuerst seine Schlüsselformel aufzuschreiben, wobei man das Kristallwasser auslässt. z B.:

Per trovare il posto di un dato composto bisogna prima scrivere la formula chiave trascurando l'acqua di cristallizzazione, p. es.

Verbindungen	Composto	Na_2SO_4	$HClO_4 \cdot 3H_2O$	$Hg(C_{18}H_{35}O_2)_2$	$2Fe_2O_3 \cdot P_2O_5 \cdot 12H_2O$	$Ni_3Pr_2(NO_3)_{12} \cdot 24H_2O$	$I_2C_6H_8SO_2H$	$(NH_4)_2CO_3$
Schlüssel-formel	Formula chiave	82-8-1	4-2-1	30-16-2-1	43-12-1	60-45-11-1	16-8-6-2-1	16-11-2-1

In die Schlüsselformel müssen die Schlüsselnummern in *absteigender* Reihenfolge geschrieben werden.

Alle chemischen Verbindungen (B-Tabellen) sind in der umgekehrten Reihenfolge der Schlüsselformeln angeordnet Z B : Um die Verbindung $Hg(C_{18}H_{35}O_2)_2 = 30\text{-}16\text{-}2\text{-}1$ zu finden, hat man zuerst den Abschnitt 30 aufzusuchen. Dann hat man den Kolonnen der chemischen Verbindungen abwärts zu folgen, bis man zuerst das Element 16 (C) antrifft, von da an setzt man weiter fort, bis das Element 2 (H) gefunden ist und dann weiter, bis das Element 1 (O) erreicht ist. Bei dieser Stelle werden alle Verbindungen gefunden, welche sich aus den 4 Elementen Hg, C, H, und O zusammensetzen. Diese Verbindungen sind in deutlicher Art, entsprechend der Bezeichnungsweise chemischer Formeln, angeordnet. Um den Gebrauch der Tafeln möglichst zu erleichtern, sind die Schlüsselnummern häufig an verschiedenen Stellen eingefügt. Sie befinden sich entweder am Kopf der Seiten, oder auf der linken Seite unten, oder an beiden Stellen.

Um eine chemische Verbindung zu suchen, *benutze man immer die B-Tabellen*: die *alle* chemischen Verbindungen enthalten, ausgenommen jene der radioaktiven Elemente. Von diesen sind

Nella formula chiave, i numeri chiave devono essere scritti in ordine decrescente.

Tutti i composti in tutte le tabelle (Tabelle B) sono disposti nell'ordine numerico inverso delle loro formule chiavi.

Supponiamo ad es. di voler trovare il composto $Hg(C_{18}H_{35}O_2)_2 = 30\text{-}16\text{-}2\text{-}1$. Prima si cerca la sezione 30 della Tabella, poi si scorre la colonna delle formule fino ad incontrare l'elemento 16 (C). Da questo punto si continua finchè si trova l'elemento 2 (H), e quindi fino a raggiungere l'elemento 1 (O). Qui si trovano tutti i composti risultanti dai quattro elementi Hg, C, H e O ordinati secondo gli indici delle formule. Per facilitare l'uso delle tabelle i numeri chiave sono inseriti ad intervalli frequenti nella testata o lungo il margine sinistro della pagina, o nell'una e nell'altro.

Per cercare un composto bisogna *sempre consultare la tabella B* che contiene *tutti* i composti tranne quelli degli elementi radioattivi; di questi sono riportati nella tabella B soltanto i composti di U, Th, Ra. Per gli altri vedi p. 364. In alcune tabelle B, laddove si trovano formule chiave che cominciano con 16, si troveranno spesso soltanto pochi composti fra i più semplici e il lettore

ℭ-Table where the remainder of such compounds will be found listed under a different arrangement known as

The ℭ-Arrangement

In this arrangement the compounds are arranged according to their empirical formulae (including water of crystallization), in the order C, H, with the remaining symbols alphabetical, e.g., C₄H₄O₄S. The ℭ-Tables, however, will not contain any carbon compound whose key-formula contains a number greater than 16.

SYSTEMS OF MORE THAN ONE COMPONENT

The components of each system are first arranged according to the standard arrangement, giving the order A, B, C, etc. The systems are then arranged, according to the standard arrangement, in the order of their A-components. All systems having the same A-component will be found (under that component) in the order of their B-components, etc.

In certain tables, the above plan will be based upon the ℭ-arrangement instead of the standard arrangement. Such cases will always be so indicated.

Name Indices

The chemical formulae of nearly all of the organic compounds and minerals whose properties are given in I. C. T. can be found with the aid of the extensive indices of names given on p. 174 and 280. If the name is not found there, other works of reference must be consulted for the formula. It should be noted, however, that the exact formula is not required. The compound can be readily located if only the elements composing it are known (in the case of inorganic compounds) or if only the number of carbon atoms are known (in the case of organic compounds) provided only that the user can recognize either name or formula when he sees it.

PHYSICAL PROPERTIES OF CHEMICAL SUBSTANCES

INTRODUCTION

The following tables (p. 96 to 314) are intended to serve as a source of ready reference for the *approximate* values of certain properties of chemical substances, displayed in such a manner as to be of the greatest utility. The values given may be uncertain by one or more units in the last significant figure. Non-significant figures are given in small type. Thus, 23oo indicates that the correct value lies between 1800 and 2800, with 2300 as most probable value.

More accurate values for these properties, if known, will be found in subsequent sections of I. C. T., together with their literature references.

A. ELEMENTARY SUBSTANCES AND ATMOSPHERIC AIR

A-Tables, p. 102. Values in parentheses are estimated, usually with the aid of the Periodic Law.

B. CHEMICAL COMPOUNDS. STANDARD ARRANGEMENT (v. p. 96)

B-Tables, p. 106

1. Formula or formula and name.
2. Gram-formula-weight. (I. C. T. atomic weights, v. p. 43.)
3. Crystal system.
   B-Table.
   Special tables.

in den ℬ-Tabellen nur die Verbindungen des U, Th und Ra enthalten. Für die anderen siehe Seite 364. In einigen ℬ-Tabellen, dort wo die Schlüsselnummern mit 16 beginnen, findet man häufig nur einige wenige einfache Verbindungen. Der Leser wird dann auf die ℭ-Tabellen verwiesen, wo die restlichen derartigen Verbindungen gefunden werden können. Diese Tabellen sind nach anderen Gesichtspunkten zusammengestellt. Es ist das die

ℭ-Anordnung (ℭ-Arrangement)

Bei dieser Anordnung sind die Verbindungen nach ihrer empirischen Formel gegeben (einschliesslich Kristallwasser) und zwar in der Ordnung C, H, die restlichen Zeichen dann in alphabetischer Ordnung, z.B. $C_6H_4I_2O_3S$. Die ℭ-Tabellen enthalten jedoch keine Kohlenstoffverbindung, in deren Schlüsselformel eine Zahl grösser als 16 vorkommt.

SYSTEME MIT MEHR ALS EINER KOMPONENTE

Die Komponenten jedes einzeln Systemes sind zuerst in der Reihenfolge A, B, C, u. s. w., entsprechend des "Standard-Arrangement" anzuordnen. Die Systeme sind dann, entsprechend des "Standard-Arrangement," in der Reihenfolge ihrer A-Komponenten angegeben. Alle Systeme, welche dieselbe A-Komponente haben, werden unter dieser Komponente in der Reihenfolge ihrer B-Komponenten gefunden.

In gewissen Tabellen wird dieser Plan entsprechend der ℭ-Anordnung, an Stelle des "Standard Arrangement," gewählt. Solche Fälle werden immer entsprechend bemerkt.

Namenverzeichnis (Englisch)

Die chemischen Formeln von so ziemlich allen organischen Verbindungen und Mineralien, deren Eigenschaften in den I. C. T. enthalten sind, können mit Hilfe des ausgedehnten Namenverzeichnisses auf Seite 174 und 280 gefunden werden. Ist der Name hier nicht auffindbar, so müssten andere Quellen für die Formel nachgesehen werden. Es soll aber bemerkt werden, dass eine genaue Formel nicht nötig ist. Die Verbindung kann bei anorganischen Verbindungen leicht aufgefunden werden, wenn nur die Elemente, die sie zusammensetzen, bekannt sind, bei organischen Verbindungen, wenn nur die Zahl der Kohlenstoffatome bekannt ist. Nötig ist es, dass der Leser entweder den Namen oder die Formel beim Ansehen erkennt.

DIE PHYSIKALISCHEN EIGENSCHAFTEN CHEMISCHER STOFFE

EINFÜHRUNG

Die folgenden Tafeln (s. 96 bis 314) sollen zur raschen Orientierung über angenäherte Werte gewisser Eigenschaften chemischer Verbindungen dienen. Sie sind in einer solchen Art angeordnet, um vom grösstmöglichem Nutzen zu sein. Die angegebenen Werte können auf einer und mehreren Stellen der letzten grossgeschriebenen Ziffer unsicher sein. Z.B. sagt die Zahl 2300 aus, dass der zwischen 1800 und 2800 liegende Wert am wahrscheinlichsten 2300 sein wird.

Genauere Werte für diese Eigenschaften können, wenn sie bekannt sind, in den weiter unten vorhandenen Abschnitten der I. C. T. zusammen mit der Literatur gefunden werden

A. ELEMENTARE STOFFE UND DIE ATMOSPHÄRISCHE LUFT

A-Tabellen, Seite 102. Werte, die in den Klammern sich befinden, sind geschätzt gewöhnlich nach dem periodischem System der Elemente.

ℬ. CHEMISCHE VERBINDUNGEN. NORMAL-ANORDNUNG [STANDARD-ARRANGEMENT] (siehe S. 97)

ℬ-Tabellen, Seite 106
1. Formel oder Formel und Name.
2. Gramm-Formel-Gewicht (Atomgewichte der I. C. T. siehe S. 43.)

sarà rimandato a una tabella ℭ dove si troveranno gli altri disposti con criterio differente che viene chiamato

La Disposizione ℭ

Secondo questa i composti sono disposti in base alle formule empiriche (compresa l'acqua di cristallizzazione) nell'ordine C, H e con i rimanenti simboli ordinati alfabeticamente P. es. $C_6H_4I_2O_3S$. Le tabelle ℭ non comprendono però composti del carbonio che hanno un numero chiave più grande di 16.

SISTEMI DI PIU' D'UN COMPONENTE

I componenti di ciascun sistema sono dapprima disposti secondo la disposizione tipo, nell'ordine A, B, C, etc. I sistemi sono quindi disposti, secondo la disposizione tipo, nell'ordine dei loro componenti A. Tutti i sistemi aventi lo stesso componente A verranno trovati, sotto questo componente, nell'ordine dei loro componenti B, etc.

In alcune tavole il piano sarà basato sulla disposizione ℭ in luogo della disposizione tipo. Di ciò verrà sempre fatta menzione.

Indici Per Nome (Inglese)

Le formule chimiche di quasi tutti i composti organici e minerali di cui sono riportate le proprietà nelle T. C. I. si possono trovare con l'aiuto di estesi indici di nomi dati a p. 174, e 280. Se negli indici non si trova il nome bisogna consultare altre opere per trovare la formula. Deve tuttavia notarsi che non è necessaria la formula esatta. Il composto può essere facilmente ritrovato se si conoscono solo gli elementi componenti (nel caso di composti inorganici) o se si conosce solo il numero di atomi di carbonio (nel caso di composti organici) purché il lettore sia in grado di riconoscerne il nome o la formula quando li vede.

PROPRIETA' FISICHE DELLE SOSTANZE

INTRODUZIONE

Le tabelle seguenti (p. 96 a 314) hanno lo scopo di fornire per una serie di sostanze valori approssimati di certe proprietà disposti in modo da essere della più grande utilità. I valori riportati pòs sono essere incerti per una o più unità nelle ultime cifre significative. Le cifre non significative sono indicate in caratteri piccoli. Così 2300 indica che il valore esatto si trova fra 1800 e 2800, e che 2300 è il valore più probabile.

Valori più precisi di queste proprietà quando sono conosciuti, sono riportati nelle sezioni successive delle T. C. I. insieme con le relative indicazioni bibliografiche.

A. SOSTANZE ELEMENTARI ED ARIA ATMOSFERICA

Tabelle A, p. 102. I valori fra parentesi sono calcolati generalmente con l'aiuto della legge periodica.

ℬ. COMPOSTI, DISPOSIZIONE STANDARD (v. p. 97)

Tabelle ℬ, p. 106

1. Formula oppure formula e nome.
2. Peso della formula in grammi. (T. C. I. pesi atomici v. p. 43.)
3. Sistema cristallino.
   Tabella ℬ.
   Tabelle speciali.
4. Punto di fusione. (Alla pressione di una atmosfera, tranne che non sia diversamente indicato dalla soprascritta; così $125^{17atm.}$ fonde a 125° alla pressione di 17 atmosfere.)
   Tabella ℬ

4. Melting point. (Under 1 atm. unless otherwise indicated by superscript, thus $125^{17atm.}$ melts at 125° under 17 atm.)

ẞ-Table.

5. Boiling point. (Under 760 mm Hg unless otherwise indicated by superscript, thus 321^{125} = boils at 321° under 125 mm Hg.)

ẞ-Table.

6. Density, g cm⁻¹. (At 20° unless otherwise indicated by superscript, thus 1.853^{40} = 1.853 g cm⁻¹ at 40°C.)

ẞ-Table.

7. Refractive index and dispersion, (n_D and H_β − H_α) for 20° unless otherwise indicated.

ABBREVIATIONS AND CONVENTIONS

at. or atm.	atmosphere
C.	cubic or regular
d.	decomposes, e.g., d. 335 = decomposes at ca. 335°; 335 d. = melts (resp. boils) at 335° with decomposition
diss.	a dissociation temperature
exp.	explodes
l.	liquid
H.	hexagonal
M.	monoclinic
P.	under pressure
s.	sublimation
s. d.	slight decomposition
R.	rhombic or orthorhombic
Tet.	tetragonal
Tr.	transition temperature
Tri.	triclinic
Trig.	trigonal
vac.	in vacuo
var.	variable

THE PROPERTY-SUBSTANCE TABLES

Following the General Tables will be found (p. 306) the Property-substance Tables, in each of which the substances, identified by Index Number, are arranged in ascending order of the values of the property, the intervals on the scale of values of the property being given in black-face type.

To Identify a Substance by Means of Its Properties.—*Example:* A liquid is found to have the following properties: B. P. = 81 1° at 745 mm, d = 0 783, n_D = 1 347. What is the substance ? With the aid of Craft's rule, first correct the boiling point to 760 mm. If the general nature of the substance is unknown, put $c = 10^{-4}$ in the Craft's equation, $\Delta t = cT_B(760 − P)$. Thus in the present instance, we should have $\Delta t = 10^{-4} \times (81.1 + 273)(760 − 745) = 0.3°$, and $t_B = 81.1 + 0.3° = 81.4°$. Next turn to the special B. P. (p. 310), d (p. 313), and n (p 276) tables and read off from these tables the index numbers of substances having values of the above properties in the neighborhood of those for the unknown substance. Thus, for the present example, the following index numbers will be obtained· For B. P , 130, 758, 727, 1612, 168, 277, 1535, 506, 792; for d, 208, 168, 395, 506, 3320, 1049, 262, 792, 5156; for n_D, 141, 168, 213. The only index number common to each of these properties is 168; and on turning to this index number in the General **C**-Table, we can readily identify our substance as acetonitrile. The identification can then be further checked by appropriate chemical tests, if desired.

3. Système cristallin.

Table **ẞ**.

Tables spéciales.

4. Point de fusion. (Sous 1 atm. à moins d'une indication par exposant, ainsi $125^{17atm.}$ = fond à 125° sous 17 atm.)

Table **ẞ**.

5. Point d'ebullition. (Sous 760 mm Hg à moins d'une indication par exposant, ainsi 321^{125} = boût à 321° sous 125 mm Hg.)

Table **ẞ**.

6. Densité, g cm⁻¹. (A 20° à moins d'une indication par exposant, ainsi $1,853^{40}$ = g cm⁻¹ à 40°C.)

Table **ẞ**.

7. Indice de réfraction, et dispersion (n_D et H_β − H_α) à 20° à moins d'une indication.

ABRÉVIATIONS ET CONVENTIONS

at. ou atm.	atmosphère
C.	cubique ou régulier
d.	Se décompose, par ex., d. 335 = se décompose à environ 335°; 335 d. = fond (resp. bout) à 335° avec décomposition
diss.	une température de dissociation
exp.	exploser
l.	liquide
H.	hexagonal
M.	monoclinique
P.	sous pression
s.	sublimation
s d.	légère décomposition
R.	rhombique ou orthorhombique
Tet.	tétragonal ou quadratique
Tr.	température de transition
Tri.	triclinique
Trig.	trigonal
vac.	dans le vide
var.	variable

TABLES DES PROPRIÉTÉS DES SUBSTANCES

On trouvera (p. 306) à la suite des Tables générales, les Tables des Propriétés des Substances, dans chacune desquelles, les substances identifiées par leur Nombre-Index, sont arrangées dans l'ordre ascendant des valeurs de la propriété; les intervalles de l'échelle des valeurs de la propriété sont donnés en caractères gras.

Pour identifier une substance au moyen de ses propriétés.— *Exemple:* On a trouvé qu'un liquide a les propriétés suivantes: P.E. = 81,1° à 745 mm, d = 0,783, n_D = 1,344. Quelle est la substance? Au moyen de la règle de Craft, on corrige premièrement le point d'ébullition à 760 mm. Si la nature générale de la substance est inconnue, on pose $c = 10^{-4}$ dans l'équation de Craft, $\Delta t = cT_B$ $(760 − P)$. Ainsi dans le cas présent, nous aurions $\Delta t = 10^{-4} \times$ $(81 1 + 273)(760 − 745) = 0 3°$, et $t_B = 81.1° + 0.3° = 81.4°$. Ensuite on cherche dans les tables spéciales des P.E. (p. 310), des d (p 313) et des n (p. 276) et on note les nombres-index des substances ayant les valeurs des propriétés ci-dessus dans le voisinage de celles de la substance inconnue. Ainsi, pour l'exemple présent, les nombres-index suivants seront obtenus; Pour le P.E., 130, 758, 727, 1612, 168, 277, 1535, 506, 792; pour d, 208, 168, 395, 506, 3320, 1049, 262, 792, 5156; pour n_D, 141, 168, 213. Le seul nombre-index commun à chacune de ces propriétés est 168; en revenant à ce nombre-index dans la Table générale **C**, et en notant les autres propriétés, on peut rapidement identifier notre substance comme étant acétonitrile. L'identification peut être alors poussée plus loin au moyen d'essais chimiques appropriés, si on le désire.

3. Kristall-System
ℬ-Tabellen.
Besondere Tabellen.

4. Schmelzpunkt. (Bei 1 Atmosphäre: wird dem Werte eine Zahl rechts hinaufgesetzt, so bedeutet diese den Druck unter welchem der Schmelzpunkt angegeben ist. Es bedeutet 125^{17atm}, der Schmelzpunkt ist bei einem Druck von 17 Atm. bei 125°)
ℬ-Tabellen.

5. Siedepunkt. (Unter 760 mm Quecksilber: wird dem Werte eine Zahl rechts hinaufgesetzt, so bedeutet diese Zahl den Druck, unter welchem der Siedepunkt angegeben ist. Es bedeutet 321^{125}; der Siedepunkt liegt bei einem Druck von 125 mm Hg bei 321°)
ℬ-Tabellen.

6. Dichte, g cm^{-3}. (Bei 20°C: wird dem Wert eine Zahl rechts hinaufgesetzt, so bedeutet diese Zahl die Temperatur, für welche die Dichte angegeben ist. Es bedeutet 1.853^{40}: die Dichte bei 40' beträgt 1.853).
ℬ-Tabellen.

7. Brechungs-Index und Dispersion, (n_D und $H_\beta - H_\alpha$) für 20°, wenn nichts anderes angegeben ist.

ABKÜRZUNGEN UND ZEICHEN

at. oder atm.	Atmosphäre
C.	kubisch oder regulär
d.	zersetzt sich, z. B. d335 bedeutet, zersetzt sich bei *ungefähr* 335°; 335d bedeutet, schmilzt (oder siedet) bei *ungefähr* 335° unter Zersetzung
diss.	Dissoziations Temperatur
exp.	explodiert
l.	flüssig
H.	hexagonal
M	monoklin
P.	unter Druck
s.	Sublimation
s.d.	schwache Zersetzung
R.	rhombisch oder orthorhombisch
Tet.	tetragonal
Tr.	Umwandlungstemperatur
Tri.	triklin
vac.	im Vacuum
var.	variabel

STOFF-EIGENSCHAFTS TAFELN

Den Haupttabellen folgend, findet man Seite 306 Stoff-Eigenschafts Tafeln. In jeder dieser Tafeln, in welcher die Stoffe durch ihre Indexzahlen bezeichnet sind, werden die Stoffe in aufsteigender Ordnung der Werte dieser Eigenschaften dargestellt. Die Intervalle an der Scala der Eigenschaftswerte sind in fettgedruckten Ziffern angegeben.

Die Erkennung eines Stoffes mit Hilfe seiner Eigenschaften. *Beispiel:* Es ist eine Flüssigkeit gefunden, welche folgende Eigenschaften hat: Siede-Punkt 81.1° bei 745 mm, $d = 0.783$, $n_D = 1.344$. Welcher Stoff ist das? Mit Hilfe der Regel von Craft corrigiere man zuerst den Siede-Punkt auf 760 mm. Ist die allgemeine Natur des Stoffes nicht bekannt, setze man $c = 10^{-4}$ in die Gleichung von Craft ein: $\Delta t = cT_B(760 - P)$. Im gegenwärtigen Falle ist also $\Delta t = 10^{-4} \times (81.1 + 275)(760 - 745) = 0.3°$, wonach dann der Siede-Punkt $t_B = 81.1° + 0.3° = 81.4°$ sich ergibt. Dann verwende man die Sd.P. Tabellen (Seite 310), die d-Tabellen (Seite 313) und die n-Tabellen (Seite 276), suche in diesen die Indexzahlen jener Stoffe heraus, deren oben genannte Eigenschaften solche Werte haben, die in der Nähe der Eigenschafts Zahlen des unbekannten Stoffes liegen. So erhält man für das gewählte Beispiel, folgende Indexnummern: für Sd. P. 130, 758, 727, 1612, 168, 277, 1535, 506, 792, für d, 208, 168, 395, 506, 3320, 1049, 262, 792, 5156; für n_D 141, 168, 213. Die einzige Index-Nummer, die alle drei Eigenschaften vereinigt, ist 168. Diese Index-Nummer wird in der Haupt ℭ-Tabelle aufgesucht; mit Beachtung noch anderer Eigenschaften kann man leicht die Flüssigkeit als Azetonitril erkennen. Die Identifizierung kann dann noch weiter durch eine chemische Untersuchung, wenn nötig, bestätigt werden.

---

5 Punto di ebollizione. (Alla pressione di 760 mm Hg tranne che non sia altrimenti indicato dalla soprascritta; così 321^{125} = bolle a 321° alla pressione di 125 mm Hg.)
Tabella ℬ.

6. Densità, g cm^{-3} (A 20°, tranne che non sia altrimenti indicato dalla soprascritta; così $1.853^{40} = 1.853$ g cm^{-3} a 40°C.)
Tabella ℬ.

7 Indice di rifrazione e dispersione (n_D e $H_\beta - H_\alpha$) per 20° tranne che non sia altrimenti indicato.

ABBREVIAZIONI E CONVENZIONI

at. oppure atm.	atmosfera
C.	cubico o regolare
d.	si decompone; per es. d335 = si decompone a *ca.* 335°; 355d = fonde (o bolle) a 335° con decomposizione
diss.	una temperatura di dissociazione
exp.	esplode
l.	liquido
H.	esagonale
M.	monoclino
P.	sotto pressione
s.	sublimazione
s d	leggera decomposizione
R.	rombico od ortorombico
Tet.	tetragonale
Tr.	temperatura di trasformazione
Tri.	triclino
Trig.	trigonale
vac.	nel vuoto
var.	variabile

LE TABELLE DELLE PROPRIETA' DELLE SOSTANZE

Seguendo le tabelle generali si troveranno (p. 306) le tabelle delle proprietà in ciascuna delle quali le sostanze, indicate col numero indice, sono disposte secondo l'ordine ascendente dei valori della proprietà. Gli intervalli nella scala dei valori della proprietà sono indicati in grassetto.

Identificazione di una sostanza a mezzo delle sue proprietà.— *Esempio:* si supponga che un liquido abbia le seguenti proprietà: B.P. = 81.1° a 745 mm, $d = 0.783$, $n_D = 1.344$. Che sostanza è? Con l'aiuto della regola di Craft, bisogna anzitutto ridurre il punto di ebollizione a 760 mm. Se non si conosce la natura della sostanza bisogna mettere, nella equazione di Craft, $c = 10^{-4}$, $t = cT_B(760 - P)$. Così, nel caso nostro, si avrebbe $t = 10^{-4} \times (81.1 + 273)(760 - 745) = 0.3°$, e $t_B = 81.1° + 0.3° = 81.4°$. Dopo bisogna guardare alle tabelle speciali per il B. P. (p. 310), per d (p. 313) e per n (p. 276), e ricavare da queste tabelle i numeri indici delle sostanze aventi valori delle suddette proprietà vicini a quelli della sostanza sconosciuta. Così, per il nostro esempio, si otterranno i seguenti numeri indici: per B.P., 130, 758, 727, 1612, 168, 277, 1535, 506, 792; per d, 208, 168, 395, 506, 3320, 1049, 262, 792, 5156; per n_D 141, 168, 213. L'unico numero indice comune a ciascuna di queste proprietà è 168; tornando a questo numero indice nella Tabella Generale ℭ, e osservando le altre proprietà, si può prontamente identificare la sostanza nell'acetonitrile.

La identificazione può quindi essere ulteriormente comprovata da appropriati saggi chimici, se si desidera.

ELEMENTARY SUBSTANCES AND ATMOSPHERIC AIR. λ-TABLE

THE GASEOUS STATE

Chem. symb.	Standard density (0°, 1A., g l⁻¹) d_0	Density of the saturated vapor at the normal boiling point g l⁻¹ d_s	Critical constants t_c °C	p_c atm	d_c g cm⁻¹	Specific heat joules per gram atom at 15° C_p	Viscosity η = A × 10⁻⁸ poises A	t
A	1.7824	5 89	-122 4	48 0	0 531	20 2	221	20
As			>1400					
Br			302		1 18		155	20
Cl	3.214		144	76	0 573	17 2	132	20
F	1.695							
H	0.08987	1 33	-239.9	12 8	0 0310	14 5s	88.7	20
He	0 1785	(11 2)	-267.9	2 26	0 069	20 9	197	20
Hg		0 020 at 320°	1650	3500	5		494	273
I			553.				184	124
Kr	3 708	(8.3)	- 62 6	54.2			248	20
N	1.2506	4 61	-147 1	33.5	0.311	14 56	176 5	23
Ne	0.9002	9 46	-228.7	26 9	0.484		312	20
O	1.4290	4 74	-118 8	49.7	0 430	14 60	203 9	23
O₃	3 0₃ at -80°		- 5 0	(67)	0 54			
P			721	100				
Rn	9.73	(12 6)	104 4	62 4			229	20
S			1040					
Tl		14 8						
Xe	5 851	(9 7)	16 6	58 2	1 15		225	20
Air	1 2930						284 2	20

THE LIQUID STATE

Chem. symb.	Density g cm⁻¹ d		Thermal expansion $\frac{1}{d_0}\frac{dd}{dt}$ = A × 10⁴ A	at t°	Normal boiling point (s = "solid") t_B	Latent heat of vaporization at t_B Kilo-joules per gram atom (s = "solid") L_v
	d	t	A	at t°	t_B	L_v
A	1.402	-185 7	4500	-183	-185 7	6 3
Ac					(>1700)	
Ag	9 4	960	11o	960-1200	1950	249
Al	2.40	658	11₃	658 1100	1800	225.
As					615 s	139 s
Au	17.	1063			2600	368
B					(2550)	
Ba					1140.	361.
Be					(1500)	
Bi	10.1	270.	12₂	270-630	1150	193.
Br	3.119	20	11oo	0 30	58 7s	15 0
C					4200	600
Ca					1170	399.
Cb					(>3300)	
Cd	8 0	320	15o	320 540	767	107.
Ce					1400	
Cl	1.557	- 33 6	1500	-34	-34 6	10.0
Co					2900	380
Cr					2200	320
Cs	1 84	26	370	27-123	670	73.
Cu	8.3	1083.	19o	1083-1295	2300.	467.

THE LIQUID STATE.—(Continued)

Chem. symb.	d	t	A	at t°	t_B	L_v
F	1.11	-187.	3000	-200	-187.	(6.)
Fe	6 9	1530			3000.	380.
Ga	6 095	29.7			>1600	
Ge					(2700.)	(500.)
H	0 0709	-252 7	13000	-255	-252.7	0.450
	0 126	-268 9				
He	0.147	-270 8			-268.9	0.10
	d_{max}					
Hf					(>3200.)	
Hg	13 546	20	182	20	356 9o	59. 3
I	4.00	107.	80o	107-150	184.3s	22.0
In					>1450.	
Ir					(>4800)	
K	0.83	62.	29o	62-150	760.	84.
Kr	2 6	146.			-151.8	(9.4)
La					1800.	
Li			18o	186-230	>1200.	(170.)
Mg	1 57	650	38o	650-800	1110.	262.
Mn					1900.	240.
Mo					3700.	710.
N	0 808	-195 8	6000	-195	-195 8	2.80
Na	0.93	97.5	28o	100-200	880	105.
Ne	1 204	-245 9			-245.9	1.74
Ni					2900.	380.
O	1 14	-183.	4100	-195	-183 0o	3.41s
O₃	1.71	-183.	2000	-183	-112.	4.8s
Os					(>5300.)	
P	1 745	44.5	520	50-60	280.	
Pa					(6200.)	
Pb	10 3	327.	120	327-825	1620.	193.
Pd	11.	1550.			2200.	
Pt	19.	1755.			4300.	520.
Ra					(1140.)	
Rb	1.475	38.5	340	40-140	700.	74.
Rh					(>2500.)	
Rn	4.4	-62.			-61 8	(18 1)
Ru					(>2700.)	
S	1.808	115.	430	115	444 6	8.9s
Sb	6.55	631.	10o	630-1050	1380.	190.
Sc					(2400)	
Se					688	31.
Si					2600	170?
Sn	6 98	232.	10o	232-1600	2260	325.
Sr					1150.	383.
Ta					(>4100)	
Te					1390.	85.
Th					(>3000)	
Ti					(>3000.)	
Tl	11.0	300.	14o	300-350	1650.	120?
						256?
V					(3000.)	
W					5900.	910.
Xe	3 06	-109 1			-109.1	(13.4)
Yt					(2500)	
Zn	6.7	463	15o	419-543	907.	99.2
Zr					(>2900.)	
87					(620.)	(69.6)
85					(520)	(83.7)

Air

Mole % O₂ in liquid	d	t	A at t°	t_B	L_V
10	0.831	−195.0		−195 0	
20	.856	−194.3		−194 3	
20.94	.861	−194.2		−194 2	0 185 (per gram)
30	.893	−193.5		−193 5	
40	.932	−192.6		−192 6	
50	.974	−191 5		−191 5	

Chem. symb.	C_p	t	A	n	t
P			2 3	6	25.
Pb			98.	− 6	400.
Rb	32 .	50	23.5	− 6	50.
S	30 4	100	95	10	115.
Sb	28	630	12	− 6	860.
Se			76 6	− 9	390.
Sn	31.	232	49 .	− 6	300.
Tl			74.	− 6	300.
Zn			43	− 6	440.
Air	1 91*	−200			

* Per gram, for liquid containing 20 94 mole % O₂

Chem. symb.	C_p Specific heat joules per gram atom		Electrical resistivity ohm-cm $R = A \times 10^n$		
	C_p	t	A	n	t
A	22 4	−100.			
Ag	33 8	907–1100	17 0	− 6	1000
Al	28	660	20 1	−- 6	657
Au	27.	1100	30 8	− 6	1063
Bi	31.	400	127		269 .
Br	36 .	13–45	7 8	12	17
Cd	36	321	34	− 6	400
Cl	33 5	0–24	>10	15	−70
Cs	32.	50	36 6	− 6	28
Cu	27.	1084	21 3	− 6	1083
Ga	23	119	27	− 6	30
H	0 975	−252			
Hg	27 .9	20	95 8	− 6	20
I	8 01	114–185	78	6	110 5
In			29 .	− 6	155
K	30.	63	13	− 6	62
Li			45 .	− 6	230
N	27 8	−200			
Na	32	100	9 7	− 6	100 .
Ni	33	1452	109		1500
O	26 4	−200			

Surface Tension

Chem symb	γ dyne cm⁻¹	t	Chem. symb	γ dyne cm⁻¹	t
A	12 5	−185 8	N	8 85	−195.8
Al	520	75 0	O	13 2	−183.
Bi	376	300	Pb	442 .	350.
Br	36	58 6	S	60.	120.
Cd	628 .	350	Se	72 .	217.
Cl	27	− 34 5			
Ga	358	30(CO₂)	Air, with 50 mole % O₂	11 6	−190.3
H	1 91	−252 7			
Hg	476	20			

Refractive Index

Chem. symb.	n_D	t	Chem. symb.	n_D	t
B	2 5*		N	1 205₃	−190.
Br	1 661	15	Na	0.0045	
Cd	0 82*		O	1 221	−181.
Cl	1 385	20.	Pb	2 6*	
H	1 097*	−252 8	S	1.929	110.
Hg	1 6–1 9	20	Se	2 9	220.
N	1 1975*	−195 8	Sn	2 1	

* These values are for the Hg line 5790 Å.

THE CRYSTALLINE STATE

Chem. symb. (At. wt. v. p. 43)	Crystal system or form	Density, g cm⁻¹		Thermal expansion $\frac{1}{l}\frac{dl}{dt} = A \times 10^{-6}$		Melting point °C	Specific heat joules per gram atom 1 joule = 4.185 cal.		Latent heat of fusion at t_F Kilojoules per gram atom	Electrical resistivity ohm-cm $R = A \times 10^{-6}$	
		d	t	A at t°		t_F	C_p at t°		L_F	A	t
A	C.	1.65	−233			−189 2 (1800)	25 9	−223	1 12		
Ac											
Ag	C.	10 5		18 9	20	960 5	25 2	20	11	1.62	20
Al	C.	2.702	20	23 03	20	660 0	24 2	20	8.0	2.62	20
As	Met.H.	5.7		4 7	20	814³⁶atm	25 8	0–100		35	0
	Black	4.7	20				27.0	0–100			
	Yel. C.	2 0	20								
Au	C.	19 3	20	14.2	20	1063.0	25.7	18	13.3	2.4	20
B		2 .		2		2300.	14	0–100		1.8 × 10¹²	0
Ba		3 5	20			850.					
Be	H.	1 8	20			1350.	16 1	0–100	12	18.5	20
Bi	H.	9 80	20	13 3	20	271	25.6	20	10 9	115	20
Br	R.	(3 4)				−7 2	23 5	−192 to −108	5.4	>10¹⁴	
C	Dia. C.	3 51	20	0 9	20		6.1	20		5 × 10²⁰	15
Graphite	C.	2.255	20	3	20	3500.	8.5	20		1400.	20
Graphite	Single crystal									39–60	20

THE CRYSTALLINE STATE.—(Continued)

Chem. symb.	Crystal system	d	t	A at t°		tF	Cp at t°		LF	A	t
Ca	C.	1 55	20	25	0–21	810	26 0	20		4 6	20
Cb		8 4	20			1950					
Cd	H.	8 6	20	29 8	20	320 9	28	20	6 2	7 5	20
Ce	C.	6 90	20			640	24 8	0 100		78	20
	H.	(6 7)									
Cl	R.	(1 9)				−101 6	28	−113	3 40		
Co	C.	8 9	20	12 3	20	1480	24 8	20	14 4	9 7	20
Cr	C.	7 1		8 2	20	1615	23	20	6 9	2 6	0
Cs		1 90	20	97	0–26	26	29	20	2 1	20	20
Cu	C.	8 92	20	16 6	20	1083	24 5	20	11 5	1 69	20
F		(1 3)				−223			(0 8)		
Fe	C.	7 86	20	11 7	20	1535	24 9	20	11 2	10 0	20
Ga	Tet.	5 91	20	18	0 30	29 75	23	12–23	5 55	53	0
Ge	C.	5 36	20			958 5	22 3	0–100		89 × 10^7	0
H	C.	0 0808	−262			−259 14	2 4	−260 6	0 059		
He						< −272 2					
Hf						(1700)					
Hg	H.?	14 19	−38 9	90	−190 to −40	−38 87	28.0	−40	2 33	21 3	−50
I	R.	4 93	20	93	20 100	113 5	27 8	20	8.38	1 3 × 10^15	20
In	Tet.	7 3	20	33	20	155	27 3	0 100		9	20
Ir	C.	22 4	20	6 5	20	2350	26 1	0-100		6	20
K	C.	0 86	20	83	20	62 3	29	14	2 38	7 0	20
Kr		(2)				−169			(1 5)		
La		6 15	20			826	26	0 100		59	18
Li	C.	0 53	20	56	20	186	23	0	(3 5)	9 3	20
Ma						(2300)					
Mg	H.	1 74	20	25 6	20	651	25	20	7 13	4 46	20
Mn		7 2	20	23	20	1260	24 6	0	8 4	5	
Mo	C.	10 2		4	20	2620 ±10	26	20-100		4 77	20
N	C.	1 026	−252 5			−209 86	23	−212	0 356		
Na	C.	0 97	20	71	20	97 5	28 4	20	2 65	4 6	20
Nd		6 9	20			840	27	0 100		79	20
Ne		(1 0)				−248 67			(0 24)		
Ni	C.	8 90	20	12 8	20	1452	25 8	20	18 17	6 9	20
O	H.	1 426	−252 5			−218 4	22 5	−221 8	0 22		
O₃	Ozone					−251.					
Os	H	22 48	20	6 1	20	2700.	25	20 100		9	20
P	Yel H.	1 82	20	12 5	0–40	44 1	23	9	0 654	10^17	11
	Red, C.	2 20	20			590^{43atm}	24	−21 to +7			
	Black									710 × 10^3	0
Pb	C.	11 34	20	29 1	20	327 5	26 5	20	4 70	21 9	20
Pd	C.	12 0	20	11 8	20	1555	26 2	18	16	10 8	20
Po						(1800)					
Pr		6 5	20			940	27	0 100		88	18
Pt	C.	21 45	20	8 9	20	1755	26 5	20	22	10 5	20
Ra		(5)				(960)					
Rb		1 53	20	90	20	38 5	28 7	0	2 18	12 5	20
Re						(3000)					
Rh	C.	12 5	20	8 4	20	1955	25	0-100		5 1	20
Rn		(1)				−71			(3 25)		
Ru	H.	12 2	20	9 1	20	2450	26	0 100		10	20
S	R.	2 07	20	64	40	112 8	23	0 30		2 × 10^?	20
	M.	1 96	20			119 0	24	0 30	1 18		
Sa		7 7				>1300					
Sb	H.	6 684	25	11 4	20	630 5	25	20	19 5	39	20
Sc		(2 5)				1200					
Se	Gray, Trig.	4 80	25	37	40	220	28	0- 41	(2.2)	1 2	20
	Red, H.?	4 50	25							85 × 10^3	20
Si	C.	2 4	20	2 8 7 3	20	1420	20 7	20		11 4	20
Sn	White, Tet.	7 31	20	20	20	231 85	26 9	18	(7)		
	Gray, C.?	5 750	20	5	−163 to −18		25 6	20			

THE CRYSTALLINE STATE.—(Continued)

Chem. symb.	Crystal system	d	t	A at $t°$		t_F	C_p at $t°$		l_F	A	t
Sr		2 6				800				23	20
Ta	C.	16 6		7	20	2850	27	20		15	20
Te	α Met. H.?	6 24	20	16 8	40	452	25	20	3 9	[5.8 − 33 × 10³]	
	β H.?	6 00	20								
Th	C.	11 2				1845	26 8	0 100		18	20
Ti	C.	4 5	20			1800	29	0-100		3	20
Tl	Tet.	11 85	20	28	20	303 5	26 6	20	6 15	18 1	20
U		18 7				<1850	28	0 100		60	20
V	C.	5 96				1710	24 6	0 100			
W	C.	19 3		4	20	3370	26	20 100		5 48	20
Xe		(2 7)				−140			(2 05)		
Yt		5 51				1490					
Zn	H.	7 140	20	33	20	419 45	25 3	20	7.1	6	20
Zr	C.	6 4	20			1700	25 2	0 100		170	0
85						(470)					
87						(23)					

CHEMICAL COMPOUNDS

ꕔ-TABLE

Compiled with the cooperation of Raleigh Gilchrist, F. W. Smithers and Edward Wichers, Bureau of Standards, Washington D. C.; J. A. Almquist, J. M. Braham and E. W. Guernsey, Fixed Nitrogen Laboratory, Washington, D. C.; H. E. Merwin, H. S. Roberts, R. B. Sosman and E. G. Zies, Geophysical Laboratory, Washington, D. C.; John C. W. Frazer, F. O. Rice and H. C. Urey, Johns Hopkins Univ., Baltimore, Md.; Robert D. Coghill, Florence Fenwick, Donald M. Hetler, Norman W. Krase and Hugh M. Spencer, Yale Univ., New Haven, Conn. The list of minerals was supplied by E. T. Wherry, Bureau of Chemistry, Washington, D. C.

General index number	Formula	Molecular weight (I. C. T atomic weights, v. p. 43)	Crystal system	Normal melting point, °C	Specific gravity 20°/4° (or at other indicated temperature)	Refractive index finding number, v. p. 165
1	H_2O	18 0154		0	0 917° l. 0 9982	203 8
2	H_2O_2	34 0154		— 1 7	1 643$_4^{-4.44}$ l. 1 442	16
3	$H_2O_2.2H_2O$	70 0462		— 51		
4	HF	20 0077		— 83	l. 0 988[14 6]	
5	$Cl_2.8H_2O$	215 039	R.	d. 9 6	1 23	
6	ClO_2	67 4580		— 76		
7	Cl_2O	86 9160		— 20?		
7 1	Cl_2O_6	166 916		— 1	1 65	
8	Cl_2O_7	182 916				
9	HCl	36 4657		—111	l. 1 194$^{-85\ 8}$	3
10	$HCl.H_2O$	54 4811		— 15.35	1 48	
11	$HCl.2H_2O$	72 4965		— 17 7	l. 1 46$_1^{18\ 3}$	
12	$HCl.3H_2O$	92 6119		— 24 4		
13	$HClO_4$	100 466		—112	l. 1 768	
14	$HClO_4.H_2O$	118 481		50	1 88 l. 1 776$_4^{50}$	
15	$HClO_4.2H_2O$	136 497		— 17 8		
16	$HClO_4.3H_2O$	154 512		— 43 2 (α) — 37 (β)		
17	HBr	80 9237		— 86	l. 2 16^{-68}	5
18	$HBr.2H_2O$	116 955		— 11	2 11^{-15}	
19	$HBr.3H_2O$	134 970		— 47 5		
20	$HBr.4H_2O$	152 985		— 55 8		
21	HBrO	96 9237				
22	$HBrO_3$	128 924		d. 100		
23	BrF_3	136 916		5		
24	IO_2	158 932		d. 130	4 2$_{10}^{10}$	
25	I_2O_5	333 864		d. 300	4 799$_4^{24}$	
26	HI	127 940		— 50.8	l. 2 847$^{-4.7}$	27
27	$HI.2H_2O$	145 955		— 43		
28	$HI.3H_2O$	163 970		— 48		
29	$HI.4H_2O$	181 985		— 36.5		
30	HIO_3	175 940	R.	110	4 629°	
31	HIO_4	191 940				
32	$HIO_4.2H_2O$	227 971	M. ?	d. 110		
33	$I_2O_5.HIO_3$	509 804		Tr. 170		
34	IF_5	221 932		8	l. 3 5	
35	ICl (α)	162 390		27 2	l. 3 24$_{14}^{44}$ 3 182$_4^9$	
35 1	ICl (β)	162 390	R.	13 9	l. 3 24$_{14}^{44}$ l. 3 182$_4^9$	
36	ICl_3	233 306	R.	ca. 33	3 11^{15}	
37	IBr	206 848		ca. 42	4 414^{10}	

Ag Al As Au   B Ba Be Bi Br   C Ca Cb Cd Ce   Cl Co Cr Cs Cu   Dy Er Eu F Fe   Ga Gd Ge Gl H   Hf Hg Ho I In   Ir K La Li Lu
47 65 13 83   54 79 75 15 5   18 77 51 29 59   4 44 46 85 31   67 00 64 3 43   25 65 20 75 2   73 30 68 6 26   36 63 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
38	SO₂............	64 0650		− 72 7		15
39	SO₃................ .. . SO₃	80 0650		16 83	1. 1 923	
40	S₂O₇................	176 130		0		
41	H₂S...	34 0804		− 82 9	1. 0.96⁻⁶⁶	10
42	H₂S₂	66 1454		− 88	1. 1.376	65
43	H₂S₃	98 2104		− 53	1. 1 496¹⁴	
44	H₂S₄	162 340			1 1 71¹⁴	
45	H₂SO₄...	98 0804		10 49	1 1.834	18
46	H₂SO₄.H₂O	116 095		8 62	1. 1 842¹⁴	
47	H₂SO₄.2H₂O	134 019		− 38 9	1. 1 650⁰	
48	H₂SO₄.4H₂O	170 142		− 25		
49	H₂SO₅..	114 080		45		
50	H₂S₂O₇ . . .	178 145		35	1. 1 9¹⁰	
51	H₂S₃O₈ ..	194 145		<60		
52	SF₄..	146 065		− 55		
53	SOF₂	86 0650		−110		
54	SO₂F₂ .	102 065		−120⁶⁶ᵐᵐ		
55	SCl₂..	102 981		− 78	1. 1 621¹⁴	56
56	SCl₄..	173 897		− 30		
57	S₂Cl₂	135 046		− 80	1. 1 678	61
58	SOCl₂	118 981		− 80	1. 1 638	52
59	SO₂Cl₂	134 981		− 54 1	1. 1 667	22
60	SO₃.SO₂Cl₂ . .	215 046		− 37 5	1. 1 837	
61	S₂O₅Cl₄ .	253 962	R.	57 d.		
62	SO₂OHCl..	116 531		− 80	1. 1 753	20
63	S₂Br₂	223 962		− 46	1. 2 635	64
64	SOBr₂.	207 897		− 50	1. 2 68¹⁴	
65	SOClBr	163 439			1. 2 31⁰	
66	SeO₂.....	111 200		340	3 953¹⁴	
67	HSe......... ...	80 2077				
68	H₂Se	81 2154		− 64	1. 2 12⁻⁴²	
69	H₂SeO₃	129 215	H.	d.	3 004¹⁴	
70	H₂SeO₄ . .	145 215	H.	58	2 950¹⁴ 1. 2 608¹⁴	
71	H₂SeO₄.H₂O .	161 230		25	2 627¹⁴ 1. 2 356⁰	
72	SeF₄ .	155 200		− 80		
73	SeF₆. .	193 200				
74	SeCl₄	221 032				
75	Se₂Cl₂	229 316			1. 2 906¹⁷ ⁴	
76	SeOCl₂	166 116		8 5	1. 2 44	
77	Se₂Br₂ .	318 232			1. 3 604¹⁴	
78	SeOBr₂	255 032		41 7	1. 3 38⁵⁰	
79	H₂SeO₄.SO₃	225 280		6 6		
80	H₂SeO₄.2SO₃	305 345		20		
81	SO₃.SeCl₄	301 097		165		
82	TeO₂—Tellurite	159 500	Tet. P.		Tet. 5 66⁰ R. 5 89⁰	1056
83	TeO₃	175 500		d.	5 08¹⁰ ⁶	
84	H₂Te .	129 515		− 48	1. 2 57ᵢ²⁰	
85	H₂TeO₄	193 515		d. 160	3.44¹⁹ ²	
86	Te(OH)₆ (α)	229 546	C.		3 053	
86.1	Te(OH)₆ (β)	229 546	M.		3 071	
87	TeF₆	241 500				
88	TeCl₂ . .	198 416		175		
89	TeCl₄ .	269 332		214		
90	TeCl₄.HCl.5H₂O	395 875		− 20		
91	TeBr₂	287 332		ca. 280		
92	TeBr₄ . .	417 164		ca. 380	4.31¹⁴	
93	TeI₄	635 228		259	8 403¹⁴	
94	2TeO₂.SO₃	399 065	R.	d. 500	4 7	
95	NO	30 0080		−161	1. 1 269ₐ¹⁶⁰·²	7
96	NO₂.	46 0080		− 9 3	1. 1 448	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
97	N_2O.	44.0160		−102 4	l. 1 226[-68]	2
98	N_2O_3	76 0160		−102	l. 1 447[2]	
99	N_2O_4	108 016	R.	30		
100	$2N_2O_4.H_2O$	234 047		5	l. 1.682[18]	
101	N_4O_4.	152 032				
102	NH_3	17 0311		− 77 7	0.817[-79]	
					l. 0.607	6
103	H_2N NH_2	32 0468		1 4	l. 1 011[14]	28
104	$N_2H_4.H_2O$	50 0622		< − 40	l. 1 03[21]	
105	N_3H	43 0317		− 80		
106	$NH_3 HN_3$	60 0628		110		
107	$2NH_3.H_2O$	52 0776		− 78		
108	$N_2H_4.HN_3$	75 0785		65		
109	HNO_3	63 0157		− 42	l. 1 502	12
110	$HNO_3.H_2O$	81 0311		− 38		
110 1	$HNO_3.3H_2O$	117 0619		− 18 5		
111	NH_2OH	33 0311		34	1 35	
					l. 1 204[23] [6]	21
112	H_2NO_4	81 0311	R.	− 34		
113	NH_4OH ..	35 0465		− 77		
114	H_4NO_4	99 0465		− 35		
115	$(OH)_2NON(OH)_2$	180 078		− 39		
116	NH_3NO_2	62 0314		72 d.		
117	NH_4NO_2	64 0468		d.		
118	NH_4NO_3	80 0468	R.	169 6	α 1 66[26]	
					β 1 725[26]	
119	NH_4ONNOH	79 0625		65		
120	$N_2H_4.HNO_3$	95 0625		70 7		
				62 1		
121	$NH_4NO_3.HNO_3$	143 063		12		
122	$N_2H_4.2HNO_3$	158 078		104		
123	$NH_4NO_3 2HNO_3$	206 078		30		
124	$NH_4NO_3 3NH_3$	131 140		ca. − 40		
125	NOF	49 0080		−134		
126	NO_2F	65 0080		−139		
127	$NH_4F.HF$	57 0465	R.		l. 1 211 [12]	
128	$N_2H_4(HF)_2$	72 0622	C.	105		
129	NCl_3	120 382			l. 1 653	
130	$NOCl$	65 4660		− 64 5	l. 1 417[-12]	
131	NO_2Cl	81 4660		< − 30	l 1 32[14]	
132	NH_4Cl—Salammoniac	53 4968	C.	89	1 536	145
133	$N_2H_4 HCl$	68 5125		89		
134	$N_2H_4 2HCl$	104 978	C.	198	1 42	
135	$NH_4Cl 3NH_3$	104 590		10 7		
136	$NH_4Cl 6NH_3$	155 683		− 18		
137	$NH_2OH.HCl$	69 4968	M.	151	1 67[17]	
138	NH_4ClO_4	117 497	R.	d.	1 95	489
139	$N_2H_4 HClO_3$	116 513		exp. 80		
140	$N_2H_4 HClO_4.2H_2O$	168 543		132		
141	$NOBr$	109 924		− 55 5		
142	$NOBr_3$	269 756		− 40	1 2 637	
143	NH_4Br	97 9548	C.		2 548	
144	$N_2H_4 HBr$	112 971		80		
145	$HBr 2NH_3$	114 986				
146	$NH_4Br.3NH_3$	149 048	R.	13 7		
147	$NH_4Br.6NH_3$	200 141		− 20		
148	NH_4I	144 971	C.		2 563	153
149	NH_4I_2	270 895		− 2	l. 2 46[14]	
150	NH_4I_3	398 835	R.		3.749	
151	$NH_4I NH_3$	162 002				
152	$N_2H_4.HI$	159 987		exp. 127		
153	$N_2H_4.2HI$.	287 926		220		
154	$NI_3.NH_3$	411 835		d.>20	3 5	

Ag Al As Au   B Ba Be Bi Br   C Ca Cb Cd Ce   Cl Co Cr Ca Cu   Dy Er Eu F Fe   Ga Gd Ge Gl H   Hf Hg Ho I In   Ir K La Li Lu
32 55 13 33   54 79 75 15 5   16 77 51 29 59   4 44 46 85 31   67 69 64 3 43   25 65 20 75 2   73 30 68 6 26   36 83 58 51 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
155	NH₄I.3NH₃	196 064		− 8		
156	NH₄I.4NH₃	213 095		− 5 1		
157	3NH₃.2HI	352 020		90		
158	NH₄I 6NH₃	247 157		28		
159	NH₄IO₃	192 971	R.	d. 150	3 309$_4^{41}$	
160	NH₄IO₄	208 971	Tet	exp.	3 056^{45}	
161	2NH₄IO₃.H₂O	403 957	Tri.	exp. 150		
162	3NH₂OH.HI	227 033		104		
163	N₂S₅	188 341		11	1 1 901^{45}	
164	N₄S₄	184 292	R.	178	2 22^{19}	
165	N₂O₃.2SO₃	236 146		230	2 14	
166	NH₄SH	51 1115				
167	(NH₄)₂S	68 1426		d.		
168	NO₂SO₃H	127 081	R.	73 d.		
169	NH₂SO₃H	97 0961	R.	205 d.	2 03$_4^{17}$	
170	NH₄HSO₄	115 112		146 9	1 78	
171	SO₂(NH₂)₂	96 112	R	92		
172	NH₃SO₃NH₄	114 127		125		
173	N₂H₄.H₂SO₄	130 127	R.	254	1 37	
174	(NH₄)₂SO₄ —Mascagnite	132 143	R.	513 d.	1 769	602
175	(NH₂OH)₂ H₂SO₄	164 143	M.	170		
176	(NH₄)₂S₂O₃	148 208	M.	d. 150		
177	(NH₄)₂S₂O₅	180 208	R.	d.		
178	(NH₄)₂S₂O₆	196 208	M.	d. 130		
179	(NH₄)₂S₂O₈	228 208	M.	d. 120	1 982	543
181	NH(SO₂NH₄)₂	179 223				
182	NH(SO₃NH₄)₂	211 223	M.	357	1 965	
183	(N₂H₄)₃ H₂SO₄	162 174		117		
184	NH₄SO₃F	117 104		245		
185	NSe	93 2080		exp. 200		
186	SeO₂(NO₃)₂	203 216		− 13		
187	NH₄HSeO₄	162 247	R.	d.	2 162	
188	(NH₄)₂SeO₄	179 278	M.	d.	2 194	686
189	(NH₄)₂SeBr₆	594 774	C.		3 326	
190	(NH₄)₂TeO₄	227 578			3 01^{24}	
191	P₂O₃	110 048	M.	22 5	2 135^{21}	
192	P₂O₄	126 048	R.?	>100	2 537$_4^{22,4}$	
193	P₂O₅	142 048		563 var.	2 387	
194	P₄O	140 096			1 912^{26}	
195	PH₃	34 0471		−132 5	l. 0 746^{90}	4
196	P₂H	63 0557			1 83^{19}	
197	P₂H₄	66 0788			l. 1 012	
198	P₉H₂	281 231			1 95^{18}	
199	P₁₂H₆	378 334			1 83^{19}	
200	H₂PO₃	81 0394		35		
201	H₃PO₂	66 0171			1 493^{18,8}	
202	H₃PO₃	82 0471		73 6	1 651^{21,2}	
203	H₃PO₄	98 0471		42 35	1 834^{18,2}	
204	PF₃	88 0240		−160		
205	PF₅	126 024		− 83		
206	POF₃	104 024		− 68		
207	PCl₃	137 398		−111 8	l. 1 574$_4^{20,8}$	47
208	PCl₅	208 314	Tet.	148 P.		
209	P₂Cl₄	203 880		− 28		
210	POCl₃	153 398		1 25	l. 1 675	25
211	P₂O₃Cl₄	251 880		< − 50	l. 1 587	
212	PH₄Cl	70 5128		28 atm		
213	PF₃Cl₂	158 940				
214	PBr₃	270 772		− 40	l. 2 852$_4^{19}$	62
215	PBr₅	430 604	R.			
216	POBr₃	286 772		56	2 822	
217	PH₄Br	114 971				
218	POCl₂Br	197 856		13	l. 2 104	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
219	POClBr₂	242.314		30	l. 2.45⁰⁰	
220	PI₃.	411 820	H.	61		
221	P₂I₄	569 776	Tri.	110		
222	PH₄I	161 987				
223	P₂S₃	158 243		290		
224	P₂S₄	222 373		276	2.03	
225	P₄S₄	285 462		298		
226	P₄S₃	220 291		172 5	2 03¹⁷	
227	P₄S₇	348 551		310	2 19¹⁷	
228	P₄S₁₀	444 746		290		
229	P₂O₃S₃	190 243		300		
230	P₄O₆S₄	348 356		102		
231	PSF₃	120 089		3 8⁷·⁶at.		
232	PSCl₃	169 463		− 35	l. 1.635	193
233	PS₂Cl₃	272 444		< − 17		
234	PSBr₃	302 837		38	2.85¹⁷	
235	P₂SBr₄	573 609		− 5		
236	P₂S₃Br₄	477 907			l. 2 262¹⁷	
237	PSCl₂Br	213 921		− 30	l. 2 12⁰	
238	PSClBr₂	258 379		− 60	l. 2.48⁰	
239	P₂SI₂ .	347 977		75		
240	P₃N₄ .	163 112			2 51¹⁸	
241	NH₄H₂PO₃	83 0782		100		
242	NH₄H₂PO₃	99.0782		ca. 123		
243	NH₄H₂PO₄	115.078	Tet.		1 803	250
244	N₂H₄ H₃PO₄	114 094		36		
245	N₂H₆H₃PO₄	130 094		82		
246	(NH₄)₂HPO₄	118 091			1.619	
247	(N₂H₄)H₄P₂O₆	194.126		152		
248	(NH₄)₂H₃P₂O₆	196 141		170		
249	N₂H₄(H₂PO₃)₂	196 141		82		
250	P₃N₃Cl₆	347 844	R.	114	1 98	
251	P₄N₄Cl₈	463 792		123.5	2.18²⁴	
252	P₅N₅Cl₁₀	579 740		41		
253	P₆N₆Cl₁₂	695 688		91		
254	P₆N₇Cl₉	603.322		237 5		
255	P₇N₇Cl₁₄	811.636		< − 18		
256	PNBr₂	204 864	R.	190		
257	PS₂NH₄ .	145 258			l. 1.78¹⁶·⁵	
258	As₄O₆	197 920		275	3.71	
259	As₂O₃ — Arsenite	197 920	C.		3 865²⁶	
260	As₂O₃ Arsenolite	197 920	C.		3 86	160
261	As₂O₃ — Claudetite	197 920	M.	31₅	4 15	986
262	As₂O₅ .	229 920			4 086	
263	AsH₃	77 9831		−113 5		
264	AsF₃	131 960			l. 2 666⁰₉	
265	AsF₅	169 960		− 80		
266	AsCl₃	181.334		− 18	l. 2 163	191
267	AsCl₅	252 250		ca. − 40		
268	AsBr₃	314 708		32.8	l. 3 540²⁵	
269	AsI₃ .	455 756		146	4.39¹⁵	
270	AsI₅	709 620		76	3 93	
271	As₂S₂ — Realgar	214 050	M.	307 (β) Tr. 267	α 3 506¹⁹ β 3 254¹⁹	1067
272	As₂S₃ — Orpiment	246 115	M.	300 Tr. 170	3 43	1071
273	As₄S₃ . . .	396 035			3 60¹⁹	
274	2AsSCl.As₂S₃	531 081		120		
275	2AsI₃.SI₆.	1705 17		72		
276	NH₄H₂AsO₄	159 014	Tet.		2 311⁹·¹	283
277	(NH₄)₂HAsO₄ . . .	176 045	M.		1.989	
278	SbO₂ — Cervantite	153 770	C.		4 07	174
279	Sb₂O₃ — Valentinite	291.540	R.	656	5 67	1024

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
280	Sb₄O₆—Senarmontite	291 540	C.		5 2	178
281	Sb₄O₆	323 540			3 78	
282	SbH₃	124 793		− 88	1 2 26⁻⁸⁸	
283	SbF₃	178 770	R. ?	292	4 379⁹⁰ ⁹	
284	SbF₅	216 770		7	1. 2.990³³⁴ ⁴	
285	SbF₅.2SbF₃	574 310		390	4 188²¹	
286	SbCl₃	228 144		73 4	3 140¹⁴	
287	SbCl₅	299 060		2 8	1 2 336	58
288	SbOCl	173 228		170 d.		
289	Sb₄O₆Cl₂	637 996	M.	55	5 014	
290	SbF₂Cl₃	206 144		55		
291	SbBr₃	361 518		96 6	4 148²²	
					1. 3 845¹⁸ ⁶	
292	SbI₃	502 566	Trig. M. R.	167	M. 4 768²²	
				Tr 114	Trig. 4 848¹⁸	
				(R. to Trig.)		
				Tr. 125		
				(M. to Trig)		
293	SbI₅	756 430		79		
294	SbF₆I	343 702		ca. 80		
295	(SbF₅)₂I	560 472		ca. 115		
296	Sb₂S₃—Stibnite	339 735	R.	550	4 64	1032
					red 4 120⁰	
					gray 4 284⁰	
					black 4 652⁰	
297	Sb₂(SO₄)₃	531 735			3 625⁴	
298	Sb₂O₃.2Sb₂S₃—Kermesite	971 010	M.		4 6	1073
299	SbF₃S	248 835		230		
300	SbSe	200 970		542		
301	Sb₂Se₃	481 140		611		
302	SbS₃e₄	682 110		605		
303	Sb₄Se₄	883 080		590		
304	Sb₂Te₃	626 040		629		
305	BiO	225 000			7 5	
306	BiO₂	241 000			5 6	
306 1	BiO₂.2H₂O	277 031		d 110	5 6	
307	Bi₂O₃ (I)	466 000	R.	820	8 9	
308	Bi₂O₃ (II)	466 000		Tr. 704	8 20	
309	Bi₂O₃ (III)	466 000	R.	860	8 5	
310	Bi₂O₃.3H₂O—Bismite	520 046	R.	d. 415	4 36	393
311	Bi₂O₄	498 000			5 10	
312	HBiO₃	258 008		d. 120	5.75	
313	BiF₃	266 000			5 32	
314	BiOF	214 000			7 5	
315	BiCl	244 458		320		
316	BiCl₃	315 374		230	4 7	
317	BiCl₄	350 832		225		
318	BiOCl	260 458			7 72	
319	BiBr	288 916		287		
320	BiBr₃	448 748		218	5 7	
321	BiOBr	304 916			8 08	
322	BiI₃	589 796	H.	439	5 7	
323	BiOI	351 932	R.		7.92	
324	BiS	241 065		685	7.7	
325	Bi₂S₃—Bismuthinite	514 195	R.		7 39	
326	BiSe	288 200		625		
327	Bi₂Se₃—Guanajuatite	655 600	R.	710	6 82	
328	Bi₂Te₃	800 500		573	7 7	
329	Bi₂TeO₆.2H₂O—Montanite	677 531			3 79	1002
330	Bi₂Te₂S—Tetradymite	705 065	R.		7.5	
331	Bi(NO₃)₃.5H₂O	485 101	Tri.	d. 30	2.83	
332	Bi(NO₃)₃.6H₂O	503 116			2 76	
333	BiPO₄	304 024	M.		3.23	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	43	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
334	BiAsO$_4$	347 960	M.		7.14	
335	Bi$_2$AsH$_2$O$_8$—Atelestite	831 975	M.		6.4	1009
336	5Bi$_2$O$_3$.2As$_2$O$_3$.9H$_2$O?—Rhagite	2887 98			6 82	
337	CO	28 0000		−207	l. 0.8138$_4^{-195}$	
338	CO$_2$	44 0000		− 56 6$^{5.2at}$	1 53^{-79}	
					l. 1.101^{-37}	
339	C$_3$O$_2$	68 0000		−107	1.114^0	23

Compounds of C with elements of key numbers 2 to 15 in C-Table, p. 176						
340	SiO$_2$—Cristobalite	60 0600	C. Tet. ?	1710	2 32	228
341	SiO$_2$ Lechatelierite	60 0600			2 20	24
342	SiO$_2$—Quartz	60 0600	Trig.	<1470 m.	2 651	267
343	SiO$_2$ Tridymite	60 0600	R.	1670	2 26	463
344	SiO$_2$.H$_2$O Opal	60 0600			2 1 to 2 3	69, 82
345	SiH$_4$	32 0908		−185	l. 0 68^{-185}	
346	Si$_2$H$_6$	62 1662		−132 5	l. 0 69^{-25}	
347	Si$_3$H$_8$	92 2416		−117	l. 0 725^0	
348	Si$_4$H$_{10}$	122 317		− 93 5	l. 0.79^0	
349	Si$_2$H$_6$O	78 1662		−144	l. 0 881^{-30}	
350	SiF$_4$	104 060		− 77		
351	SiHF$_3$	86 0677		ca. −110		
352	SiCl$_4$	169 892		− 70	l. 1 483	192
353	Si$_2$Cl$_6$	268 868		− 1	l. 1 58^0	
354	Si$_3$Cl$_8$	367 844		− 67		
357	Si$_4$Cl$_{10}$	466 820				
358	Si$_5$Cl$_{12}$	565 796				
359	Si$_6$Cl$_{14}$	664 772		170 s. d.		
360	Si$_2$OCl$_6$	284 868		− 33		
361	Si$_3$O$_2$Cl$_8$	459 904				
362	Si$_4$O$_3$Cl$_{10}$	514 820				
363	Si$_5$O$_{10}$Cl$_{12}$	809 976				
364	SiH$_3$Cl	66 5411		−118	l. 1 145$_4^{-113}$	
365	SiH$_2$Cl$_2$	100 991		−122	l. 1 42$_4^{-122}$	
366	SiHCl$_3$	135 442		−134	l. 1 34	
367	SiBr$_4$	347 724		5	2.812$_4^0$	190
368	Si$_2$Br$_6$	535 616		95		
369	Si$_3$Br$_8$	723 508		133		
370	Si$_4$Br$_{10}$	911 400		185 d.		
371	SiH$_3$Br	110 999		− 94	l. 1 533^0	
372	SiH$_2$Br$_2$	189 907		− 77	l. 2 17^0	
373	SiHBr$_3$	268 816		<− 60	l. 2 7^{17}	
374	Si$_2$H$_5$Br	141 075		−100		
375	Si$_2$HBr$_5$	456 708		89		
376	SiCl$_3$Br	214 350		<− 60		
377	SiCl$_2$Br$_2$	258 808		<− 60		
378	SiClBr$_3$	303 266		− 39	l. 2 432	
379	SiI$_4$	535 788		120 5		
380	Si$_2$I$_6$	817 712		250		
381	SiHI$_3$	409 864		8	l. 3 314	
382	SiCl$_3$I	261 366		<− 60		
383	SiCl$_2$I$_2$	352 840		<− 60		
384	SiClI$_3$	444 314		2		
385	SiBr$_3$I	394 740		14		
386	SiBr$_2$I$_2$	441 756		38		
387	SiBrI$_3$	488 772		ca. 53		
388	SiS	60 1250			1 853$_4^{15}$	
389	SiSCl$_2$	131 041		75		
390	SiCl$_3$.SH	167 507				
391	SiSBr$_2$	219 957		93		
392	SiN	42 0680			3 17	
393	Si$_2$N$_3$	98 1440			3 64	
394	Si$_3$N$_4$	140 212			3.44	
395	Si$_3$N$_3$H	99 1517			2 015^{17}	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
37 53 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
396	Si_3H_8N	107 257			1. 0 895[108]	
397	$N_2H_6.H_2SiF_6$	176.122		186 d.		
398	$(NH_4)_2SiF_6$—Cryptohalite	178.138	C.		2.01	08
399	$SiBr_4.6NH_3$	449 911			2 307[17]	
400	$SiO_2.P_2O_5$....	202 108			3.1	
401	$3SiO_2.2Bi_2O_3$—Agricolite	1112 18	M.		6	994
402	$3SiO_2.2Bi_2O_3$—Eulytite	1112 18	C.		6 11	175
403	SiC—Carborundum	40 0600	H.	>2700	3 17	110
404	$Si(CH_3)H_3$..	46 1062		−156.4	1. 0 62[67]	
405	$Si(CH_3)_2H_2$	60 1216		−149 9	1. 0.68[80]	
406	$Si(CH_3)_4$	88 1524			1. 0 645[21.9]	
407	$Si(CH_3)_3C_2H_5$	102.168			1. 0.684	
408	$Si(C_2H_5)_3H$.	116 183			1. 0.751[9]	
409	$Si(CH_3)_2[(C_2H_5)_2]$.	116 183			1. 0.7168	
410	$Si(CH_3)_3C_3H_7$	116.183			1. 0 701[21]	
411	$Si(CH_3)_2[(CH_2)_4]$.	128 183			1. 0 804	439
412	$Si(CH_3)_2(C_2H_5)(C_3H_7)$	130.199			1. 0.732[17] e	
413	$Si(CH_3)_3(C_4H_9)$	130 199			1. 0.721[17]	
414	$Si(CH_3)_3(iso-C_4H_9)$	130 199			1. 0.717[18]	
415	$Si(CH_3)_2(C_3H_7)_2$	144 214			1. 0 741[17] e	
416	$Si(CH_3)_2(C_2H_5)(iso-C_4H_9)$	144 211			1. 0.743	
417	$Si(CH_3)_3(iso-C_5H_{11})$	144 214			1. 0 731[18]	
418	$Si(C_2H_5)_4$	144 214			1. 0.766[19] e	1036
419	$Si(C_2H_5)_3H$	158 229			1. 0 762[18]	
420	$Si(C_2H_5)_3(C_3H_7)$	158 229			1. 0 774[17]	
421	$Si(C_2H_5)_3(C_4H_9)$	172 245			1. 0 779[18]	
422	$Si(C_2H_5)_3(iso-C_4H_9)$	172 245			1. 0 781[18.6]	
423	$Si(C_2H_5)_3(iso-C_5H_{11})$	186 260			1. 0.782[19]	
424	$Si(C_6H_5)_4$	336 214		233		
425	$Si_2(CH_3)_6$	146 259			1. 0 725[23] e	
426	$Si(OCH_3)_4$.	152 152			1. 1 028[22]	9
427	$Si(C_2H_5)_3OH$	132 183			1. 0 871[0]	
428	$Si(C_2H_5)_3OC_2H_5$	160 214			1. 0 840[0]	
429	$Si(OC_3H_7)_4$.	264 276			1. 0.915	1034
430	$Si(C_6H_5)_3OH$.	276 183			1 178	
431	$Si(C_6H_5CH_2)_3OH$.	318 229		106	1 177	
432	$Si_2O(OC_2H_5)_6$	426 443			1. 0 977[22] e	1035
433	$Si(CH_3)H_2Cl$. .	80 5565		−134 1	1. 0 935[80]	
434	$Si(CH_3)HCl_2$	115 007		− 93	1. 0 93[0]	
435	$Si(C_2H_5)Cl_3$.	163 473			1. 1 239[0] e	
436	$Si(C_3H_7)Cl_3$	177 488			1. 1 210[0]	1
437	$Si(C_2H_5)_2Cl_2$	157 053			1. 1 106[15]	
438	$Si(C_4H_9)Cl_3$	191 503			1. 1 162[18] e	
439	$Si(iso-C_4H_9)Cl_3$	191 503			1. 1 154	
440	$Si(C_2H_5)(C_4H_9)Cl_2$	185 084			1. 1 042	
441	$Si(C_6H_5)Cl_3$	211 473			1. 1 326[18] e	
442	$Si(C_6H_5CH_2)Cl_3$	225 488			1. 1 289[19] e	
443	$Si(C_2H_5)(C_6H_5)Cl_2$	205 053			1. 1.159[18]	
444	$Si(SCN)_4$	260 352		143 8		
445	TiO_2—Anatase	79 9000	Tet.		3.84	407
446	TiO_2—Brookite	79 9000	R.		4.17	1028
447	TiO_2—Rutile	79 9000	Tet.	1640 d.	4 26	409
448	Ti_2O_3	143 800	Trig.		4 6	
449	TiF_4 .	123 900			2 798[20] b	
450	$TiCl_4$	189 732		− 30	1. 1 726	59
451	$TiBr_4$	367 564		39		
452	$TiBrCl_3$	234 190				
453	TiI_2 .	301 764			4 30	
454	TiI_4	555 628		150		
455	$TiCl_4.SCl_4$	363 629		64		
456	Ti_2N_2 .	123 816		2930	5 18[18]	
457	TiP	78 9240			3 95[20]	
458	$TiCl_4.PCl_5$	327 130		85 5		

Mg Mn Mo N	Na Nb Nd Ni O	Os P Pb Pd	Pr Pt Ra Rb	Rh Ru S Sa	Sb Sc Se Si Sn	Sr Ta Tb Te Th	Ti Tl Tm U V	W Y Yb Zn Zr
76 42 47 11	82 51 61 45 1	35 12 23 41	60 37 80 84	40 39 8 63	14 56 9 18 22	78 52 66 10 24	19 27 70 49 50	48 57 71 28 21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
459	$TiCl_4.POCl_3$	343 130		110		
460	$TiCl_4.2POCl_3$	496 528		107		
461	TiC	59 9000		3180	4.25	
462	$Ti_{10}C_3N_4$	615 064			5.29	
463	Ti_2Si	123 860			4 02	
464	GeO_2	104 380	R.		4.703	
465	GeH_4	76 4108		-165	l. 1.523⁻¹⁴⁷	
466	Ge_2H_6	150 806		-109	l. 1.98⁻¹⁰⁹	
467	Ge_3H_8	225 202		-105 6	l. 2 20⁻¹⁰⁵	
468	$GeCl_4$	214 212		-49 5	l. 1 874	
469	$GeHCl_3$	179 762				
470	$GeBr_4$	392 044		26 1	l. 3 132	
471	GeI_4	580 108		144	4 322	
472	$Ge(C_2H_5)_4$	188 534		-90	0 991	13

All Zr salts probably contaminated with 1-5% Hf

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
473	ZrO_2 Baddeleyite	123 000	M.	2700	5 49	1012
473 I	ZrO_2 (free from Hf)	123 000			5.73	
474	ZrF_4	167 000			4 43	
475	$ZrCl_4$	232 832				
475 5	$ZrOCl_2\ 8H_2O$	322 039				274.5
476	$ZrOS$	139 065			4 87	
477	$4ZrO_2.3SO_3$	732 195			4 1	
478	$4ZrO_2.3SO_3.15H_2O$	1002 43	M.		2 5	
478 5	$(NH_4)_3ZrF_7$	278 034	C.			70 2
479	ZrP_2	153 048			4 77	
480	$2ZrCl_4.PCl_5$	673 978		164 5		
481	ZrC_2	115 000				
482	$ZrSi_2$	147 120			4 88²²	
483	$ZrO_2.SiO_2$ Zircon	183 060	Tet.	2500	4 5	382, 387
484	SnO	134 700	C.		6 95	
485	SnO_2 Cassiterite	150 700	Tet. H. R.		7 0	391
486	SnF_4	194 700			4 78	
487	$SnCl_2$	189 616		246 8		
488	$SnCl_4$	260 532		-30 2	l. 2 226	
489	$H_2SnCl_6.6H_2O$	441 556			1 925²⁷	
490	$SnBr_2$	278 532		215 5	5 12¹⁷	
491	$SnBr_4$	438 364		31 0	1 3 34³⁶	
492	$SnCl_2Br$	304 990		-31	l. 2 5¹³	
493	$SnCl_2Br_2$	349 448		-20	l. 2 8¹³	
494	$SnClBr_3$	393 906		1	l. 3 1¹³	
495	SnI_2	372 564		320		
496	SnI_4	626 428		143 5	4 46	
497	$SnCl_2I_2$	443 480			1 3 29	
498	$SnBr_2I_2$	532 396		50 d.	3 6	
499	SnS	150 765		880	5 080⁰	
500	SnS_2	182 830			4 5	
501	$SnSe$	197 900		861	6 18⁰	
502	$SnSe_2$	277 100			5 0	
503	$SnTe$	246 200		780	6 48	
504	$SnCl_4.2NOCl$	391 464		180	2 6	
505	$2NH_4Cl\ SnCl_4$	367 526			2 4	
506	$(NH_4)_2SnBr_6$	634 274			3.50	
507	Sn_4P_3	567 872			5 18	
508	$SnCl_4\ POCl_3$	413 930		58		
509	Sn_3As_2	462 280			6 56	
510	SnC_2O_4	206 700			3.56¹⁸	
512	$Sn(C_2H_5)_2$	176 777			l. 1 654	
513	$Sn(CH_3)_4$	178 792			l. 1 314⁰	
514	$Sn(CH_3)_2(C_2H_5)_2$	206 823			l. 1 232	50
515	$Sn(C_2H_5)_4$	234 854			l. 1 187²³	
516	$Sn(C_3H_5)_3$	272 777		225 7		44

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
517	Sn(C₂H₅)₄	426 854		226		
518	Sn₂(C₂H₅)₆	411 631			1. 1 412⁰	
519	Sn(C₂H₃O₂)₂	236 746		182		
520	SnCl(C₂H₅)₃	241 274			1. 1 428⁰	
521	SnBr(C₂H₅)₃	285 732			1. 1 630	
522	SnI(CH₃)₃	290 701			1. 2 100¹¹	
523	SnI(C₂H₅)₃.	332 748			1. 1 833¹¹	
524	PbO—Litharge	223 200	Tet.	88₈	9 5₃	423
525	PbO—Massicotite	223 200	R.		8 0	1068
526	PbO₂—Plattnerite	239 200	Tet.		9 375	417
527	Pb₃O₄—Minium	685 600			9.1	
528	PbF₂ . . .	245 200		855	8 2₄	
529	PbCl₂—Cotunnite	278 116	R.	50₁	5 8₅	1016
530	PbCl₄ .	349 032		– 15	1. 3 18⁰₆	
531	Pb(ClO₂)₂.	342 116		exp. 126		
532	Pb(ClO₃)₂	374 116			3 89	
533	Pb(ClO₃)₂.H₂O	392 131	M.	d. 110		
534	Pb(ClO₄)₂.3H₂O	460 162	R.	d. 100	2 6	
535	PbO.PbCl₂—Matlockite	501 316	Tet.	524 d.	7 21	1008
536	2PbO.PbCl₂—Mendipite	724 516	R.	69₃	7 0₈	1022
537	PbO.2PbCl₂—Penfieldite	779 432	H			398
538	6PbO.PbCl₂—Lorettoite.	1617 32	Tet.		7 6	418
539	PbCl₂.PbO.H₂O—Laurionite	519 331	R	d. 142	6 2₄	1006
540	PbCl₂.PbO.H₂O—Paralaurionite	519 331	M	d. 150	6.0₅	
541	2PbCl₂.PbO.H₂O—Fiedlerite .	797 447	M	d. 150	5 8₈	1005
542	PbFCl	261 658	Tet.	60₁		
543	PbBr₂	367 032	R.	37₃	6 6₆	
544	Pb(BrO₃)₂.H₂O	481 047	M.	d. 180	5.5₃	
545	PbO.PbBr₂.H₂O	608 248	R.		6.7₂	
546	PbClBr.	322 574			5 7₄	
547	PbI ..	334 132		d. 300		
548	PbI₂	461 064	H.	40₂	6 16	
549	Pb(IO₃)₂	557 064		d. 300		
550	PbO.PbI₂...	684 264		300 d.		
551	PbI₂.PbO.H₂O	702 280	R.	d. <100	6 8₃	
552	PbS—Galena	239 265	C.	111₄	7 5	189
553	PbSO₄—Anglesite .	303 265	R. M.	117₀ Tr. 86₄	6 2	981
554	PbS₂O₃	319 330			5 18	
556	Pb₂S₂O₆.4H₂O	439 392			3 2₂	311
557	Pb₂O(SO₄)—Lanarkite	526 465	M	977	6 9₂	995
558	PbSe—Clausthalite	286 400	C.	106₅	8 10	
559	PbSeO₄	350 400	R	d.	6 3₇	
560	PbTe—Altaite	334 700	C.	917	8 16	
561	PbN₆	291 248		exp. 350		
562	Pb(NO₃)₂	331 216	C M.	47₀	4 5₃	162
563	2PbO.N₂O₅.1.5H₂O	581 439	M.	d. 100		
564	4PbO.N₂O₅.N₂O₅ 2H₂O	1112 86	R.	d. 100		
565	2PbO.N₂O₅.H₂O	572 431	R	d. 180	5 9₃	
566	(NH₄)₂PbCl₆	456 026	C	d. 120		
567	Pb(PO₃)₂.	365 248		80₀		
568	Pb₂P₂O₇. .	588 448	R.	821	5.8	
569	3PbO.P₂O₅ .	811 648		1014 Tr. 782		389
570	4PbO.P₂O₅	1034 85		982		
571	5PbO.2P₂O₅	1400 10		946		
572	8PbO P₂O₅	1927 65		860		
573	PbCl₂.3Pb₃(PO₄)₂—Pyromorphite .	2713 06	H.	115₆	6 8	1000
574	Pb(AsO₂)₂	421 120			5 8₅	
575	Pb(AsO₃)₂	453 120	H.		6 4₂	
576	Pb₂As₂O₇	676 320		802	6 85	998
577	Pb₃(AsO₄)₂	899 520		1042	7 30	
578	Pb₃(AsO₄)₂.0.5H₂O......	908 528			7.00	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
579	5PbO.Pb₃(AsO₄)₂	2015 52		862		
580	5PbO Pb₃(AsO₄)₂.0.5H₂O	2024 53	R.		8 04	
581	10PbO.3As₂O₄.3H₂O	2975 81	H.		6 86	179
582	PbHAsO₄	347 168	M.	d. >200	5 79	1054
583	Pb(H₂AsO₄)₂	489 151	Tri.	d. 140	4 46	963
584	Pb₄(PbOH)₂(AsO₄)₄	2040 26			7 08	
585	2Pb₃(AsO₄)₂.2Pb(OH)₂.10H₂O	2461 62			7.1	
586	65PbO 21As₂O₄ 12H₂O	19552 5		d. >200	7.1o	
587	9PbO.3As₂O₄.PbCl₂ Mimetite	2976 68	H.	1140 Tr. 395	7 13	399
588	4PbO As₂O₄.2PbCl₂ Ecdemite	1646 15	R.		7 0	
589	3PbCl₂.3PbO.As₂O₄ Georgiadesite	1733 87	R.	d.	7 1	
590	5PbO.2PbCl₂ As₂O₄	1870 15	Tet.		7 14	
591	PbS As₂S₃ Sartorite	485 380	R.	<700 d.	4 6	
592	2PbS.As₂S₃ Dufrenoysite	724 645	R.		5 50	
593	3PbS 2As₂S₃ Rathite	1210 03	R.		5 41	
594	4PbS.As₂S₃ Jordanite	1203 18	M.		6 1o	
595	4PbS 3As₂S₃ Baumhauerite	1695 41	M.		5 33	
596	7PbS 2As₂S₄ Lengenbachite	2167 09	Tri.		5 8	
597	10PbS 3As₂S₃ Guitermanite	3131 00			5 94	
598	3PbO.Sb₂O₄ Monimolite	1236 68	C.		6 58	
599	PbO.PbCl₂.Sb₂O₃ Nadorite	792 856	R.		7 02	1059
600	PbS.Sb₂S₃ Zinkenite	579 000	R.		5 3	
601	2PbS.Sb₂S₃ Plumosite	818 265	M.		5 62	
602	3PbS.Sb₂S₃ Dürfeldtite	1057 53			5 9	
603	3PbS.2Sb₂S₃ Domingite	1397 27			5 62	
604	4PbS Sb₂S₃ Meneghinite	1296 80	R.		6 30	
605	5PbS.Sb₂S₃ Geocronite	1536 06	R.		6 4	
606	5PbS.2Sb₂S₃ Boulangerite	1875 80	R.		6 18	
607	5PbS 2Sb₂S₃ Mullanite	1875 80	R.		6.3	
608	5PbS.4Sb₂S₃ Plagionite	2555 27	M.		5 47	
609	6PbS.Sb₂S₃ Kilbrickenite	1775 33			6 5	
610	PbS.Bi₂S₃ Galenobismutite	753 460			6 9	
611	2PbS.Bi₂S₃ Cosalite, Bjelkite	992 725	R.		6 6	
612	2PbS.3Bi₂S₃ Chiviatite	2021 12			6 92	
613	3PbS.Bi₂S₃ Lillianite	1231 99	R.		7 0	
614	4PbS 5Bi₂S₃ Rezbanyite	3528 04			6 2	
615	6PbS Bi₂S₃ Beegerite	1949 79	C.		7 27	
616	2BiSCl PbS.Bi₂S₃	1306 51			6 42	
617	PbCO₃ Cerussite	267 200	R.	500 d. d. 315	6 6	1001
618	PbC₂O₄	295 200			5 28	
619	Pb(CH₃)₄	267 292		— 27 5	l. 1 995	42
621	Pb(CH₃)₃(C₂H₅)	281 308			l. 1 889	43
622	Pb(CH₃)₂(C₂H₅)₂	295 323			l. 1 790	48
623	Pb(CH₃)₃(C₃H₇)	295 323			l. 1 760$_4^{22}$	37
624	Pb(C₂H₅)₃(CH₃)	309 339			l. 1 712$_4^{21}$	46
625	Pb(CH₃)₃(C₄H₉)	309 339			l. 1 674$_4^{24}$	34
626	Pb(CH₃)₃(iso-C₄H₉)	309 339			l. 1 668$_4^{23.5}$	32
627	Pb(CH₃)₂(C₃H₇)₂	323 354			l. 1 623$_4^{24.4}$	35
628	Pb(C₂H₅)₄	323 354			l. 1 659$_4^{18}$	51
629	Pb(CH₃)₃(iso-C₅H₁₁)	323 354			l. 1 524$_4^{21.4}$	30
630	Pb(C₂H₅)₃(C₃H₇)	337 369			l. 1 595$_4^{22.5}$	49
631	Pb(C₂H₅)₂(C₃H₇)₂	351 385			l. 1 529$_4^{25.4}$	41
632	Pb(C₂H₅)₂(iso-C₄H₉)₂	351 385			l. 1 504$_4^{20 6}$	33
633	Pb(C₂H₅)₃(iso-C₄H₉)	351 385			l. 1 530$_4^{22.6}$	40
634	Pb(CH₃)₂(iso-C₅H₁₁)₂	379 416			l. 1 430	31
635	Pb(C₂H₅)₂(iso-C₄H₉)₂	379 416			l. 1 456$_4^{22}$	36
636	Pb(C₂H₅)₃(C₅H₁₁)	365 400			l. 1 482	38
637	Pb(C₂H₅)₃(iso-C₅H₁₁)	365 400			l. 1 506$_4^{21.8}$	39
638	Pb(C₆H₅)₄	515 354		227 7		
639	Pb(CHO₂)₂	297 215	R.	d. 190	4.63	973
640	Pb(dl-C₄H₄O₆)	355 231			2 530¹⁹	
641	Pb(d-C₄H₄O₆)	355 231	R.		3 871¹⁹	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ge Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
22 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
642	Pb(C₂H₃O₂)₂ ..	325 246		280	3 251	
643	Pb(C₂H₃O₂)₂.3H₂O .	379 292	M.	75	2 55	710
644	Pb(C₂H₃O₂)₂.10H₂O	505 400	R.	22	1 689	
645	Pb(C₃H₅O₂)₄	459 292		180	2 23¹⁸	
646	Pb(C₃H₅O₂)₄	515 354		132		
647	Pb(C₆H₁₁O₂)₂.	437 369		74		
648	Pb(C₇H₁₃O₂)₂	465 400		91 5		
649	Pb(C₈H₁₅O₂)₂	493 431		84 5		
650	Pb(C₉H₁₇O₂)₂	521 416		95		
651	Pb(C₁₀H₁₉O₂)₂	549 493		100		
652	Pb(C₁₂H₂₃O₂)₂	605 554		104		
653	Pb(C₁₄H₂₇O₂)₂	661 616		107		
654	Pb(C₁₆H₃₁O₂)₂	717 677		112		
655	Pb(C₁₈H₃₅O₂)₂	769 708		ca 80		
656	Pb(C₁₈H₃₃O₂)₂	773 739		125		
657	3PbO.2CO₂.H₂O —Hydrocerusite	775 615	H.	d. 400	6 14	395
658	PbCl₂.PbCO₃—Phosgenite	545 316	Tet	d.	6 13	396
659	PbBr₂.PbCO₃	634 232	Tet.	d.	6 55	
660	Pb(OH)₂.PbSO₄.2PbCO₃—Leadhillite	1078 88	M.		6 5	996
661	Pb(OH)₂.PbSO₄.2PbCO₃—Maxite	1078 88	R.		6 9	
662	Pb(SCN)₂	323 346	M.		3 82	
663	PbSiO₃—Alamosite	283 260	M.	766	6 49	992
664	2PbO.SiO₂ ?	506 460		746		
665	3PbO.SiO₂ ?	729.660		717		
666	3PbO.2SiO₂ —Barysilite	789 720	Trig.		6 72	304
667	SnPbS₂—Teallite	390 030	R		6 4	
668	ThO₂—Thorianite	264 150	C.	>2800	9 69	182
669	ThCl₄	373 982	R	820	4 59	
670	ThBr₄	551 814			5 67	
671	ThS₂	296 280		d.	6 8	
672	ThOS	280 215		d.	6 14	
673	Th(SO₄)₂.9H₂O	602 419	M.	d.	2 77	
674	Th(PO₃)₄	548 246	R.		4 08	
675	ThC₂.	256 150			8 96	
676	ThSi₂	288 270			7 96¹⁶	
677	ThO₂.SiO₂—Thorite	324 210	Tet.	175	5 3	
678	GaCl₂	140 636		175		
679	GaCl₃	176 094		75 5	1 2 36²⁰	
680	(NH₄)₂Ga₂(SO₄)₂.24H₂O	992 147			1 77	89
681	In₂O₃	277 600	Trig.		7 179	
682	InCl₃	221 174			4 0	
683	In(ClO₄)₃.8H₂O	557 297		80		
684	InI	241 732		351		
685	InI₂ .	368 664		212		
686	InI₃	495 596		199		
687	In₂(SO₄)₃	517 795			3 438	
688	(NH₄)₂InCl₅.H₂O	346 483	R.		2 281	
689	(NH₄)₂InBr₅.H₂O	568 473	R.		3 167	
690	(NH₄)₃In(SO₄)₃.12H₂O	541 154			2 011	88
691	Tl₂O	424 800		300		
692	Tl₂O₃	456 800		759	brown 9 65²¹	
					black 10 19²²	
693	TlOH	221 408				
694	Tl(OH)₃	255 423		>340		
695	TlF	223 400				
696	TlCl	239 858		430	7.00	
697	TlCl₃.4H₂O	382 836		37		
698	TlClO₃	287 858			5 047₉	
699	TlClO₄	303 858		501	4 89	
700	TlBr	284 316		460	7 557⁷ ⁸	
701	TlBr₃.4H₂O.	516 210		40		
702	TlBr₂Cl.4H₂O...	471 752		40 d.		
703	TlI	331 332		440	7.09¹⁴ ⁷	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
704	Tl₂S ...	440 865		44₈	8.0	
705	Tl₄S₄ ...	569 125		12₅		
706	Tl₆S₇	185.966		12₇		
707	Tl₂SO₄	504 865	R.	63₂		975
708	Tl₂S₂O₆	568 930	M.		5.57	
709	TlHSO₄	301 473		12o d.		
710	Tl₂Se	488 000		340		
711	Tl₂Se.Tl₂Se₃	1134 40		33₈		
712	Tl₂SeO₄	552 000	R.		6 87₅	991
713	Tl₂Te	536 300		412		
714	Tl₂TeO₄	600 300			5 712	
715	TlN₃	246 424		33₄		
716	TlNO₃	266 408	γ R. β Trig. α C.	206 Tr. 75 (γ to β) Tr. 145 (β to α)	$5\ 556^{21}_{4}\ ^{4}$	1053
717	(NH₄)₂TlCl₅.2H₂O	507 295			2 38⁹	
718	Tl₃PO₄	708 224			6 8⁹	
719	Tl₄P₂O₇	991 648	M.	>12o	6 78₆	
720	TlH₂PO₂	269 439	M.	19o		
721	TlH₂PO₄	301 439	M.	19o	4 72₃	
722	Tl₂H₂P₂O₇	584 863		27o		
723	Tl₃S.As₂S₃ - Lorandite	686 980	M.		5 53	1072
724	TlSbAs₂S₄ - Vrbaite	636 415	R.		5 30	
725	Tl₂CO₃ ...	468 800			7 11	
726	Tl(C₂H₃O₂)	263 423		110	3 68 1 3 9	
727	Tl(CHO₂)₃	339 423	M.	95		
728	Tl(C₃H₅O₂)	277 439		140	2 8	
729	Tl(d-C₄H₅O₆)	353 439	R.		3 496	
730	Tl(dl-C₄H₅O₆)	353 439	Tri.		3 494	
731	Tl(meso-C₄H₅O₆) 0.5H₂O	362 446	Tri.		3 518	
732	TlH(C₄H₄O₆)₂	323 454		64		
733	Tl₂(d-C₄H₄O₆)	556 831	Trig.		4 80	558
734	Tl₂(meso-C₄H₄O₆)	556 831	Tri.		5 110	899
735	Tl₂(dl-C₄H₄O₆)	556 831	M.	165	4 66	957
736	Tl₂(d-C₄H₄O₆).0 5H₂O ...	565 838	M.		4 60	
738	TlH(Cl₂CCO₂)₂	530 156	Tet.		$2\ 822^{18}_{4}$	
739	TlH(CBr₂CO₂)₂	796 904	M.		$3\ 923^{18}$	
740	TlOC₆H₂(NO₂)₃ - Picrate	432 440	M. (red) Tri. (yellow)		$3\ 164^{17}_{4}$ $2\ 993^{25}_{4}$	
741	Tl(SbO)(d-C₄H₄O₆).H₂O	508 216	R.		3 990	
742	TlCl 2PbCl₂	796 090	C.	435		
743	TlGa(SO₄)₂.12H₂O	682 435			2 477	110
744	ZnO - Zincite	81 3800	H.	>1800	5 60₆	392
745	ZnO	81 3800			5 4₇	
746	Zn(OH)₂	99 3954	R.	d. 125	3 053	
747	ZnF₂	103 380	M. Tri. ?	87₂	$4\ 84^{18}_{4}$	
748	ZnF₂.4H₂O ...	175 442	R.	Tr. 100	$2\ 535^{12}_{4}$	
749	ZnCl₂	136 296	C.	365	2.91^{25}_{4}	
750	Zn(ClO₃)₂.4H₂O	304 357			2 15	
751	Zn(ClO₄)₂.6H₂O	372 388			2 15	
752	ZnBr₂	225 212	R.	394	4 219	
753	ZnI₂ ...	319 244	C.	446	$4\ 666^{14\ 2}_{4}$	
754	Zn(IO₃)₂	415 244		d.	4 98	
755	ZnS(α) - Würzite	97 4450	H.	1850¹⁵ᵒᵃᵗ	4 08₇	404
756	ZnS(β) - Sphalerite	97 4450	C.	Tr. 102o	$4\ 102^{25}_{4}$	187
757	ZnSO₄ - Zinkosite	161 445	R.	d. 74o	$3\ 74^{18}_{4}$	860
758	ZnSO₄.H₂O	179 460		d. 238	$3\ 28^{16}$	
759	ZnSO₄.6H₂O	269 537	M.	Tr. 70 0	$2\ 072^{15}_{4}$	
760	ZnSO₄.7H₂O - Goslarite	287 553	R.	Tr. 39 0	1 97	490
761	ZnS₂O₆.6H₂O	333 602	Tri.		1 915	
762	ZnSe	144 580	H.		$5\ 42^{15}$	188.1

```
Ag Al As Au   B Ba Be Bi Br   C Ca Cb Cd Ce   Cl Co Cr Cs Cu   Dy Er Eu F Fe   Ga Gd Ge Gl H   Hf Hg Ho I In   Ir K La Li Lu
51 53 13 33   54 79 75 15 5    16 77 51 29 59    4 44 46 85 31   67 69 64 3 43   25 65 20 75 2   73 30 68 6 26   36 83 55 81 72
```

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
763	$ZnSeO_4.5H_2O$	298 657	Tri.	d. >50	2 591	
764	$ZnSeO_4.6H_2O$	316 672	Tet.	d.	2 325	252
765	$ZnTe$	192 880	C.	1238 s	5 54[18]	188.2
766	$Zn(NO_3)_2$	189 396		44 07		
767	$Zn(NO_3)_2.3H_2O$	243 442		45 5		
768	$Zn(NO_3)_2.6H_2O$	297 488	Tet.	36 4	2 065[14]	
769	$ZnCl_2.NH_3$	153 377				
770	$ZnCl_2.2NH_3$	170 358	R.	210 8		
771	$ZnCl_2.2NH_4Cl$.	243 290	R.		1 82	
772	$Zn(ClO_3)_2.4NH_3$ . .	300 420		exp. 205	1 84	
773	$ZnBr_2.2NH_4Br$	421 122			2 625	
774	$Zn(BrO_3)_2.4NH_3$	389 336		exp. 169	2 27	
775	$Zn(IO_3)_2.4NH_3$.	483 368		exp. 215	2 82	
776	$ZnSO_4.(NH_4)_2SO_4$	293 588			2 2s	
777	$ZnSO_4.(NH_4)_2SO_4.6H_2O$	401 680	M.	d.	1 931	516
778	$Zn(SeO_4).(NH_4)_2SO_4.6H_2O$	495 950	M.		2 20	620
779	Zn_3P_2	258 188	C.	>420	1 55[18]	
780	$Zn_3(PO_4)_2$.	386 188	R.	900	3 998[16]	
781	$Zn_3(PO_4)_2.4H_2O$—α Hopeite	458 250	R.	Tr. >105	3 04	734
782	$Zn_3(PO_4)_2.4H_2O$—β Hopeite	458 250	R.	Tr. >140	3 03	720
783	$Zn_3(PO_4)_2.4H_2O$—Parahopeite	458 250	Tri.	Tr. >163		793
784	$ZnH_4(PO_4)_2.2H_2O$	295 490	Tri.	100 d.		
785	$Zn_2(OH)PO_4$—Tarbuttite	242 792	Tri.		4 1s	898
786	$Zn_3(PO_4)_2 Zn(OH)_2.3H_2O$—Spencerite	539 630	M.	d. 100	3 14	755
787	$Zn_3P_2S_8$	385 198	H.		2 2	
788	$ZnAs_2$	215 300		771		
789	Zn_3As_2	346 060	C.	1015		
790	$Zn_2As_2O_7$.	392 680			4 701[11]	
791	$Zn_3As_2O_8$	474 060	R.		4 913[14]	
792	$Zn_3(AsO_4)_2.8H_2O$—Koettigite	618 183	M.	d. 100	3 309[16]	881
793	$4ZnO.As_2O_5.H_2O$—Adamite	573 455	R.	d. >100	4 345	918
794	$ZnCO_3$—Smithsonite	125 380	Trig.	d. 300	4 44	369
795	ZnC_2O_4	153 380			2 58[17.8]	
796	$ZnC_2O_4.2H_2O$	189 411		d. 100	2 562	
797	$Zn(CH_3)_2$	95 4262		— 40	1 1 386[10]	
798	$Zn(C_2H_5)_2$	123 457		— 28	1. 1 182[16]	
799	$Zn(C_3H_7)_2$	151 488				
800	$Zn(iso-C_5H_{11})_2$	207 549			1 1 022[6]	
801	$Zn(CHO_2)_2$	155 395			2 3s	
802	$Zn(CHO_2)_2 2H_2O$	191 426	M.		2 205	
803	$Zn(C_2H_3O_2)_2$	183 126		142	1 840	
804	$Zn(C_2H_3O_2)_2.2H_2O$	219 457	M.	237	1 735	518
805	$Zn(l-C_4H_5O_5)_2.2H_2O$—l-Malate	367 488	Tet.		1 701[20]	
806	$Zn(C_3H_7CO_2)_2$.	239 488	M.			535
807	$5ZnO 2CO_2.3H_2O$—Hydrozincite	548 947	M ?		3 7	920
808	$Zn(CH_3SO_3)_2.3H_2O$—Ethane disulfonate	307 587	Tri.		2 043	
809	$ZnC_{10}H_6O_6S_2.6H_2O$—1, 5-Naphthalene disulfonate	459 649	M.		1 793	791
810	$Zn(CN)_2$	117 396	R.	d. 800		
811	$ZnO.SiO_2$	141 440		1437	3 52 1 3 86 gls	
812	$2ZnO.SiO_2$—Willemite	222 820	Trig.	1509	3 9	341
813	$2ZnO.SiO_2.H_2O$—Calamine	240 835	R.		3 45	780
814	$ZnSiF_6.6H_2O$ . .	315 532	H.		2 104	209
815	$ZnSiS$	125 505			3 41	
816	$ZnO.TiO_2$	161 280			3 17	
817	$ZnO.3TiO_2$	321 080			4 92[16]	
818	$3ZnO 2TiO_2$	403 940			3 83	
819	$4ZnO.5TiO_2$ . . .	725 020			3.68[16]	
820	$Tl_2Zn(SO_4)_2.6H_2O$.	774 402	M.	d. 120	3 72o	771
821	CdO .	128 410	C.		8 1s	
822	Cd_2O .	240 820		d.	8 192[18]	
823	$Cd(OH)_2$	146 425	Trig.	d. 300	4 79[15]	

Mg Mn Mo N	Na Nb Nd Ni O	Os P Pb Pd	Pr Pt Ra Rb	Rh Ru S Sa	Sb Sc Se Si Sn	Sr Ta Tb Te Th	Ti Tl Tm U V	W Y Yb Zn Zr
76 42 47 11	82 51 61 45 1	35 12 23 41	60 37 80 84	40 39 8 63	14 56 9 18 22	78 52 66 10 24	19 27 70 49 50	48 57 71 28 21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
824	CdF$_2$	150 410	C.	1100	6.64	
825	CdCl$_2$	183 326	C.	568	4 047$_4^{16}$	
826	CdCl$_2$.2.5H$_2$O	228 364	M.	Tr. 34	3 327	829
827	Cd(ClO$_3$)$_2$.2H$_2$O	315 357		80		
828	CdCl$_2$.CdO H$_2$O	329 751	H.	d. 280	4 56$_4^{16}$	
829	CdBr$_2$	272 242		583	5.192$_4^{24}$	
830	Cd(BrO$_3$)$_2$.2H$_2$O	404 273	R.		3.758	
831	CdO CdBr$_2$ H$_2$O	418 667			4.87$_4^{16}$	
832	CdI$_2$(α)	366 274	H.	388	5 670$_4^{20}$	
832 1	CdI$_2$(β)	366 274			5 305$_4^{20}$	
833	Cd(IO$_3$)$_2$	462 274			6 48	
834	Cd(IO$_3$)$_2$.H$_2$O	480 289		Tr. 160	6 43	
835	CdS –Greenockite	144 475	H.	1750^{100} at	4 820	406
836	CdSO$_4$	208 475	R.	1000	4 691$_4^{24}$	
837	CdSO$_4$ H$_2$O	226 490	M.	Tr. 108	3 786	
838	CdSO$_4$.2 66H$_2$O	256 583	M.	Tr. 41 5	3 090	688
839	CdSO$_4$.7H$_2$O	334 583	M.	Tr. 4	2 48	
840	CdS$_2$O$_6$.6H$_2$O	380 632	Tri.	d.	2 272	
841	CdSe	191 610	H.		5 81$_4^{16}$	
842	CdSeO$_4$.2H$_2$O	291 641	R.	d. 100	3 632	
843	CdTe	239 910	C.	1041	6 20^{16}	
844	Cd(NO$_3$)$_2$	236 426		350		
845	Cd(NO$_3$)$_2$.4H$_2$O	308 488		59 4	2 455$_4^{17}$	
846	CdCl$_2$ NH$_4$Cl	236 823	R.		2 93	
847	CdCl$_2$.4NH$_4$Cl	397 313	Trig.	Tr – 20	2 01	296
848	CdCl$_2$.2NH$_4$OH	249 388		d. 130	2 72$_4^{16}$	
849	Cd(ClO$_3$)$_2$.6NH$_3$	381 513		exp. 184	1 78	
850	Cd(BrO$_3$)$_2$.4NH$_3$	436 366		exp. 192	2 53	
852	Cd(IO$_3$)$_2$.4NH$_3$	530 398		exp.	3 23	
853	CdSO$_4$.(NH$_4$)$_2$SO$_4$	340 618		d.	3 11	
854	CdSO$_4$.(NH$_4$)$_2$SO$_4$.6H$_2$O	448 710	M.	d. 100	2 067	500
855	CdSeO$_4$.(NH$_4$)$_2$SeO$_4$.2H$_2$O	470 918	Tri.		3 376	
856	CdSeO$_4$.(NH$_4$)$_2$SeO$_4$.6H$_2$O	542 980	M.	d. 20	2 307	
857	Cd$_2$P$_2$O$_7$.2H$_2$O	434 899		900	4 965$_4^{16}$	
858	Cd$_3$(PO$_4$)$_2$	527 278		1500		
859	5CdO.2P$_2$O$_5$.5H$_2$O	1016 22	M.	d. 550	4 13$_4^{16}$	
860	Cd(H$_2$PO$_4$)$_2$.2H$_2$O	342 520	Tri.	d. 100	2 742$_4^{15}$	
861	Cd$_3$(PO$_4$)$_2$.2CdHPO$_4$.4H$_2$O	1016 22	M.	d. 600	4 06	
862	3Cd$_3$(PO$_4$)$_2$.CdCl$_2$	1765 16			5 46$_4^{16}$	
863	Cd$_3$As$_2$	487 150	C.		6 211	
864	Cd$_2$As$_2$O$_7$	486 740			5 974	
865	CdHAsO$_4$ H$_2$O	270 393		d. >120	4 164$_4^{16}$	
866	Cd(H$_2$AsO$_4$)$_2$.2H$_2$O	430 392	Tri.	d. 75	3 241$_4^{16}$	
867	CdSb...	234 180		455		
868	CdCO$_3$	172 410	Trig.	d. <500	4 258	
869	CdC$_2$O$_4$	200 410		d. 340	3 32^{18}	
870	Cd(CH$_3$)$_2$	142 456				
871	Cd(CHO$_2$)$_2$.2H$_2$O	238 456	M.		2 44	
872	Cd(C$_2$H$_3$O$_2$)	171 433		256	2 341	
873	Cd(C$_2$H$_3$O$_2$) 2H$_2$O	207 464	M.		2 01	
874	Cd(CH$_3$SO$_3$)$_2$.2H$_2$O	336 602	Tri.		2 570	
875	Cd(CN)$_2$	164 426		d. >200		
876	CdO SiO$_2$	188 470		1242	4.93	
877	2CdO SiO$_2$	316 880		1243		
878	HgO–Montroydite	216 610	R.	d. 100	11 14	1027
879	Hg$_2$O	417 220		d. 100	9.8	
880	HgF	219 610	C. ?	570	8 73	
881	HgF$_2$	238 610	C.	645 d.	8 95	
882	HgCl –Calomel	236 068	Tet.	302	7.150	390
883	HgCl$_2$—Corrosive sublimate	271 526	R.	277	5 44 1. 4.44^{280}	
884	HgClO$_3$....	284 068	R.	d. 250		

Ag Al As Au
21 55 13 33
B Ba Be Bi Br
54 79 75 15 5
C Ca Cb Cd Ce
16 77 51 29 59
Cl Co Cr Cs Cu
4 44 46 85 31
Dy Er Eu F Fe
67 69 64 3 43
Ga Gd Ge Gl H
25 65 20 75 2
Hf Hg Ho I In
73 30 68 6 26
Ir K La Li Lu
36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{15}	Ref. ind. finding No.
885	HgClO₄.6H₂O	408 160		d. 150	4 28	
886	Hg(ClO₄)₂.7H₂O	525 634		34 d.	2 78	
887	Hg₂ClO—Terlinguaite	452 678	M.	d.	8 725	1070
888	HgCl₂.2HgO	704 746	H	d.	red 8 3	
			M.	d.	black 8 5	
889	HgO.2HgCl₂	759 662			6 42	
890	Hg₂O.2HgCl—Eglestonite	889 356	C.		8 33	195
891	HgCl₂.3HgO—Kleinite	921 356	H.		7 93	
892	HgCl₂.4HgO	1137 97	H.	d. 260	9 10	
893	HgBr	280 526			7 307	
894	HgBr₂	360 412	R.	237	6 05s	
					1. 5 12²⁴⁰	
895	HgBr₂.4HgO	1226 88	R.	d. 230	8 73	
896	HgI	327 512	Tet.	290 d	7 70	
897	HgI₂ (red)	454 174	Tet	Tr 127	6 283	
898	HgI₂ (yellow)	454 174	R	259	6 271	
					1. 5 24²⁴⁰	
899	Hg₂Cl₂I₂	726 000	R.	153		
900	HgS—Metacinnabarite	232 675	C		7 5o	
901	HgS (α)—Cinnabarite	232 675	H.		8 1o	411
902	HgS (β)	232 675	H		7 7₃	
903	HgSO₄	296 675	R	d	6 47	
904	Hg₂SO₄	497 285	M.	d.	7 56	
904 1	Hg₂SO₄Cl₂	568 201		270		
904 2	Hg₂SO₄Br₄	816 949		d 125		
904 3	Hg₂SO₄I₂	751 149		248		
905	HgSO₄.3HgS	994 700		d. 120	6 41₆	
906	Hg₂SeO₃	528 120		180 d		
907	HgNO₂	246 618		d. 140	5 925	
908	HgNO₃.H₂O	280 633	M.	70	4 785³˙⁷	
909	Hg(NO₃)₂.0.5H₂O	333 634		79	4 3	
910	Hg₂(NO)₂...	461 236		d. 100	7 33	
911	(HgOH)₂.NH₄OH.	468 267			4 083	
912	HgCl₂.N₂H₄.HCl	340 039		157		
913	HgCl₂.2NH₄Cl H₂O..	396 535	R.		2 84	
914	HgCl₂.12NH₃	475 899		— 9 P.		
914 1	Hg₂(NO₃)₂Cl₄	667 068		d. 100		
915	HgBr₂ 2N₂H₄ HBr.H₂O	603 475		73		
916	NHg₂Br.3NH₄Br	789 008	R.	180 d.		
916 1	Hg₂(NO₃)₂I₄	1032 96		250		
917	HgS 2Sb₂S₃—Livingstonite	942 145	R.		4 81	1029
918	Hg(CH₃)₂	230 656			1. 3 069	53
919	Hg(C₂H₅)₂	258 687			1. 2 444	54
920	Hg(C₃H₇)₂	286 718			1. 2 124¹⁸	
921	Hg(iso-C₄H₉)₂	314 748			1 1 835¹⁵	
922	Hg(C₆H₅)₂	354 687		121 8	2 318	
923	Hg(C₁₀H₇)₂—Mercury α-naphthyl	454 718		188	1 929	
924	Hg(C₂H₃O₂)₂	318 656		d.	3 27o	
925	Hg(C₃H₅O₂)₂.	346 687		110		
926	Hg(C₇H₅O₂)₂...	442 687		165		
927	Hg(C₁₈H₃₃O₂)₂—Oleate	763 118		103		
928	Hg₂(C₂H₃O₂)₂.	547 297		225 d.		
929	HgCH₃Cl	251 091		170	4 063	
930	HgC₂H₅Cl	265 107		193	3 482	
931	HgCH₃I	342 565		143		
932	Hg(C₂H₅S)₂	322 817		77		
933	Hg(CN)₂	252 626	Tet.		4 00	
934	CuO—Paramelaconite	79 5700			6 4	
935	CuO—Tenorite	79 5700	C.	d. 1026¹⁵² ᵐᵐ O₂	6 40	1078
936	Cu₂O—Cuprite	143 140	C.	1235⁰⁴ ᵐᵐ O₂	6 0	188
937	CuF	82 5700		908		
938	CuF₂.5HF.6H₂O.	309 701	M.	d.	2 405	
939	CuCl—Nantokite	99 0280	C.	422	3 5₃	173

Mg Mn Mo N Na Nb Nd Ni O Os P Pb Pd Pr Pt Ra Rb Rh Ru S Sa Sb Sc Se Si Sn Sr Ta Tb Te Th Tl Ti Tm U V W Y Yb Zn Zr
76 42 47 11 82 51 61 45 1 35 12 23 41 60 37 80 84 40 39 8 63 14 56 9 18 22 78 52 66 10 24 19 27 70 49 50 43 57 71 28 21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
940	CuCl₂	134 486		49s	3.054	
941	CuCl₂.2H₂O	170 517	R	110 d.	2 390[22 4]	883
942	Cu(ClO₂)₂.6H₂O	338 578	C. ?	65		
943	Cu(ClO₄)₂.7H₂O	388 594			1 955	
944	3CuO.CuCl₂.3H₂O — Atacamite	427 242	R.	d. 200	3 94	1033
945	3CuO.CuCl₂.3H₂O—Paratacamite	427 242	Trig.	d. 200	3 74	172
946	4CuO.Cl₂O₄.3H₂O	523 242	R. M. ?	d.	3 55	
947	CuBr	143 486	C.	504	4 72	
948	CuBr₂	223 402	M.	498		
949	CuBr₂.4H₂O	295 464	R.	Tr. 30		
950	Cu(BrO₃)₂.6H₂O	427 494	C.	d. 180	2 583	
951	CuI— Marshite	190 502	C. Tet.	605	5 62	186
952	Cu(IO₃)₂	413 434	M.	d.	5 241[15]	
953	Cu(IO₃)₂.H₂O	431 449	Tri.	d. 240	4 876[15]	
954	Cu(IO₃)OH	255 510	R.	d. 290	4 878[16]	
955	CuS—Covellite	95 6350	H. M. ?	Tr. 103	4 6	
956	Cu₂S—Chalcocite	159 205	R.	1100	5 6	
957	Cu₂S	159 205	C.	1130	5 783	
958	CuSO₄—Hydrocyanite	159 635	R.	200	3 6	
959	CuSO₄.H₂O	177 650		d. 221	3 17	
960	CuSO₄.3H₂O	213 681	M.		2 663	
961	CuSO₄.5H₂O—Chalcanthite	249 712	Tri.	d. 20	2 286[15 6]	641
962	CuSO₄.7H₂O—Boothite	285 743	M.		1 944[21]	
963	Cu₂SO₃.H₂O	225 220	H.		3 83[15]	
964	3CuO.SO₃.2H₂O—Antlerite	354 806	R.		3 9	921
965	Cu₂SO₄.CuSO₄.2H₂O	386 871		d. 150	3 57	
966	4CuO.SO₃.3H₂O— Brochantite	452 391	R.		3 907	944
967	4CuO.SO₃.4H₂O—Langite	470 407	R.		3 49	939
968	7CuO.2SO₃.5H₂O	807 197	R.		3 85	
969	20CuO.SO₃.2CuCl₂.20H₂O—Connellite	2300 75	H.		3 4	350
970	Cu₂Se	206 340	C.	1113	6 749[20]	
971	Cu₂Se₂—Umangite	349 110			5 620	
972	CuO.SeO₂.2H₂O—Chalcomenite	226 801	M. R. ?		3 76	916
973	CuSeO₄.5H₂O	296 847	Tri.		2 559	
974	Cu(NO₃)₂.3H₂O	241 631		114 49	2 047	
975	Cu(NO₃)₂.6H₂O	295 678		26 4 d.		
976	4CuO.N₂O₅.3H₂O—Gerhardite	480 342	R.		3 43	903
977	CuCl₂.2NH₄Cl..... .	241 480			1 905[11 6]	
978	CuCl₂.2NH₄Cl.2H₂O	277 510	Tet.	d. 110	1 98	354
979	CuCl.3NH₃	150 121		123		
980	2CuCl.NH₃	215 087		162		
981	2CuCl.3NH₃ . .	249 149		144		
982	3CuCl₂.10NH₃	573 769		270		
983	Cu(ClO₃)₂.4NH₃ . .	298 610		d. 90	1 81	
984	CuBr₂.2NH₃	257 464		d. 200		
985	CuBr.3NH₃	194 579		115		
986	2CuBr.3NH₃	338 065		135		
987	Cu(BrO₃)₂.4NH₃ . .	387 526		exp. 140	2 31	
988	CuI.3NH₃ .	241 595		105		
989	2CuI.3NH₃ . .	432 097		117		
990	Cu(IO₃)₂.5NH₃	498 590		exp. 215	2 72	
991	(NH₄)₂SO₄.CuSO₄	291 778			2 348	
992	(NH₄)₂SO₄.CuSO₄.6H₂O.	399 870	M.	d. 120	1 87	538
993	(NH₄)₂SeO₄.CuSeO₄.6H₂O	494 140	M.		2 22	639
994	CuP .	94 5940			5 14	
995	Cu₂P	158 164		d.	6 4	
996	Cu₃P₂	252 758		d.	6 67	
997	4CuO.P₂O₅.H₂O—Libethenite .	478 343	R.		3 7	932
998	4CuO.P₂O₅.2H₂O—Pseudolibethenite .	496 359			4 0	
999	4CuO.P₂O₅.3H₂O—Tagilite .	514 374			4 08	968
1000	5CuO.P₂O₅.2H₂O—Dihydrite	575 929	M. Tri.		4 2	940
1001	6CuO.P₂O₅.3H₂O—Phosphochalcite ..	673 514			4.4	
1002	Cu(H₂PO₄)₂	193 649		exp. 90		

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1003	CuPO₄.CuOH.............	239 172	R.			931
1004	Cu₃As—Domeykite....	265 670	H.	830	8 0o	
1005	3CuO.As₂O₅.5H₂O—Trichalcite	558 707	R.			885
1006	4CuO.As₂O₅.H₂O—Olivenite	566 215	R.		4 3	951
1007	4CuO.As₂O₅.3H₂O—Leucochalcite	602 246	R.			960
1008	4CuO.As₂O₅.7H₂O—Euchroite	674 308	R.		3 4o	891
1009	5CuO.As₂O₅.H₂O—Erinite	645 785			4 04	964
1010	6CuO.As₂O₅.3H₂O—Clinoclasite	761 386	M.		4 37	976
1011	7CuO.As₂O₅.14H₂O—Chalcophyllite	1039 12	Trig.		2 66	306
1012	5CuO.As₂O₅.9H₂O—Tyrolite	789 909	R.		3 0₅	912
1013	2Cu₂S.As₂S₃...	564 525			4 289	
1014	3Cu₂S.As₂S₅—Enargite	787 860	C.		4 40	
1015	3Cu₂S.2As₂S₃—Binnite	969 845	C.		4 48	
1016	Cu₃(AsO₄)₂.3NH₃.4H₂O	591 785	Tri		3 05	
1017	Cu₃Sb (β).....	312 480		687 Tr. 407 (β to α)	8 51 (β) 8 48 (α)	
1018	Cu₂Sb₂	561 390		830		
1019	Cu₂S.Sb₂S₃—Chalcostibite	498 940	R.		4 932	
1020	Cu₂S.2Sb₂S₃—Guejarite	838 675	R.		4 811	
1021	3Cu₂S.Sb₂S₃—Stylotypite	817 350			5 147	
1022	Cu₂S.Bi₂S₃—Emplectite	673 400	R.		6 10¹⁴	
1023	5Cu₂S.2Bi₂S₃—Wittichenite	1824 42			5 9¹⁴	
1024	2Cu₂S.Bi₂S₃.2BiSCl	1385 7			6 78	
1025	2Cu₂S.Bi₂S₃.2BiSBr	1474 6			6 41	
1025 1	20CuO.Bi₂O₃.5As₂O₅.22H₂O—Mixite	3603 34			3 79	352
1026	2CuO.CO₂—Mysorine	203 140			4 398	
1027	2CuO.CO₂.H₂O—Malachite	221 155	M.		4 0	977
1028	3CuO.2CO₂.H₂O—Azurite	344 725	M.	d. 220	3 88	938
1029	Cu(CHO₂)₂.... ..	153 585			1 831	
1030	Cu(CHO₂)₂.4H₂O	225 647	M		1 795	652
1031	Cu(C₂H₃O₂)₂.	181 616			1 930	
1032	Cu(C₂H₃O₂)₂.H₂O	199 632		115	1 882	667
1033	Cu(C₂H₃O₂)₂.2H₂O	217 647		d. 240	1 9	
1034	Cu(CH₂SO₃)₂.4H₂O—Ethane disulfonate	323 790	Tri.		2 061	
1035	CuC₁₀H₆O₆S₂.6H₂O—1, 5-Naphthalene disulfonate	457 839	M.		1 783	792
1036	CuCN.	89 5780	M.	474 5		
1037	CuC₂O₄.2NH₃	185 632			2 305⁷ (α) 2 225⁷ (β)	
1038	CuSCN	121 613			2 846¹⁵	
1039	Cu₂(NH₃)₂(SCN)₂	277 348	R.	d. 20	1 021¹⁵	
1040	Cu₂Si	155 200			6 9¹⁸	
1041	Cu₄Si	282 340		850	7 53	
1042	Cu₆Si₂	373 970		775		
1043	CuO.SiO₂.H₂O—Bisbeeite	157 645	R.			783
1044	CuO.SiO₂.H₂O—Dioptase	157 645	Trig.		3 05	319
1045	2CuO.2SiO₂.H₂O—Shattuckite	297' 275	M.			948
1046	6CuO.5SiO₂.2H₂O—Plancheite	813 751	M.		3 36	320
1047	CuSiF₆.6H₂O...... ..	313 722	R.		2 158¹⁹	211
1048	CuCl₂.PbO.H₂O—Percylite	375 701	C.		4.67¹⁶ ⁷	176
1049	2CuO.5PbO.3SO₃.CO₂.3H₂O—Linarite	1613 38	M.		5 4	967
1050	CuO.4PbO.P₂O₅—Tsumebite	1114 42	R.		6	987
1051	Cu₂S.2PbS.Bi₂S₃—Aikinite	1151 93	R.		6 45	
1053	5Cu₂S.2ZnS.2As₂S₃—Tennantite	1483 14	C.		4 4	198
1054	Cu₂HgI₄.	835 478			6 096⁷	
1055	CuCl.HgS.	331 703			6 29	
1056	Ag₂O..	231 760	C.	d. 3oo	7 143¹⁶ ⁶	
1057	Ag₂O₂.	247 760		d. >100	7 44	
1058	AgF.	126 880	C.	435	5 852¹⁶ ⁵	
1059	AgCl—Cerargyrite..	143.338		455	5.56	177
1060	AgClO₃ .	191 338	Tet.	23o	4.43o	
1061	AgClO₄	207 338		d. 486		
1062	AgBr—Bromyrite	187 796	C.	43₄	6.474	185

Mg	Mn	Mo	N	Na	Nb	Nd	Ni	O	Os	P	Pb	Pd	Pr	Pt	Ra	Rb	Rh	Ru	S	Sa	Sb	Se	Se	Si	Sn	Sr	Ta	Tb	Te	Th	Tl	Tl	Tm	U	V	W	Y	Yb	Zn	Zr
75	42	47	11	82	51	61	45	1	35	12	23	41	60	37	80	84	40	39	8	63	14	56	9	18	22	78	52	66	10	24	19	27	70	49	50	45	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1063	AgBrO₃	235 796	Tet	d.	5.206	372
1064	AgI—Iodyrite	234 812	H.	d.552	5.67	400
1065	AgIO₃	282 812	R.	>200	5 525	
1066	Ag₂S Acanthite	247 825	R.	825	7 326	
				Tr. 175		
1067	Ag₂S Argentite	247 825	C.	Tr. 175	7.317	
1068	Ag₂SO₄	311 825	R.	652	5.45₄²⁹ ¹	
1069	Ag₂S₂O₈ 2H₂O	411 921	R.		3 61	844
1070	Ag₂Se Naumannite	294 960		880	8 0	
1071	Ag₂SeO₃	342 960			5 929	
1072	Ag₂Te Hessite	343 260	C.	955	8 ₅	
1073	AgN₃	149 904		exp. 251 5		
1074	AgNO₂	153 888	R.	d 140	4.453²⁶	
1075	AgNO₃	169 888	R.	212	4.352⁴⁹	1050
1076	Ag₂(NO₃)	275 776		d. 110	5 75²⁰	
1077	AgNO₃.NH₃	170 919	Tet.	70 d		
1078	NH₄NO₃.AgNO₃	249 935	R.	109 6		
1079	Ag(NH₃)₂NO₂	203 950	R.	170 d		
1080	AgCl AgNO₃	313 226		160		
1081	2AgCl 3NH₃	337 769	R.	68 d.		
1082	AgI.AgNO₃	404 700	R.	94		
1083	AgI.2AgNO₃	574 588	R.	119 1		
1084	AgBr.NH₄Br.4(NH₄)₂S₂O₃	878 580	Tet.			336
1085	Ag₃P₃	308 832		d.	4 63	
1086	AgPO₃	186 904		482	6 370	
1087	Ag₃PO₄	418 664	C.	849	6 370²⁵	
1088	Ag₄P₂O₇	605 568		585	5 306⁷ ⁵	
1089	Ag₂HPO₄	311 792	Trig.	d. 110		366
1090	Ag₃AsO₃	446 600		150 d		
1091	Ag₃AsO₄	462 600	C.		6 657²⁵	
1092	Ag₃AsBr₄	638 348		d.	5 55²⁵	
1093	Ag₂S.As₂S₃ Smithite	493 940	M.		4 700	1066
1094	Ag₂S.As₂S₃—Trechmannite	493 940	Trig.		4 700	422
1095	3Ag₂S.As₂S₃ Proustite	989 590	Trig.		5 49	412
1096	3Ag₂S.As₂S₅—Xanthoconite	1053 72	R.		5 2	1030
1097	Ag₂S.Sb₂S₃ Miargyrite	587 560	M.		5 36¹⁷	
1098	3Ag₂S Sb₂S₃—Pyrargyrite	1083 21	Trig.		5 7₆	425
1099	3Ag₂S.Sb₂S₃—Pyrostilpnite	1083 21	M. Tri.		5 790¹⁷	
1100	5Ag₂S Sb₂S₃—Stephanite	1578 86	R.		6 ₃	
1101	8Ag₂S.Sb₂S₃—Polybasite	2322 34	R.		6 1	1031
1102	12Ag₂S Sb₂S₃—Polyargyrite	3313 64	R.		6 50	
1103	Ag₂S.Bi₂S₃ Matildite	762 020	R.		6 9	
1104	AgNO₃ Bi(NO₃)₃ 2NH₄NO₃	629 006			3 055¹⁵	
1105	Ag₂CO₃	275 760		218 d.	6 077	
1106	Ag₂C₂O₄	303 760		exp. 140	5 029⁴	
1107	AgC₂H₃O₂	166 903		d.	3 259¹⁶	
1108	AgC₃H₅O₃.0 5H₂O Lactate	205 995		100		
1109	Ag₂(d-C₄H₄O₆)	363 791		d	3 432¹⁶	
1110	Ag₂(dl-C₄H₄O₆)	363 791			3.775¹⁶	
1111	AgCN	133 888		320 d.	3 95	
1112	AgCNO	149 888		d.	4 00	
1113	AgCN.NH₃	150 919	M.	102 d		
1114	Ag(SbO)(d-C₄H₄O₆).H₂O	364 886	R.		3 481¹⁸ ²	
1115	4Ag₂S GeS₂ Argyrodite	1127 81	C.		6 085¹⁵	
1116	4Ag₂S.SnS₂ Canfieldite	1174 13	C.		6 28	
1117	Ag₂S 2As₂S₃.6PbS Lengenbachite	2175 65	Tri.		5 8	
1118	3Ag₂S.4PbS.3Sb₂S₃—Diaphorite	2719 74	R.		5 9	
1119	3Ag₂S.4PbS 3Sb₂S₃—Freieslebenite	2719 74	M.		6 3	
1120	AgNO₃.2TlNO₃.Bi(NO₃)₃	1001 73			4 87¹⁴	
1121	AgCl.HgCl	379 406			6 495	
1122	2AgI HgI₂	924 098		Tr. 45	5 998⁰	
1123	4AgI.CuI Miersite	1129 75			5 64	183
1124	Ag₂S.Cu₂S—Stromeyerite	407 030	R.		6 ₂	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
47 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1125	Au₂O............	410.400		d. 20₅		
1126	Au₂O₂....	426 400		d. 18₀		
1127	Au₂O₃.....................	442 400		d. 16₀		
1128	AuCl	232 658		d. 289 5	7 4	
1129	AuCl₃	303 574		254 d	3 9	
1130	Au₂Cl₄	536 232		d. 25₀	5 1	
1131	AuBr	277 116		d. 115		
1132	AuBr₃	436 948		16₀ d.		
1133	Au₂Br₄	714 064		d. 115		
1134	AuHBr₄.5H₂O...... .	607 949		27		
1135	AuI	324 132		d. 120		
1136	Au₂S₃	458 530		d. 14₀		
1137	Au₂S₃	490 595		d 19₇	8 754	
1138	Au₂Se₃	632 000			4 65''	
1139	AuTe—Calaverite .	324 700	Tri.		9 04	
1140	Au₂Te₄.......	904 400		47₂		
1141	HAu(NO₃)₄.3H₂O	500 286		72 d.	2 81	
1142	Au₂O₃.4NH₃....	510 524		exp. 143		
1143	Au₂P₃....	487 472			6 67	
1144	Au(CN)₃.3H₂O.	329 270		d. 50		
1145	4AuCl₃.3AgCl.8NH₄Cl	2072 28	R.			159
1146	OsO₂ .	222 800			7 91	
1147	OsO₄ (yellow) . .	254 800	M.	41	4 91	57
					1 4 41⁴⁰ ¹	
1147 5	OsO₄ (white)	254 800		39 5		
1148	OsF₆	304 800				
1149	OsF₈	342 800		34 5		
1150	(NH₄)₂OsCl₆ ..	439 626	C.		2 93	
1151	(NH₄)₂OsBr₆	706 374			4 09	
1152	IrCl.. .	228 558		d. 798	10 18	
1153	IrCl₂	264 016		d. 773		
1154	IrCl₃..... . . .	299 474		d 763	5 30	
1155	(NH₄)₂IrCl₆ . ..	441 926	C.		2 856	
1156	IrCl.4NH₃.H₂O .	314 698	Trig.			327
1157	[Ir(NH₃)₅Cl]Cl₂	384 630	R		2 675	
1158	[Ir(NH₃)₅Br]Br₂	518 004	R.		3 215⁴⁰ ᵇ	
1159	[Ir(NH₃)₅Cl]Br₂	473 546	R.		3 01	
1160	[Ir(NH₃)₅I]I₂	659 052	R.		3 586⁴⁰ ᵇ	
1161	[Ir(NH₃)₅Cl]I₂	567 578	R.		3 12	
1162	Ir₂(SO₄)₃.(NH₄)₂SO₄.24H₂O	1238 91	C.	106		
1163	PtCl₂ .	266 146		d. 5₅1	5 87	
1164	PtCl₂.8H₂O	481 185			2 43	
1165	H₂PtCl₆.6H₂O	518 086		60	2 431	
1166	PtBr₄ .	514 894		d. 180		
1167	H₂PtBr₆.9H₂O	838 880	M.	<100 d.		
1168	PtI₄ .	702 958		d 100		
1169	PtS	227 295			8 897	
1170	PtSe₂	353 630			7 65	
1171	PtSe₃ .	432 830			7 15	
1172	Pt(NH₃)₄(OH)₂ .	297 370		110 d.		
1173	Pt(NH₃)₂Cl₂ . .	300 208	R.	d 270		
1174	(NH₄)₂PtCl₆ ..	441 056	C.		3 065	
1175	[Pt(NH₃)₄]Cl₂.H₂O .	352 286	Tet.	d. 110	2 737	
1176	(NH₄)₂PtBr₆ . .	710 804	C.		4 265	
1177	(NH₄)₂PtI₆ .	992 900	C.		4 61	
1178	PtP₂O₇ . . .	369 278		d >600	4 856	
1179	PtAs₂—Sperrylite	345 150	C.	>800	10 60	
1180	[Pt(CO)Cl₂]₂	588 292		195		
1181	2PtCl₂.3CO	616 292	M.	130		
1182	[Pt(CO)Br₂]₂.	766 124	M.	182		
1183	[Pt(CO)I₂]₂	954 188		ca. 150 d.		
1184	[CH₃(C₇H₅)₂SCl]₂PtCl₄	618 308	M.	210		888
1185	[(C₂H₅)₂SCl]₂PtCl₄	646 339	M.			811

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1186	$[C_3H_5NH_3]_2H_2PtCl_6$	500 117		218 d.	2 275[18]	
1187	$[(CH_3)_2N]_2H_2PtCl_6$	528 148		245 d.	2 015	139
1188	$[CH_3(C_2H_5)NH]_2H_2PtCl_6$	528 148		208	2 115[18]	
1189	$[C_3H_7NH_3]_2H_2PtCl_6$	528 148		214	2 218	
1190	$[(iso-C_3H_7)NH_3]_2H_2PtCl_6$	528 148		228	2 229	
1191	$[(CH_3)_4N]_2PtCl_6$	556 179	C.	278 d.	1.811[16]	
1192	$[CH_3(C_4H_9)NH_2]_2H_2PtCl_6$	556 179		200 d.	1 068[16]	
1193	$[(CH_3)_3C_3H_5N]_2PtCl_6$	584 210	C.	266 d.	1.762[17]	
1194	$[(C_2H_5)C_3H_7NH]_2H_2PtCl_6$	584 210		199	1.89	
1195	$[C_2H_5(iso-C_3H_7)NH]_2H_2PtCl_6$	584 210		180	1 885	
1196	$[C_2H_5(iso-C_4H_9)NH]_2H_2PtCl_6$	612 240		201 d.	1 804	
1197	$[(C_3H_5)_3N]_2H_2PtCl_6$	612 240		100	1 903	
1198	$[(C_3H_7)_2NH]_2H_2PtCl_6$	612 240		175 d.	1 704[16]	
1199	$[(CH_3)_3C_4H_7N]_2PtCl_6$	612 240	C.	252 d.	1 821	
1200	$[(CH_3)_3(iso-C_4H_7)N]_2PtCl_6$	612 240	C.	237	1 871[16]	
1201	$[(C_3H_7)(iso-C_4H_9)NH]_2H_2PtCl_6$	640 271		188	1 702[16]	
1202	$[(CH_3)_3(C_3H_5)_3N]_2PtCl_6$	640 271	C.	250 d.	1 731	
1203	$[(CH_3)_2(C_2H_5)(C_3H_7)N]_2PtCl_6$	640 271	C.	256 d.	1 812	
1204	$[(CH_3)_3(C_4H_9)N]_2PtCl_6$	640 271	C.	259 d.	1 795	
1205	$[(CH_3)_3(iso-C_4H_9)N]_2PtCl_6$	640 271	C.	220	1 751[17]	
1206	$[(CH_3)(C_3H_7)_2N]_2H_2PtCl_6$	640 271		>200	1 737	
1207	$[(C_2H_5)_4N]_2PtCl_6$	668 302	C.	250 d.	1 776	
1208	$[(iso-C_4H_9)_2NH]_2H_2PtCl_6$	668 302		213	1.62[16]	
1209	$[(C_2H_5)(C_3H_7)_2N]_2H_2PtCl_6$	668 302		175	1.726	
1210	$[(CH_3)_2(C_3H_7)_2N]_2PtCl_6$	668 302	Tet.	250	1.745	
1211	$[(C_2H_5)_3(C_3H_7)N]_2PtCl_6$	696 333	C.	235 d.	1 710	
1212	$[(CH_3)(C_2H_5)(C_3H_7)_2N]_2PtCl_6$	696 333	C.	228 d.	1 712	
1213	$[(C_2H_5)_2(C_3H_7)_2N]_2PtCl_6$	724 364	C.	220 d.	1 677	
1214	$[(CH_3)(C_2H_5)(C_3H_7)(iso-C_4H_9)N]_2PtCl_6$	724 364		236 d.	1 637	
1215	$[(C_2H_5)_3(C_4H_9)N]_2PtCl_6$	724 364	C.	220	1 629[16]	
1216	$[(C_2H_5)_3(iso-C_4H_9)N]_2PtCl_6$	724 364	M.	215	1 602	
1217	$[(C_2H_5)(C_3H_7)_3N]_2PtCl_6$	752 394	Tri.	212	1 571[17]	
1218	$[(C_3H_7)_4N]_2PtCl_6$	780 424	Tri.	199	1 515	
1219	$[(CH_3)(iso-C_4H_9)_3N]_2PtCl_6$	808 456	R. ?	174	1 696	
1220	$[(C_2H_5)(iso-C_4H_9)_3N]_2PtCl_6$	836 487	Tet.	170	1 562[17]	
1221	$[(C_3H_7)(iso-C_4H_9)_3N]_2PtCl_6$	864 518	C.	168	1 509	
1222	$Pt_x(NO_2)_y(C_nH_bS_c)_z$	Tschugaeff and Chlopin, 93, 82 : 402 ; 12.				
1223	PtSi	223 290		1100	11 63[16]	
1224	Pt_2Si	418 520			13 8[16]	
1225	Pt_3Si_2	641.810			14 1	
1226	$PtPbCl_6.4H_2O$	687 240	C.		3 681	
1227	$PtPbBr_6$	881 926		d. >120	6 025	
1228	$PtZnCl_6.6H_2O$	581 450	Trig.		2 717	
1229	$PtZnBr_6.12H_2O$	956 291	Trig.		2 877	
1230	$PtZnI_6.9H_2O$	1184 34	Trig.		3 689	
1231	$PtCdCl_6.6H_2O$	628 480	Trig.		2 882	
1232	$PtCuCl_6.6H_2O$	579 964	Trig.		2 734	
1233	RuO_2	133 700	Tet.		7 2	
1234	RuO_4	165 700		25 5	5 77[100]	
1235	Ru_2S_3 Laurite	299 595	C.		6 99	
1236	RuSi	129 760			5 4	
1237	$[Rh_2(NH_3)_{10}Cl_2]Cl_4$	588 879	R.	d. 200	2 079[18]	
1238	$[Rh(NH_3)_5Br]Br_2$	427.814	R.		2 65	
1239	$[Rh(NH_3)_5I]I_2$	568 862	R.		3.12[16]	
1240	$NH_4Rh(SO_4)_2.12H_2O$	529 264	C.	103		115
1241	$TlRh(SO_4)_2.12H_2O$	715 625	C.			130
1242	$RbRh(SO_4)_2.12H_2O$	596 665	C.	109		
1243	PdO	122 700		d. 877		
1244	$PdCl_2$	177 616		500		
1245	PdI_2	360 564		d. 350		
1246	PdS	138 765		950		
1247	Pd_2S	245 465		800 d.	7.3	
1248	PdSe	185 900		<960		

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1249	Pd(NH₃)₂Cl₂ . . .	211 678	Tet.		2 5	
1250	(NH₄)₂PdCl₄	284 610	Tet.		2 17	
1251	(NH₄)₂PdCl₆	355 526	C.		2 418	
1252	(NH₄)₂PdSO₂Cl₄.H₂O	365 268	Trig.			316
1253	Pd(CO)Cl₂	205 616		197		
1254	Pd(CO)₂Cl₂	233 616		142		
1255	2PdCl₂.3CO	439 232		132		
1256	PdSi	134 760			7 31[14]	
1257	ZnPdCl₄.6H₂O	492 920	H.		2 359	
1258	MnO—Manganosite	70 9300	C.	1650	5 18	180
1259	MnO.H₂O—Pyrochroite	88 9454	Trig.		3 258[14]	349
1260	MnO₂—Polianite, Pyrolusite	86 9300	R.		5 026	
1261	MnO₂.H₂O	104 945	C.			171
1262	Mn₂O₃ . . .	157 860	C.		4 50	
1263	Mn₂O₃.H₂O—Manganite	175 875	R.		3 258	1058
1264	Mn₃O₄—Hausmannite	228 790	Tet.		4 700	121
1265	MnF₂	92 9300		856	3 98	
1266	MnF₃	111 930			3 54	
1267	MnF₃.5HF.6H₂O	301 061			1 921	
1268	MnCl₂—Scacchite	125 846	C.	650	2 977[b]	
1269	MnCl₂.4H₂O	197 908	M.	58 01	2 01	
1270	Mn(ClO₄)₂.8H₂O	397 969			1 99	
1270.1	MnCl₂.3MnO₂.3H₂O—Kempite	440 682	R.		2 94	889
1271	MnBr₂	214 762			4 385[b] fused	
1272	MnBr₂.4H₂O	285 820	M.	64 3d		
1273	MnS—Alabandite	86 9950	C.	d.	3 99	197
1274	MnS₂—Hauerite	119 060	C.		3 463	106
1275	MnSO₄	150 995		700	3 25	
1276	MnSO₄.H₂O—Szmikite	169 010	M. ?		2 954	742
1277	MnSO₄.2H₂O	187 026			2 526	
1278	MnSO₄.3H₂O	205 041			2 356	
1279	MnSO₄.4H₂O	223 057	M. R.		2 107	
1280	MnSO₄.5H₂O	241 072	Tri.		2 103	
1281	MnS₂O₆.6H₂O	323 152	Tri		1 757	
1282	MnSe	134.130	C.		5 59[b]	
1283	MnSeO₄.2H₂O	234 161	R		2 949	
1284	MnSeO₄.5H₂O	288 207	Tri.		2 334	
1285	Mn₃N₂ . . .	302 666			6 63	
1286	Mn(NO₃)₂.3H₂O	232 992		34 81		
1287	Mn(NO₃)₂.6H₂O	287 038		25 8	1 82	
1288	NH₄MnO₄	136 969	R.		2 208[10 a]	
1289	(NH₄)₂SO₄.MnSO₄.6H₂O	391 229	M.		1 83[1]	484
1290	(NH₄)₂SO₄.2MnSO₄	434 133	C.		2 56[14]	
1291	(NH₄)₂SO₄.Mn₂(SO₄)₃	530 196			2 40[13]	
1292	(NH₄)₂SeO₄.MnSeO₄.6H₂O	485 500	M.		2 093	
1293	Mn₃P₂ . . .	391 628			4 94	
1294	Mn₂P₂O₇ . . .	283 908	M.		3 707[15]	897
1295	3MnO.P₂O₅.3H₂O—Reddingite	408 884	R.		3 1	842
1296	3MnO.P₂O₅.4H₂O ?—Stewartite	426 898	Tri.		2 94	846
1297	5MnO 2P₂O₅.4H₂O—Palaite	710 808	M.		3 17	843
1298	5MnO.2P₂O₅.5H₂O—Hureaulite	728 823	M.		3 18	835
1299	3MnO.As₂O₅—Armangite	442 710	H. R.		4 23	
1300	4MnO.As₂O₅.H₂O—Sarkinite, Polyarsenite	531 655	M.		4 15	954
1301	Mn₂O₃.4MnO.As₂O₅.4H₂O—Flinkite	743 562	R.		3 87	959
1302	6MnO.As₂O₅.5H₂O—Hemafibrite	745 577	R.		3 6	980
1303	7MnO.As₂O₅.4H₂O—Allactite . . .	798 492	M.		3 84	945
1304	MnSb	176 700			5 6[17]	
1305	10MnO.Sb₂O₄—Manganostibite	1032 84	M.			989
1306	Mn₃C	176 790			6 89[17]	
1307	MnCO₃—Rhodochrosite	114 930	Trig.		3 125	368
1308	MnC₂O₄	142 930			2 43[21 7]	
1309	Mn(CHO₂)₂	144 945			2 205	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1310	$Mn(CHO_2)_2.2H_2O$	180 976	R.		1 953	
1311	$Mn(C_2H_3O_2)_2$	172 976			1 74	
1312	$Mn(C_2H_3O_2)_2.4H_2O$	245 038	M.		1 589	
1313	$MnCl_2.2C_2H_4N.HCl$	320 405		175		
1314	$MnSi$	82 9900		1280	$5\ 90^{15}$	
1315	$MnSi_2$	111 050			$5\ 24^{15}$	
1316	Mn_2Si	137 920		1316	6.20^{15}	
1317	$MnO.SiO_2$	130 990		1273	3.48_4^{25}	63
1318	$MnO.SiO_2$—Rhodonite	130 990	Tri.	1323	$3\ 72_4^{25}$	929
1319	$2MnO.SiO_2$—Tephroite	201 920	R.	1300	$4\ 043_4^{25}$	949
1320	$3Mn_2O_3.MnO\ SiO_2$ Braunite	604 570	Tet.		4 78	
1321	$8MnO\ 7SiO_2\ 5H_2O$—Bementite	1077 94	R.		2 90	803
1322	$12MnO\ 8SiO_2\ 7H_2O$—Ectropite	1457 75	M. ?		2 46	1044
1323	$MnSiF_6.6H_2O$	305 082	Trig.	d.	$1\ 904^{17\ 5}$	206
1324	$5MnO.SiO_2\ As_2O_3.H_2O$—Dixenite	630 645	H.		4 2	385
1324 I	$12MnO\ 9SiO_2.As_2O_3.7H_2O$ Schallerite	1747 73			3 368	344
1325	$MnO.TiO_2$—Pyrophanite	150 830	Trig.	1404	4 54	405
1326	$2MnO\ 6PbO.3As_2O_3.H_2O$—Trigonite	2188 84	M.		8 28	1004
1327	$2Mn_2O_3\ 3PbO.3SiO_2$—Kentrolite	1165 44	R.		6 19	1014
1328	$2Mn_2O_3\ 3CuO$—Crednerite	554 430			5 0	
1329	$MnPtCl_6.6H_2O$	571 000	Trig.	d.	2 692	
1330	$MnPtCl_6\ 12H_2O$	679 093	Trig.		2 112	
1331	$MnPtBr_6.12H_2O$	945 841	Trig.		2 759	
1332	$MnPtI_6.9H_2O$	1173 89	Trig.	d	3 604	
1333	FeO	71 8400		1420		
1334	Fe_2O_3—Hematite	159 680	Trig.	1560 d	5.12	424
1335	$Fe_2O_3.H_2O$—Goethite	177 695	R.		4 28	1026
1336	$Fe_2O_3.H_2O$—Lepidocrocite	177 695	R.		4 09	1013
1337	Fe_3O_4 Magnetite	231 520	C.	1538 d	5 2	
1338	FeF_2	93 8400			4 09	
1339	FeF_3	112 840			3 18	
1340	$FeCl_2$—Lawrencite	126 756	H.		2 7	280
1341	$FeCl_2\ 4H_2O$	198 818			1 93	
1342	$FeCl_3$ Molysite	162 214	H.	282	2 8	
1343	$2FeCl_3\ 2HCl\ 4H_2O$	469 421		45 7		
1344	$FeBr_2$	215 672			$4\ 636_4^{25}$	
1345	$FeBr_3\ 6H_2O$	403 680		27		
1346	FeI_2	309 704		177		
1347	$FeI_2\ 4H_2O$	381 764			2 87	
1348	FeS—Troilite	87 9050	H.	1194	4 8	
1349	FeS_2—Marcasite	119 970	R.	Tr 450	4 87	
1350	FeS_2 Pyrite	119 970	C.		5 0	
1351	Fe_2S_3	207 875			4 3	
1352	Fe_3S_4	295 780			4 55	
1353	Fe_7S_8—Pyrrhotite	647 400	H.	d. >700	4 6	
1354	$FeSO_4.H_2O$—Szomolnokite	169 920	M.		3 08	
1355	$FeSO_4.5H_2O$—Siderotilate	241 982	Tri.		2 2	642
1356	$FeSO_4.7H_2O$—Melanterite	278 012	M.		1 89	471
1357	$Fe_2O_3.2SO_3.7H_2O$—Amarantite	445 918	Tri.		2.11	762
1358	$Fe_2O_3.2SO_3.10H_2O$ Fibroferrite	499 964	R.		1 86	255
1359	$Fe_2O_3\ 3SO_3.9H_2O$—Coquimbite	562 014	Trig.		2 1	270
1360	$Fe_2O_3.4SO_3.9H_2O$—Rhomboclasite	642 079	R.			675
1361	$FeO.Fe_2O_3.4SO_3.24H_2O$—Bilinite	984 150			1 87	530
1362	$2Fe_2O_3.SO_3.6H_2O$—Glockerite	507 517				958
1363	$2Fe_2O_3.5SO_3.18H_2O$—Copiapite	1043 96	R.		2 1	654
1364	$3Fe_2O_3.4SO_3.10H_2O$—Carphosiderite	979 454	Trig.		2 6	371
1365	$Fe_2O_3.3TeO.4H_2O$—Durdenite	662 242	R.			990
1366	Fe_3N	125 688		d.	6 35	
1367	$Fe(NO_3)_3.6H_2O$	349 956		35		
1368	$(NH_4)_2SO_4.FeSO_4.6H_2O$	392 140	M.		1 864	513
1369	$(NH_4)_2SO_4.Fe_2(SO_4)_3\ 24H_2O$	964 387	C.		1 71	102
1370	$(NH_4)_2SeO_4\ FeSeO_4.6H_2O$	486 410	M.		2 160	612
1371	FeP	86 8640			5 2	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
32 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1372	Fe₃P	142.704		1290	5 7	
1373	Fe₂P₃	204.752			4 5	
1374	Fe₃P	198.544		1110	6 74	
1375	Fe₃P₄	291.616			5 04	
1376	Fe(PO₃)₃	292.912			3 02	
1377	Fe₂O₃.P₂O₅.4H₂O—Strengite	373.790	R.		2 87	917
1378	3FeO.P₂O₅.8H₂O—Vivianite	501.691	M.		2 58	757
1379	2Fe₂O₃.P₂O₅.12H₂O—Cacoxenite	677.593	H.		3 38	285
1380	3Fe₂O₃.2P₂O₅.8H₂O—Beraunite	907.259	M.		2 9	950
1381	7FeO.2P₂O₅.9H₂O—Ludlamite	949.115	M.		3 72	873
1382	2Fe₂O₃.P₂O₅.2SO₃.2H₂O—Destinezite	657.569	Tri		2 1	794
1383	2Fe₂O₃.P₂O₅.2SO₃.2H₂O—Diadochite	657.569			2 0	142
1384	FeAs	130.800		1020	7 83	
1385	FeAs₂—Arsenoferrite	205.760	C.	990	7 4	
1386	FeAs₂—Löllingite	205.760	R.		7	
1387	FeAsO₄.4H₂O—Scorodite	266.862	R.		3 2	941
1388	3FeO.As₂O₅.8H₂O—Symplesite	589.563	M.		2 96	857
1389	3Fe₂O₃.2As₂O₅.13H₂O—Pharmacosiderite	1109.08	M. ?, C		3	874
1390	FeS₂.FeAs₂—Arsenopyrite	325.730	R.		6 2	
1391	2FeO.Sb₂O₅—Tripuhyite	467.220			5 82	1015
1392	FeS.Sb₂S₃—Berthierite	427.640	R.		4 0	
1393	Fe₃C	179.520		1837	7 4	
1394	FeCO₃.H₂O—Siderite	133.855	Trig.		3 8	377
1395	FeC₂O₄.2H₂O	179.871	R.	d 160	2 28	
1396	Fe(CO)₄	167.840		d 140	1 996¹⁸	
1397	Fe(CO)₅	195.840		−21	1 1 157	
1398	Fe₂(CO)₉	363.680		d. 100	2 085¹⁸	
1399	FeC₂₀H₁₄O₆S₂.6H₂O—Naphthalene-β-sulfonate	578.170				1039
1400	(NH₄)₄Fe(CN)₆.2NH₄Cl.3H₂O	415.083	Trig.		1 490	301
1401	Fe₄(NO)₇S₃N(C₂H₅)₄	659.773			1 883¹⁷	
1402	FeSi	83.9000			6 1	
1403	FeSi₂	111.960			5 4	
1404	Fe₂Si	139.740			7 0	
1405	Fe₃Si₂	223.640			6 7	
1406	FeO.SiO₂—Gruenerite	131.900	M.	1550	3 5	890
1407	2FeO.SiO₂—Fayalite	203.740	R.	1255		978
1408	2Fe₂O₃.2SiO₂.3H₂O—Iddingsite	493.526	R.		2 8	928
1409	FeSiF₆.6H₂O	305.992	Trig.			207
1410	FeO.TiO₂—Ilmenite	151.740	Trig.		4 76	
1411	Fe₂O₃.3TiO₂—Arizonite	399.380	M. ?		4 25	1069
1412	2Fe₂O₃.3TiO₂—Pseudobrookite	559.060	R.		4 7	1061
1413	6FeO.Sb₂O₃.5TiO₂—Derbylite	1122.08	R.		4 53	420
1414	2Fe₂O₃.PbO.3SO₃.4H₂O—Vegasite	854.817	H.			555
1415	3Fe₂O₃.PbO.4SO₃.6H₂O—Plumbojarosite	1130.59	Trig.		3 63	378
1416	3Fe₂O₃.2PbO.P₂O₅.2SO₃.6H₂O—Corkite	1335.71	Trig.		4 2	383
1417	5Fe₂O₃.3PbO.6As₂O₄—Carminite	2847.52			4 4	
1418	FeS.3Sb₂S₃.4PbS—Jamesonite	1967.98	M.		5 7	
1419	3Fe₂O₃.2PbO.As₂O₄.2SO₃.6H₂O—Beudantite	1423.58	Trig.		4 1	386
1420	9Fe₂O₃.4PbO.6As₂O₅.4SO₃.33H₂O—Lossenite	4622.21	R.			952
1421	2Fe₂O₃.3PbO.3SiO₂—Melanotekite	1169.14	R.		5 73	1010
1422	TlFe(SO₄)₂.12H₂O	668.555	C.		2 38	124
1423	Zn(FeO₂)₂	241.060			5 33	
1424	Fe₂O₃.CuO	239.250		1458		
1425	FeS.CuS—Chalcopyrite	183.540	Tet.		4 2	
1426	FeS.2Cu₂S.CuS—Bornite	501.950	C.		5 0	
1427	2FeS.CuS—Cubanite	271.415	R.		4 0	
1428	4FeS.Cu₂S.2CuS	702.095			5 0	
1429	4FeS.3Cu₂S.3CuS	1116.14			4 85	
1430	3Fe₂O₃.CuO.2P₂O₅.8H₂O—Chalcosiderite	986.829	Tri.		3 1	969
1431	Fe₂O₃.2CuO.As₂O₅.2H₂O—Chenevixite	584.771			3 93	379

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1432	$FeS.Cu_2S.SnS_2$—Stannite	429 940	Tet.		4.4	
1433	$Fe_2O_3.CuO.PbO.2SO_3.4H_2O$—Beaverite	694 642	H.		4.36	373
1434	$2Ag_3Fe(CN)_6.3NH_3$	1122 15			2.45	
1435	$FePtCl_6.6H_2O$	571 910			2.7	
1436	$FePtI_6.9H_2O$	1174 80			3.45	
1437	$FeO.MnO_2$—Bixbyite	158 770	C.		4.95	
1438	$Fe_2O_3.MnO$—Jacobsite	230 610	C.		4.75	
1439	$Fe_2O_3.9MnO.4P_2O_5.14H_2O$—Salmonsite	1618 46	R.		2.88	848
1439 1	$9(MnFe)O.8SiO_2.MnCl_2.7H_2O$—Friedelite		Trig.		3 1	329
1440	CoO	74 9700	C.	d. 80o	5 6s	
1441	Co_2O_3	165 940			5 1s	
1442	Co_3O_4	240 970			6 07з	
1443	$Co(OH)_2$	92 9854		d.	3 597¹⁵	
1444	CoF_2	96 9700	M.		4 43	
1445	$CoF_2.3H_2O$	151 016			2 583²¹	
1446	$CoF_3.5HF.6H_2O$	305 101	Trig.		2 045	
1447	$CoCl_2$	129 886			3 356	
1448	$CoCl_2.2H_2O$	165 917			2 477²⁵	
1449	$CoCl_2.6H_2O$	237 978	M.	86	1 924²⁵	
1450	$Co(ClO_3)_2.6H_2O$	333 978		61	1.92	
1451	$Co(ClO_4)_2.6H_2O$	365 978	H.	143		131
1452	$Co(ClO_4)_2.7H_2O$	383 994			2 075	
1453	$CoBr_2$	218 802			4.909¹⁵	
1454	$CoBr_2.6H_2O$	326 894		100 d.		
1455	CoI_2	312 834			5 68	
1456	$Co(IO_3)_2.6H_2O$	516 926			3.689²¹	
1457	CoS—Syeporite	91 0350		>1100	5 45	
1458	Co_3S_4—Linnaeite	305 170	C.		4.9	
1459	$CoSO_4$	155 035			3 710²⁵	
1460	$CoSO_4.H_2O$	173 050		d.	1 92	
1461	$CoSO_4.4H_2O$	227 096			2 368²⁵	
1462	$CoSO_4.6H_2O$	263 127	M.		2 029²⁵	
1463	$CoSO_4.7H_2O$—Bieberite	281 143	M. ?		1.948²⁵	481
1464	$CoSe$	138 170			7 65	
1465	$CoSeO_4.5H_2O$	292 247	Tri.	d.	2 512	599
1466	$CoSeO_4.6H_2O$	310 262	M.		2 32	
1467	$CoSeO_4.7H_2O$	328 278	M.		2.135	
1468	$Co(NO_3)_2.3H_2O$	237 032		91		
1469	$Co(NO_3)_2.6H_2O$	291 078	M.	<100	1 883²⁵	
1470	$Co(NO_3)_2.3NH_3$	248 087			2 001¹⁷	
1471	$[Co(NH_3)_4(NO_2)_2]NO_2$	281 118	R.		1.922¹⁷	
1472	$Co(NO_3)_2.6NH_3$	285 173			1.473²¹	
1473	$CoF_2.6NH_3$	199 157			1.744²⁵	
1474	$CoCl_2.NH_3$	146 917		ca. 321		
1475	$CoCl_2.2NH_3$ (α)	163 948		27з	2 097²⁵	
1476	$CoCl_2.2NH_3$ (β)	163 948			2 073²⁵	
1477	$CoCl_2.4NH_3$	198 010		d.	1.593²⁵	
1478	$CoCl_2.5NH_3$	215 042			1 580²⁵	
1479	$[Co(NH_3)_5Cl]Cl_2$	250 500	R.		1 819²⁵	
1480	$CoCl_2.6NH_3$	232 073		d.	1 497²⁵	
1481	$CoCl_3.6NH_3$	267 531	M.		1 744²⁵	
1482	$CoCl_3.10NH_3$	300 197			1.71²⁵	
1483	$[Co(NH_3)_4(OH_2)Cl]Cl_2$	251 484	R.		1.847	
1484	$[Co(NH_3)_5(NO_2)]Cl_2$	261 050	M.		1.698¹⁸	
1485	$[Co(NH_3)_5(NO_2)](NO_3)Cl$	287 500	R.		1.800	
1486	$CoBr_2.2NH_3$	252 864		26o		
1487	$[Co(NH_3)_5Br]Br_2$	383 874		d.	2.483¹⁷·⁸	
1488	$CoBr_2.6NH_3$	320 989			1.955	
1489	$[Co(NH_3)_5Br]Cl_2$	294 958			2 095¹⁶·⁸	
1490	$CoI_2.2NH_3$	346 896		22₂		
1491	$(NH_4)_2SO_4.CoSO_4.6H_2O$	395 270	M.		1.901	521
1492	$Co(SO_4)_2.4NH_3.2H_2O$	355 255			1 804²⁵	
1493	$Co(SO_4)_2.5NH_3$	336 256			1.703²⁵	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1494	[Co(NH$_3$)$_6$(SO$_4$)]SO$_4$.2H$_2$O	373 294	R.		1 828[18]	
1495	[Co(NH$_3$)$_6$(OH$_2$)]$_2$(SO$_4$)$_3$.3H$_2$O	666 523	Tet.		1 854	
1496	[Co(NH$_3$)$_6$]Cl(SO$_4$).3H$_2$O	346 726	R.		1 765	
1497	(NH$_4$)$_2$SeO$_4$.CoSeO$_4$.6H$_2$O	489 540	M.	d.	2 212	623
1498	Co(NH$_3$)$_6$Cl(SeO$_4$).3H$_2$O	393 861	R.		1.937	
1499	Co(H$_2$PO$_2$)$_2$.6H$_2$O	297 141			1.809[18,a]	
1500	CoAs$_2$—Safflorite	208 890		d.	6.97[6]	
1501	CoAs$_2$—Smaltite	208.890		d.	6 5	
1502	CoAs$_3$—Skutterudite	283 850			6.79	
1503	Co$_2$As$_5$	342 820		d.	7.35[6]	
1504	Co$_5$As$_2$	326 830		d.	7.82[6]	
1505	Co$_3$(AsO$_4$)$_2$.8H$_2$O—Erythrite	598 953	M.		2 9	850
1506	CoAsS—Cobaltite	165 995	C.	d.	6.2	
1507	CoCO$_3$—Spherocobaltite	118 970	Trig.		2 818[3]	375
1508	CoC$_2$O$_4$	146 970			2.325[1]	
1509	Co(CO)$_4$	170 970		51	1 73[18]	
1510	Co(CHO$_2$)$_2$.2H$_2$O	185 016			2 129[22]	
1511	CoC$_3$H$_2$O$_4$.2H$_2$O—Malonate	197 016			2 279	
1512	Co(C$_2$H$_3$O$_2$)$_2$.4H$_2$O	249 078	M.		1 7[18,7]	651
1513	Co(C$_5$H$_7$O$_2$)$_2$—Acetylacetonate	356 132				
1514	CoC$_{10}$H$_6$O$_6$S$_2$.6H$_2$O—1, 5-Naphthalene-disulfonate	453 239	M.		1 77	799
1515	Co(CO)$_3$NO	172 978		--1 05	1. 1 513[14]	
1516	[Co(NH$_3$)$_4$(C$_2$O$_4$)]NO$_3$.HNO$_3$	357 149			1 264[15]	
1517	CoSi	87 0300		1393	6 30	
1518	CoSi$_2$	115 090		1277	5 3[6]	
1519	CoSi$_3$	143 150		1307		
1520	Co$_2$Si	146 000		1327	7 1[17]	
1521	Co$_2$SiO$_4$	210 000			4.63	
1522	CoSiF$_6$.6H$_2$O	309 122	Trig.		2.087	413
1523	CoSnCl$_6$.6H$_2$O	498 510	R. Trig.		2.699	
1524	CoPtCl$_6$.6H$_2$O	575 040	Trig.	d.	2.699	
1525	CoPtBr$_6$.12H$_2$O	949.881	Trig.		2.762	
1526	CoPtI$_6$.9H$_2$O	1177.93	Trig.		3 618	
1527	CoPtI$_6$.12H$_2$O	1231.98	Trig.		3 018	
1528	NiO—Bunsenite	74 6900	C.		7.45	201
1529	Ni$_2$O$_3$	165 380			4 83	
1530	Ni$_3$O$_4$.2H$_2$O	258 085			3 412[32]	
1531	NiF$_2$	96 6900			4.63	
1532	NiF$_2$.3H$_2$O	150 736			2.014[19]	
1533	NiF$_2$.5HF.6H$_2$O	304 821	Trig.		2.132	
1534	NiCl$_2$	129 606			3 544	
1535	Ni(ClO$_3$)$_2$.6H$_2$O	333 698		80 d.	2 07	
1536	Ni(ClO$_4$)$_2$.6H$_2$O	365 698	H.	149		132
1537	Ni(ClO$_4$)$_2$.7H$_2$O	383 714			2 15	
1538	NiBr$_2$	218 522			4 64[18]	
1539	Ni(IO$_3$)$_2$	408 554			5 07	
1540	Ni(IO$_3$)$_2$.4H$_2$O	480 616	H.	d. ca. 100		
1541	NiS—Millerite	90 7550	Trig.	797	4 60	
1542	Ni$_3$S	149 415			5 52	
1543	Ni$_3$S$_2$	240 200		794 Tr. 545		
1544	Ni$_3$S$_4$—Polydymite	304 330	C.		4.7	
1545	NiSO$_4$	154 755			3.6$_8$	
1546	NiSO$_4$.H$_2$O	172 770			1.98	
1547	NiSO$_4$.6H$_2$O	262 847	Tet. M.	Tr. 53 3	2 07	246
1548	NiSO$_4$.7H$_2$O—Morenosite	280 863	R.		1 948	501
1549	NiS$_2$O$_6$.6H$_2$O	326 912	Tri.	d.	1.908	
1550	NiSe	137 890			8.46	
1551	NiSeO$_4$.6H$_2$O	309 982	Tet.		2 31	262
1552	Ni(NO$_3$)$_2$.6H$_2$O	290 798	M.	56.7	2.05	
1553	NH$_4$Cl.NiCl$_2$.6H$_2$O	291 195	M.		1 645	
1554	Ni(ClO$_3$)$_2$.6NH$_3$	327 793		180	1.52	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1555	$Ni(BrO_3)_2.6NH_3$	416 709		exp. 195	1.99	
1556	$Ni(IO_3)_2 5NH_3$	493 710			2.97	
1557	$(NH_4)_2Ni(SO_4)_2.6H_2O$	394 990	M.		1 923	539
1558	$(NH_4)_2Ni(SeO_4)_2.6H_2O$	489 260	M.	d.	2 22	643
1559	NiP_2	120 738			4 62[18]	
1560	NiP_2	151 762			4 19[18]	
1561	Ni_2P	148 404		1112	6.3[15]	
1562	Ni_3P_2	238 118			5 99	
1563	$Ni(H_2PO_2)_2.6H_2O$	296 861		d.	1 824	
1564	$NiAs$—Nicollite	133 650	H.	968	7.57⁰	
1565	$NiAs_2$—Rammelsbergite	208 610	R.		7 1	
1566	$Ni_{11}As_8$—Maucherite	325.990	Tet.		7 86⁰	
1567	$Ni_{12}As_2$	443 370		998		
				Tr. 970		
1568	$Ni_3(AsO_4)_2$	453 990			4 982	
1569	$3NiO.As_2O_5.8H_2O$—Annabergite	598 113	M.		3 0	845
1570	$NiAsS$ Gersdorffite	165 715			6 3	
1571	$NiSb$ Breithauptite	180 460	H.	1158	7.70⁰	
1572	Ni_5Sb_2	536 990		1170		
1573	$NiSbS$—Ullmannite	212 525	C.		6.6	
1574	NiC_2O_4	146 690			2.235	
1575	$Ni(CO)_4$	170 690		−25	l. 1.310	
1576	$3NiO.CO_2.H_2O$ Zaratite	286 085			2 6	136, 143
1577	$Ni(CHO_2)_2.2H_2O$	184 736			2.154	
1578	$Ni(C_2H_3O_2)_2.$	176 736			1.798	
1579	$Ni(C_2H_3O_2)_2.4H_2O$	248 798			1 744[167]	
1580	$NiC_{10}H_6O_6S_2.6H_2O$ 1, 5-Naphthalene disulfonate	452 959	M.		1 79	808
1581	Ni_2Si	145 440		1309	7 2[17]	
1582	$2NiO_2.3SiO_2 2H_2O$ Connarite	397 590	H.		2 5	292
1583	$NiSiF_6.6H_2O$	308 842	Trig.	d.	2 134	210
1584	$NiPdCl_6.6H_2O$	486 230	H.		2 353	
1585	$3NiO.6CuO.2As_2O_5.SO_3.7H_2O$—Landuc- kerite	1367 50	M. ?		2 2₅	851
1586	$NiPtCl_6.6H_2O$	574 760	Trig.		2 798	
1587	$NiPtBr_6.6H_2O$	841 508	Trig.		3 715	
1589	CrO_3	100 010	R.	190 d.	2 7	
1590	Cr_2O_3	152 020	H.	1900	5 21	
1591	$Cr_4O_3.3H_2O$	310 086			2 90	
1592	Cr_3O_9	404 050			4	
1593	CrF_2	90 0100		1100	4 1₁	
1594	CrF_3	109 010	R.	>1000	3 8	
1595	$CrCl_2$	122 926			2 75	
1596	$CrCl_3$	158 384			2 7	
1597	CrO_2Cl_2	154 926		− 96 5	l. 1 836	
1598	$(CrO_2)_4Cl_6$	632 798			2 5	
1599	CrS	84 0750			4 1	
1600	Cr_2S_3	200 215			3 7	
1601	$Cr_2(SO_3)_3$	344 215			2 2	
1602	$Cr_2(SO_4)_3$	392 215			3 0	
1603	$Cr_2(SO_4)_3.17H_2O$	698 476			1 7	
1604	H_2CrSO_7	198 090		190 d.		
1605	H_2CrSeO_7	245 225		200		
1606	$(NH_4)_2CrO_4$	152 088	M.		1 8	
1607	$CrO_3.3NH_3$	167 103	R.		1 96	
1608	$(NH_4)_2Cr_2O_7$	252 098	M.		2 1₅	
1609	$(NH_4)_2Cr_3O_{10}$	352 108	R.		2 33	
1610	$(NH_4)_2Cr_4O_{13}$	452 117		170	2 3₄	
1611	$NH_4IO_3.CrO_3$	292 981	R.		3 . 5	
1612	$(NH_4)_2CrSO_7$	232 153		160		
1613	$Cr_2(SO_4)_3.(NH_4)_2SO_4.24H_2O$	950 727	C.	100 d.	1 72	101
1614	CrP	83 0340			5 7	
1615	$Cr(PO_3)_3$	289 .082			2.9₇	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
21 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 60 64 3 43	25 65 29 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1616	$Cr_4(P_2O_7)_3$	730 184	M.		3 2	
1617	Cr_2As_3	328 900			6 2	
1618	$4CrO_3.As_2O_5.2(NH_4)_2O.H_2O$	752 131		d 175	1 83	
1619	Cr_3C_2 . . .	180 030		1890	6 68	
1620	Cr_4C	220 040			6 75	
1621	Cr_5C_2	284 050		1665	6 92	
1622	$CrC_2O_4.H_2O$ . .	158 025			2 46	
1623	$Cr(d-C_4H_4O_6)$	200 041			2 33¹⁰	
1624	$Cr[CH(COCH_3)_2]_3$—Acetylacetonate	349 172		211		
1625	$[Cr(CON_2H_4)_6]Cl_3.3H_2O$	572 711		150		
1626	$[Cr(CON_2H_4)_6](CN)_3.5.5H_2O$	589 400		75		
1627	$[Cr(CON_2H_4)_6](SCN)_3$	586 510		90 d.		
1628	$CrSi_2$.	108 130			4 4	
1629	Cr_2Si . . .	184 090			6 52	
1630	Cr_3Si_2	212 150			5 5	
1631	$PbCrO_4$—Crocoitite	323 210	M.	844	6 3	1000
1632	$3PbO.2CrO_3$—Phoenicochroite	869 620			5 75	
1633	$TlCr(SO_4)_2.12H_2O$	664 725	C.		2 38	122
1634	$ZnCr_2O_4$	233 400			5 3	
1635	$(NH_4)_2Cr_2O_7.HgCl_2$	523 624	M		3 11	
1636	Ag_2CrO_4 . .	331 770			5 625	
1637	$Ag_2Cr_2O_7$	431 780			4 770	
1638	$MnO.Cr_2O_3$	222 950			4 87	
1639	$FeCr_2O_4$—Chromite	223 860	C.		4 5	181
1640	$NiCr_2O_4Cl_2.9H_2O$	491 765		47		
1641	MoO_2	128 000	Tet.		4 516¹⁹ ᵇ	
1642	MoO_3	144 000	R.	795	4 50¹⁹ ᵇ	
1643	$Mo_5O_{14}.6H_2O$	812 092			3 6¹⁸	
1644	H_2MoO_4 .	162 015	H	d 115		
1645	H_4MoO_5	180 031	M Tn ?		3 124¹⁵	
1646	MoF_6	210 000		17		
1647	MoO_2F_2	166 000			3 494	
1648	$MoOF_4$ . .	188 000		98	3 001	
1649	$MoCl_5$	273 290		194		
1650	MoI_2 .	349 864			4 3	
1651	MoS_2—Molybdenite	160 130	H	1185	4 8	
1652	Mo_2S_3	288 195			5 9¹⁵	
1653	$(NH_4)_2MoO_4$	196 078	M.		2 270	
1654	$18MoO_3.14NH_3.3H_2O_2.18H_2O$	3256 76	M.		2 975	
1655	Mo_2P_2 . .	254 018			6 17	
1656	$Mo(PO_3)_3$ . . .	333 072			3 28⁰	
1658	$MoCl_5.POCl_3$	426 688		127		
1659	$18MoO_3.As_2O_5.28H_2O$	3326 35	Tri.		3 088	
1660	$18MoO_3.As_2O_5.38H_2O$	3506 51	Tri.	d.	2 822	
1661	$Bi_2O_3.MoO_3$—Koechlinite	610 000	R.			1065
1662	MoC	108 000		2570	8 40	
1663	Mo_2C	204 000		2380	8 9	
1664	$Mo(CO)_6$ . .	264 000			1 95	
1665	$3C_2H_4(NH_2)_2.HSCN.Mo(OH)(SCN)_3$	462 447		128 d.		
1666	$MoSi_2$	152 120			6 1	
1667	$TiO_2.12MoO_3.22H_2O$	2204 24	Tet.	60		
1668	$PbMoO_4$—Wulfenite	367 200	Tet.	1068	6 7	419
1669	$2PbO.MoO_3$	590 400		951		
1670	$Fe_2O_3.3MoO_3.7.5H_2O$—Molybdite	774 796	R.		4 5	919, 936, 953
1671	$WO_3.H_2O$—Tungstite . .	250 015	R.	1473	5 5 ?	1018
1672	WF_6	298 000		2 5		
1673	WOF_4	276 000		110		
1674	WCl_5	361 290		248		
1675	WCl_6	396 748		275		
1676	WO_2Cl_2	286 916				
1677	$WOCl_4$	341 832		211		
1678	WBr_6	583 580		276		

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1679	WOBr₄	519 664		27₇		
1680	WCl₄.3WBr₄	2387 24		232	6 9¹⁸	
1681	WI₂	437 864			5 2¹⁸	
1682	WI₄	691 728			7 5¹⁰	
1683	WS₂	248 130			8 5	
1684	WP	215 024			5 8	
1685	WP₂	246 048			5 21	
1686	W₄P₃	798 048			4 68	
1687	24WO₃.P₂O₅.45H₂O	6520 74	C.		6 9¹⁸	
1688	WAs₂	333 920			15 7¹⁸	
1689	WC	196 000		2777	16 06¹⁸	
1690	W₂C	380 000		2877		
1691	W₄C	564 000		>2700		
1692	WSi₂	240 120			9 3⁰	
1693	W₅Si₃	452 180			10 9	
1694	PbO.WO₃—Raspite	455 200	M.	1123		1023
1695	PbO.WO₃ Stolzite	455 200	Tet.		8 23	401
1696	CuO.WO₃ Cuprotungstite	311 570	Tet.			1007
1697	MnO WO₃ Hübnerite	302 930	M.		7 2	1017
1698	FeO.WO₃ Ferberite	303 845	Tet.		6 64	1062
1699	Fe₂O₃.WO₃.6H₂O—Ferritungstite	499 772	H.			364
1700	NiO.WO₃	306 690	R.		6 88²⁰ ⁵	
1701	3Cr₃C₂.W₂C	920 090			8 4²²	
1702	UO₂—Uraninite	270 170	R.		10 5	
1703	UO₃	286 170			5 9₂	
1704	UO₄.2H₂O	338 201		d. 11₅		
1705	U₃O₈—Pitchblende	842 510			7 3₁	
1706	UF₄	352 170	M.		4 68	
1707	(UO₂)(ClO₄)₂.4H₂O	541 148		11₀ d.		
1708	(UO₂)(ClO₄)₂.6H₂O	577 178		9₀		
1709	UBr₄	557 834			4 8₄	
1710	UI₄	745 898		50₀	5 6	
1711	UO₂(IO₃)₂	620 034	R.	d. 25₀	5 2	
1712	UO₂(IO₃)₂ H₂O	638 049			5 0₅	
1713	UO₂SO₄.3H₂O	420 281		d. 10₀	3 2₈	
1714	UO₂NO₃.6H₂O	440 270	R.	59	2 742	
1715	UO₂(NO₃)₂ 3H₂O	448 232		12₀		
1716	UO₂(NO₃)₂.6H₂O	502 278	R.	d. 10₀	2 8₁	525
1717	(NH₄)₂(UO₂)(NO₃)₄.2H₂O	590 310			2 78	
1718	(NH₄)₂(UO₂)(SO₄)₂ 2H₂O	534 408			3 0₁	
1719	UO₃.2P₂O₅	554 266	R.		3 9	
1720	3UO₂.P₂O₅.6H₂O—Phosphuranylite	1060 65	C.			906
1721	3UO₃.As₂O₅.12H₂O—Troegerite	1304 61	M.		3 3	802
1722	Bi₂O₃.2UO₂ 3H₂O—Uranosphrite	1060 39	R.		6 36	993
1723	5Bi₂O₃.3UO₃.2As₂O₅.12H₂O—Walpurgite	3816 53	Tri.		5 7₆	997
1724	UC₂	262 170		226₀	11 3¹⁸	
1725	U₂C₃	512 340		240₀	11 28	
1726	UO₃.CO₂ Rutherfordine	330 170	Tet.		5 6	935
1727	UO₂C₂O₄	358 170			2 98	
1728	UO₂(CHO₂)₂.H₂O	378 201		d. 110	3 69¹⁰	
1729	UO₂(C₂H₃O₂)₂.2H₂O	424 247	R.	d. 275	2 89¹⁵	
1730	(NH₄)₄(UO₂)(CO₃)₃.2H₂O	558 356			2 7₇	
1731	UO₂(C₂H₃O₂)₂ NH₄C₂H₃O₂	465 278	Tet.			223
1732	USi₂	294 290			8 0	
1733	12U₂O₃.5SiO₂.14H₂O Soddite	6844 60	R.		4 627	
1734	U₃Pb₂O₁₇.4H₂O—Curite	1949 31			7 1₉	
1735	8UO₃.4PbO.3P₂O₅.12H₂O Dewindtite	3824 49			4 8	
1736	UPbSiO₆.1.33H₂O—Kasolite	593 450	M.		5 9₆	
1737	Cu(UO₂)₂P₂O₈.8H₂O—Metatorbernite I	938 081	Tet.		3 ₅	303
1738	CuO.2UO₃.P₂O₅.8H₂O—Torbernite	938 081	Tet.		3.₅	737
1739	CuO.2UO₃ As₂O₅.8H₂O—Zeunerite	993.953	Tet.		3 ₂	317
1740	VO	66.9600			5 758¹⁴	
1741	VO₂	82 9600		>1755	4.399	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1742	V_2O_3	133 920			3 64	
1743	V_2O_4	149 920		1970	4 87$\frac{1}{4}$	
1744	V_2O_5	181 920		800	3 357	
1745	VF_3	107 960	R.		3 363[19]	
1746	VF_4	126 960		d. 325	2 975[13]	
1747	VF_5	145 960			2 177[19]	
1748	VOF_2	104 960		d.	3 396[19]	
1749	VOF_3	123 960		300	2 459	
1750	VCl_2	121 876	H.		3 23[18]	
1751	VCl_3	157 334			3 00[18]	
1752	VCl_4	192 792		—109	1. 1 816[39]	
1753	$VOCl$	102 418			2 824	
1754	$VOCl_2$	137 876			2 88[13]	
1755	$VOCl_3$	173 334		< —15	1. 1 829	
1756	V_2O_3Cl	201 378			3 64	
1757	$VOBr$	146 876		d. 480	4 00[18]	
1758	$VOBr_3$	306 708			2 933[14,b]	
1759	V_2S_2	166 050			4 200	
1760	V_2S_3	198 115			4 7[21]	
1761	V_2S_5	262 245			3 000	
1762	$V_2O_4.3SO_3.16H_2O$—Minasragrite	694 361	M. Tri.			619
1763	VN	64.9680		2050	5 630	
1764	$(NH_4)_3VS_4$	233 336			1 620	
1765	$(NH_4)_4V_2S_6O$	382 465			1 716	
1766	$Bi_2O_3.V_2O_5$—Pucherite	647 920	R.		6 25[74,b]	1064
1767	VC	62 9600		2830	5 4	
1768	V_4C_3	239 840		2750[smm]		
1769	$(NH_4)_2VO(CNS)_4.5H_2O$	425 407	R.	58		
1770	VSi_2	107 080			4 42	
1771	V_3Si	129 080			5 48[17]	
1772	$PbO.V_2O_5$	405.120		849		
1773	$2PbO.V_2O_5$	628 320		722		
1774	$3PbO.V_2O_5$	851 520		952		
1775	$8PbO.V_2O_5$	1907 52		794		
1776	$9PbO.3V_2O_5.PbCl_2$—Vanadinite	2832 68	H.	424	6.863	403
1777	$TlVO_3$	303 360		424		
1778	Tl_3VO_4	728 160		566		
1779	$Tl_4V_2O_7$	315 200		454		
1780	$Tl_6V_4O_{13}$	1638 24			8 59[17,b]	
1781	$4(PbZn)O.V_2O_5.H_2O$—Descloizite		R.		6 0	1021
1782	$Cd_{10}V_4Cl_2O_{24}$	1884 78	H.		5 204[16]	
1783	$Cd_{10}V_6Br_2O_{24}$	1973 69	H.		5 456[16]	
1784	$2PbO.2CuO.V_2O_5.H_2O$—Cuprodescloizite	805 475	R.		6.1	1020
1785	$Ag_4V_2O_7$	645 440		383		
1786	$5(NH_4)_2O.P_2O_5.3V_2O_5.15MoO_3.39H_2O$	3810 80			2 410	
1787	$6(NH_4)_2O.P_2O_5.6V_2O_5.12MoO_3.41H_2O$	4012 67			2 411	
1788	$3(NH_4)_2O.SiO_2.V_2O_5.9MoO_3.20H_2O$	2054 52			2 802[16]	
1789	$3(NH_4)_2O.SiO_2.V_2O_5.10MoO_3.21H_2O$	2216 54			2 804[16]	
1790	$3(NH_4)_2O.SiO_2.V_2O_5.11MoO_3.27H_2O$	2468 63	M. ?		2 807	
1791	$3(NH_4)_2O.SiO_2.V_2O_5.15MoO_3.24H_2O$	2990 58			2 816	
1792	$3(NH_4)_2O.SiO_2.V_2O_5.9WO_3.24H_2O$	2918 58			3 40	
1793	$3(NH_4)_2O.SiO_2.V_2O_5.10WO_3.21H_2O$	3096 53			3 43	
1794	$2UO_3.3V_2O_5.15H_2O$—Uvanite	1388 33	R.			979
1795	Cb_2O_5	266 200		1520	4 60$\frac{51}{4}$,[2]	
1796	CbF_5	188.100		75 5	3 29	
1797	$CbCl_5$	270 390		194	2 75	
1798	$CbOCl_3$	215 474				
1799	CbC	105 100				
1800	Cb_2FeO_6—Ferroniobite	338 040	R.		6 26	1063
1801	Ta_2O_5	443 000	R.	1470 d.	8 736$\frac{51}{4}$,[2]	
1802	TaF_5	276 500		96 8	4 74	
1803	$TaCl_5$	358 790		221	3 68[27]	
1804	$TaBr_5$	581 080		240		

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	43	47	11		82	51	61	45	1		25	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No
1805	TaC	193 500				
1806	TaSi₂	237 620			8.83°	
1807	Ta₂O₆.MnO — Manganotantalate	513 930	R.		7.03	1019
1808	B₂O₃	69 6400			l. 1.85 glass	26
1809	B₂O₃.3H₂O — Sassolite	123 686	Tri.	d.	1.49	448
1810	B₂H₆	27 6862		−169		
1811	B₄H₁₀	53 3570		−112		
1812	B₁₀H₁₄	122 308		99.5	0.94	
1813	BF₃	67 8200		−127		
1814	BCl₃	117 194		−107	l. 1.434°	
1815	BBr₃	250 568		− 45	l. 2.60	
1816	B₂HBr	102 564		−104		
1817	BI₃	391 616		43	l. 3.3⁵⁰	
1818	B₂S₃	117.835		310	1.55	
1819	BN₃	38 8360				
1820	NH₄BF₄	104 859			1.851¹⁷	
1821	CB₄	76 9200		23⁵⁰	2.6	
1822	B(CH₃)₃	55.8893		56		
1823	B(C₂H₅)₃	97.9355			l. 0 696²³	
1824	B(OCH₃)₃	103 889			l. 0 915	
1825	B(OC₂H₅)₃	145 936			l. 0.864²⁶ ᵇ	11
1826	B(OC₃H₅)₃	187 982			l. 0.867¹⁶	
1827	B(OC₄H₉)₃ Isobutyl	230 028			l. 0.864⁰	14
1828	B(OC₅H₁₁)₃ — Isoamyl	272 074			l. 0 872⁰	17
1829	SiB₃	60 5200			2 52	
1830	SiB₆	92 9800			2 47	
1831	Zr₂B₄	316 280			3 7	
1833	ThB₄	275 430			7 5	
1834	ThB₆	297 070			6 4	
1835	TlBO₂	247 220		472		
1836	Tl₃BO₃	672 020		370 d.		
1837	Tl₄B₂O₅	919 240		434		
1838	B₂O₃.CdO	198 050		87⁵		
1839	B₂O₃.CuO	149 210		d. 875	3 86	
1840	MnB₂	76 5700			6.9	
1841	Mn₃B₄O₉	352 070	Tri.		3 61	923
1842	FeB	66 6600			7.1₅	
1843	Fe₂B	122 500			7 4	
1844	FeB₂	77 4800			5 0	
1845	Fe₂B₃	165 780		13⁴⁰		
1846	Fe₆B₇	300 840		135¹		
1847	CoB	69 7900			7.2₅	
1848	Co₂B	112 740			7.9	
1849	NiB	69.5100			7.4	
1850	Ni₃B	128 200		122⁵	8.0	
1851	Ni₂B₃	197.710		116⁰		
1852	CrB	62 8300			5 ₅	
1853	Cr₃B₄	177.670			6.7¹⁵	
1854	Mo₃B₄	331 280			7	
1855	WB₂	205 640			10.8	
1857	B₂O₃.9WO₃.2NiO.18H₂O	2631 30	M.	80	l. 3 6⁸⁰	
1858	Al₂O₃ — Corundum	101.920	Trig.	205⁰	4 00	359
1859	Al₂O₃.H₂O — Diaspore	119 935	R.	d. 360	3 413	911
1860	Al₂O₃.3H₂O — Gibbsite	155 966	M.	d. 2⁰⁰	2.423	692
1861	Al(OH)₃	77 9831	M.			632
1862	AlF₃	83.9600	Tri.	104⁰	3.0₇	
1863	AlF₃.H₂O — Fluellite	101 975	R.		2 17	507
1864	AlCl₃	133 334	H.	194	2.44²⁵	
1865	AlBr₃	266 708	Trig.	97.5	l. 1.31²⁰⁰ / 3 01²⁵	
1866	AlBr₃.15H₂O	536 939		− 7 5 m	l. 2 64¹⁰⁰	
1867	Al(BrO₃)₃.9H₂O	572 847		62.3		

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1868	AlBrCl₂..............	177.792		143		
1869	AlI₃..............	407 756		191	3.98	
					l. 3 20$_4^{100}$	
1870	Al₂S₃..........	150 115	H.	110o	2 02	
1871	Al₂O₃.SO₃.9H₂O—Aluminite	344 124	M.	d.	1 705⁰	453
1872	Al₂O₃.2SO₃—Alumian...	262 050	Trig.		2.7₄	286
1873	Al₂O₃.3SO₃. ...	342 115		d. 77o	2 71	
1874	Al₂O₃.3SO₃.18H₂O—Alunogenite	630 361	M		1 691¹⁷	468
1875	2Al₂O₃.SO₃.10H₂O—Felsoebanyite	464 059	R		2 33	587
1876	2Al₂O₃.SO₃.15H₂O—Paraluminite	554 136				462
1877	AlN	40 9680	R.	215o		
1878	Al(NO₃)₃.9H₂O..	375 123	R.	73		
1879	AlCl₃.NH₄Cl..	186 831		304		
1880	AlCl₃.3NH₃	184 427		280 d.		
1881	Al₂(SO₄)₃.(NH₄)₂SO₄.	474.258			2 039	
1882	Al₂O₃.(NH₄)₂O.4SO₃.24H₂O Tschermigite...........	906 628	C.	9.₃ ₅	1 64	81
1883	AlPO₄........	121 984	H.		2 59	
1884	Al₂O₃.P₂O₅.4H₂O—Metavariscite	316 030	R.	>150o	2 54	680
1885	Al₂O₃.P₂O₅.6H₂O— Lucinite	352 060	R		2 566	724
1886	Al₂O₃.P₂O₅.6H₂O —Zepharovichite	352 060		>150o	2 37	664
1887	Al₂O₃.3P₂O₅.	528 064			2 779	
1888	2Al₂O₃.P₂O₅.3H₂O—Augelite.	399 934	M	d.	2 77	712
1889	5Al₂O₃.2P₂O₅.9H₂O—Spherite	955 835	R.	d.	2 536	711
1890	Al(AsCl)₃...	358 214			2 85²²	
1891	Al₄C₃	143 840			2 36	
1892	Al₂O₃.C₁₂O₉.18H₂O—Mellite	714 197	Tet.		1 64	260
1893	Al(CH₃)₃	72 0293				19
1894	Al(C₂H₅)₃.	114 076				20
1895	Al(C₅H₇O₂)₃— Acetylacetonate	324 122		194		
1896	Al(OC₆H₅)₃	306 076		ca. 265	1 23	
1897	NH₄(CH₃)Al(SO₄)₂.12H₂O	467 329	C.		1 508	75
1898	Al₂O₃.SiO₂—Andalusite	161 980	R.	d.	3 2	815
1899	Al₂O₃ SiO₂—Cyanite	161 980	Tri.	d.	3 6	907
1900	Al₂O₃.SiO₂—Sillimanite	161 980	R.	d. <155o	3 23	819
1901	Al₂O₃ 2SiO₂.2H₂O —Kaolinite	258 071	M		2 6	690
1902	Al₂O₃.2SiO₂.4H₂O—Newtonite	294 102	Tet		2 37	274
1903	Al₂O₃.4SiO₂.H₂O— Pyrophyllite	360 175	R.		2 8₅	727
1904	3Al₂O₃.2SiO₂— Mullite . .	425 880	R	1810 d.	3 15o	
1905	2(AlF)O.SiO₂ —Topaz .		R		3.58	784
1906	Al₃Ti₂	176 680	Tet.		3 348	
1907	3Al₂O₃.2PbO.2P₂O₅.7H₂O- P l u m b o-gummite . . .	1162 36	H	d.	4 014	325
1908	3Al₂O₃ 2PbO.2SO₃.P₂O₅.6H₂O—Hinsdalite	1162 43	H.		3 65	805
1909	2Al(OH)₃ Pb(HCO₃)₂ -Dundasite	485 182			3 25	
1910	Al₂(SO₄)₃.Tl₂SO₄.24H₂O	1279 35	C.	91	2 320	107
1911	Al₂O₃.ZnO—Automolite (Gahnite)	183 300	C.		4 58	161
1912	3Al₂O₃.2ZnO.2SO₃.18H₂O —Zincaluminite	1278 45	H.	d	2 26	256
1913	Al₂O₃.4CuO.SO₃.8H₂O —Cyanotrichite	644 388	R		2 737	779
1914	(AlCl)O.6CuO.SO₃.9H₂O—Spangolite		Trig.	d	3 14	340
1915	3Al₂O₃.CuO.2P₂O₅.9H₂O- Turquoise..	831 565	Tri	d. 300	2 67	782
1916	4Al₂O₃.18CuO.5As₂O₅.55H₂O —Liroconite	3980 39	M	d.	2 96	830
1917	Al₂O₃.MnO. . .	172 850	C		4 12	
1918	Al₂O₃.MnO.4SO₃.24H₂O—Apjohnite.	925 480	M.		1 782	477
1919	Al₂O₃.2MnO.P₂O₅.4H₂O—Eosphorite..	457 890	R.		3 1₃	837
1920	Al₂O₃.MnO.2SiO₂.2H₂O—Carpholite	329 004	R.		2 94	801
1921	Al₂O₃.3MnO.3SiO₂—Spessartite......	494 890	C.		4 180	167
1922	Al₂O₃.7MnO.8SiO₂.6H₂O—Ganophyllite.	1187 00	M.		2 84	914
1923	Al₂O₃.FeO—Hercynite	173 760	C.		3 9₃	165
1924	Al₂O₃.FeO.4SO₃.24H₂O—Halotrichite .	926 390	M.		2 04	505
1925	Al₂O₃.FeO.P₂O₅.11H₂O —Paravauxite ..	513 977	Tri.	d.	2 3	681
1926	Al₂O₃.2FeO.P₂O₅.4H₂O—Childrenite....	459 710	R.	d.	3.2₃	876

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	60		43	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1927	$2Al_2O_3.4FeO.3P_2O_5.24H_2O$—Vauxite	1349.71	Tri.		2.45	677
1928	$Al_2O_3.3FeO 3SiO_2$—Almandite	497 620	C.		4.04	166
1929	$Al_2O_3.3FeO 2SiO_2.3H_2O$—Daphnite	491 606	M.			826
1930	$5Al_2O_3.2FeO 4SiO_2.H_2O$—Staurolite	910.528	R.		3.7	930
1931	$Al_2O_3.CoO$	176.890	C.		$4 37_4^{18}$	
1932	$3Al_2O_3.4CoO$	605 640			4 80	
1933	AlB_{12}	156 800	M.		2.5	
1934	$Al_2O_3.B_2O_3$—Jeremejevite	171 560	H.		3 3	313
1935	$BO_3(AlO)_3$	187.700	R.			758
1936	$C_2B_{13}.3AlB_{12}$	624 240	Tet		$2 61_5$	
1937	$8Al_2O_3.B_2O_3 6SiO_2 H_2O$—Dumortierite	1263 38	R.		3 3	886
1938	Se_2O_3	138 200			$3 86_4$	
1939	$SeCl_4$	151 474		93_0		
1940	$SeBr_4$	284 848			3 91	
1941	$Se_2(SO_4)_3$	378.395			$2 57_9$	
1942	$Se(NO_3)_3$	231 124		15_0		
1943	$Se(NO_3)_3.4H_2O$	303 186		d. 10_0		
1944	$Se_2O_3.2SiO_2$—Thortveitite	258 320	R.		3 57	946
1945	Yt_2O_3	226 000		241_0	4 84	
1946	$YtCl_3$	195 374		$<68_6$	$2 8_4^{14}$	
1947	$YtCl_3.H_2O$	213 389		16_0		
1948	$Yt(BrO_3)_3.9H_2O$	634 887		74		
1949	$Yt_2(SO_4)_3$	466 195			$2 61_2$	
1950	$Yt_2(SO_4)_3.8H_2O$	610 318	M.		2.55_8	661
1951	$Yt_2O_4.P_2O_5$ Xenotime	368 048	Tet.		4 6	348
1952	$Yt_4(P_2O_7)_3$	878 144			3.059	
1953	YtC_2	113 000			4 13	
1954	$Yt(CH_3CO_2)_3.4H_2O$	338 131	Tri.		1 696	
1955	$Yt(C_2H_5SO_4)_3.18H_2O$	1163 90	H.		1.764_4^{25}	238
1956	$2Yt_2O_3.4SiO_2.H_2O$ Thalenite	710 255	M.		4.23	925
1957	$Yt_2Pt_3(CN)_{12}.21H_2O$	1453 90	R.		2 376	
1957 1	$Yt_2(MoO_4)_3$	658 000		1347	$4 79^{16}$	415
1958	La_2O_3	325 820		>2000	6 51	
1959	$LaCl_3$	245 284		90_7	$3 94_4^{18}$	
1960	$LaCl_3 7H_2O$	371 392		d. 91		
1961	$La(BrO_3)_3 2H_2O$	558 689		d. 15_0		
1962	$La(BrO_3)_3 9H_2O$	684 797		37 5		
1963	LaS_2	203 040		d. 65_0		
1964	La_2S_3	374 015			$4 911^{11}$	
1965	$La_2(SO_4)_3$	566 015			$3 60_0$	
1966	$La_2(SO_4)_3 9H_2O$	728 154			$2 82_1$	
1967	$(NH_4)_2La_2(SO_4)_4 8H_2O$	842 281	M.		$2 51_6$	
1968	$La_2O_3 5P_2O_5$	1036 06	M.		3.241	
1969	LaC_2	162 910			$5 0_2$	
1970	$La(C_2H_5SO_4)_6 18H_2O$	1213 81	H.		$1 845_4^{25}$	224
1971	$Tl_2La(NO_3)_5.4H_2O$	929 812		72 d.	$3 318_4^{0}$	
1972	$Zn_3La_2(NO_3)_{12} 24H_2O$	1650 43		98 0	$2 161_4^{0}$	
1973	$La_2Pt_3(CN)_{12}.18H_2O$	1499 88	M.		2 626	
1974	$Mn_3La_2(NO_3)_{12} 24H_2O$	1619 08		87 2	$2 080_4^{0}$	
1975	$Co_3La_2(NO_3)_{12} 24H_2O$	1631 20		101 8	$2 131_4^{0}$	
1976	$Ni_3La_2(NO_3)_{12}.24H_2O$	1630 36		110 5	$2 146_4^{0}$	
1976 1	$La_2(MoO_4)_3$	757 820	Tet.	1181	4.77^{14}	
1977	CeO_2	172 250	C.	195_0	7 3	
1978	CeF_3 Fluocerite	197 250	H	1324	5 8	298
1979	$CeCl_3$	246 624		84_8	$3 92_4^{0}$	
1980	$Ce(BrO_3)_3.9H_2O$	686 137	H	49		
1981	Ce_2S_3	376 695			$5 02_0^{11}$	
1982	$Ce_2(SO_4)_3$	568 695			$3 91_2$	
1983	$Ce_2(SO_4)_3 5H_2O$	658 772	M.		3 17	
1984	$Ce_2(SO_4)_3 8H_2O$	712 818	Tri.	630	$2 886^{17}$	
1985	$Ce_2(SO_4)_3.9H_2O$	730 834	H.		$2 83_1$	
1986	$Ce_2(S_4O_6)_3.15H_2O$	1031 12	Tri.		2 288	560
1987	Ce_2SeO_4	423.700	R.		4 456	748

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
1988	$(NH_4)_2Ce(NO_3)_6.4H_2O$	558.429	M.	74		
1989	$(NH_4)_2SO_4.Ce_2(SO_4)_3.8H_2O$	844.961	M.		2.52₃	
1990	$CePO_4$	235.274			5.2₂	
1991	$Ce(PO_3)_3$	377.322			3.27	
1992	CeC_2	164.250			5.2₃	
1993	$Ce(C_2H_3O_2)_3$	258.296		308 d.		
1994	$CeOF.CO_2$—Bastnäsite	219.250	H		5.0	346
1995	$Ce(C_2H_5SO_4)_6.18H_2O$	1215.15	H.		1.930_4^{16}	225
1996	$CeSi_2$	196.370			5.67^{17}	
1997	$Tl_2Ce(NO_3)_6.4H_2O$	931.152		61.5 d.	3.326_4^0	
1998	$Zn_3Ce_2(NO_3)_{12}.24H_2O$	1653.11	Trig	92.8	2.188_4^0	
1999	$Ce_2Pt_3(CN)_{12}.18H_2O$	1502.56	M		2.657	
2000	$Mn_3Ce_2(NO_3)_{12}.24H_2O$	1621.76		83.7	2.102_4^0	
2001	$Co_3Ce_2(NO_3)_{12}.24H_2O$	1633.88		98.5	2.157_4^0	
2002	$Ni_3Ce_2(NO_3)_{12}.24H_2O$	1633.04		108.5	2.173_4^0	
2002 1	$Ce_2(MoO_4)_3$	760.480	R. Tet.	973	4.83	416
2003	$Ce_2(WO_4)_3$	1024.50	Tet.	108₉	$6.77^{16.5}$	
2004	$Ce_2O_3.3Al_2O_3.2P_2O_5.6H_2O$—Florencite.	1026.45	Trig		3.59	337
2005	Pr_2O_3	329.840			6.87	
2006	Pr_4O_7	675.680			6.71₃	
2007	$Pr_{10}O_{18}$	1697.20			6.70₄	
2008	$PrCl_3$	247.294		81₈	4.020_4^{20}	
2009	$Pr(BrO_3)_3$	524.668		d. 150		
2010	$Pr(BrO_3)_3.9H_2O$	686.807	H.	56.5		
2011	Pr_2S_3	378.035			5.04_2^{11}	
2012	$Pr_2(SO_4)_3$	570.035			3.72^{016}	
2013	$Pr_2(SO_4)_3.5H_2O$	660.112	M.		3.17₃	
2014	$Pr_2(SO_4)_3.8H_2O$	714.158	M.		2.82	663
2015	$Pr_2(SeO_4)_3$	711.440			4.30^{15}	
2016	$Pr_2(SeO_4)_3.8H_2O$	855.563			$3.094^{11.5}$	
2017	PrC_2	164.920			5.1	
2018	$Pr(C_2H_5SO_4)_6.18H_2O$	1215.82	H		1.876_4^{26}	226
2019	$Zn_3Pr_2(NO_3)_{12}.24H_2O$	1654.45	Trig.	91.5	2.202_4^0	
2020	$Mn_3Pr_2(NO_3)_{12}.24H_2O$	1623.10		81.0	2.109_4^0	
2021	$Co_3Pr_2(NO_3)_{12}.24H_2O$	1635.22		97.0	2.176_4^0	
2022	$Ni_3Pr_2(NO_3)_{12}.24H_2O$	1634.38		108.0	2.195_4^0	
2023	Nd_2O_3	336.540			7.24	
2024	$NdCl_3$	250.644		78₄	4.134_4^{26}	
2025	$NdCl_3.6H_2O$	358.736		124	$2.282_4^{16.5}$	
2026	$Nd(BrO_3)_3.2H_2O$	564.049		d. 150		
2027	$Nd(BrO_3)_3.9H_2O$	690.157	H	66.7		
2028	Nd_2S_3	381.735			$5.179^{11.7}$	
2029	$Nd_2(SO_4)_3.8H_2O$	720.858	M.		2.850	668
2030	NdC_2	168.270			5.1₅	
2031	$Nd(C_2H_5SO_4)_6.18H_2O$	1219.17	H.		1.883_4^{26}	227
2032	$Zn_3Nd_2(NO_3)_{12}.24H_2O$	1661.15		88.5	2.215_4^0	
2033	$Mn_3Nd_2(NO_3)_{12}.24H_2O$	1629.80		77.0	2.114_4^0	
2034	$Co_3Nd_2(NO_3)_{12}.24H_2O$	1641.92		95.5	2.195_4^0	
2035	$Ni_3Nd_2(NO_3)_{12}.24H_2O$	1641.08		105.6	2.202_4^0	
2035 1	$Nd_2(MoO_4)_3$	768.510	Tet.	1176	5.14^{16}	414
2036	$(NdPr)_2(SO_4)_3.8H_2O$		M.			658
2037	Sa_2O_3	348.860			7.43	
2038	$SaCl_2$	221.346			3.69^{22}	
2039	$SaCl_3$	256.804		68₆	4.46_4^{18}	
2040	$SaCl_3.6H_2O$	364.896	Tri.		2.383	
2041	$SaOCl$	201.888			7.02	
2042	$SaBr_3.6H_2O$	498.270			2.971	
2043	$Sa(BrO_3)_3.2H_2O$	570.209		d. 150		
2044	$Sa(BrO_3)_3.9H_2O$	696.317	H.	75		
2045	Sa_2S_3	397.055			3.7	
2046	$Sa_2(SO_4)_3.8H_2O$	733.178	M.		2.930	670
2047	$Sa(NO_3)_3.6H_2O$	441.546	Tri.		2.375	
2048	$SaPO_4$	245.454			$5.83^{17.5}$	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Tl	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	26	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2049	SnC₂	174 430			5.8₆	
2050	Sn(CHO₂)₂	285 453			3.733	
2051	Sn(C₂H₃O₂)₂.4H₂O	399 561			1 94	
2052	Sn(C₃H₅O₂)₂	369 546			1.894	
2053	Sn(C₃H₅O₂)₂.3H₂O	423 592			1.786	
2054	Sn(C₂H₅SO₄)₂.18H₂O	1225 33	H.		1 904₄²⁵	234
2055	Zn₃Sn₂(NO₃)₁₂.24H₂O	1673 47		76 ₅	2 283₄⁰	
2056	Mn₃Sn₂(NO₄)₁₂.24H₂O	1642 12		70 ₂	2 188₄⁰	
2057	Co₃Sn₂(NO₄)₁₂.24H₂O	1654 24		83 ₂	2 237₄⁰	
2058	Ni₃Sn₂(NO₄)₁₂.24H₂O	1653 40		92 ₂	2 272₄⁰	
2059	Sn₂O₃B₂O₃	386 500			6 05	
2060	Eu₂O₃	352 000			7 42	
2061	Eu(C₂H₅SO₄)₃.18H₂O	1226 90	H.		1 909₄²⁵	239
2062	Gd₂O₃	362 520			7 40₇	
2063	GdCl₃	263 634		62₈	4 52₄⁰	
2064	GdCl₃.6H₂O	371 726			2 424₄¹⁰	
2065	GdBr₃.6H₂O	505 100			2 844¹⁵	
2066	Gd₂(SO₄)₃	602 715			4 139¹⁴ ⁶	
2067	Gd₂(SO₄)₃.8H₂O	746 838	M		3 010¹⁴ ⁶	
2068	Gd(NO₃)₃.5H₂O	433 361		92	2 406¹⁵	
2069	Gd(NO₃)₃.6H₂O	451 376	Tri	91	2 332	
2070	Gd₂(C₂O₄)₃.10H₂O	758 674	Tri	110		
2071	Gd(C₂H₃O₂)₃.4H₂O	406 391	Tri		1 611	
2072	Gd(C₂H₅SO₄)₃.18H₂O	1232 16	H.		1 919₄²⁵	235
2073	Zn₃Gd₂(NO₃)₁₂.24H₂O	1687 13		56 ₅	2 351₄⁰	
2074	Gd₂Pt₃(CN)₁₂.21H₂O	1590 63	R		2 563	
2075	Co₃Gd₂(NO₄)₁₂.24H₂O	1667 90		63 ₂	2 315₄⁰	
2076	Ni₃Gd₂(NO₄)₁₂.24H₂O	1667 06		72 ₅	2 356₄⁰	
2077	TbCl₃	265 574		58₈	4 35₄⁰	
2078	Tb(NO₃)₃.6H₂O	453 316	M	89 3		
2079	Dy₂O₃	373 040			7 81	
2080	DyCl₃	268 894		68₀	3 67₄⁰	
2081	Dy(C₂H₅SO₄)₃.18H₂O	1237 42	H.		1 492²⁵	240
2082	Er₂O₃	383 400			8 64₀	
2083	Er₂(SO₄)₃	623 595			3 67₈	
2084	Er₂(SO₄)₃.8H₂O	767 718			3 18₀	
2085	Er(C₂H₃O₂)₃.4H₂O	416 831	Tri.		2 114	
2086	Er(C₂H₅SO₄)₃.18H₂O	1242 60	H.		1 907₄²⁵	233
2087	Yb₂O₃	395 200			9 1₇	
2088	YbCl₃.6H₂O	388 066			2 57₅	
2089	Yb₂(SO₄)₃	635 395			3 79₃	
2090	Yb₂(SO₄)₃.8H₂O	779 518			3 28₆	
2091	Yb₂(SeO₄)₃	776 800			4 14₀	
2092	Yb₂(SeO₄)₃.8H₂O	920 923			3 3₀	
2093	Yb(NO₃)₃.4H₂O	431 686			2 68₂	
2094	Yb₂(CO₃)₃.4H₂O	599 262			3 67	
2095	Yb(C₂O₄)₃	437 600			2 439	
2096	Yb(C₂O₄)₃.10H₂O	617 754			2 644	
2097	Yb(C₂H₃O₂)₃.4H₂O	422 731			2 09	
2098	LuCl₃	281 374		>916	3.98	
2099	HfO₂	211 000		281₂	9 68	
2099 5	HfOCl₂.8H₂O	410 039				270.5
2099 6	(NH₄)₃HfF₇	366 034	C.			70.1
2100	BeO	25 0200	H.	240₀	3 02₅	347
2101	BeF₂	47 0200			1. 2 1¹⁵	
2102	2BeO.5BeF₂	285 140			2 3	
2103	BeCl₂	79 9360		14₀	1 89₉₄²⁵	
2104	BeBr₂	168 852		49₀		
2105	BeI₂	262 884		51₀	4 20¹⁵	
2106	BeSO₄	105 085			2 44₃	
2107	BeSO₄.4H₂O	177 147	Tet.		1 713¹⁰ ⁵	219
2108	BeSeO₄.4H₂O	224 282	R.		2.03	537
2109	Be₃N₂	55 0760		2200		

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
22 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 58 81 72

Index No.	Formula	Mol. wt	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2110	Be(NO₃)₂.3H₂O	187 082		60		
2111	Be₂C	30 0400			1 9¹⁵	
2112	Be(C₂H₅)₂	67 0970				
2113	Be(C₃H₇)₂	95 1278				
2114	Be(C₅H₇O₂)₂—Acetylacetonate	207 128	M	108	1 168⁴	
2115	BeO.3Be(C₂H₃O₂)₂	170 126		284	1 36⁴	
2116	BeO.3Be(C₃H₅O₂)(C₃H₅O₂)	448 265		127		
2117	BeO.3Be(C₃H₅O₂)₂	490 311		120		
2118	BeO.3Be(C₄H₇O₂)₂	574 403				
2119	BeO.Be(C₂H₅SO₄)₂.4H₂O	356 309	Tet			220
2120	BeO.SiO₂	85 0800		>1755		
2121	2BeO.SiO₂—Phenacite	110 100	Tri.		3 0	326
2122	4BeO.2SiO₂.H₂O—Bertrandite	238 215	R.		2 6	764
2123	BeOH.BeBO₃—Hambergite	93 8677	R.		2 35	733
2124	BeO.Al₂O₃—Chrysoberyl	126 940	R.		3 76	933
2125	3BeO.Al₂O₃.6SiO₂—Beryl.	537 340	H.	1410	2 66	284
2126	2BeO.Al₂O₃.2SiO₂.H₂O—Euclase	290 095	M		3 1	839
2127	2BeO.Yt₂O₃.FeO.2SiO₂ Gadolinite	468 000	M.		4 3	947
2128	MgO—Periclase	40 3200	C.	2800	3 65	158
2129	MgO.H₂O—Brucite	58 3354	Trig		2 4	272
2130	MgF₂—Sellaite	62 3200	Tet	1396	3 0	208
2131	MgCl₂—Chloromagnesite	95 2360	H	712	2 325	335
2132	MgCl₂ 6H₂O—Bischofite	203 328	M.	118 d	1 56	562
2133	Mg(ClO₃)₂.6H₂O	299 328		35	1 80	
2134	Mg(ClO₄)₂	223 236		d. 251	2 60⁴⁴	
2135	Mg(ClO₄)₂.6H₂O	331 328		147	1 970¹⁵	
2136	MgBr₂	184 152		700	3 72	
2137	Mg(BrO₃)₂ 6H₂O	388 244	C			117
2138	MgI₂	278 184			4 25	
2139	Mg(IO₄)₂.4H₂O	446 216	M.		3 3¹³⁵	
2140	MgS	56 3850			2 80	
2141	MgSO₄	120 385		1185	2 66	
2142	MgO SO₃.H₂O—Kieserite	138 400	M.		2 57	637
2143	MgSO₄.5H₂O	210 462	Tri.		1 718	511
2144	MgSO₄.6H₂O—Hexahydrite	228 477	M.		1 76	
2145	MgO.SO₃.7H₂O—Epsomite	246 493	R.		1 68	447
2146	MgS₂O₆ 6H₂O	292 542	Tri.		1 666	
2147	MgSeO₄.6H₂O	275 612	M.		1 928	503
2148	MgO.N₂O₅.H₂O—Nitromagnesite	166 351				558
2149	Mg(NO₄)₂.6H₂O	256 428		95	1 464	
2150	(NH₄)₂O.MgO 2SO₃.6H₂O—Boussingaultite	360 620	M.	>120	1 70	464
2151	(NH₄)₂O.MgO.2SeO₃.6H₂O	454 890	M.		2 04	568
2152	Mg₂P₂O₇	222 688			2 598⁴⁴	761
2153	2MgO.P₂O₅.7H₂O—Newberyite	348 796	R.		2 10	585
2154	3MgO.P₂O₅.8H₂O—Bobierite	407 431	M.		2 41	595
2155	Mg(H₂PO₂)₂.6H₂O	262 491	Tet		1 59¹⁵	
2156	3MgO.P₂O₅.MgF₂—Wagnerite	325 328	M.		3 12	701
2157	(NH₄)₂O.2MgO.P₂O₅.12H₂O—Struvite	490 950	R.		1 72	522
2158	3MgO.(NH₄)₂O.2P₂O₅.10H₂O—Hannayite	637 288	Tri		1 89	703
2159	3MgO.As₂O₅.8H₂O—Hoernesite	495 003	M.		2 60	702
2160	(NH₄)MgAsO₄.6H₂O	289 111			1 932¹⁵	
2161	Mg₃Sb₂	316 500		961		
2162	Mg₃Bi₂	490 960		715		
2163	MgO.CO₂—Magnesite	84 3200	Trig		3 037	342
2164	MgO.CO₂.3H₂O—Nesquehonite	138 366	R		1 850	542
2165	MgO.CO₂.5H₂O—Lansfordite	171 397	M		1 73	459
2166	2MgO.CO₂.4H₂O—Artinite	196 702	R.		2 02	630
2167	4MgO.3CO₂.4H₂O—Hydromagnesite	365 342	R		2 16	622
2168	Mg(d-C₄H₅O₅).5H₂O	262 128	M.		1 67	
2169	Mg(d-C₄H₅O₆)₂.4H₂O	394 459	R.		1 72	
2170	Mg(C₂H₃O₂)₂	142 366		323	1 42	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2171	Mg(C₂H₃O₂)₂.4H₂O	214.428	M.		1.454	512
2172	Mg(CH₃SO₃)₂.4H₂O —Ethane disulfonate	284.542	Tri.		1.727	
2173	MgC₁₀H₆O₆S₂.6H₂O —1, 5-Naphthalene disulfonate	418.589	M.		1.64	777
2174	Mg₂Si	76.7000		1102		
2175	MgO.SiO₂—Clinoenstatite.....	100.380	M.	1557 d.	3.28	836
2176	MgO.SiO₂ - Enstatite... ..	100.380	R.	d.	3.19	832
2177	2MgO.SiO₂—Forsterite	140.700	R.	1890	3.2₆	828
2178	2MgO.3SiO₂.4H₂O -Parasepiolite	332.882	R.			557
2179	3MgO.2SiO₂.2H₂O—Chrysotile ...	277.111	R.		2.5	647
2180	3MgO.3SiO₂.2H₂O -Antigorite....	337.171	R.		2.62	545
2181	3MgO.4SiO₂.H₂O - Talc ..	379.215	M.		2.7₆	728
2182	MgSiF₆.6H₂O ...	274.472	Trig.			204
2183	2MgO.SiO₂.Mg(F,OH)₂ - Prolectite ..		M.		3.1	861
2184	4MgO.2SiO₂.Mg(F,OH)₂ -Chondrodite.		M.		3.1₅	781
2185	6MgO.3SiO₂.Mg(F,OH)₂ -Humite ...		R.		3.1₅	790
2186	8MgO.4SiO₂.Mg(F,OH)₂ -Clinohumite.		M.		3.1	863
2187	MgO.TiO₂ -Geikielite	120.220	Trig.		3.98	402
2188	MgSnCl₆.6H₂O	463.860	Trig.		2.08	289
2189	2(MgPb)O.SiO₂.H₂O Molybdophyllite		H.		4.7₂	367
2190	MgCl₂.2CdCl₂.12H₂O	678.073	R.			629
2191	MgHg₂I₄.7H₂O	1313.24			3.8⁰	
2192	MgPtCl₆.6H₂O	540.390	Trig.		2.437	
2193	MgPtBr₆.12H₂O	915.231	Trig.		2.802	
2194	MgPdCl₆.6H₂O	451.860	H.		2.12	
2195	Mg₂MnCl₆.12H₂O	532.503	H.		1.802	
2196	MgO.Fe₂O₃ Magnesioferrite	200.000	C.		4.6	194
2197	MgO.Fe₂O₃.3SO₄.13H₂O - Quetenite...	674.395	M.		2.12	626
2198	2MgO.Fe₂O₃.4SO₃.15H₂O--Botryogenite.	830.811	M.		2.1	660
2199	6MgO Fe₂O₃ CO₂.12H₂O—Pyroaurite	661.785	H.		2.07	275
2200	6MgO.Fe₂O₃ CO₂ 12H₂O Brugnatellite .	661.785	H.		2.07	264
2201	3(Fe, Mg)O Fe₂O₃ 2SiO₂.3H₂O - Cronstedtite		Trig. ?		3.34	363
2202	MgO.CoO₂	131.290			5.06	
2203	Mg₂Ni₂O₃ 3SiO₂.6H₂O -Genthite	486.292	R. ?		2.5	
2204	MgCrO₄.7H₂O	266.438	R.		1.695	665
2205	MgO.Cr₂O₃	192.340			4.5₀	
2206	MgCrO₄ (NH₄)₂CrO₄.6H₂O	400.510	M		1.84	813
2207	6MgO Cr₂O₃ CO₂.12H₂O—Stichtite	654.125	H.		2.16	265
2208	MgW₂O₁₁ 8H₂O .. .	1112.44	M.			926
2209	3MgO 5V₂O₅.28H₂O	3407.09	Tri.		2.18₀	
2210	4MgO.Cb₂O₅	427.480	H		1.4	
2211	MgO B₂O₃.3H₂O Pinnoite	164.006	Tet.		2.30	277
2212	2MgO.B₂O₃.H₂O —Ascharite	168.295			2.7	666
2213	2MgO B₂O₃.H₂O -Camsellite	168.295	R. ?			1041
2214	3MgO.B₂O₃	190.600	R.		2.99	833
2215	6MgO 8B₂O₃.MgCl₂ -Boracite impure	894.276	R. C.	Tr. 265 R. to C.	2.9	856
2216	10MgO.4B₂O₃ 3H₂O -Szaibelyite	735.806			3	321
2217	6MgO.2B₂O₃ 2SO₃.9H₂O -Sulfoborite	703.469	R.		2.4	650
2218	3MgO B₂O₃.P₂O₅.8H₂O- Lueneburgite	476.771	M		2.1	649
2219	3MgO.B₂O₃.MnO Mn₂O₃--Pinakiolite	419.390	R.		3.9	999
2220	3MgO.B₂O₃.FeO Fe₂O₃— Ludwigite	422.120	R.		4.0	972
2221	4MgO.B₂O₃.Fe₂O₃ —Magnesioludwigite	390.600	R		4.0	971
2222	MgO.Al₂O₃ --Spinel	142.240	C	2135	3.6	156
2223	MgO Al₂O₃.4SO₃ 22H₂O--Pickeringite	858.839	M.		1.85	473
2224	6MgO.Al₂O₄.CO₂.12H₂O--Hydrotalcite	604.025	H.		2.06	247
2225	3MgO.Al₂O₃.3SiO₂ —Pyrope	403.060	C.		3.5	154
2226	4MgO.Al₂O₃ 2SiO₂.5H₂O -Coleramite.	473.397	H.		2.51	273
2227	5MgO.Al₂O₃.3SiO₂ 4H₂O---Leuchtenbergite	555.762	M.		2.7	726
2228	5MgO.Al₂O₃.6SiO₂.4H₂O - Zebedassite	735.942			2.19	590
2229	5MgO 6Al₂O₃ 2SiO₂—Sapphirine	933.240	M.		3.45	900
2230	(FeMg)O.Al₂O₃.P₂O₅.H₂O —Lazulite ...		M.		3.1	804

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2231	Mg₃Gd₂(NO₃)₁₂.24H₂O	1563.95	Trig.	77.5	2.163	
2232	CaO—Lime	56.0700	C.	2572	3.40	168
2233	CaH₂	42.0854		d. 675	1.7	
2234	Ca(OH)₂	74.0854	R. Trig.		2.343	318
2235	CaF₂—Fluorite	78.0700	C.	1360	3.180	71
2236	CaCl₂—Hydrophyllite	110.986	C.	772	2.152 fused	120
2237	CaCl₂.6H₂O	219.078	Trig.	29.92	1.68	212
2238	CaF₂.CaCl₂	189.056		d. 737	3.07	
2239	CaBr₂	199.902		765	3.353	
2240	CaBr₂.3H₂O	253.948	R	80.5		
2241	CaBr₂.6H₂O	307.994	H	38.2		
2242	Ca(BrO₃)₂.H₂O	313.917	M	d.	3.329	
2243	CaF₂.CaBr₂	277.972			3.15	
2244	CaI₂	293.934		575	3.956	
2245	CaI₂.6H₂O	402.026		42		
2246	Ca(IO₃)₂—Lautarite	389.934	Tri.		4.591	
2247	CaS—Oldhamite	72.1350	C		2.8	
2248	CaSO₄—Anhydrite	136.135	R. M.	Tr. 1193 (R. to M.) M. 1450	2.96	708
2249	CaSO₄.2H₂O—Gypsum	172.166	M		2.32	600
2250	CaS₂O₆.4H₂O	272.262	Trig.		2.176	269
2251	CaSeO₄	183.270			2.93	
2252	CaSeO₄.2H₂O	219.301	M.		2.676	
2253	Ca₃N₂	148.226		900	2.63	
2254	Ca(NO)₂	100.086			2.53	
2255	Ca(NO₂)₂.H₂O	150.101	H.		2.23	
2256	Ca(NO₂)₂.4H₂O	204.148			1.674	
2257	Ca(NO₃)₂—Nitrocalcite	164.086	C.	561	2.36	
2258	Ca(NO₃)₂.3H₂O	218.132		51.1		
2259	Ca(NO₃)₂.4H₂O (α)	236.148	M.	42.7	1.82	526
2260	Ca(NO₃)₂.4H₂O (β)	236.148		39.7		
2261	Ca₃P₂	182.258		>1600	2.51	
2262	CaP₂O₆	198.118		975	2.82	
2263	Ca₂P₂O₇	254.188		1230	3.09	
2264	2CaO.P₂O₅.H₂O—Monetite	272.204	Tri.	d.	2.75	586
2265	2CaO.P₂O₅.5H₂O—Brushite	344.265	M.		2.25	656
2266	Ca₃(PO₄)₂	310.258		1670	3.14	
2267	Ca₄P₂O₉	366.328	M.	1630	3.06	148
2268	4CaO.P₂O₅.5H₂O—Isoclasite	456.405	M.		2.92	698
2269	5CaO.2P₂O₅.1.5H₂O—Martinite	591.469	M. ?		2.89	765
2270	10CaO.3P₂O₅	986.844		1540	2.89	
2271	Ca(H₂PO₄)₂	234.149	Tri.	d.	2.546	
2272	Ca(H₂PO₄)₂.H₂O	252.164	Tri.	d.	2.220	
2273	CaF₂.3Ca₃P₂O₈—Fluoroapatite	1008.84	H.	1630	3.18	309
2274	Ca₅P₃ClO₁₂—Chloroapatite	520.880		1530	3.17	331
2275	3Ca₃(PO₄)₂.CaFCl—Apatite	1025.30		1270	3.11	308
2276	(NH₄)CaPO₄.7H₂O	279.241	M.	d.	1.561	
2277	Ca₃As₂	270.130			2.5	
2278	2CaO.As₂O₅.3H₂O—Haidingerite	396.106	R.		2.907	756
2279	2CaO.As₂O₅.5H₂O—Pharmacolite	432.137	M.		2.535	730
2280	2CaO.As₂O₅.8H₂O—Wapplerite	486.183	Tri.		2.48	621
2281	9CaO.3As₂O₅.CaF₂—Svabite	1272.46	H.		3.80	345
2282	5CaO.3Sb₂S₃—Romeite	1491.95	C.		5.04	169
2283	CaC₂	64.0700		2300	2.22	
2284	CaCO₃—Aragonite	100.070	R.		2.93	880
2285	CaCO₃—Calcite	100.070	H.	1339	2.711	328
2286	CaCO₃.6H₂O	208.162	M.			633
2287	CaC₂O₄	128.070			2.24	
2288	CaO.C₂O₃.H₂O—Whewellite	146.085	M.		2.23	674
2289	Ca(CHO₂)₂	130.085	R.	d.	2.015	577
2290	CaC₄H₂O₄.H₂O—Maleate	172.101	R.			706
2291	CaC₄H₂O₄.2H₂O—Fumarate	190.116	R.			754

Mg Mn Mo N	Na Nb Nd Ni O	Os P Pb Pd	Pr Pt Ra Rh	Rh Ru S Sn	Sb Se Si Sn	Sr Ta Tb Te Th	Ti Tl Tm U V	W Y Yb Zn Zr
76 42 47 11	82 51 61 45 1	35 12 23 41	60 37 80 84	40 39 8 63	14 56 9 18 22	78 52 66 10 24	19 27 70 49 50	48 57 71 23 21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No
2292	$CaC_4H_4O_5.3H_2O$ —Malate	194 147	R.			676
2293	$CaC_4H_4O_4.3H_2O$—Succinate	210 147				648
2294	$Ca(meso-C_4H_4O_6).3H_2O$.	242 147	Tri.			609
2295	$Ca(d-C_4H_4O_6).4H_2O$?	260 162	R.			638
2296	$Ca(C_3H_3O_2)_2$	158 116				683
2297	$Ca(C_3H_6O_3)_2\ 3H_2O$ —Lactate	218 147		100		
2298	$Ca(C_4H_5O_2)_2$ —Crotonate	210 147				695
2299	$CaC_4H_4O_5.6H_2O$ Acid malate	414 239	R.			561
2300	$Ca(C_6H_5CO_2)_2.3H_2O$	336 193	R.		1 436	
2301	$CaH_2(C_4H_4O_6)_2.2C_4H_6O_6$ —d-Tetratartrate	638 239	R.		1 851[19]	
2302	$Ca_3C_{12}H_6O_{12}$ Aconitate	462 256				636
2303	$Ca_3C_{12}H_{10}O_{14}.2H_2O$ - Citrate	534 318		130		
2304	$Ca_3C_{12}H_{10}O_{14}.4H_2O$ - Citrate	570 349				618
2305	$Ca(C_4H_2O_5NO_2)_2$ xH_2O Nitrotetronate		M.		1.745	822
2306	$Ca(C_9H_8NO_3)_2.3H_2O$ —Hippurate	450 255	R ?		1.318	
2307	$7CaO.CO_2\ 2P_2O_5$ —Dahllite	720 586	H.		3 08	310
2308	$10CaO\ CO_2\ 3P_2O_5$ - Podolite	1030 84	H.		3 077	807
2309	$10CaO\ CaF_2\ CO_2\ 3P_2O_5\ H_2O$ Francolite	1126 92	H.		3 1	304
2310	$CaSi$	68 1300			2 35[16]	
2311	$CaSi_2$	96 1900			2 5	
2312	Ca_3Si_2	176 330			1 64	
2313	Ca_6Si_{10}	521 020		1200		
2314	$CaSiO_3$	116 130	H.		2 89	299
2315	$CaO\ SiO_2$ -Pseudowollastonite	116 130	M.	1540		773
2316	$CaO.SiO_2$ Wollastonite	116 130	M.	Tr. 1200	2 9	800
2317	$CaO\ 2SiO_2\ H_2O$ Okenite	194 205	R.		2 3	578
2318	$2CaO\ SiO_2\ (\alpha)$	172 200	M. Tri.	2130		908
2319	$2CaO\ SiO_2\ (\beta)$	172 200	M. R	Tr. 1420 β to α		1049
2320	$2CaO.SiO_2\ (\gamma)$	172 200	M	Tr. 675 γ to β		824
2321	$2CaO.SiO_2\ H_2O$- Hillebrandite	190 215	R. ?		2 69	772
2322	$2CaO.2SiO_2.3H_2O$ Riversideite	286 306			2 61	751
2323	$3CaO.2SiO_2$	288 330	R.	1475 d.		1046
2324	$4CaO.4SiO_2\ 7H_2O$ Crestmonite	590 628			2 22	759
2325	$CaSiF_6.2H_2O$	218 161	Tet.		2 25	
2326	$3CaO.CaF_2\ 3SiO_2\ 2H_2O$ Zeophyllite	462 191	Trig		2 76	276
2327	$3CaO.CaF\ 2SiO_2\ H_2O$ - Custerite	365 415	M.		2 96	732
2328	$5CaO\ SiO_2\ P_2O_5$	482 458		1760	3 01	
2329	$3CaO.SiO_2.CO_2\ SO_3\ 15H_2O$ Thaumasite	622 566	H.		1 87	243
2330	$5CaO\ 2SiO_2\ CO_2$ Spurrite	414 170	M. ?		3 01	867
2331	$CaO.TiO_2$ Perovskite	135 970	R.		4 10	1025
2332	$CaTi(SO_4)_3$	376 165	C.			91
2333	$5CaO.2TiO_2\ 3Sb_2O_5$ -Lewisite	1410 77	C.		4 95	184
2334	$CaO.TiO_2\ SiO_2$ -Titanite	196 030	M.	1142	3 5	983
2335	$CaO\ SnO_2\ 3SiO_2\ 2H_2O$ Stokesite	422 981	R.		3 2	776
2336	$Ca_3PbC_{18}H_{30}O_{12}$ -Propionate	725 571	Tet.			251
2337	$2CaO.PbO\ 3SiO_2$	515 520			3 99	955
2338	$1CaO\ 6PbO\ 6SiO_2\ H_2O$ Ganomalite	1902 86	Tet.		5 74	985
2339	$4CaO.5PbO\ PbCl_2\ 6SiO_2$ - Nasonite	1978 76	H.		5 7	380, 384
2340	$CaO\ ZnO\ SiO_2\ H_2O$—Clinohedrite	215 525	M.		3 33	862
2341	$2CaO\ ZnO\ SiO_2$ Hardystonite	253 580	Tet.		3 4	332
2342	$CaHgI_4$	748 408			3 30[6]	
2343	$CaHg_6I_{12}\ 8H_2O$	2710 43			4 69[6]	
2344	$Ca_3Hg_4I_{14}\ 2H_2O$	3132 07			3 61[0]	
2345	$CaSO_4.3Cu(OH)_2\ CuSO_4\ 3H_2O$ — Urvolgyite	574 542	R.		3 132	
2346	$2CaO.2CuO\ As_2O_5\ H_2O$ Higginsite	519 215	R.		4 33	965
2347	$CaCu(C_2H_3O_2)_4.6H_2O$.	357 748	Tet.		1 42	213
2348	$CaPt(CN)_4.5H_2O$	429 409	R.			1045
2349	$2CaO.MnO.P_2O_5\ 2H_2O$—Fairfieldite	361.149	Tri.		3 07	823
2350	$2CaO.MnO.As_2O_5\ 2H_2O$ —Brandtite.	449 021	Tri.		3 671	902
2351	$CaO.MnO.SiO_2$—Glaucochroite.	187.060	R.		3.41	910

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2352	4CaO.2Mn₂O₃.5SiO₂.4H₂O—Orientite	912 362	R.		3 1	943
2353	4CaSiO₃.3MnSiO₃—Bustamite	857 490	Tri.			868
2354	CaO.Fe₂O₃	215 750		1216 d.		408
2355	2CaO.Fe₂O₃	271 820		1436 d		1057
2356	2CaO.FeO.P₂O₅.4H₂O—Anapaite	398 090	Tri.		2 82	778
2357	6CaO.3Fe₂O₃.4P₂O₅.19H₂O—Calcioferrite	1725 94	M.		2 53	282
2358	3CaO.2Fe₂O₃.2As₂O₅.6H₂O— Arseniosiderite	1055 50	R.		3 36	376
2359	FeCa₃(CN)₆.12H₂O	508 212	Tri.			718
2360	CaO.FeO.2SiO₂—Hedenbergite	248 030	M.	1100	3 7	922
2361	2CaO.4FeO.Fe₂O₃.4SiO₂.H₂O –Ilvaite	817 435	R.		4 0	984
2362	CaO.Cr₂O₃	208 090			4 8¹⁸	
2363	15CaO.8CrO₃.7I₂O₅— Dietzeite	397 818	M.		3 70	970
2364	3CaO.Cr₂O₃.3SiO₂—Uvarovite	500 110	C.		3 42	170
2365	CaMoO₄—Powellite	200 070	Tet.		4 35	388
2366	CaO.WO₃ —Scheelite	288 070	Tet.		6 06	381
2367	CaO.8UO₃.2SO₃ 25H₂O—Uranopilite	2505 56	Tri. ?		3 8	788
2368	CaO.2UO₃.P₂O₅ 8H₂O—Autunite	914 581	R.		3 1	707
2369	CaO.2UO₃.P₂O₅.8H₂O— Bassetite	914 581	M.		3 10	705
2370	CaO.2UO₃.As₂O₅.8H₂O—Uranospinite	1002 45	R.		3 45	719
2371	2CaO.UO₂.4CO₂.10H₂O—Uranothallite	738 464	R.		2 8	547
2372	CaO.2UO₃ 2SiO₂.6H₂O— Uranophane	856 622	Tri. ?		3 9	855
2373	CaV₆O₁₁	419 910		637		
2374	CaO.3V₂O₅.9H₂O—Hewettite	763 969	R.		2 554	1011
2375	CaO.3V₂O₅.9H₂O—Metahewettite	763 969	R.		2 51	1003
2376	2CaO.3V₂O₅.11H₂O—Pascoite	856 069	M.		2 46	961
2377	CaCl₂.Ca₃(VO₄)₂	461 116	R.		4 01	
2378	CaB₆	104 990			2 3	
2379	CaO.B₂O₃	125 710	R.	1100		841
2380	2CaO.B₂O₃	181 780		1304		
2381	2CaO.3B₂O₃.5H₂O—Colemanite	411 437	M.	d.	2 43	739
2382	2CaO.3B₂O₃.7H₂O—Meyerhofferite	447 468	Tri.	d.	2 12	635
2383	2CaO.3B₂O₃.13H₂O—Inyoite	555 260	M.	d.	1.875	570
2384	4CaO.5B₂O₃.9H₂O—Pandermite	734 619	M.	d.	2 43	738
2385	5CaO.6B₂O₃.9H₂O—Priceite	860 329	Tri.		2 4	735
2386	CaO.2SiO₂.B₂O₃—Danburite	245 830	R.		3 0	806
2387	2CaO.2SiO₂.B₂O₃.H₂O—Datolite	319 915			3 0	831
2388	4CaO 5B₂O₃ 2SiO₂.5H₂O—Howlite	782 677	M.		2 6	746
2389	8CaO.5B₂O₃.6SiO₂.6H₂O –Bakerite	1265 21			2 8	721
2390	CaO.B₂O₃.SnO₂—Nordenskioeldine	276 410	Trig.		4 2	
2391	CaO.Al₂O₃	157 990	M. ?, Tri.	1600		838
2392	3CaO.Al₂O₃	270 130	C.	1535 d		155
2393	3CaO.5Al₂O₃	677 810	Tet. ?, R.	1720		300
2394	5CaO.3Al₂O₃	586 110	C.	1455		141
2395	CaF₂.Al(F, OH)₃.H₂O—Gearksutite		M.		2 77	445
2396	CaF₂ 2Al(F, OH)₃.H₂O—Prosopite		M Tri.		2 88	548
2397	6CaO.Al₂O₃.3SO₃.33H₂O—Ettringite	1273 04	H.		1 75	231
2398	CaO.2CaF₂.2Al(F, OH)₃.SO₃ 2H₂O— Creedite		M.		2 73	470
2399	CaO.2Al₂O₃.P₂O₅.5H₂O— Crandallite	492 035	R.		3 5	294
2400	CaO.Al₂O₃.2SiO₂ —Anorthite	278 110	Tri.	1551	2 765	723
2401	CaO.Al₂O₃.2SiO₂.2H₂O –Hibschite	311 141	C.		3 05	149
2402	CaO.Al₂O₃.2SiO₂.2H₂O—Lawsonite	314 141	R.		3 09	869
2403	CaO.Al₂O₃.3SiO₂.5H₂O—Levynite	428 247	Trig.		2 1	241
2404	CaO.Al₂O₃.4SiO₂.4H₂O—Gismondite	470 292		1550	2 3	644
2405	CaO.Al₂O₃.4SiO₂.4H₂O—Laumontite	470 292	M.		2 3	605
2406	CaO.Al₂O₃.6SiO₂.5H₂O- -Epistilbite	608 427	M.		2 25	572
2407	CaO.Al₂O₃.6SiO₂.5H₂O—Heulandite	608 427	M.		2 2	528
2408	CaO.Al₂O₃.7SiO₂.7H₂O—Stellerite	704 518	R.		2.12	509
2409	CaO.2Al₂O₃ 2SiO₂.H₂O—Margarite	398 045	M.		3 0	820
2410	2CaO.Al₂O₃.SiO₂—Velardenite	274 120	Tet.	1590	3.04	333
2411	2CaO.Al₂O₃.3SiO₂.H₂O—Prehnite	412 255	R.		2.9	796
2412	2CaO.Al₂O₃.5SiO₂.6H₂O—Laubanite	622 452	M. ?		2.2	221

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Tl	Ti	Tm	U	Y		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2413	$2CaO.3Al_2O_3.9SiO_2$ —Didymolite	958.440	M.		2.71	540
2414	$3CaO.Al_2O_3.SiO_2$ ..	330.190	R.			1048
2415	$3CaO.Al_2O_3.3SiO_2$ —Grossularite	450.310	C.		3.530	157
2416	$3CaO.Al_2O_3.6SiO_2.H_2O$ —Bavenite	648.505	M.		2.72	717
2417	$4CaO.3Al_2O_3.6SiO_2$ —Meionite.	890.400	Tet.		2.74	295
2417.1	$4CaO.3Al_2O_3.6SiO_2.H_2O$ —Clinozoisite	908.415	M.		3.36	915
2418	$4CaO.3Al_2O_3.6SiO_2.H_2O$ —Zoisite	908.415	R.		3.3	896
2419	$3CaO.5Ce_2O_3.6P_2O_5.24H_2O$ —Churchite	3095.37	M.		3.14	785
2420	$CaO.2CeOF.3CO_2$ —Parisite	538.570	Trig.		4.32	279
2421	$CaPO_4.BeOH$ —Hydro-herderite	161.122	R.		3.0o	774
2422	$CaCl_2.2MgCl_2.12H_2O$ —Tachyhydrite	517.643	H.	> 168 d.	1.66s	249
2423	$2CaO.2MgO.As_2O_5.H_2O$ —Adelite	440.715	M.		3.76	909
2424	$2CaO.MgO.As_2O_5.MgF$ —Tilasite	425.700	M.		3.28	847
2425	$CaO.MgO.2CO_2$ —Dolomite	184.390	Trig.		2.872	339
2426	$CaO.MgO.SiO_2$ —Monticellite	156.450	R.	d. 1498	3.2	852
2427	$CaO.MgO.2SiO_2$ —Diopside	216.510	M.	1391	3.3	864
2428	$CaO.3MgO.2SiO_2$ —Merwinite	297.150	M.		3.15	901
2429	$CaO.3MgO.4SiO_2$ —Tremolite	417.270	M.		3.0	786
2430	$2CaO.MgO.2SiO_2$ —Åkermannite	272.580	Tet.	1458	2.944	307
2431	$5CaO.2MgO.6SiO_2$	721.350		d. 1365		797
2432	$CaO.MgO.3B_2O_3.6H_2O$ —Hydroboracite	413.402	M.		2.0	631
2433	$CaO.MgO.Al_2O_3.SiO_2$ —Gehlenite	258.370	Tet.		3.04	330
2434	SrO	103.620	R.	2430	4.7	
2435	$Sr(OH)_2$	121.635			3.625	
2436	$Sr(OH)_2.8H_2O$	265.758	Tet.		1.9o	242
2437	SrF_2	125.620	C.	1190	2.44	
2438	$SrCl_2$	158.536	C.	873	3.052	140
2439	$SrCl_2.6H_2O$	266.628	Trig.	d. 61	1.93	257
2440	$Sr(ClO_3)_2$	254.536	R.	120 d.	3.152	763
2441	$SrF_2.SrCl_2$	284.156	Tet.	962	4.18	324
2442	$SrBr_2$ . .	247.452		643	4.216_4^{24}	
2443	$SrBr_2.6H_2O$	355.544		d. 20	2.358^{18}	
2444	$Sr(BrO_3)_2.H_2O$	361.467	M.	d.	3.773	
2445	$SrBr_2.SrF_2$	373.072			4.06	
2446	SrI_2	341.484		402	4.549_4^{25}	
2447	$Sr(IO_3)_2$	437.484	Tri.		5.045^{15}	
2448	$SrI_2.SrF_2$	467.104			4.5	
2449	SrS .	119.685	C.		3.70^{15}	
2450	$SrS_4.6H_2O$	323.972		25		
2451	$SrO.SO_3$ —Celestite	183.685	R.	1580 d.	3.96	789
2452	$SrS_2O_3.5H_2O$	289.827	M.	d.	2.17^{17}	
2453	$SrS_2O_6.4H_2O$	319.812	Trig.		2.373	253
2454	$Sr(NO)_2$	147.636			2.683	
2455	$Sr(NO)_2.5H_2O$	237.713			2.173_4^{50}	
2456	$Sr(NO_2)_2$	179.636			2.867^{27}	
2457	$Sr(NO_2)_2.H_2O$	197.651		d.	2.408_0^{0}	
2458	$Sr(NO_3)_2$	211.636	C.	570	2.986	135
2459	$Sr(NO_3)_2.4H_2O$	283.698	M.		2.2	
2460	Sr_3P_2	324.908			2.68	
2461	$SrHPO_4$	183.652	R.		3.544	
2462	SrC_2	111.620			3.2	
2463	$SrO.CO_2$ —Strontianite	147.620	R.	1497^{60} at	3.70	853
2464	$Sr(CHO_2)_2$ . .	177.635	R.	71.9	2.69	704
2465	$Sr(CHO_2)_2.H_2O$	195.651	R.		2.25	
2466	$Sr(CHO_2)_2.2H_2O$	213.666	R.		2.69_5	597
2467	$Sr(C_2H_3O_2)_2$	205.666			2.099	
2468	$Sr(CH_3SO_3)_2.H_2O$ —Ethane disulfonate.	293.796	M.		2.355 (α) 2.453 (β)	
2469	$Sr(C_2H_5O_4S)_2.2H_2O$ —Ethylsulfate	373.858	M.		2.032	554
2470	$Sr(C_4H_2O_5NO_2)_2.xH_2O$ —Nitrotetronate		M.		2.043	812
2471	$Sr(SbOC_4H_4O_6)_2$	627.222	H.			426
2472	$SrSiO_3$ . .	163.680		1580	3.65	60
2473	$2SrO.SiO_3$	267.300		> 1700	3.84	

Ag	Al	As	Au	B	Ba	Be	Bi	Br	C	Ca	Cb	Cd	Ce	Cl	Co	Cr	Cs	Cu	Dy	Er	Eu	F	Fe	Ga	Gd	Ge	Gl	H	Hf	Hg	Ho	I	In	Ir	K	La	Li	Lu
32	55	13	33	54	79	75	15	5	16	77	51	39	59	4	44	46	85	31	67	60	64	3	43	25	65	20	75	2	73	30	68	6	26	36	83	56	81	72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2474	$SrSiF_6.2H_2O$.............	265.711	M.		$2\ 991^{7.8}$	
2475	$SrCl_2.2CdCl_2.7H_2O$....... . ..	651 296	M.		$2\ 718^{24}$	
2476	$SrHg_4I_{12}.8H_2O$......... .	2757.98			4.66^{5}	
2477	$Sr_2Cu(CHO_2)_6.8H_2O$..	562.964	Tri.			593
2479	$SrCrO_4$............. .	203.630	M.		$3\ 895^{15}$	
2480	$SrCr_2O_7.3H_2O$.... .	357 686	M.			905
2481	$Sr(OCrO_2Cl)_2.4H_2O$	430 618		72		
2482	$SrMoO_4$.	247 620			$4\ 145$	
2483	$SrWO_4$.	335 620			$6\ 184$	
2484	$Sr_2W_{12}SiO_{40}.16H_2O$.. . . .	3339 55	M.			934
2485	SrB_6......	152 540			$3\ 3$	
2486	$SrO.B_2O_3$.	173.260		1100		
2487	$SrO.2B_2O_3$...	242 900		930		
2488	$2SrO.B_2O_3$	276 880		1130		
2489	$2SrO.3Al_2O_3.2P_2O_5.7H_2O$—Goyazite	923.204	Trig		$3\ 2$	305
2490	$2SrO.3Al_2O_3.P_2O_5.2SO_3.6H_2O$—Svanbergite.	923 270	Trig		$3\ 5$	314
2491	$SrO.Al_2O_3.2SiO_2$...... .	325 660		>1700		
2492	$3SrO.2Ce_2O_3.7CO_2.5H_2O$—Ancylite	1365 94	R.		$3\ 95$	974
2493	$SrCa_2C_{18}H_{30}O_{12}$—Propionate	605 991	Tet.			230
2494	BaO......... .	153 370	C.	1923	$5\ 72$	
2495	BaO_2........	169 370			4.96	
2496	BaH_2........	139 385		d. 675	$4\ 21^{0}$	
2497	$Ba(OH)_2$ ...	171 385	M.		$4\ 495$	
2498	$Ba(OH)_2.8H_2O$.	315 509	M.	77.9	$2\ 1_8$	544
2499	BaF_2 .	175 370	C.	1280	$4\ 83$	
2500	$BaCl_2$.	208 286	M. C.	Tr. 925 962	$3\ 856^{7.4}$	
2501	$BaCl_2.2H_2O$	244 317	R.		$3\ 097^{7.4}$	825
2502	$Ba(ClO)_2$... .	240 286		d. 235		
2503	$Ba(ClO_2)_2$ ...	304 286		41_4		
2504	$Ba(ClO_3)_2.H_2O$	322 301	M.	d. 120	$3\ 179$	713
2505	$Ba(ClO_4)_2$	336 286		505		
2506	$Ba(ClO_4)_2.3H_2O$	390 332	H.		$2\ 74$	
2507	$BaClF$ .. .	191 828	Tet.	1008	$5\ 931$	315
2508	$BaCl_2.BaF_2$	383 656			$4\ 51^{18}$	
2509	$BaBr_2$.. .. .	297 202		847	$4\ 781^{7.4}$	
2510	$BaBr_2.2H_2O$.	333 233	M.		$3.582^{7.4}$	913
2511	$Ba(BrO_3)_2.H_2O$	411 217	M.		$3\ 99^{18}$	
2512	$BaBr_2.BaF_2$.	472 572			$4\ 96^{18}$	
2513	BaI_2	391 234		740 d.	$5\ 15_1$	
2514	$BaI_2.6H_2O$.	499 326	H.	25.7		
2515	$BaI_2.7H_2O$.	517 342			$3\ 67$	
2516	$Ba(IO_3)_2$..... .	487 234	M.		$5\ 2_3$	
2517	$Ba(IO_3)_2.H_2O$.	505 249	M.		$5\ 0^{15}$	
2518	$BaI_2.BaF_2$	566 604			$5\ 21^{18}$	
2519	BaS.... .	169 435	C.		$4\ 25^{15}$	
2520	$BaS_4\ 2H_2O$	301 661	R.	d.	$2\ 988$	
2521	$BaO.SO_3$—Barite	233.435	R.	Tr. 1149 to M. ? 1580	$4\ 499^{15}$	816
2522	$BaS_2O_3.H_2O$	267 515	R.		3.45^{18}	
2523	$BaS_2O_6.2H_2O$.	333 531	R. M.		$4\ 536^{13.5}$	744
2524	$BaS_2O_6.4H_2O$	369 562	M.		$3\ 142$	1076
2525	$BaSeO_4$... .	280 570	R.	d.	$4\ 75$	
2526	$BaTeO_4$..	328 870			$4\ 48^{16}$	
2527	BaN_6.... .	221.418	R.	d. 21_9		
2528	$Ba(NO_2)_2$.. .	197 386			3.891^{23}	
2529	$Ba(NO_2)_2$. . .	229 386		217	$3\ 23^{24}$	
2530	$Ba(NO_2)_2.H_2O$	247 401			$3\ 173^{29}$	
2531	$Ba(NO_3)_2$—Nitrobarite...	261 386	C.	592	3.244^{23}	137
2532	$Ba(NH_2)_2$	169 417		280		
2533	$Ba_2P_2O_7$	448.788	R.		4.1^{16}	
2534	$Ba_3(PO_4)_2$	602 158	C.		4.1^{16}	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh		Ru	S	Sb		Sb	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40		39	5	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2535	BaHPO₄	233 402	R.		4 165[15]	
2536	BaH₄(PO₄)₂ H₂O	285 464	M.		2 90[17]	
2537	BaF₂.3Ba₃P₂O₈	1981 84	H.	1670		334
2538	BaCl₂.3Ba₃P₂O₈	2014 76	H.	1584	5 949	343
2539	Ba₃As₂	562 030			4 1[15]	
2540	BaHAsO₄ H₂O	295 353	R M.		3.93[15]	
2541	BaC₂	161 370			3.75	
2542	BaCO₃ Witherite	197 370	R.	Tr. 811 to α	4.43	875
2543	BaCO₃ (α)	197 370	H.	Tr. 982 to β		
2544	BaCO₃ (β)	197 370		1740[90 at]		
2545	BaC₂O₄	225 370			2 658	
2546	Ba(CHO₂)₂	227 385	R.		3 21	745
2547	BaC₃H₂O₄ Malonate	239 385			2 147[18]	
2548	Ba(meso-C₄H₄O₆) H₂O	303 416			2 98	
2549	Ba(dll-C₄H₄O₆) 5H₂O	375 478	M.			1051
2550	Ba(C₂H₃O₂)₂	255 416			2 468	
2551	Ba(C₂H₃O₂)₂ H₂O	273 432	Tri.		2 19	582
2552	Ba(C₂H₃O₂)₂ 3H₂O	309 462	Tri.		2 021	
2553	Ba(C₃H₅CO₂)₂ H₂O	301 462	R.			584
2554	Ba(CH₃SO₄)₂ Ethane disulfonate	325 531	R.		2 779	
2555	BaC₆H₄O₇S₂ 1H₂O —Phenol-2, 4-disulfonate	461 592	M.			767
2556	BaC₁₀H₆O₆S₂ H₂O —Naphthalene-1, 5-disulfonate	441 562	R.		2 282	904
2557	BaSiO₃	213 430		1604	4 399	872
2558	BaSiO₃ 6H₂O	321 522	R.		2 59	659
2559	BaO 2SiO₂	273 490	R.	1420	3.73	775
2560	2BaO SiO₂	366 800		>1755		1052
2561	2BaO 3SiO₂	486 920		1450	3 93	795
2562	BaSiF₆	279 430			4 279[15]	
2563	BaO.TiO₂ 3SiO₂ Benitoite	413 450	H.		3 7	356
2564	BaCdCl₄ 4H₂O	463 674	Tri.		2 968	827
2565	BaCdBr₄ 4H₂O	641 506	Tri.		3 687	894
2566	BaCd(CHO₂)₄ 2H₂O	465 842	M.			627
2567	BaHg₄I₁₂ ...	2692 60			4 63[6]	
2568	Ba₃Hg₄I₁₆.16H₂O	3734 32			4 06	
2569	BaPtBr₆.10H₂O	992 250	M.		3 713	
2570	BaPt(CN)₄.4H₂O	508 694	M.		3 05	1047
2571	BaO.MnO₂	240 300			5 85	
2572	BaO.FeO.4SiO₂ —Gillespite	465 450	Trig.		3 33	302
2573	4BaO.FeO.2Fe₂O₃.10SiO₂ Taramellite	1605 28	R.		3 92	942
2574	BaN₁₂O₆	334 750			4 8	
2575	BaCrO₄	253 380			4 498[15]	
2576	Ba₃[Cr(C₂O₄)₃]₂	1044 13			2 57	
2577	Ba₃[Cr(C₂O₄)₃]₂ 7H₂O	1170 24			2 896[28]	
2578	Ba₃[Cr(C₂O₄)₃]₂.12H₂O	1260 31			2 372[27]	
2579	BaMoO₄.	297 370			4 65	
2580	BaWO₄	385 370			6 35	
2581	BaO 4WO₃.9H₂O	1243 51	R		4 30	
2582	Ba₄W₁₂SiO₄₀.16H₂O	3439 05	M			962
2583	BaO 2UO₃.P₂O₅.8H₂O —Uranocircite	1011 88	R.		3 53	787
2584	Ba₃V₂O₇	488 660		ca. 863		
2585	3BaO.10WO₃.V₂O₅ SiO₂ 28H₂O	3526 52			3 66	
2586	BaB₆	202 290			4 36	
2587	BaO B₂O₃..	223 010		1060		
2588	2BaO B₂O₃..	376 380		1002		
2589	3BaO B₂O₃	529 750		1315		
2590	BaCl₂ 2AlCl₃ .	474 954		290		
2591	BaO Al₂O₃ 2SiO₂ —Celsian	375 410	M.	>1700	3 37	727
2592	BaO Al₂O₃.3SiO₂ 3H₂O Edingtonite	435 470	R.		2 7	662
2593	4BaO.Al₂O₃.7SiO₂ —Barylite	1135 82	R.		4 03	884
2594	BaF₂.Ce₂O₃ 3CO₂ —Cordylite	635 870	H.		4 31	357
2595	BaO.CaO.2CO₂ —Barytocalcite	297 440	M.		3 6s	828

Ag Al As Au B Ba Be Bi Br C Ca Cb Cd Ce Cl Co Cr Cs Cu Dy Er Eu F Fe Ga Gd Ge Gl H Hf Hg Ho I In Ir K La Li Lu
47 55 13 33 54 79 75 15 5 16 77 51 29 59 4 44 46 85 31 67 60 64 3 43 25 65 20 75 2 73 30 68 6 26 36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
						73
2596	BaCa₂C₁₈H₃₀O₁₂—Propionate	655 741	C.			338
2597	BaO.2CaO.3SiO₂	445 690	H. ?	1320 d.		
2598	RaCl₂	296 866	M.	1000 Tr. 870	4 9₁	
2599	RaBr₂	385 782	M.	728	5 79	
2600	Li₂O	29 8780		>1700	2 013²⁴·ᵃ	
2601	LiH	7 94670	C.	680	0 820	
2602	LiOH	23 9467		450	2 54	
2603	LiOH.H₂O	41 9621			1 83	
2604	LiF	25 9390	C	870	2 295²¹·ᵇ / l. 1 789⁸⁷⁰	
2605	LiCl	42 3970	C.	613	2 068²⁸/₄	
2606	LiClO₃	90 3970		129		
2607	LiClO₃.0.5H₂O	99 4047		65		
2608	LiClO₄	106 397		236	2 429	
2609	LiClO₄.3H₂O	160 443	H.	95	1 841	
2610	LiBr	86 8550	C.	517	3 464²⁵ᵇ	
2611	LiBr 2H₂O	122 886		44		
2612	LiBr.3H₂O	140 901		3 5		
2613	LiI	133 871		446	4 061⁷⁵ / l. 2 827⁶²·⁵·⁴	
2614	LiI.3H₂O	187 917		73		
2615	Li₂S	45 9430			1 66	455
2616	Li₂SO₄	109 943	M.	860	2 221 / l. 2 001⁸⁶⁰	469
2617	Li₂SO₄ H₂O	127 958	M		2 06 / 2 158	684
2618	Li₂S₂O₆.2H₂O	210 039	R		2 123¹³	
2619	LiHSO₄	104 012			1 615⁰	
2620	LiNO₂.H₂O	70 9624			l. 1 77²⁷²	
2621	LiNO₃	68 9470	Trig.	255	2 38	353
2622	LiNO₃.3H₂O	122 993		d 29 6		
2623	LiNH₂	22 9624		390	1 178¹⁷ᵇ	
2624	Li₂NH	28 8937			1 303¹⁹	
2625	LiBr.NH₃	103 886		97		
2626	LiNH₄SO₄	121 043	M (α) H (β) M (γ ?)		1 204	
2627	LiPO₃	85 963			2 461 / 2 537¹⁷·ᵇ	
2628	Li₃PO₄	115 841	R	837	1 645	
2629	Li₃PO₄.12H₂O	332 026	Trig.	100	2 461	
2630	LiH₂PO₄	103 978		>100	3 07	
2631	Li₃AsO₄	159 777		>950	3 2¹⁷	
2632	Li₃Sb	142 587			1 65¹⁸	
2633	Li₂C₂	37 8780			2 111¹⁷·ᵇ	694
2634	Li₂CO₃	73 8780	M.	618	l. 1 765⁹⁰⁰ / 2 121¹⁷·ᵇ	
2635	Li₂C₂O₄	101 878			1 46	
2636	LiCHO₂.H₂O	69 9621	R.			682
2637	LiHC₄H₄O₅ 6H₂O—Malate	248 070	M	70		533
2638	LiC₂H₃O₂.2H₂O	101 993	R.		1 817	
2639	Li₂(CH₂SO₃)₂.2H₂O—Ethane disulfonate	238 070	M.			
2640	Li₂C₁₀H₆O₆S₂.2H₂O—Naphthalene 1, 5-disulfonate	336 085	M.		1 664	814
						614
2641	LiNH₄(dl-C₄H₄O₆).H₂O	191 024	M.			693
2642	LiNH₄(d-C₄H₄O₆).H₂O	191 024	R.		1 12	
2643	Li₆Si₂	97 7540		1201	l. 2 33²⁵/₄	55
2644	Li₂O.SiO₂	89 9380	R.		2 52²⁵/₄ / 2 454²¹/₄	322, 1042
2645	Li₂O.2SiO₂	149 998		1032 d. 1256	2 28	1043
2646	2Li₂O.SiO₂	119 816			2 3	
2647	Li₂SiF₆.2H₂O	191 969	M.		3.144	
2648	TlLi(dl-C₄H₄O₆).2H₂O	395 401	Tri.			

Rh 40   Ru S Sa 39 8 63   Sb Sc Se Si Sn 14 56 9 18 22   Sr Ta Tb Te Th 78 82 66 10 24   Tl Ti Tm U V 19 27 70 49 50   W Y Yb Zn Zr 48 57 71 28 21

Mg Mn Mo N 76 42 47 11   Na Nb Nd Ni O 82 51 61 45 1   Os P Pb Pd 35 12 23 41   Pr Pt Ra Rb 60 37 80 84

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2649	$2LiI.HgI_2.6H_2O$..	830.308			3.26⁰	
2650	$2LiI.HgI_2.8H_2O$....	866 339			2.95⁰	
2651	$Li_2O.2MnO.P_2O_5$—Lithiophilite	313 786	R.		3.5	878
2652	$Li_2O 2FeO.P_2O_5$—Triphylite	315 606	R.		3.55	895
2653	$Li(UO_2)(C_2H_3O_2)_3.3H_2O$	508 224	M.		2 280[15]	
2654	$Li_2O.B_2O_3$ ..	99 5180		843		
2655	$Li_2O.B_2O_3.16H_2O$	387 764	Trig.	47	1.38	
2656	$Li_2O 2B_2O_3$.	169.158		900		
2657	$Li_2O.Al_2O_3$	131.798		>1625	2.554[25] [1]	
2658	$2LiF Al_2O_3.P_2O_5$—Amblygonite	295.846	Tri.		3.05	740
2659	$Li_2O.Al_2O_3.2SiO_2$—Eucryptite	251.918	H.	1388	2.67	268
2660	$Li_2O.Al_2O_3.4SiO_2$—Spodumene.	372 038	M.	1400	3.2	854
2661	$Li_2O.Al_2O_3 5SiO_2$	432 098			2.40	
2662	$Li_2O Al_2O_3 6SiO_2$	492 158			2 41	
2663	$Li_2O.Al_2O_3.8SiO_2$—Petalite	612 278	M.	137⁰	2 4	573
2664	$2Li_2O 7Al_2O_3.2B_2O_3 6SiO_2 12H_2O$ - Manandonite.	1489 02	H.		2 89	749
2665	Na_2O	61.9940			2 27	
2666	$Na_2O_2 8H_2O$......	222 117	H.	d. 30		
2667	NaH	24 0047			0 92	
2668	$NaOH$	40 0047		318 4	2 130	
2669	$NaOH 3.5H_2O$	103.059		15.5		
2670	NaF Villiaumite	41.9970	Tet.	980	2 79	66
2671	$NaCl$ Halite...	58 4550	C.	804	2.163	129
2672	$NaOCl 2.5H_2O$.	119.494		57 5		
2673	$NaOCl.5H_2O$	164 532		24 5		
2674	$NaClO_3$	106 455	C. Trig.	248	2.490[15]	110
2675	$NaClO_4$	122 455	R.	482 d.		
2676	$NaClO_4 H_2O$..	140 470	H.	d. 130	2.02	
2677	$NaBr$	102 913	C.	75₅	3.205	
2678	$NaBr.2H_2O$......	138.944	M.	50.7	2 176	
2679	$NaBrO_3$	150 913	C.	38₁	3 339[17] [5]	138
2680	NaI	149 929	C.	651	3 667	
2681	$NaIO_3$	197.929	R.	d.	4.277	
2682	$NaIO_4$	213 929	Tet.	d. 300	3 865[16]	
2683	$NaIO_4 3H_2O$	267.975	Trig.		3 219[15]	
2684	Na_2S	78.0590			1.856	
2685	Na_2S_2	110 124		445		
2686	Na_2S_3	142 189		223 5		
2687	Na_2S_4	174 254	C.	275		
2688	$Na_2S_4.6H_2O$....	282 346		25		
2689	Na_2S_5	206 319		251 8		
2690	$Na_2SO_3.7H_2O$.	252 167	M.		1 561	
2691	$Na_2SO_4 (\alpha)$ Thenardite...	142 059	R.	Tr. 100	2 69	466
2692	Na_2SO_4	142 059	R.	Tr. 100 to M.	2 698	
			M.	Tr. 500 to H		
			H.	88₄		
2693	$Na_2SO_4.10H_2O$—Glaubers salt	322 213	M.	d. 32 4	1 464	434
2694	$Na_2SO_4.10H_2O$- Mirabilite	322 213	M.		1 48	428
2695	$Na_2S_2O_3$....	158 124	M.		1 667	
2696	$Na_2S_2O_3 5H_2O$	248 201	M.	d. 48 0	1 68₅	564
2697	$Na_2S_2O_6 2H_2O$	242 155	R.		2 189	520
2698	$NaHS.3H_2O$.	110 116	R	22		
2699	$NaHSO_4$	120 070	Tri.	>315	2 742	
2700	$2Na_2O NaCl.NaF.2SO_3$—Sulphohalite.	384.570	C.		2 49	76
2701	Na_2Se_4	362 794		−55		
2702	Na_2SeO_4.	189 194	R.		3.098	
2703	$Na_2SeO_4.10H_2O$....	369 348	M.		1 58	
2704	$NaNO_2$..... . .	69 0050	R.	271	2 168⁰	
2705	$NaNO_3$ Soda-niter	85 0050	Trig.	308	2 257	288
2706	$Na_2(NO)_2$..	106 010		300 d.	2 466[30]	
2707	$NaNH_2$	39.0204		210		
2708	$3Na_2O.N_2O_5 2SO_3.2H_2O$—Darapskite	490.159	M.		2 2	475

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
32 55 13 83	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 83 85 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2709	6NaNO₃.2Na₂SO₄.3H₂O—Nitroglauberite	848 194	R.			534
2710	NaNH₄SO₄.2H₂O—Lecontite	173 132	R.	d.	1 63	443
2711	NaPO₃	102 021		616 d.	2 476	
2712	Na₃PO₄	164 015		1340	2.537[17] ⁵	
2713	Na₃PO₄.12H₂O	380 200	Trig.	d. 73 4	1 62	214
2714	(NaPO₃)₃.2H₂O	342 094	Tri.	d.	2.476	
2715	Na₄P₂O₆.10H₂O	430 190	M.		1 832	480
2716	Na₄P₂O₇	266 036		988	2 45	
2717	Na₄P₂O₇.10H₂O	446 190	M.	d.	1 82	444
2718	NaH₂PO₂.2.5H₂O	149.075	M.	42		432
2719	NaH₂PO₄.H₂O	138 052	R.	d. 190	2 040	487
2720	NaH₂PO₄.2H₂O	156 067	R.	ca. 60	1 91	450
2721	Na₂HPO₃.5H₂O	216 103	R.			438
2722	Na₂HPO₄.2H₂O	178.057	H.		1 848	
2723	Na₂HPO₄.7H₂O	268 134	M.	d.	1.679	437
2724	Na₂HPO₄.12H₂O	358 211	R. M.	34 6	1.52	433
2725	Na₂H₂P₂O₆.6H₂O	314 150	M.		1.849	504
2726	Na₂H₂P₂O₇	222 057	M.	d. 220	1.862	
2727	Na₂H₂P₂O₇.6H₂O	330 150	M.		1.848	454
2729	Na₂HP₂O₆.9H₂O	390 185	M.	d. 100	1 743	465
2730	Na₃PO₄.H₃PO₄.15H₂O	532 293		55		
2731	Na₃PO₄.NaF.12H₂O	422 197	C.		2 216	
2732	2Na₃PO₄.NaF.19H₂O	712 320	C.		2.217	74
2733	NH₄NaHPO₄.4H₂O—Microcosmic salt, Stercorite	209 129	M.	ca 79 d.	1 574	436
2734	Na₃AsO₄	207 951			2 835	
2735	Na₃AsO₄.12H₂O	424 136	Trig.	86 3	1 759	216
2736	NaH₂AsO₄.H₂O	181 988	R.		2 535	672
2737	NaH₂AsO₄.2H₂O	200 003	R.		2 309	546
2738	Na₂HAsO₄.7H₂O	312 070	M.		1 871	556
2739	Na₂HAsO₄.12H₂O	402 147	M.	28	1 72	441
2740	2Na₂AsO₄.NaF.19H₂O	800 192	C.		2 85[25]	90
2741	Na₃AsS₄.8H₂O	416 334	M.	d.		879
2742	2Na₂O.As₂O₃.2SO₃	514 038			2 425[21]	
2743	(NH₄)NaHAsO₄.4H₂O	253 065	M.		1 845[17]	457
2744	NaSb	144 767		465		
2745	Na₃Sb	190 761		856		
2746	NaSbO₂ 3H₂O	230 813	R.	d.	2 864	
2747	Na₃SbS₄.9H₂O	481 160	C.		1 839	
2748	Na₃Bi	277 991		775		
2749	Na₂C₂	69 9940			1.575[15]	
2750	Na₂CO₃	105 994		851	2 533	
2751	Na₂CO₃.H₂O—Thermonatrite	124 009	R.		1 55	
2752	Na₂CO₃.7H₂O	232 102	R. Trig.	d. 35 1	1 51	
2753	Na₂CO₃.10H₂O—Natron	286 148	M.		1 46	431
2754	NaCHO₂	68.0017	M.	253	1 92	
2755	NaHCO₃	84 0047	M.		2 20	
2756	NaC₂H₃O₂	82 0201		324	1 528	
2757	NaC₂H₃O₂.3H₂O	136 063	M.	58; 78	1 45	452
2758	NaHC₃H₂O₄.H₂O—Acid malonate	144 036	R.			604
2759	NaH(d-C₄H₄O₆).H₂O	190 051	R.			628
2760	NaC₄H₇O₄—Diacetate	142 051	C.			79
2761	NaC₁₆H₃₁O₂—Palmitate	278 236		ca. 270		
2762	NaC₁₈H₃₃O₂—Elaidate	304 251		227		
2763	NaC₁₈H₃₃O₂—Oleate	304 251		235		
2764	Na₂(d-C₄H₄O₆).2H₂O	230.056	R.		1.818	
2765	Na₂CO₃.NaHCO₃.2H₂O—Tronite	226 030	M.		2 147[21] ⁷	563
2766	Na₃C₆H₅O₇.5H₂O—Citrate	348 107	R.		1 857[22] ⁵	
2767	NaC₁₀H₆S₂O₆.2H₂O—Naphthalene 1, 5-disulfonate	345 040	M.		1 777	809
2768	Na₂(CH₂SO₃)₂.2H₂O—Ethane disulfonate	270 186	M.		1.939 (α) 1.880 (β)	
2769	NaCN	49 0050		563 7		

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Se	Si	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2770	NaNH₄(meso-C₄H₄O₆).H₂O	207 082	M.		1.740	1074
2771	NaNH₄(d-C₄H₄O₆).4H₂O.	261 128	R.		1.587	527
2772	NaC₅H₈NO₄ – Glutamate	169 067	M.			574
2773	NaSCN	81 0700	R.	562 3		
2774	NaC₆H₄(NH₂)SO₃.2H₂O—Sulfanilate	231 147	R.			696
2775	NaC₁₀H₆NO₃S 4H₂O 1, 4-Naphthyl-amine sulfonate	317 193	M.			747
2776	Na₂O.SiO₂	122 054		1088		1040
2777	Na₂O.2SiO₂	182 114	R.	874		571
2778	Na₂SiF₆	188 054	H.		2 679	202
2779	Na₂O.3TiO₂	301 694	M.		3 5¹⁹	
2780	Na₂O.ZrO₂.6SiO₂.3H₂O —Elpidite	599 400	R.		2 58	689
2781	Na₂O.Pb(OH)Cl.SO₃—Caracolite	401 725	R.		4 5	937
2782	TlNa(dl-C₄H₄O₆).2H₂O	411 459	Tri.		3 289	
2783	TlNa(meso-C₄H₄O₆) 2 5H₂O	420 466	Tri.		3.120	
2784	TlNa(d-C₄H₄O₆).4H₂O	447 489	R.		2 580	
2785	NaTl₃(d-C₄H₄O₆)₂	932 259	R.		4 145	
2786	ZnNaPO₄	183 401	R.		3 3	
2787	Zn(Na₂PO₄)₂	347 416	C.		2 8	
2788	Na₂SO₄.CdSO₄	350 534		551		
2789	Na₂SO₄.CuSO₄.2H₂O Kroehnkite	337 725	M.		2 06¹	715
2790	Na₂SO₄.Cu(OH)₂.3CuSO₄.3H₂O Natrochalcite	772 596	M.	d. 350	2 33	840
2791	NaCu(CN)₂	138 583		d. 100	1 013	
2792	Na₃IrCl₆ 12H₂O	691 024		50		
2793	Na₂PtCl₄.4H₂O	455 118		100 d.		
2794	Na₂PtCl₆ 6H₂O	562 064	Tri.		2 50	
2795	Na₂PtBr₆ 6H₂O	828 812	Tri.		3 323	
2796	Na₂PtI₆ 6H₂O	1110 91	M. ?		3 707	
2798	Na₂Ru(NO₂)₅.2H₂O	413 765	M.		2 9	741
2799	Na₄MnP₂O₇	274 972			2 9	
2800	Na₂O.2MnO.P₂O₅ Natrophilite	345 902	R.		3 41	871
2801	Na₄Mn(PO₄)₂	336 966			2 7	
2802	Na₂O.3Fe₂O₃.4SO₃.6H₂O –Natrojarosite	969 386	R.		3 2	966
2803	2Na₂O Fe₂O₃.4SO₃.7H₂O– Sideronatrite	684 042	R.		2 2	725
2804	3Na₂SO₄ Fe₂(SO₄)₃ 6H₂O—Ferrinatrite	934 144	Trig.		2 55	271
2805	Na₆Fe₂(C₂O₄)₆ 10H₂O	957 816	M.		1 973¹⁷ ⁵	
2806	Na₂Fe(CN)₅NO 2H₂O	297 913	R.		1 7₂	
2807	Na₄Fe(CN)₆ 12H₂O	520 061	M.		1 458	616
2808	Na₂O Fe₂O₃.4SiO₂ –Aegirite	461 914	M		3 5	956
2809	Na₂O.Fe₂O₃ FeO.5SiO₄ –Riebeckite	593 814	M.		3 44	887
2810	Na₂O 2FeO Fe₂O₃ 6SiO₂ –Crocidolite	725 714	M.		3 2	893
2811	Na₂CrO₄	162 004	R.	39₂	2 723	
2812	Na₂CrO₄.4H₂O	234 066	M.	d. 64 s		
2813	Na₂CrO₄.6H₂O	270 096	Tri.	d. 25 9		
2814	Na₂CrO₄.10H₂O	342 158	M.		1 483	
2815	Na₂Cr₂O₇ 2H₂O	298 045	M	320	2 52¹⁴	892
2816	Na₂O 2CrO₃ I₂O₅.2H₂O	631 909			3 21	
2817	Na₂Cr₂S₄	278 274	H	d	2 55¹⁴	
2818	NH₄NaCrO₄ 2H₂O	193 077	R	d.	1 842¹⁵	
2819	NaCrP₂O₇	249 055	R.		3	
2820	Na₂MoO₄	205 994		687	1. 2 590¹·²⁶	
2821	Na₂Mo₂O₇	349 994		612		
2822	3Na₂O.7MoO₃ 22H₂O	1590 32	M.	ca 700		
2823	3Na₂O.5MoO₃ P₂O₅.14H₂O ...	1300 25	R.			818
2824	Na₂WO₄	293 994	R.	698	4 179 1. 3 613²⁶·⁶	
2825	Na₂WO₄.2H₂O	330 025	R.		3 245	
2826	Na₂W₂O₆	509 994			7 28	
2827	Na₂W₃O₉	741 994		d.	6 617	
2828	Na₂W₄O₁₂	973 994			7 195⁴	
2829	Na₂O.4WO₃ 10H₂O	1170 15	C.	706 6	3 847¹³	
2830	Na₂W₅O₁₅	1205 99			7 283¹⁷	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2831	$4Na_2O.10WO_3.23H_2O$	2982 33	M.	680.8	4 3	
2832	$5Na_2O.12WO_3.28H_2O$	3598 40	Tri.	705.8		
2833	$9Na_2O.22WO_3.51H_2O$	6580 73		683.3		
2834	$Na_2O.3UO_3$...	920 504	R ?		6 912	
2835	$NaU(C_2H_3O_2)_3$	438 236	Tet.		2 56	109.1
2836	$NaVO_3$ ...	121 957	M "	562	2 79	
2837	$Na_2O.V_2O_4.5V_2O_5$	1137 51	R "	ca 800 d.		
2838	Na_3VO_4 .	183 951		ca. 866		
2839	$Na_3VO_4.10H_2O$	364 105	C H			127, 263
2840	$Na_3VO_4.12H_2O$	400 136	Trig			245
2841	$Na_4V_2O_7$ ..	305 908	H	654		
2842	$2Na_3VO_4.NaF.19H_2O$	752.192	C			123
2843	$Na_4VSO_3.10H_2O$	380 170		18	1 773	
2844	$3Na_2O.V_2O_5.10WO_3.SiO_2.29H_2O$	3270 41	C.		3 344	
2845	Na_2CbO_3	187.094			4 19	
2846	$Na_2O B_2O_3$	131 634		966		
2847	$Na_2O 2B_2O_3$	201 274		711	1 2 5 glass	45
					2 37	
2848	$Na_2B_4O_7.10H_2O$ — Borax	381 428	M.	75	1 73	460
2849	$Na_2O.4B_2O_3$	340 554		783		
2850	$NaAlO_2$..	81 9570		1650		
2851	$2NaF.AlF_3$—Chiolite	167 954	Tet.		3 0	205
2852	$3NaF.AlF_3$—Cryolyte	209 950	M	1000	2 90	427
					1 2.10^{1000}	
2853	$Na_2O.Al_2O_3.4SO_3.12H_2O$ —Tamarugite	700 359	M. Tri.		2.03	494
2854	$Na_2O Al_2O_6.4SO_3 22H_2O$ Mendozite	880 513	M. ?		1 88	449
2855	$Na_2SO_4.Al_2(SO_4)_3 24H_2O$	916 514	C.	61	1 675	72
2856	$Na_2O.3Al_2O_3.4SO_3 6H_2O$— Natroalunite	796 106	Trig. C.		2 6	287
2857	$Na_2O.Al_2O_3.P_2O_5 H_2O$—Fremontite	323 977	M ?		3 04	760
2858	$Na_2O.2AlOF.As_2O_5$ –Durangite	396 834	M	4 0		866
2859	$Na_2O.Al_2O_3.2CO_2.2H_2O$ –Dawsonite	287 944	R		2 4	653
2860	$Na_2O.Al_2O_3.2SiO_2$– Carnegieite	284 034	Tri. ?	1526	2 57	596
2861	$Na_2O Al_2O_3 2SiO_2$—Nephelite	284 034	H	Tr 1248	2 67	266
2862	$Na_2O.Al_2O_3.3SiO_2.2H_2O$— Natrolite	380 125	R		2 25	478
2863	$Na_2O.Al_2O_3.4SiO_2$—Jadeite	404 154	M	1050	3 34	834
2864	$Na_2O.Al_2O_3.4SiO_2.2H_2O$—Analcite	440 185	C.		2 25	229
2865	$Na_2O.Al_2O_3.6SiO_2$—Albite....	524 274	Tri	1100	2 61	615
2866	$Na_2O.Al_2O_3.9SiO_2.2NaF$—Leifite	788 448	H.		2 57	248
2867	$Na_2O.3Al_2O_3.6SiO_2.2H_2O$—Paragonite	764 145	M		2 8	750
2868	$2Na_2O.Al_2O_3.6SiO_2.H_2O$—Ussingite	604 283	Tri.		2 50	565
2869	$2Na_2O.3Al_2O_3.6SiO_2.7H_2O$— Hydronephelite	916 216	H		2 3	236
2870	$3Na_2O 3Al_2O_3 6SiO_2.2NaCl$—Sodalite	969 012	C.		2 2	99
2871	$3Na_2O.3Al_2O_3.18SiO_2.2NaCl$— Marialite.	1689 73	Tet		2 56	261
2872	$3Na_2O.3Al_2O_3.6SiO_2.2Na_2S$ –Lazurite	1008 22	C.		2 4	108
2873	$5Na_2O 3Al_2O_3.6SiO_2.2SO_3$–Noselite.	1136 22	C.		2 3	105
2874	$Na_2La(NO_3)_5.H_2O$	512 959	M		2 63$^4_4$	
2875	$Na_2Ce(NO_3)_5.H_2O$	514 299			2 65$^7_4$	
2876	$Na_2O 2BeO P_2O_5$–Beryllonite	251 082	R.		2 85	679
2877	$Na_2O.2BeO 6SiO_2.H_2O$ –Epididymite	490 409	R		3 55	700
2878	$Na_2O.2BeO.6SiO_2.H_2O$ –Eudidymite.	490 409	M.		2 55	657
2879	$Na_2SO_4 MgSO_4$	262 444	R.		2 729	
2880	$Na_2O.MgO 2SO_3.2.5H_2O$–Loeweite	307 483	Trig.	Tr. 71	2 37	232
2881	$Na_2O MgO 2SO_3.4H_2O$ –Bloedite	334 506	M.		2 23	498
2882	$3Na_2O MgO.4SO_3$—Vanthoffite	546 562	M. ?		2 69	497
2883	$NaMgPO_4$	142 344			2 5	
2884	$Na_2MgP_2O_7$...	244 362	C. ?		2 2	
2885	$Na_2Mg(CO_3)_2$	190 314	Tet.		2 729^{16}	
2886	$NaCl Na_2CO_3.MgCO_3$—Northrupite	248 769	C.		2 377^{16}	118
2887	$3Na_2O.2MgO 4CO_2 SO_3$—Tychite	522 687	C.		2 52	113
2889	$Na_2O.CaO.2SO_3$–Glauberite	278 194	M.		2 83	625
2890	$Na_2O.CaO.2SO_3.4H_2O$—Wattevilhte	350 257	M.		1 81	446
2891	$3Na_2O.3CaO.2P_2O_5$...	638 288	M.		2 4	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2893	$Na_2O.CaO.2CO_2.2H_2O$—Pirssonite	242.095	R.	813	2.35	567
2894	$Na_2O.CaO.2CO_2.5H_2O$—Gaylussite	296.141	M.		1.94	580
2895	$Na_2O.4CaO.6SiO_2.H_2O$—Pectolite	664.650	M.		2.73	766
2896	$Na_2O.2CaO.5B_2O_3.16H_2O$ Ulexite	810.580	M.	d.	1.95	551
2897	$NaF.CaF_2.AlF_3.H_2O$ Pachnolite	222.042	M.		2.98	429
2898	$NaF.CaF_2.AlF_3.H_2O$—Thomsenolite	222.042	M.		2.98	430
2899	$Na_2O.CaO.2Al_2O_3.10SiO_2.20H_2O$ Faujasite	1282.81	C.		1.92	92
2900	$Na_2O.2CaO.3Al_2O_3.9SiO_2.8H_2O$ Mesolite	1164.56	Tri.		2.27	555
2901	$Na_2O.2CaO.3Al_2O_3.9SiO_2.8H_2O$ Pseudomesolite	1164.56	Tri.		2.22	531
2902	$5(Na_2, Ca)O.3Al_2O_3.6SiO_2.2SO_4$ Hauynite		C.		2.4	106
2903	$NaF.CaO.BeO.2SiO_2$ Leucophanite	243.207	R.		2.96	743
2904	$NaF.2CaO.2BeO.3SiO_2$ Meliphanite	384.357	Tet.		3.01	297
2905	$NaCaMgAlSi_4O_{12}$—Tuxtlite	418.587	M.		3.27	870
2906	Na_2SrSO_7	277.679		280		
2907	$Na_2Sr(CO_3)_2$	253.614		750		
2908	$Na_4SrCa(CO_3)_4$	459.678		720		
2909	$Na_2Ba(CO_3)_2$	303.364		740		
2910	$2Na_2O.BaO.2TiO_2.10SiO_2$ Leucosphenite	1037.76	M.		3.1	849
2911	$Na_4BaCa(CO_3)_4$	509.428		660		
2912	$NaLi(dl-C_4H_4O_6).2H_2O$	213.998	M.			506
2913	$3NaF.3LiF.2AlF_3$ Cryolithionite	371.728	C.		2.78	67
2914	K_2O	94.1900			2.32	
2915	K_2O_4	142.190		>280		
2916	KH	40.1027		d.	0.80	
2917	KOH	56.1027		Tr. 260	2.044	
				380	l. 1.87$_4^{130}$	
2918	KF	58.0950		880	2.48	
					l. 1.869$_4^{212}$	
2919	$KF.2HF$	98.1104		105		
2920	$KF.3HF$	118.118		100		
2921	KCl Sylvite	74.5530	C.		1.988	103
2922	$KClO_3$	122.553	M.	368.4	2.32	579
2923	$KClO_4$	138.553	R.	d. 400	2.52	
2924	KBr	119.011		730	2.75	134
2925	$KBrO_3$	167.011	Trig.	370 d.	3.27$^{17.5}$	
2926	KI	166.027	C.	773	3.123	150
2927	KI_3	419.891	M.	45	3.498	
2928	KIO_3	214.027	M.	560	3.89	
2929	KIO_4	230.027	Tet.	582	3.618	
2930	$K_2H_3IO_6.3H_2O$	358.191	Tri.			541
2931	$KICl_2$	236.943	M.	60		
2932	$KIBr_2$	325.859	R.	60		
2933	K_2S	110.255		471	1.805	
				Tr. 146.4		
2934	$K_2S.5H_2O$	200.332		60		
2935	K_2S_2	174.385		252.0		
2936	K_2S_4	206.450		>145		
2937	K_2S_5	238.515		206.0		
2938	K_2SO_4 Arcanite	174.255	R.	Tr. 588	2.662	519
				1067		
2939	$K_2S_2O_7$	190.320	C.	d. 400		
2940	$K_2S_2O_7.0.33H_2O$	196.325	M.		2.23	
2941	$K_2S_3O_6$	238.320	Trig		2.278	215
2942	$K_2S_3O_7$	254.320		>300	2.277	
2943	$K_2S_4O_6$	270.320	Tri.			458
2944	$K_2S_2O_6$	270.385	R.		2.304	472
2945	$K_2S_4O_6$	302.450	M.		2.296	
2946	$K_2S_5O_6.1.5H_2O$	361.538			2.112	

Ag Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
23 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 36	36 83 56 81 73

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
2947	KSH..............	72 1677		455		
2948	KHSO₄—Misenite.....	136 168	R. M	210	2 35	
2949	KHS₂O₇	216 233		168		
2950	K₂SO₄.KHSO₄......	310 423	M.		2 59[18]	508
2951	4K₂SO₄.3H₂SO₄.....	991 261		d. <25	2 277[18]	
2952	KSO₂F	138 160		311		
2953	KI.4SO₂ . .	422 287		0 26		
2954	K₂Se	157 390			2 851	
2955	K₂SeO₄ .	221 390	R.	3 066		646
2956	K₂SeSO₇ ...	301 455		120		
2957	K₄H₂TeI₂O₁₀.2H₂O	657 600	Trig.			307
2958	KNO₂........ ..	85.1030		297	1 915	
2959	KNO₃—Niter	101 103	R. Trig.	Tr. 129 R. to Trig 333	2 11[10 6]	556
2960	KNH₂	55 1184		338		
2961	KNO₃.2HNO₃..	227 134		22		
2962	KBr.4NH₃.... .	187 135		45		
2963	KNO₃.KHSO₄ .	237 271			2 38	
2964	5K₂O.(NH₄)₂O.6SO₃—Taylorite	1003 42				440
2965	KPO₃.	118.119		Tr. 450 810	2 258[18 6] 1 2 068[200]	
2966	K₃PO₄.. .	212 309		1340		
2967	K₄P₂O₇	330.428		Tr. 278 1090	2 33	
2968	KH₂PO₄..	136 134	Tet.	96	2 338	244
2969	K₂H₂P₂O₆.2H₂O	274 284	M.	d.		624
2970	K₂H₂P₂O₆.3H₂O	292 300	R.	d.		483
2971	KH₂AsO₄.... ..	180 070	Tet.	288	2 867	278
2972	5K₂O.As₂O₅.8SO₃.6H₂O	1449 48			2 289	
2973	KSb.	160 865		605		
2974	K₃Sb	239 055		812		
2975	K₂CO₃. . .	138 190		891	2 29	
2976	(KCO)₂ .	134 190		78		
2977	K₂C₂O₄.H₂O .	184 205	M.		2 13	486
2978	K₂O.2CO₂.H₂O—Kalicinite	200 205	M.	d. <200	2 17	476
2979	2K₂CO₃.3H₂O.......	330 426	M.		2 013	
2980	KCHO₂........ .	84 1027		167 5	1 91	
2981	KHC₂O₄........	128 103	M.		2 0	655
2982	KHC₂O₄.H₂O.....	146 118			2.044	
2983	KC₂H₃O₂.........	98 1181		292	1 8	
2984	KC₄H₅O₄—Acid succinate	156 134	M.	242 d.	1 767	
2985	KC₄H₅O₄.2H₂O—Acid succinate	192 164	R.		1 616	617
2986	KH(d-C₄H₄O₆)	188 134	R.		1 956	
2987	KH(dl-C₄H₄O₆) . . .	188 134	M.		1.954	
2988	KH(C₂H₃O₂)₂.	158 149		142		
2989	KC₆H₇O₇—Citrate .	230 149	Tri.		1 906	
2990	KC₂H₃O₂.2C₂H₄O₂..... ..	218 180		112	1 47	
2991	KHC₈H₄O₄—Acid phthalate	204 134	R.		1 636	
2992	KH(C₄H₅O₄)₂—Disuccinate	274 180	M.	162	1 56	
2993	KC₉H₇O₄.2H₂O—Acetylsalicylate	254 180		65		
2994	KC₁₈H₃₃O₂—Oleate	320 349				1037
2995	K₂C₄H₄O₄.3H₂O—Succinate.	248 267	R.		1.564	
2996	K₂(d, l-C₄H₄O₆).	226 221	M.		1.984	
2997	K₂(d-C₄H₄O₆).0.5H₂O . .	235 229	M.		1.98	610
2998	2K₂C₂O₄.H₂C₂O₄.2H₂O—Tetraoxalate.	458 426	R.		1 213[22]	592
2999	KH(CCl₂CO₂)₂.	364 851	Tet.		2 005[18]	
3000	KC₂H₅O₄S—Ethyl sulfate...... ..	164 199	M.		1.843	
3001	KC₆H₅O₄S—p-Phenolsulfonate..... ..	212 199	R.	>260	1 87	770
3002	KC₆H₅O₄S.2H₂O—o-Phenolsulfonate. .	248 229	R.		1 734	697
3003	KC₆H₄O₇S₂.H₂O—2, 4-Phenoldisulfonate	309 271	R.			768
3004	CH₂(SO₃K)₂—Methane disulfonate ..	252 335	M.		2 376	645
3005	K₂C₁₀H₆O₆S₂.2H₂O—Naphthalene 1, 5-disulfonate....................	336 397	M.		1.797	859

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Ba		Sb	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Tl	Ti	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	60		48	57	71	28	31

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
3006	KCN	65 1030		634.5	1.52[16]	
3007	KCNO	81 1030			2.048	
3008	KNH₄(d-C₄H₄O₆).0.5H₂O	214 172			1.700	
3009	KC₅H₂N₄O₄ - Acid uroxanate	253 142				1038
3010	KC₆H₂O₇N₃ Picrate	267 134	R		1.852	982
3011	KCNS	97 1680		173.2	1 886	
3012	K(SbO)(d-C₄H₄O₆)0.5H₂O Tartar emetic	333 904	R.		2 607	810
3013	K₂O SiO₂	154 250		976		
3014	K₂O 2SiO₂	214 310	R ?	1041		532
3015	K₂O 4SiO₂ H₂O	352 445	R.	d. 100	2 417	634
3016	K₂SiF₆ Hieratite	220 250	C.		2 665	
3017	K₄Ti₂O₅	253 990		980		
3017 5	K₂ZrF₆	283 190	M.			1037 2
3017 6	K₃ZrF₇	341 285	C.			68.2
3018	K₂Sn(OH)₆	298 936	Trig.		3.197	
3019	K₂SnCl₆	409 638	C.		2.71	147
3020	K₂SnBr₆	676 386			3.783	
3021	K₂SnS₃.3H₂O	347 131			1 847[15]	
3022	KPb₂Cl₅	630 785	R.	440		
3023	K₂PbCl₄	498 138	C.	d 190		
3024	KC₂H₃O₂ PbI(C₂H₃O₂)	491 273		208 5		
3025	KGa(SO₄)₂ 12H₂O	517 130			1 895	86
3026	K₃InCl₆ 2H₂O	480 864	Tet		2 483	
3027	K₃InBr₆ 2H₂O	747 612	Tet		3 140	
3028	K₃TlCl₆ 2H₂O	570 464	Tet		2 859	
3029	K₂SO₄.ZnSO₄.6H₂O	443 792	M	d. 121	2 24₅	482
3030	K₂Zn(SeO₄)₂ 2H₂O	466 001	Tri.		3 21	
3031	K₂Zn(SeO₄)₂ 6H₂O	538 062	M.		2 554	588
3032	K₂Zn(CN)₄	247 602	C	d. 150		70
3033	4KCl.CdCl₂	481 538	Trig.		2 5	293
3034	K₄Cd(NO₂)₄	374 632	R.			691
3035	CdKPO₄	246 529	R.		3.8	
3036	KCl 2HgCl₂ 2H₂O	653 636	R		4 11[15]	
3037	2KCl HgCl₂ H₂O	438 647	R		3 58[15]	877
3038	KBr.HgBr₂	479 153			4 40	
3039	KBr.HgBr₂ H₂O	497 468			3 865	
3040	KI HgI₂ H₂O	638 516		104		
3041	2KCN Hg(CN)₂	382 832	Tet.		2 447[21 3]	
3042	2KCl CuCl₂ 2H₂O	319 623	Tet		2 41	312
3043	K₂O.CuO 2SO₃ 6H₂O Cyanochroite	441 982	M		2 2₂	491
3045	K₂SeO₄ CuSeO₄.6H₂O	536 252	M		2 527	603
3046	K₂CO₃ CuCO₃	261 760			1 35[65]	
3047	K₂Cu(CN)₄	284 887	Trig.			121
3048	KNO₃ AgNO₃	270 991	M.	125	3 219	
3049	2KNO₂ AgNO₂ Bi(NO₂)₃	671 118			3 33	
3050	KAgCO₃	206 975		d.	3 769	
3051	KAuCl₄	378 127	M.	357		
3052	K₄Os(CN)₆.3H₂O	557 274	M.			769
3053	K₂IrCl₆	484 038	C.	d.	3 546	
3054	K₂SO₄ Ir₂(SO₄)₃ 24H₂O	1281 02	C	10₃		
3055	K₃Ir(C₂O₄)₃ 4H₂O	646 417	Tri.		2 510[15]	
3056	K₃IrCl₂(C₂O₄)₂ H₂O Chloroxalate	615 316	M.			736
3057	K₃IrCl₂(NO₂)₂C₂O₄ 2H₂O Dichloro dinitro oxalate	597 348	R.			716
3058	K₂PtCl₄	415 252	Tet.		3 30	
3059	K₂PtCl₆	486 168	C.	d 250	3 499	
3060	K₂PtBr₆	752 916	C.	>400 d.	4 66	
3061	K₂PtI₆	1035.01	C.		5.18	
3062	K₂S 3PtS.PtS₂	1051 50		d.	6 44[15]	
3063	[Pt(NH₃)Cl]K.H₂O	375 746	R.			709
3064	K₂Pt(NO₂)₂Br₂.H₂O	543 283	Tri.			858
3065	K₂Pt(NO₂)₂I₂ 2H₂O	655 331	Tet.			362

Ag Al As Au   B Ba Be Bi Br   C Ca Cb Cd Ce   Cl Co Cr Cs Cu   Dy Er Eu F Fe   Ga Gd Ge Gl H   Hf Hg Ho I In   Ir K La Li Lu
32 55 13 33   54 79 75 15 5   16 77 51 29 59   4 44 46 85 31   67 69 64 3 43   35 65 20 75 2   73 30 68 6 36   36 83 58 81 72

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
3066	$K_2Pt(C_2O_4)_2.2H_2O$ ..	485.451	M.		3.03	
3067	$K_2Pt(CN)_4$......	377.452	R.		2 45	
3068	$K_2Pt(NO_2)_2C_2O_4.H_2O$	471 451	M.			
3069	$K_2Pt(SCN)_6$...	621.858	H.			817
3070	$K_2Pt(SCN)_6.2H_2O$	657.889	M R		3.70[19]	
3071	$K_2Pt(SeCN)_6$.	904 668	R.	d 80	2 342[18]	
3072	$KRuO_4.H_2O$.	222.810	Tet.	d. 400 [vac]	3 378[12 3]	
3073	$K_4Ru(CN)_6.3H_2O$	468.174	M			722
3074	$K_3Rh(CN)_6$..	376 243	M			669
3075	K_2PdCl_4....	326 722			2 67	
3076	K_2PdCl_6..	397 638	C.		2 738	
3077	$KMnO_4$.. .	158 025	R.	d <240	2 703	291
3078	$K_2MnCl_4.2H_2O$.........	310 983	Tri.		2 221	
3079	K_4MnCl_6—Chloromanganokalite	424.058	Trig.		2 31	
3080	$K_2SO_4.MnSeO_4.2H_2O$.....	408 416	Tri.		3 07	
3081	$K_3Mn(CN)_6$	328 695	M.			1055
3082	$K_2Fe(SO_4)_2$.....	326.160			2 177	
3083	$K_2Fe(SO_4)_2.6H_2O$..	434.252	M.		2 169	479
3084	$K_2Fe_2(SO_4)_4.24H_2O$.	1006 50	C.	33	1 831	97
3085	$K_2O.3Fe_2O_3.4SO_3.6H_2O$—Jarosite	1001 58	R.		3 2	370
3086	$K_2Fe_2(CrO_4)_4.6H_2O$	806 342	M.		1 448[17 5]	678
3087	$K_3Fe(CN)_6$	329 173	M		1 894[17]	699
3088	$K_4Fe(CN)_6$	368 268			1 898[19]	
3089	$K_4Fe(CN)_6.3H_2O$..	422 314	M.			714
3090	$2KF.CoF_2$......	213 160	M.		3 22	
3091	$K_2SO_4.CoSO_4.6H_2O$	437 382	M		2 218	492
3092	$K_2SeO_4.CoSeO_4.6H_2O$	531 652	M.		2 514	589
3093	$[Co(NH_3)_2(NO_2)_4]K$.	316 159	R		2.076	
3094	$K_2Co(C_3H_2O_4)_2$—Malonate	341 191			2.234	
3095	$K_3Co(CN)_6$	332 303	M		1 906	
3096	$K_2SO_4.NiSO_4.6H_2O$	437 102	M	d. <100	2 237	514
3097	$K_2Ni(SeO_4)_2.6H_2O$.	531.372	M.	d <100	2 539	608
3098	$K_2Ni(COS)_4$.	377 140	M.		2.132[5 4]	125
3099	$2KCN.Ni(CN)_2 H_2O$	258 927	M.		1 871[14 5]	
3100	$K_2O.CrO_3$—Tarapacaite	194 200	R.	975	2 732[18]	927
3101	$K_2Cr_2O_7$	294 210	Tri	398	2 69	924
3102	$K_2Cr_3O_{10}$	394 220	M.	250	2 648	
3103	$K_2Cr_4O_{11}$	494 230	M.	215	2 649	
3104	$KCrClO_3$	174 563	M.	d.	2 497[19]	
3105	$K_2O 2CrO_3.I_2O_5$	628 074			3 66	
3106	K_2CrSO_7 .	274 265		350		
3107	$K_2SO_4.Cr_2(SO_4)_3.24H_2O$	998 840	C		1 83	95
3108	K_2CrSeO_7	321 400		120		
3109	$3K_2CrO_4.2(NH_4)_2CrO_4$	886 775			2 403[16]	
3110	$K_2O.Cr_2O_3.2P_2O_5$	530 306	M.		3 5[20]	
3111	$K_3Cr(CN)_6$...	325 343	M.	150 d	1.71	607
3112	$K_3Cr(SCN)_6.4H_2O$	589 795	R.		1 711[16]	
3113	$K_2Cr_2O_7.HgCl_2$	565 736	R.		3 531[19]	
3114	$K_2Cr_2O_7.Hg(CN)_2.2H_2O$	582 867	R			1077
3115	K_2MoO_4.... . . .	238 190		919	l. 2 342[m1]	
3116	K_2WO_4...	326 190	M	921	3 120[m1]	
				Tr. 388		
3117	$K_2W_2O_7$.	558 190		555		
3118	$K_2O.8WO_3$ ...	1950 19			6.53	
3119	$K_2SeO_4.Cr_2(SeO_4)_3.24H_2O$	1187 38			2 078[17 5]	
3120	$K_4U(C_2O_4)_4.5H_2O$	772 627	M		2 563	
3121	$KUO_2(C_2H_3O_2)_3.H_2O$	504 350	Tet.		2 396	
3122	$KV(SO_4)_2.12H_2O$	498 370			1.782	
3123	$K_4V_2S_4.3H_2O$	520 736			2 144	
3124	$K_2O.2UO_3.V_2O_5.8H_2O$—Carnotite	960 573	H. R.			988
3125	$3K_2O.SiO_2.V_2O_5.10WO_3.22H_2O$	3240 89	C.		3 664	
3126	$7K_2O.2SiO_2 3V_2O_5.18WO_3.42H_2O$	6257 86	M. Tri.		3.537	
3127	$NH_4K_4O_3.SiO_2.V_2O_5.10WO_3.23H_2O$.....	3237 85			3.74	

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
3128	2KF.TaF₄	392 690	R.		4.56	
3129	K₂O.B₂O₃	163 830	M.	947		
3130	KBF₄	125 915	C. R.	500 d.	2.50	
3131	KBO₂.KPO₃	200 034		872		
3132	3KF.AlF₃	258 245		1035		
				Tr. 300		
3133	K₂O.Al₂O₃.4SO₃.24H₂O — Kalinite	948 740	M. C.		1.75	77, 442
3134	K₂O.3Al₂O₃.4SO₃.6H₂O— Alunite	828 302	Trig.		2.60	281
3135	KAl(SeO₄)₂.12H₂O	568 640	C.		2 001	93
3136	K₂O.Al₂O₃.2SiO₂ —Kaliophilite	316 230	H.	>1745	2 6	258
3137	K₂O.Al₂O₃.4SiO₂ —Leucite	436 350		>1800	2.47	114
3138	K₂O.Al₂O₃.6SiO₂—Microcline	556 470	Tri.	1150	2.56	613
3139	K₂O.Al₂O₃.6SiO₂ Orthoclase	556 470	M.	1170 d.	2 56	606
3140	K₂O.3Al₂O₃.6SiO₂.2H₂O—Muscovite	796 341	M.	d.	2 9	731
3141	2Al₂O₃.3B₂O₃.K₂O—Rhodizite	506 950	C.		3 4	151
3142	K₂La(NO₃)₅.5H₂O	554 163	R.	d. 60	2 54⁰	
3143	K₂Ce(NO₃)₅.2H₂O	564 511	R.	d. 180		
3143.5	K₂HfF₆	371 19	M.			1037 1
3143 6	K₃HfF₇	429 285	C.			68.1
3144	KMgF₃	120 415			2 8	
3145	K₂MgF₄	178 510			2 7	
3146	KCl.MgCl₂.6H₂O—Carnallite	277 881	R.	167	1 60	467
3147	KI.MgI₂.6H₂O	552 303			2 547	
3148	K₂SO₄.MgSO₄.4H₂O Leonite	366 702	M.		2 25	493
3149	K₂O.MgO.2SO₃.6H₂O Picromerite	402 732	M.	d. 72	2 13	451
3150	K₂SO₄.2MgSO₄ Langbeinite	415 025	C.		2 83	128
3151	KCl.MgSO₄.3H₂O —Kainite	248.984	M.		2 13	553
3152	K₂Mg(SeO₄)₂.6H₂O	497.002	M.		2 31	527
3153	KMgPO₄	158 439	R.		2 6	
3154	K₂Mg(P₂O₆)₃	576 654	M.		2 4	
3155	KHMg(CO₃)₂.4H₂O	256 484	Tri.	d. 100	1 98	
3156	K₂Mg(CrO₄)₂.2H₂O	370 561	Tri.		2 60¹⁵	
3157	K₄.4MgO.11B₂O₃.18H₂O —Heintzeite	1345 79	M.		2 1	611
3158	KCl.CaCl₃ —Chlorocalcite	185 539	C.	754		591
3159	K₂O.CaO.2SO₃.H₂O—Syngenite	289 310	M.		2 60	581
3160	K₂CaP₂O₇	292 308	H.		2 7	
3161	K₂Ca(CO₃)₂	238 260	R.	790		
3162	K₂O.8CaO.16SiO₂.16H₂O Apophyllite	1791 96	C.		2 35	259
3163	K₂CrO₄.CaCrO₄.2H₂O	386 311	Tri.		2 502	
3164	K₂O.4CaO.2Al₂O₃.24SiO₂.H₂O—Milarite	1981 77	H.		2 57	254
3165	K₂O.2CaO.MgO.4SO₃.2H₂O —Polyhalite	602 941	R.		2 7s	685
3166	K₂SO₄.4CaSO₄.MgSO₄.2H₂O— Krugite	875 211			2 801	
3167	KCl.2SrCl₂	391 625		638		
3168	2KCl.SrCl₂	307 642	R.	597		
3169	K₂SrP₂O₇	339 858	H.		2 9	
3170	KSrCr(C₂O₄)₃.6H₂O	550 817			2 155¹²⁸	
3171	K₂Ba(CO₃)₂	335 560		800		
3172	K₄BaCa(CO₃)₄	573 820		758		
3173	LiKSO₄	142 099	H.		2 393	218
3174	2KNO₂.LiNO₂.Bi(NO₂)₃	570 177			3 21¹⁵	
3175	LiKCO₃	106 034		515		
3176	LiK(d-C₄H₄O₆).H₂O	212 080	R.			601
3177	KLi(dl-C₄H₄O₆).H₂O	212 080	M.		1 610	1075
3178	KLiPt(CN)₄.3H₂O	399 342	R.			798
3179	K₂Li₂Fe(CN)₆.3H₂O	358 002	M.			753
3180	KLiMoO₄.H₂O	224 049	R.		2 696	
3181	K₃Na(SO₄)₂—Glaserite	332 412	Trig.	<1000	2 696	237
3182	KNaHAsO₄.7H₂O	328 168			1.884	
3183	KNa(dl-C₄H₄O₆).3H₂O	264 169	M.		1.783	
3184	KNaC₄H₄O₆.4H₂O—Rochelle salt	282 184	R.		1.790	517
3185	KCl.11Na₂O.9SO₃.2CO₂—Hanksite	1565 07	H.		2.56	222
3186	3KCl.NaCl.FeCl₃—Rinneite	408 870	Trig.		2 35	290
3187	K₃Na(CrO₄)₂	372 302	Trig.		2.767	351

Index No.	Formula	Mol. wt.	Crystal system	M. P	d_4^{20}	Ref. ind. finding No.
3188	5K₂W₄O₁₃.2Na₄W₄O₁₃	7534 93				
3189	(CaK₂Na₂)O.Al₂O₃.6SiO₂.6H₂O— Erionite		R.		7 117 2 0	435
3190	Rb₂O	186 880		d. 400	3 72	
3191	Rb₂O₂	202 880			3 65	
3192	Rb₂O₃	218 880			3 53	
3193	Rb₂O₄	234 880		280	3 05⁰	
3194	RbH	86 4477		d 300	2	
3195	RbOH	102 448		300	3 203¹¹	
3196	RbF	104 440		760	1 2 88⁰²⁰	
3197	RbCl	120 898		715	2 76 1. 2 088⁴⁷⁸⁰	104
3198	RbClO₃	168 898			3 19	
3199	RbClO₄	184 898	R		2 9	
3200	RbBr	165 356	C	682	3 35 1. 2 795⁴⁷⁸⁰	133
3201	RbBr₃	325 188	R	d. 140		
3202	RbBrO₃	213 356		430	3 68	
3203	RbBrCl₂	236 272	R	d. 110		
3204	RbBr₂Cl	280 730	R.	76		
3205	RbI	212 372	C	642	3 55 1. 2 873⁴⁷⁵	146
3206	RbI₃	466 236	R	190		
3207	RbIO₃	260 372	M ?, C	d	1 33¹⁹·⁵	
3208	RbIO₄	276 372	Tet.		3 918¹⁶	
3209	RbICl₂	283 288	R	190		
3210	RbIBr₂	372 204	R.	225		
3211	RbIBrCl	327 746	R	205		
3212	Rb₂S	202 945			2 912	
3213	Rb₂S₃	267 075		213		
3214	Rb₂S₅	331 205		225	2 618¹⁵	
3215	Rb₂SO₄	266 945	R.	1060 Tr. 653	3 613 1. 2 529¹¹⁰⁰	576
3216	Rb₂S₂O₆	331 010	H.			217
3217	Rb₂S₂O₈	363 010	M.			502
3218	RbHSO₄	182 513			2 892¹⁶	
3219	RbI.4SO₂	468 632		13 5		
3220	Rb₂SeO₄	314 080	R		3 90	673
3221	RbNO₃	147.448	H	Tr. 161 4 to C.	3 11	594
			C	Tr. 219 to R.	1. 2 395⁴⁰⁰	
	,		R Tri	310		
3222	RbNO₃.HNO₃	210 464	Tet	62		
3223	RbNO₃.2HNO₃	273 479		45		
3224	Rb₂CO₃	230 880		837		
3225	RbH₃(C₂O₄)₂.2H₂O	300 494	Tri.		2 125¹⁸	
3226	Rb(dl-C₄H₅O₆)	234 479	Tri		2 282	
3227	Rb(meso-C₄H₅O₆).0.5H₂O	243 486	Tri.		2 399	
3228	RbHC₈H₄O₄—Phthalate	250 479	R		1 933	
3229	Rb₂(d-C₄H₄O₆)	318 911	Trig.		2 692	
3230	Rb₂(meso-C₄H₄O₆).H₂O	336 926	Tri		2 584	569
3231	Rb₂(meso-C₄H₄O₆).2H₂O	354 942	M.			496
3232	Rb₃C₆H₅O₇—Citrate	360 926		212 d.		
3233	RbH(CCl₃CO₂)₂	411 196	M.		2 150¹⁸	
3234	RbSCN	143 513		195		
3235	Rb₂SiF₆	312 940			3 332	
3236	RbTi(SO₄)₂.12H₂O	541 655	C.			199
3237	RbPbCl₃	399 014	R.	440		
3238	RbPb₂Cl₅	677 130	R.	423		
3239	RbGa(SO₄)₂.12H₂O	563 475	C.		1 962	87
3240	Rb₂InCl₅.H₂O	480 985	R.		3 087	
3241	Rb₂InBr₅.H₂O	703 275			3 409	
3242	RbIn(SO₄)₂.12H₂O	608 555	C.	42	2 065	83
3243	Rb₂TlCl₅.H₂O	570 585			3 513	

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	23		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

Index No.	Formula	Mol. wt	Crystal system	M. P.	d_4^{20}	Ref. ind. finding No.
3244	Rb₃TlBr₆·2H₂O.	976 247			4.077	
3245	Rb₂Zn(SO₄)₂·6H₂O	536 482	M.		2.591	499
3246	Rb₂Zn(SeO₄)₂·6H₂O	630 752	M.		2.860	598
3247	Rb₂Cd(SO₄)₂·6H₂O	583 512			2.695	485
3248	2RbCl·CuCl₂·2H₂O	412 313			2.895	
3249	Rb₂Cu(SO₄)₂·6H₂O	534 672	M.		2 57	510
3250	Rb₃AgBi(NO₂)₆	763 808			3 67¹⁵	
3251	Rb₂SO₄·Ir₂(SO₄)₄·24H₂O	1373 71	C	109		
3253	RbRh(SO₄)₂·12H₂O	596 665	C			109
3254	RbMnO₄	204 370			3 235¹⁰ ⁴	
3255	Rb₂Mn(SO₄)₂·6H₂O	526 632	M		2 46	474
3256	RbFeCl₄·2H₂O	283 685			2 711	
3257	Rb₂FeCl₄·2H₂O	404 583			2 850	
3258	Rb₂Fe(SO₄)₂·6H₂O	526 912	M.		2 518	405
3259	RbFe(SO₄)₂·12H₂O	549 595	C.		1 92	98
3260	Rb₂FeSe₂O₆·6H₂O	621 212			2 819	
3261	Rb₂SeO₄·Fe₂(SeO₄)₃·24H₂O	1287 73	C	45	2 131¹⁵	111
3262	Rb₂Co(SO₄)₂·6H₂O	530 072	M		2 567	515
3263	Rb₂Co(C₃H₂O₄)₂·4H₂O--Malonate	505 942			2 131	
3264	Rb₂SO₄·NiSO₄·6H₂O	529 792	M.		2 586	523
3265	Rb₂SO₄·Cr₂(SO₄)₃·24H₂O	1091 53	C.	107	1 946	96
3266	RbV(SO₄)₂·12H₂O	544 715			1.915⁴	
3267	3RbF·AlF₃	397 280		985		
3268	Rb₂SO₄·Al₂(SO₄)₃·24H₂O	1041 43	C.		1.867⁰	78
3269	Rb₃La(NO₃)₅·4H₂O	691 892	M.	86	2 497⁰	
3270	Rb₂Ce(NO₃)₅·4H₂O	693 232	M.	70	2 497⁰	
3271	Rb₂Pr(NO₃)₅·4H₂O	693 902		63 5	2 50⁰	
3272	Rb₂Nd(NO₃)₅·4H₂O	697 252		47	2 56⁰	
3273	Rb₂Mg(SO₄)₂·6H₂O	495 422	M.		2 40	461
3274	Rb₂Mg(SeO₄)₂·6H₂O	589 692	M		2 684	549
3275	Rb₂Mg(CrO₄)₂·6H₂O	535 312	M		2 466	805
3276	RbLa(d-C₄H₄O₆)·H₂O	258 425	R		2 281	671
3277	RbNa(meso-C₄H₄O₆)·2.5H₂O	301 506	Tn		2 20	
3278	Cs₂O	281 620			4 36	
3279	Cs₂O₃	313 620		400	4 25⁰	
3280	Cs₂O₄	329 620		600· 515 (in O₂)	3 68⁰	
3281	CsH	133 818			2 7	
3282	CsOH	149 818		Tr. 223 272 3	3 675	
3283	CsF	151 810		683	3 586⁷⁵⁰, l. 2 549	
3284	CsCl	168 268	C	646	3 97, l. 2 732⁷⁰⁰	144
3285	CsClO₃	216 268			3 57¹⁹ ⁵	
3286	CsClO₄	232 268			3 327	
3287	CsBr	212 726	C.	636	4 44, l. 3 038⁷⁰⁰	152
3288	CsBr₃	372 558	R	180		
3289	CsBrO₃	260 726		420	4 10¹⁹ ⁵	
3290	CsBrCl₂	283 642		205		
3291	CsBr₂Cl	328 100		191		
3292	CsI	259 742	C	621	4 51, l. 3 114⁶⁹⁹	163
3293	CsI₃	513 606	R.	207 5		
3294	CsIO₃	307 742	M.		4 85	
3295	CsIO₄	323 742	R		4 259	
3296	CsICl₂	330 658	R.	230	3 86	
3297	CsIBr₂	419 574		248		
3298	CsI₂Br	466 590		195 5		
3299	CsIBrCl	375 116		235		
3300	Cs₂S₂	329 750		460		
3301	Cs₂S₃	361.815		217		

Index No.	Formula	Mol. wt.	Crystal system	M. P.	d_4^0	Ref. ind. finding No.
3302	Cs₂S₄	393.880		160		
3303	Cs₂S₅	425.945		210	2.806¹⁴	
3304	Cs₂S₆	458.010		186		
3305	Cs₂SO₄	361.685	R.	Tr. 660 to H. 10 to	4.243 / 3.034¹⁰⁴⁰	687
3306	CsHSO₄	229.883	R.	d.	3.352¹⁰	
3307	Cs₂SeO₄	408.820	R			752
3308	Cs₂(SeO₄)₂	552.020	R		4.453	
3309	CsN₃	174.834		315		
3310	CsNO₃	191.818	H	Tr. 161 to C. 414	3.685 / 2.713₄⁰⁰	
3311	CsNH₂	148.833		260		
3312	CsNO₃.HNO₃	257.834		100		
3313	CsNO₃.2HNO₃	320.849		35		
3314	CsHC₈H₄O₄—Phthalate	297.849	R		2.178	
3315	CsH(CCl₃CO₂)₂ . .	458.566	M		2.143	
3316	Cs₂SiF₆ . . .	407.680			3.372¹⁰	
3317	CsGa(SO₄)₂.12H₂O	610.845	C.		2.113	84
3318	Cs₂InCl₅.H₂O .	575.725			3.350	
3319	Cs₂InBr₅.H₂O	798.015			3.776	
3320	CsIn(SO₄)₂.12H₂O	655.925	C.		2.241	85
3321	Cs₂TlCl₅.H₂O	665.325			3.879	
3322	Cs₃Tl₂Cl₉ .	1126.35	H.			361
3323	Cs₂Zn(SO₄)₂.6H₂O	631.222	M.		2.875	552
3324	Cs₂Zn(SeO₄)₂.6H₂O	725.492	M.		3.115	640
3325	Cs₂Cd(SO₄)₂.6H₂O	678.252	M		2.957	536
3326	CsCd(CNS)₃ .	419.139		213		
3327	CsCl.HgCl₂	439.794	C. R.			164
3328	Cs₂HgI₄ . .	973.958	M.		4.806	
3329	Cs₂Hg₃I₈	1882.91	M.		5.14	
3330	Cs₃HgI₅ .	1233.70	R.		4.605	
3331	Cs₂Cu(SO₄)₂.6H₂O	629.412	M.		2.858	559
3332	2CsNO₂.AgNO₂.Bi(NO₂)₃	858.548			3.88¹⁵	
3333	CsSO₄.Ir₂(SO₄)₃.24H₂O	1335.64	C.	110		
3334	CsRh(SO₄)₂.12H₂O.	644.035	C	111		112
3335	CsMnO₄ . .	251.740			3.597¹⁰·⁴	
3336	CsMn(SO₄)₂.12H₂O	596.055	C.			200
3337	Cs₂Mn(SO₄)₂.6H₂O	620.772	M.		2.740	524
3338	CsFeCl₃.2H₂O. . . .	331.055			2.907¹⁷	
3339	Cs₂FeCl₄.2H₂O	499.323			3.275	
3340	CsFe(SO₄)₂.12H₂O	596.965	C.		2.061	100
3341	Cs₂Fe(SO₄)₂.6H₂O	621.682	M.		2.796	550
3342	Cs₂FeSe₂O₈.6H₂O	715.952	M.		3.694	
3343	Cs₂SeO₄.Fe₂(SeO₄)₃.24H₂O	1382.47	C.	60	3.618¹⁶	116
3344	Cs₂Co(SO₄)₂.6H₂O	624.812	M.		2.844	566
3345	Cs₂Co(C₃H₂O₄)₂.4H₂O—Malonate	600.682			2.682	
3346	Cs₂Ni(SO₄)₂.6H₂O. .	624.532	M.		2.872	575
3347	CsCr(SO₄)₂.12H₂O	593.135	C.	116	2.043	94
3348	CsV(SO₄)₂.12H₂O	592.085			2.033⁴	
3349	3CsF.AlF₃	539.390		823		
3350	Cs₂SO₄.Al₂(SO₄)₃.24H₂O	1136.17	C.		1.867⁰	80
3351	2Cs₂O.2Al₂O₃.9SiO₂.H₂O—Pollucite	1325.64	C.		2.9	126
3352	Cs₂La(NO₃)₅.2H₂O.	750.601	M.		2.827⁰	
3353	Cs₂Mg(SO₄)₂.6H₂O	590.162	M.		2.676	488
3354	Cs₂Mg(SeO₄)₂.6H₂O . .	684.432	M.		2.94	583
3355	Cs₂Mg(CrO₄)₂.6H₂O . . .	630.052	M.		2.747	821
3356	Cs₃Cu₂Sr(SCN)₇	1019.69	Tet.		2.882	374
3357	Cs₃Cu₂Ba(SCN)₇ . .	1069.45	Tet.		2.92	365
3358	Cs₃BaAg₂(SCN)₇	1158.07	Tet.		3.026	360
3359	CsLiCl₂	210.665		356.5		

Mg	Mn	Mo	N		Na	Nb	Nd	Ni	O		Os	P	Pb	Pd		Pr	Pt	Ra	Rb		Rh	Ru	S	Sa		Sb	Sc	Se	Si	Sn		Sr	Ta	Tb	Te	Th		Ti	Tl	Tm	U	V		W	Y	Yb	Zn	Zr
76	42	47	11		82	51	61	45	1		35	12	23	41		60	37	80	84		40	39	8	63		14	56	9	18	22		78	52	66	10	24		19	27	70	49	50		48	57	71	28	21

BOILING POINTS

General index No	Boiling point under 1 atm. (or mm of Hg indicated by superscript)	General index No	Boiling point under 1 atm. (or mm of Hg indicated by superscript)	General index No.	Boiling point under 1 atm. (or mm of Hg indicated by superscript)	General index No.	Boiling point under 1 atm. (or mm of Hg indicated by superscript)
1	100	89	444	204	$-$ 95	294	d. <260
2	152 1	91	339	205	$-$ 75	316	4₄7
4	19 4	92	424	206	$-$ 40	320	45₃
6	9 9·?	95	151 0	207	73 5	322	500 d.
7	3 8·??	96	21 3	208	16₂	337	-192.0
8	82	97	$-$ 89 5	209	180	338	s. $-$ 78.5
9	$-$85 0	98	3 5	210	107 23	339	6 3
13	16??	99	4?	211	21₂	341	22₃₀
17	$-$67 0	101	12 5	213	$-$ 8	345	-112.0
21	40??	102	$-$ 33 35	214	172 9	346	$-$ 15
23	13?	103	113 5	215	10₆	347	53
26	$-$ 35 5??	104	118 5??	216	193	348	80
31	s. 110	105	37	217	s. 38 8??	349	$-$ 15 2
34	97	109	86	218	137 6	350	$-$ 65??
35	ca 97	111	56 ?	219	ca. 165	351	$-$ 80 2
36	ca 77 diss	114	diss 44??	222	s. 61 8??	352	57 57
	s. 101??	118	d 210	223	490	353	139
37	ca. 116	120	s ca. 140	224	51₄	356	213
38	$-$ 10 0	125	$-$ 56	226	407 5	357	150??
39	44 6	126	$-$ 63 5	227	523	358	190??
40	s. 10	128	s. 105	228	515	360	137 0
41	$-$ 59 6	129	<71	230	295	361	200
42	74 5		exp 93	232	125	362	15₃
44	60??	130	$-$ 5 5	233	ca. 118	363	ca. 300
46	290	131	5	235	205 s. d.	364	$-$ 30
47	16₇	132	s. 520	237	150 d.	365	8
53	$-$ 30	139	d. <100	238	95??	366	33
54	$-$ 52	140	exp. 240	250	127??	367	153
55	59	141	$-$ 2	251	328 5	368	ca. 240
57	138	142	ca. 32	252	224??	371	2
58	78 8	143	s. 542	253	262??	372	66
59	69 4		235 vac	254	257??	373	109
60	153·??	148	s. 551	255	291??	374	0??
62	151.5??		220 vac	256	s. 150 vac.	376	80
63	54° ??	149	d. 15	263	-55	377	10₄
64	68??	164	s. 135	264	63??	378	140.5
65	115 d.	165	357.3	265	$-$ 53	379	290
66	s. 317	166	s 120	266	122	381	220
67	$-$ 41.2	170	490	268	221	382	113 5
68	$-$ 42	172	d. 160	269	40₈	383	172
72	100	177	s. 140	271	565	384	23₅
73	s. $-$ 39	181	s. 80 d.	272	707	385	192
74	d. 288	186	d. > $-$ 13	274	ca. 300 d.	386	230 5
76	176 4	191	90??	282	$-$ 17	387	255
77	22₇	192	s ca. 180	284	149 5	388	s. 940??
81	183	193	s. 347 (α)	285	390	389	92??
82	s. 450		600 (β)	286	220 2	390	96
84	$-$ 1 8	195	$-$ 87.4	287	92??	391	150??
87	$-$ 35 5	197	57.5??	291	28₀	403	s. 221o diss.
88	324	198	s. 2?0 d.	292	400.6	404	31

No.	B. P.	No.	B. P.	No.	B. P.	No.	B. P.
406	27	488	114 1	716	43o	1515	78.6
407	63.5	490	620	749	732	1534	973
408	107	491	202	752	650	1552	136.7
409	96 2	492	50^{4}	753	624	1556	78 d.
410	90	493	65^{50}	755	s 1185	1575	43^{741}
411	134$^{752\,9}$	494	65^{30}	760	d. 280	1593	>1300
412	122	495	720	769	500	1597	170
413	115 5	496	340	770	d. 271	1610	d. 175
414	108	497	191 d	779	110o	1619	3800
415	142	499	1230	797	46	1624	340
416	139 5	508	180	798	118	1646	35
417	132	513	78	799	160	1647	к. 270
418	153 7	514	146	800	220	1648	180
419	171	515	181	825	970	1649	268
420	172 5	517	> 420	829	963	1658	170 d.
421	191,	518	270 d.	832	71з	1664	3s (in H₂)
422	187	519	240	845	132	1672	19 5
423	205^{756}	520	210	870	105	1673	187
425	114 3^{756}	521	224	881	650	1674	275.6
426	122	522	170	882	383 7	1675	346 7
427	154	523	231	883	304	1676	266
428	153	528	129o	893	s. 345	1677	227 5
429	227	529	95o	894	322	1678	33з
432	195^{20}	530	exp 1o5	896	310 d.	1679	327
435	100 5$^{765\,7}$	543	916		s. 110	1689	6000
436	125	548	954	898	354	1690	6000
437	130	600	s. 475	901	s. 580	1706	69$^{zat.}$
438	149$^{754\,3}$	619	110	915	d. 150		s. 56
439	141 5	621	130^{761}	918	96	1714	118
440	154 5	622	53^{14}	919	159	1724	4100
441	201 5$^{739\,4}$	623	152^{755}	920	191	1747	111 2
442	107^{18}	624	70 5^{16}	921	135^{90}	1749	480
443	230	625	64 5^{14}	922	> 306 d.	1752	148 5^{744}
444	314 2	626	166^{760}	939	13$_{66}$	1753	127
449	. 284	627	78^{11}	940	9$_{93}$	1755	127.19
450	136 4	628	83^{14}	947	13$_{45}$	1758	13o
451	230	629	70^{11}	951	129o	1767	3900
452	154	630	99 5^{16}	958	d. 400	1796	219
454	>360	631	105^{13}	974	170 d.	1797	240.5
459	140	632	96^{13}	1032	240 d.	1798	s. 4oo
460	138	633	108 2^{16}	1059	155o	1799	4зoo
461	4зoo	634	123^{14}	1075	444 d.	1802	229.5
465	− 90	635	124^{14}	1129	s. 265	1803	242
466	29	636	121^{15}	1147	134	1804	32o
467	110 5	637	144 5^{13}	1148	20з	1805	5500
468	86 5	670	s. 610	1149	47 3	1810	87.5
469	72		725	1180	s. 240	1811	17
470	185 9	675	5000	1234	100 8^{182}	1812	d. 200
471	375	678	535	1268	119o	1813	−101
472	163 5	679	217	1334	s. 120o diss.	1814	12.5
480	416	693	139 diss.	1342	315	1815	90 6
481	5100	695	3oo	1397	102.8^{749}	1817	210
485 5	− 52	696	80$_{6}$	1447	104$_{9}$	1819	123o^{94}
486	705	700	81$_{5}$	1509	d. 52	1821	>3500
487	623	703	82$_{4}$	1513	240	1822	110

No.	B. P.	No.	B. P.	No.	B. P.	No.	B. P.
1823	95	2010	d. 100	2500	1560	2921	1416
1824	65	2044	d. 100	2601*		2924	1380
1825	120	2105	590	2604	1670	2926	1330
1826	175	2112	188	2605	1353	2927	d. 225
1827	212	2113	245	2606	d. 270	2931	d. 215
1828	255	2114	270	2608	d. 410	2932	d. 180
1858	2210	2115	331	2610	1265	2936	d. 850
1864	182 7^(42)	2116	330	2613	1190	2958	d. 350
	≮ 177 8	2117	341	2625	d. > 170	2959	d. 400
1865	268	2118	239^(10)	2668	1390	3196	1410
1866	d 7	2131	1112	2670	1700	3197	1390
1869	382	2232	2850	2671	1413	3200	1340
1870	≮ 1550 (in N₂)	2234	450 diss	2677	1390	3205	1300
1879	600 (in H₂)	2236	> 1600	2680	1300	3283	1230
1893	130	2244	718	2769	1496	3284	1290
1894	194	2285	≮ 898 6	2846	> 1400	3287	1300
1895	315	2495	795 diss.	2917	1320	3292	1280
1953	1600	2499	1400	2918	1500		

* Hüttig, Bs, 141: 133, 24

REFRACTIVE INDICES

A. LIQUIDS

Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D
1	436	1.833[10.2]	18	45	1 429	34	625	1 5035[22 b]	50	513	1 5201
2	97	1 193[16]	19	1893	1 432[12]	35	627	1 5062[23 1]	51	628	1 5218[16]
3	9	1.256	20	62	1 437[14]	36	635	1 5081[22]	52	58	1.527[10]
4	195	1.317[17 6]	21	111	1 440[22 b]	37	623	1 5082[21]	53	918	1 5327[22 1]
5	17	1.325[10]	22	59	1 444	38	636	1 5097	54	919	1 5399[22 1]
6	102	1.325[16 b]	23	339	1 454	39	637	1 5118[71]	55	2644	1 548[16]
7	95	1 330[-90]	24	341	1.46	40	633	1 5120[21 b]	56	55	1 557[14]
8	1	1 333	25	210	1 460[25 1]	41	631	1 5127[25]	57	1147	1 56[44]
9	426	1.368	26	1808	1 464	42	619	1 5128	58	287	1 601[14]
10	41	1 374	27	26	1.466[12]	43	621	1.5132[19]	59	450	1 61[10.b]
11	1825	1 381	28	103	1 470[22]	44	515	1 5143	60	2472	1.618
12	109	1 397[16 4]	29	1894	1.480[6 b]	45	2847	1 515	61	57	1 660[14]
13	472	1 400	30	629	1 4926	46	624	1 5158[24 3]	62	214	1.697[26 6]
14	1827	1 408	31	634	1.5005	47	207	1 516[14]	63	1317	1 700
15	38	1 410	32	626	1 5021[21 2]	48	622	1 5174	64	63	1 736
16	2	1 414[22]	33	632	1 5023	49	630	1 5175[19 7]	65	42	1 885
17	1828	1 421									

B. SOLIDS

I. Isotropic Group. m. = mean value

Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D	Serial No.	Gen. index No.	Refractive index n_D
66	2670	1 336	95	3107	1 4814	127	2839	1 5305	160	260	1 7550
67	2913	1.339	96	3265	1 4815	128	3150	1 5329	161	1911	1 780
68	398	1.370	97	3084	1 4817	129	2671	1 5442	162	562	1 782
68 1	3143 6	1 403	98	3259	1 4823	130	1211	1 548	163	3292	1 7876
68 2	3017.6	1 408	99	2870	1 483	131	1451	1 55 (m.)	164	3327	1.792
69	344	1 41	100	3340	1 4839	132	1536	1 55 (m.)	165	1923	1 800
70	3032	1 4115	101	1613	1 4842	133	3200	1 5530	166	1928	1 801
70 1	2099 6	1 426	102	1369	1 4854	134	2924	1 5590	167	1921	1 811
70 2	478 5	1.433	103	2921	1 4903	135	2458	1.5667	168	2232	1.83
71	2235	1 4339	104	3197	1 493	136	1576	1.57	169	2282	1 83
72	2855	1.4388	105	2873	1 495	137	2531	1 5717	170	2364	1.838
73	2596	1 444	106	2902	1 496	138	2679	1 5943	171	1261	1 862?
74	2732	1 452	107	1910	1 4976	139	1187	1.6000	172	945	1.864 (m.)
75	1897	1 454	108	2872	1 50	140	2438	<1 6	173	939	1 93
76	2700	1 454	109	3253	1 5004	141	2394	1 608	174	278	2.0
77	3133	1 4562	109 5	2835	1 501	142	1383	1.61	175	402	2 05
78	3268	1.4566	110	743	1 5066	143	1576	1 61	176	1048	2 05
79	2760	1 457	111	3261	1 5070[18]	144	3284	1 6418	177	1059	2 0710
80	3350	1.4587	112	3334	1 5077	145	132	1 642	178	280	2 087
81	1882	1 4594	113	2887	1 508	146	3205	1 6474	179	581	2 09?
82	344	1.46	114	3137	1 509	147	3019	1 6574	180	1258	2.16
83	3242	1 4638	115	1240	1 5103	148	2267	1 660 (m.)	181	1639	2.16
84	3317	1 4649	116	3343	1 5116[18]	149	2401	1 67	182	668	2 20
85	3320	1 4652	117	2137	1 514	150	2926	1 6770	183	1123	2 20
86	3025	1 4653	118	2886	1 5141	151	3141	1.69	184	2333	2 20
87	3239	1 4658	119	2674	1 5151	152	3287	1 6984	185	1062	2 253
88	690	1 4664	120	2236	1 52	153	148	1 7031	186	951	2 346
89	680	1 4684	121	3047	1 522 (m)	154	2225	1.705	187	756	2 3682
90	2740	1 4693	122	1633	1 5228	155	2392	1 710	188	936	2 705
91	2332	1.4736	123	2842	1 5230	156	2222	1.723			
92	2899	1.48	124	1422	1 5236	157	2415	1.735	188 1		2 89
93	3135	1 4801	125	3098	1 54 (m.)	158	2128	1 7364	188 2		3 56
94	3347	1 4810	126	3351	5 521	159	1145	1.74 (m.)	189	552	3 912

MISCELLANEOUS

Serial No	Gen index No.	Refractive index n	Serial No	Gen index No	Refractive index n	Serial No.	Gen. index No	Refractive index n	Serial No.	Gen. index No.	Refractive index n
190	367	1 579⁵⁵ (F)	193	232	1 563⁴¹ (C)	196	1274	2 69 (Li)	199	3236	1.46 (red)
191	266	1 621⁴⁴ (F)	194	2196	2 35 (Li)	197	1273	2 70 (Li)	200	3336	1.48 (red)
192	352	1 412 (C)	195	890	2 49 (Li)	198	1053	>2 72 (Li)	201	1528	2 18 (red)

II. Uniaxial Group

Serial No.	Gen. Index No	Refractive index ω	Refractive index ε	Serial No.	Gen. index No.	Refractive index ω	Refractive index ε
202	2778	1 300	1 296	247	2224	1 512	1 498
203	1	1 309	1 313	248	2866	1 518	1 522
204	2182	1 3439	1 3602	249	2422	1 522	1.513
205	2851	1 349	1 342	250	243	1 5246	1 4792
206	1323	1 3570	1 3712	251	2336	1.527	1 539
207	1409	1 3638	1 3848	252	764	1 5291	1 5039
208	2130	1 378	1 390	253	2453	1 5296	1 5252
209	814	1 3824	1 3992	254	3164	1 532	1 529
210	1583	1 3910	1 4066	255	1358	1 533	1 575
211	1047	1 4092	1 4080	256	1912	1 534	1 514
212	2237	1 417	1 393	257	2439	1 5364	1 4866
213	2347	1 436	1 478	258	3136	1 537	1 533
214	2713	1 4458	1 4524	259	3162	1 537	1 535
215	2941	1 455	1 515	260	1892	1 539	1 511
216	2735	1 4567	1 4662	261	2871	1 539	1.537
217	3216	1 4574	1 5078	262	1551	1 5393	1 5125
218	3173	1 4715	1 4721	263	2839	1 5398	1 5475
219	2107	1 4720	1 4395	264	2200	1 540	1 510
220	2119	1 473	1 435	265	2207	1 542	1 516
221	2412	1 475	1 486	266	2861	1 542	1 538
222	3185	1 481	1 461	267	342	1 544	1 553
223	1731	1 481	1 493	268	2659	1 545	
224	1970	1 482	1 473	269	2250	1 5496	
225	1995	1 482	1 474	270	1359	1 5519	1 5575
226	2018	1 486	1 479	270 5	2099 5	1 557	1 543
227	2031	1 487	1 479	271	2804	1 558	1 613
228	340	1 487	1 484	272	2129	1 559	1.580
229	2864	1 487	1 486	273	2226	1 56	
230	2493	1 487	1 496	274	1902	1 560	1.580
231	2397	1 49		274 5	475 5	1 563	1 552
232	2880	1 490	1 471	275	2199	1 565	
233	2086	1 490	1 480	276	2326	1 565	1 560
234	2054	1 490	1 481	277	2211	1 565	1 575
235	2072	1 490	1 482	278	2971	1 567	1 518
236	2869	1 490	1 502	279	2420	1 5690	1 6700
237	3181	1 4901	1 4996	280	1340	1 57	
238	1955	1 493	1 480	281	3134	1 572	1 592
239	2061	1 494	1 484	282	2357	1 575	1 57
240	2081	1 495	1 480	283	276	1.5766	1 5217
241	2403	1 496	1 491	284	2125	1.581	1 575
242	2436	1 4991	1 4758	285	1379	1.582	1 645
243	2329	1 507	1 468	286	1872	1 583	1 602
244	2968	1 5095	1 4684	287	2856	1 585	
245	2840	1 5095	1 5232	288	2705	1 5874	1 3361
246	1547	1 5109	1.4873	289	2188	1.5885	1.5970

Serial No.	Gen. index No.	Refractive index ω	Refractive index ε	Serial No.	Gen. index No.	Refractive index ω	Refractive index ε
290	3186	1.589	1.590	346	1994	1 717	1 817
291	3079	1.59		347	2100	1 719	1.733
292	1582	1 59	1 56	348	1951	1 721	1.816
293	3033	1 5906	1 5907	349	1259	1 723	1 681
294	2399	1 595	1 585	350	969	1 724	1.746
295	2417	1 597	1 560	351	3187	1 7278	1 7361
296	847	1.6038	1 6042	352	1025 1	1 730	1 810
297	2904	1 612	1 593	353	2621	1 735	1 435
298	1978	1 613	1 607	354	978	1 741	1 724
299	2314	1 6150	1 6360	355	1414	1 755	1 82
300	2393	1 617	1 652	356	2563	1 757	1 804
301	1400	1 6198	1 5922	357	2594	1 760	1 577
302	2572	1 621	1 619	358	733	1 768	1 812
303	1737	1.623	1.625	359	1858	1 773	1 773
304	2309	1 625		360	3358	1 7761	1 6788
305	2489	1 629	1 639	361	3322	1 784	1 774
306	1011	1 632	1 575	362	3065	1 7909	1 6527
307	2430	1 633	1 639	363	2201	1 80	
308	2275	1 634	1 631	364	1699	1 80	1 72
309	2273	1 634	1 632	365	3357	1 8013	1 6882
310	2307	1 635	1 631	366	1089	1 8036	1.7083
311	556	1 635	1 653	367	2189	1 815	1 761
312	3042	1 636	1 615	368	1307	1 817	1 5973
313	1934	1.640		369	794	1 818	1 618
314	2490	1 64		370	3085	1 820	1 715
315	2507	1 640	1 633	371	1364	1 82	1.73
316	1252	1 6430		372	1063	1.8466	1 9200
317	1739	1 643	1 623	373	1433	1 85	
318	2234	1.644	1 446	374	3356	1 8535	1 6982
319	1044	1 044	1 697	375	1507	1.855	1 60
320	1046	1 644	1 702	376	2358	1 870	1 792
321	2216	1.65	1 59	377	1394	1 875	1 633
322	2644	1.65	1 67	378	1415	1 875	1 784
324	2441	1.651	1 627	379	1431	1 88	
325	, 1907	1.654	1 676	380	2339	1 913	1 923
326	2121	1 6542	1 6700	381	2366	1 918	1.934
327	1156	1 6576	1 6666	382	483	1 923	1.908
328	2285	1.6583	1 4864	383	1416	1 93	
329	1439	1 664	1 629	384	2339	1 945	1.971
330	2433	1 666	1 661	385	1324	1 96	
331	2274	1 667	1 666	386	1419	1 96	
332	2341	1 669	1 657	387	483	1 960	2.015
333	2410	1 669	1 658	388	2365	1 967	1.978
334	2537	1 669	1 665	389	569	1 970	1 936
335	2131	1 675	1 59	390	882	1 9733	2.6559
336	1084	1 6769	1 6294	391	485	1 997	2 093
337	2004	1 680	1 685	392	744	2 008	2 029
338	2597	1 681	1 668	393	310	2 01	1.82
339	2425	1.6817	1 5026	394	666	2 07	2.05
340	1914	1 694	1 641	395	657	2.09	1.94
341	812	1 694	1 723	396	658	2 114	2 140
342	2163	1.700	1 500	397	2957	2 12	2 00
343	2538	1.701	1 699	398	537	2 13	2.21
344	1324.1	1 704	1 679	399	587	2 135	2.118
345	2281	1.706	1.698	400	1064	2.21	2.22

Serial No	Gen. index No	Refractive index		Serial No.	Gen. index No.	Refractive index	
		ω	ε			ω	ε
401	1695	2 2685	2 182	407	445	2.554	2 493
402	2187	2 31	1 95	408	2354	2 58	2 43
403	1776	2 354	2 299	409	447	2 616	2 903
404	755	2 356	2 378	410	403	2 654	2 697
405	1325	2 481	2 210	411	901	2 854	3 201
406	835	2 506	2 529	412	1095	3 0877	2 7924

MISCELLANEOUS

413	1522	1 3817 (C)	1 3872 (C)	420	1413	2 45 (Li)	2 51 (Li)
414	2035 1	2 005 (667)	2 004 (667)	421	1264	2.46 (Li)	2 15 (Li)
415	1957 1		2 013 (667)	422	1094	2 6 (La)	
416	2002 1	2 019 (667)	2 007 (667)	423	524	2 665 (Li)	2 535 (La)
417	526	2 3 (La)		424	1334	3 01 (Li)	2 94 (La)
418	538	2 35 (La)	2 33 (La)	425	1098	3 084 (Li)	2 881 (Li)
419	1668	2 402 (La)	2 304 (La)	426	2471	1 683 (red)	1 587 (red)

III. Biaxial Group

Serial No	Gen index No	Refractive index			Serial No.	Gen. index No.	Refractive index		
		α	β	γ			α	β	γ
427	2852		1 364		462	1876	1 462	1 470	1 471
428	2694	1 394	1 396	1 398	463	343	1 469	1 47	1 473
429	2897		1 413		464	2150	1 4716	1 4730	1 4786
430	2898	1 407	1 411	1 415	465	2729	1 4653	1 4738	1 4804
431	2753	1 405	1 425	1 440	466	2691	1 464	1 474	1 485
432	2718	1 4193	1 4309	1 4493	467	3146	1 466	1 475	1 494
433	2724	1 4321	1 4361	1 4373	468	1874	1 474	1 476	1 483
434	2693		1 44		469	2617	1 460	1 477	1 488
435	3189	1 438	1 44	1 452	470	2398	1 461	1 478	1 485
436	2733	1 439	1 441	1 469	471	1356	1 4713	1 4782	1 4856
437	2723	1 4412	1 4424	1 4526	472	2948	1 475	1 480	1 487
438	2721		1 4434		473	2223	1 476	1 480	1 483
439	411	1 4368	1 4458	1 4510	474	3255	1 4767	1 4807	1 4907
440	2964	1 447	1 448	1 459	475	2708	1 391	1 481	1 486
441	2739	1 4453	1 4496	1 4513	476	2978		1 482	
442	3133	1 430	1 452	1 458	477	1918	1 478	1 482	1 482
443	2710	1 440	1 452	1 453	478	2862	1 480	1 482	1 493
444	2717	1 4499	1 4525	1 4604	479	3083	1 4759	1 4821	1 4969
445	2395	1 448	1 454	1 456	480	2715	1 4777	1 4822	1 5036
446	2890	1 435	1 455	1 459	481	1463	1 477	1 483	1 489
447	2145	1 4326	1 4554	1 4609	482	3029	1 4775	1 4833	1 4969
448	1809	1 340	1 456	1 459	483	2970	1 4768	1 4843	1 4870
449	2854	1 432	1 457	1 458	484	1289	1 4801	1 4840	1 4913
450	2720	1 4401	1 4629	1 4815	485	3247	1 4798	1 4848	1 4948
451	3149	1 4607	1 4629	1 4755	486	2977	1 440	1 485	1 550
452	2757		1 464		487	2719	1 4557	1 4852	1 4873
453	1871	1 459	1 464	1 470	488	3353	1 4857	1 4858	1 4916
454	2727	1 4599	1 4645	1 4649	489	138		1 486	
455	2616		1 465		490	760	1 4620	1 4860	1 4897
456	2738	1 4622	1 4658	1 4782	491	3043	1 4836	1 4864	1 5020
457	2743	1 4649	1 4663	1 4791	492	3091	1 4807	1 4865	1 5004
458	2943	1 4609	1 4660	1 5657	493	3148	1 483	1 487	1 490
459	2165	1 456	1.468	1 507	494	2853	1 484	1 487	1 496
460	2848	1 4468	1 4686	1 4715	495	3258	1 4815	1 4874	1 4977
461	3273	1.4672	1 4689	1 4779	496	3231		1 488	

Serial No.	Gen. index No.	Refractive index			Serial No	Gen. index No	Refractive index		
		α	β	γ			α	β	γ
497	2882	1 485	1.488	1 489	552	3323	1 5022	1 5048	1.5093
498	2881	1 486	1 488	1.489	553	3151	1.494	1 505	1.516
499	3245	1.4833	1 4884	1 4975	554	2469	1.497	1 505	1.509
500	854	1 4847	1 4887	1 4959	555	2900	1.505	1 505	1.506
501	1548	1 4669	1 4888	1 4921	556	2959	1 3346	1 5056	1 5064
502	3217	1 4812	1 4888	1.5719	557	2178		1 506	
503	2147	1 4856	1 4892	1 4911	558	2118	1 314	1 506	1 506
504	2725	1.4855	1 4897	1.5041	559	3331	1 5048	1 5081	1 5153
505	1924		1 49		560	1986		1 507	
506	2912		1 490		561	2299	1 193	1 507	1.545
507	1863	1.473	1 490	1 511	562	2132	1 495	1 507	1.528
508	2950	1 479	1 490	1 526	563	2765		1 5073	
509	2408	1 484	1 49	1 495	564	2696	1.4886	1 5079	1 5360
510	3249	1 4886	1 4906	1 5036	565	2868	1 504	1 508	1.545
511	2143		1 491		566	3344	1 5057	1 5085	1.5132
512	2171		1 491		567	2893	1 5043	1 5093	1.5751
513	1368	1 4870	1 4915	1 4989	568	2151	1 5070	1 5093	1 5169
514	3096	1 4836	1 4916	1 5051	569	3230		1 510	
515	3262	1.4859	1 4916	1.5011	570	2383	1.495	1 51	1.520
516	777	1 4888	1 4930	1 4994	571	2777	1 500	1 510	1.515
517	3184	1 492	1 493	1.496	572	2106	1 502	1 510	1.512
518	804		1 494		573	2663	1 504	1 510	1.516
519	2938	1 4935	1 4947	1 4973	574	2772		1 511	
520	2697	1 4820	1 4953	1 5185	575	3346	1 5087	1 5129	1 5162
521	1491	1 4902	1 4953	1 5032	576	3215	1 5131	1 5133	1.5144
522	2157	1 495	1 496	1 504	577	2289	1 510	1 514	1.578
523	3264	1 4895	1 4961	1 5052	578	2317	1 512	1 514	1.515
524	3337	1 4946	1 4966	1.5025	579	2922	1 440	1 515	1 525
525	1716		1 4967		580	2894	1 4435	1 5156	1.5233
526	2259	1 465	1 498	1 504	581	3159	1 500	1 5170	1 5183
527	2771	1 495	1 498	1 499	582	2551	1 500	1 517	1 525
528	2407	1 498	1 499	1 505	583	3354	1 5178	1 5179	1 5236
529	3152	1 4969	1 4991	1 5139	584	2553		1 518	
530	1361,		1 500		585	2153	1 514	1 518	1 533
531	2901		1 5		586	2264	1 515	1 518	1.525
532	3014		1 500		587	1875	1 516	1 518	1 533
533	2638	1 40	1 50		588	3031	1 5121	1.5181	1 5335
534	2709	1 418	1 500	1 543	589	3092	1 5135	1.5195	1.5358
535	806	1 180	1 500	1 530	590	2228		1.52	
536	3325	1 498	1 500	1 506	591	3158		1.52	
537	2108	1 4664	1 5007	1 5027	592	2998	1 48	1.52	1 55
538	992	1 4910	1 5007	1 5054	593	2477	1 500	1 520	1.580
539	1557	1 4949	1 5007	1 5081	594	3221	1 51	1 52	1.524
540	2413		1 501		595	2154	1 510	1 520	1.543
541	2930		1 501		596	2860	1 516	1.52	1.520
542	2164	1 495	1 501	1 526	597	2466	1 484	1 521	1 538
543	179	1 4981	1 5016	1 5866	598	3246	1.5162	1 5222	1.5331
544	2498	1 4710	1 5017	ca β	599	1466		1 5225	1 5227
545	2180	1 490	1 502	1 511	600	2249	1 5205	1 5226	1.5296
546	2737	1.4794	1 5021	1 5265	601	3176		1 523	
547	2371	1 499	1.503	1 538	602	174	1.5209	1 5230	1.5330
548	2396	1.501	1.503	1 510	603	3045	1.5096	1 5235	1.5387
549	3274	1.5011	1.5031	1.5135	604	2758	1 407	1 524	1.541
550	3341	1 5003	1.5035	1.5094	605	2405	1 513	1.524	1.525
551	2896	1.491	1.504	1.520	606	3139	1 518	1 524	1.526

Serial No.	Gen. index No.	Refractive index α	β	γ	Serial No.	Gen. index No.	Refractive index α	β	γ
607	3111	1 5221	1 5244	1 5373	662	2592	1 538	1 549	1.554
608	3097	1 5199	1 5248	1 5339	663	2014	1.5399	1.5494	1.5607
609	2294	1 470	1 525	1 555	664	1886		1 55	
610	2097		1 526		665	2204	1 5211	1 5500	1 5680
611	3157	1 508	1 526	1 550	666	2212	1 53	1.55	1 55
612	1370	1 5201	1 5260	1 5356	667	1032	1 545	1.55	
613	3138	1 522	1 526	1 530	668	2029	1 5413	1 5505	1.5621
614	2641		1 529		669	3074	1 5498	1 5513	1 5634
615	2865	1 525	1 529	1 536	670	2046	1 5427	1 5519	1 5629
616	2807	1 5193	1 5205	1 5436	671	3276		1 552	
617	2985	1 417	1 530	1 533	672	2736	1.5382	1 5535	1 5607
618	2304	1 515	1 530	1 580	673	3220	1.5515	1 5537	1 5582
619	1762	1 518	1 530	1 542	674	2288	1.491	1 555	1 650
620	778	1 5240	1 5300	1 5385	675	1360	1 533	1 555	1 635
621	2280	1 525	1 53	1 550	676	2292	1 545	1 555	1 575
622	2167	1 527	1 530	1 540	677	1927	1.551	1 555	1.562
623	1497	1 5246	1 5311	1 5396	678	3086		1 556	
624	2909	1 1893	1 5314	1 5363	679	2876	1 5520	1.5579	1.5608
625	2889	1 515	1 532	1 536	680	1884	1 551	1 558	1.582
626	2197	1 527	1 532	1 583	681	1925	1 554	1 558	1.573
627	2566		1 533		682	2637	1 530	1 560	1 590 ?
628	2759		1 533		683	2296	1 55	1 56	1.57
629	2190		1 533	1 5769	684	2618	1 5487	1 5602	1.5788
630	2166	1 189	1 531	1 557	685	3165	1 548	1 562	1.567
631	2432	1 517	1 534	1 565	686	188	1 5607	1 5630	1.5846
632	1861	1 5347	1 5347	1 5577	687	3305	1.5598	1 5644	1 5662
633	2286	1 460	1 535	1 545	688	838		1 565	
634	3015	1 495	1 535		689	2780	1 560	1 565	1.574
635	2382	1 500	1 535	1 560	690	1901	1 561	1 565	1 567
636	2302	1 515	1 535	1 575	691	3034		1 565	1 608
637	2142	1 523	1 535	1 586	692	1860	1 566	1 566	1 587
638	2205	1 525	1 535 ?	1 550	693	2642		1 567	
639	993	1 5213	1 5355	1 5395	694	2634	1 428	1 567	1 572
640	3324	1 5326	1 5362	1 5412	695	2298	1 450	1 567	1 600
641	961	1 5140	1 5368	1 5433	696	2774	1 536	1 567	1 649
642	1355	1 528	1 537	1 543	697	3002	1.527	1 568	1 647
643	1558	1 5291	1 5372	1 5466	698	2268	1 565	1 568	1 580
644	2404		1 539		699	3087	1 5660	1 5689	1 5831
645	3004		1 539		700	2877	1 565	1 569	1 569
646	2955	1 5352	1 5390	1 5446	701	2156	1 569	1 570	1 582
647	2179		1 54		702	2159	1 563	1 571	1 596
648	2293	1 460	1 540	1 610	703	2158	1.555	1 572	1 575
649	2218	1 520	1 54	1 545	704	2464	1 559	1 574	1 598
650	2217	1 527	1 540	1 544	705	2369	1.56	1 574	1 580
651	1512		1 542		706	2290	1.495	1 575	1 640
652	1030	1 413	1 542	1 557	707	2368	1 553	1 575	1 577
653	2859	1 466	1 542	1 596	708	2248	1 5693	1 5752	1 6130
654	1363	1 530	1 543	1 595	709	3063	1 5438	1 5754	
655	2981	1 415	1 545	1 565	710	643		1 576	
656	2205	1 539	1 545	1.551	711	1889	1 562	1 576	1 588
657	2878	1 545	1 546	1.551	712	1888	1 574	1 576	1.588
658	2036	1 5392	1.5479	1 5592	713	2504	1 5622	1 577	1 635
659	2558	1 542	1 548	ca. 1 548	714	3089		1 5772	
660	2198	1 544	1 548	1.572	715	2789	1 544	1 578	1.601
661	1950	1.5433	1.5490	1.5755	716	3057	1.569	1 579	1.669

Serial No.	Gen. index No.	Refractive index α	Refractive index β	Refractive index γ	Serial No.	Gen. index No.	Refractive index α	Refractive index β	Refractive index γ
717	2416	1.578	1 579	1 583	772	2321	1 605	1 61	1 612
718	2359	1.5700	1 5818	1 5961	773	2315	1 610	1 611	1.654
719	2370	1 560	1 582	1 587	774	2421	1 592	1 612	1 621
720	782	1.574	1 582	1 582	775	2559	1 597	1 612	1 621
721	2389		1 583		776	2335	1 609	1 6125	1.619
722	3073		1 5837		777	2173	1 520	1 613	1 639
723	2400	1.576	1 584	1 588	778	2356	1 602	1 613	1 649
724	1885	1 563	1 585	1 592	779	1913	1 588	1 617	1.655
725	2803	1 508	1 586	1.525	780	813	1 614	1 617	1.636
726	2227	1.585	1.586	1.596	781	2184	1.607	1 619	1 639
727	1903	1.552	1 588	1.600	782	1915	1 61	1 62	1 65
728	2181	1.539	1 589	1 589	783	1043	1 61	1 62	1 71
729	2591	1.584	1.589	1 594	784	1905	1 619	1 620	1 627
730	2279	1.5825	1 5891	1 5937	785	2419	1 620	1 620	1.654
731	3140	1.561	1.590	1 594	786	2429	1 609	1 623	1 635
732	2327	1.586	1.59	1 598	787	2583	1.610	1 623	1 623
733	2123	1 5595	1 5908	1 6311	788	2367	1 621	1 623	1 631
734	781	1 572	1.591	1 59	789	2451	1 6220	1 6237	1.6300
735	2385	1.572	1 591	1 594	790	2185	1 617	1 624	1 652
736	3056		1 592		791	809	1.531	1 625	1.659
737	1738	1 582	1.592	1 592	792	1035	1 541	1 625	1 660
738	2384	1 582	1 592	<1 606	793	783	1 614	1 625	1 637
739	2381	1 5863	1.5920	1 6139	794	1382	1 615	1 625	1 665
740	2658	1.579	1 593	1 597	795	2561	1 620	1 625	1.645
741	2798	1 5889	1.5043	1.7163	796	2411	1 616	1.626	1.640
742	1276	1 562	1 595	1 632	797	2431	1.621	1.627	1 635
743	2903	1 571	1 595	1 598	798	3178	1.6237	1 6278	2.2916
744	2523	1 5860	1 5951	1.6072	799	1514	1 532	1 628	1 605
745	2546	1.573	1 597	1.636	800	2316	1 616	1 629	1 631
746	2388	1 586	1.598	1 605	801	1920		1 63	
747	2775	1 573	1.599	1 657	802	1721	1.585	1 630	1 630
748	1987	1.5989	1 5999	1 6003	803	1321	1 602	1 632	1.632
749	2664		1 6		804	2230	1.603	1 632	1.639
750	2867		1 60		805	3275	1 622	1 633	1.644
751	2322	1 595	1.60	1 603	806	2386	1.632	1 634	1 636
752	3307	1 599	1 600	1.600	807	2308		1 635	
753	3179	1 5883	1 6007	1 6316	808	1580	1.541	1 636	1.669
754	2291	1 413	1 602	1 611	809	2767	1 577	1 636	1 639
755	786	1.586	1.602	1 608	810	3012	1 620	1 636	L.638
756	2278	1.590	1.602	1.638	811	1185		1 037	
757	1378	1.579	1 603	1 633	812	2470	1 453	1 637	1 707
758	1935	1 586	1.603	1 623	813	2206	1.636	1 637	1.653
759	2324	1.593	1 603	1 607	814	2640	1 507	1 638	1.698
760	2857	1.594	1 603	1 615	815	1898	1 632	1 638	1.643
761	2152	1.602	1 604	1 615	816	2521	1.6369	1 6381	1 6491
762	1357	1.51	1 605	1 611	817	3068	1 545	1.641	1 760
763	2440	1 567	1 605	1 626	818	2823	1 596	1 641	1.652
764	2122	1.591	1 605	1 614	819	1900	1 638	1 642	1.653
765	2269		1.606		820	2409	1 632	1 643	1.645
766	2895	1.595	1.606	1 634	821	3355	1.637	1 643	1.655
767	2555		1.607		822	2305	1.462	1.643	1.722
768	3003		1.607		823	2349	1.636	1 644	1.654
769	3052		1.6071		824	2320	1.642	1 645	1.654
770	3001	1 571	1.608	1.694	825	2501	1.635	1.646	1.660
771	820	1.617	1.609	1.593	826	1929	1.643	1.649	1.649

Serial No.	Gen. index No	α	β	γ	Serial No.	Gen. index No.	α	β	γ
827	2564		1 651		882	2595	1.525	1.684	1.686
828	2177	1 635	1 651	1 670	883	941		1.685	
829	826		1 8513		884	2593	1.681	1.685	1 695
830	1916	1 612	1 652	1 675	885	1005	1 67	1 686	1 698
831	2387	1 625	1 653	1 669	886	1937	1.678	1 686	1 689
832	2176	1 650	1 653	1 658	887	2809		1.687	
833	2214	1 6527	1 6537	1 6748	888	1184	1 687	1 687	1 704
834	2863		1 654		889	1270 1	1.684	1 695	1.698
835	1298	1 647	1 654	1 660	890	1406	1 672	1 697	1 717
836	2175	1 651	1 654	1 660	891	1008	1 695	1 698	1 733
837	1919	1 633	1 655	1 662	892	2815	1 6610	1 6994	1 7510
838	2391	1 643	1 655	1 663	893	2810		1.70	
839	2126	1 652	1 655	1 671	894	2565		1.702	
840	2790	1 6491	1 6555	1 7143	895	2652		1.702	
841	2379	1 540	1 656	1 682	896	2418	1 700	1.702	1.706
842	1295	1 651	1 656	1 683	897	1294	1 695	1 704	1.710
843	1297	1 652	1 656	1 660	898	785	1 660	1 705	1 713
844	1069	1 6272	1 6573	1 6601	899	734		1.707	
845	1569	1 622	1 658	1 687	900	2229	1 705	1.709	1 711
846	1296	1 63	1 66	1 69	901	2428	1 708	1.711	1 718
847	2424	1 640	1 660	1 675	902	2350	1 709	1 711	1.724
848	1439	1 655	1 66	1 670	903	976	1.703	1 713	1 722
849	2910	1 645	1 661	1 688	904	2556	1 614	1 714	1.729
850	1505	1 6263	1 6614	1 6986	905	2480	1 7146	1 7174	1 812
851	1585	1 629	1 662	1 727	906	1720	1 691	1 720	1.720
852	2426	1 651	1 662	1 668	907	1899	1 712	1 720	1 728
853	2463	1 5155	1 664	1 666	908	2318	1 715	1 720	1 737
854	2600	1 660	1 666	1 676	909	2423	1 712	1 721	1 731
855	2372	1 642	1 667	1 669	910	2351	1 686	1 722	1 735
856	2215	1 662	1 667	1.673	911	1859	1.702	1.722	1 750
857	1388	1 635	1 668	1 702	912	1012	1 694	1 726	1 730
858	3064	1 626	1 6684	1 757	913	2510	1 7129	1 7266	1.7441
859	3005	1.485	1 669	1 697	914	1922	1 705	1 729	1 730
860	757	1 658	1 669	1 670	915	2417.1	1 724	1.729	1 734
861	2183		1 670		916	972	1.710	1 731	1 732
862	2340		1 670		917	1377	1 730	1 732	1.762
863	2186	1 668	1 670	1 690	918	793	1 708	1 733	1 758
864	2427	1 664	1 671	1 694	919	1670	1 720	1 733	1 935
865	1908	1 670	1 671	1 689	920	807	1 640	1 736	1 750
866	2858	1 634	1 673	1 685	921	964	1 730	1 737	1 785
867	2330	1 640	1 674	1 679	922	2360	1.732	1 737	1 751
868	2353	1 662	1 674	1 676	923	1841	1.617	1.738	1.776
869	2402	1 665	1 674	1 684	924	3101	1 7202	1 7380	1 8197
870	2905	1 666	1 674	1 688	925	1956	1 731	1 738	1 744
871	2800	1 671	1 674	1 684	926	2208		1.74	
872	2557	1 673	1 674	1 678	927	3100		1 74	
873	1381	1 653	1 675	1 697	928	1408	1.71	1.74	1 76
874	1389		1 676		929	1318	1 733	1.740	1 744
875	2542	1 529	1 676	1 677	930	1930	1.736	1.741	1.746
876	1926	1 643	1 678	1 684	931	1003		1 743	
877	3037	1 648	1 678	1 699	932	997	1.702	1 745	1.789
878	2651	1 676	1 679	1 687	933	2124	1.717	1 748	1 757
879	2741		1 6802		934	2484		1 749	
880	2284	1 5299	1 6809	1 6854	935	1726	1.72	1 75	1.80
881	792	1.662	1 683	1.717	936	1670	1.74	1 75	1.95

Serial No.	Gen. index No.	α	β	γ	Serial No.	Gen. index No.	α	β	γ
937	2781	1.743	1.754	1 764	985	2338	1 910	1 91	1.945
938	1028	1.730	1.758	1 838	986	261	1 871	1 92	2.01
939	967	1.708	1.760	1 798	987	1050	1 885	1 920	1.956
940	1000	1.719	1.762	1 805	988	3124	1 750	1 925	1 95
941	1387	1.765	1.774	1.797	989	1305	1 92	1 95	1.96
942	2573	1.770	1.774	1.783 ?	990	1365	1 702	1 955	1.965
943	2352	1.758	1.776	1.795	991	712	1 9493	1 9592	1 9640
944	966	1.730	1.778	1 803	992	663	1 947	1 961	1 968
945	1303	1.760	1.779	1 779	993	1722	1 955	1 985	2 05
946	1944	1 757	1 78	1 803	994	401		1 99	
947	2127	1 78	1 78	1.785	995	557	1.93	1 99	2 02
948	1045	1.752	1 782	1 815	996	660	1 87	2 00	2 01
949	1319	1.759	1.786	1.797	997	1723	1 90	2 00	2 05
950	1380	1 775	1.786	1.815	998	576		2 03	
951	1006	1.747	1.788	1 829	999	2219	1 908	2 05	2 065
952	1420	1.783	1 788	1 818	1000	573	2 042	2 050	2 050
953	1670	1 78	1 79	2 04	1001	617	1 8037	2 0763	2 0780
954	1300	1.780	1 793	1.802	1002	329		2 09	
955	2337		1 795		1003	2375	1.70	2 10	2 23
956	2808	1 763	1 799	1.813	1004	1326	2 08	2 1	2.16
957	735		1 80		1005	511	1 816	2 102	2.126 ?
958	1362	1 76	1 8	1 81	1006	539	2 0767	2.1161	2 1580
959	1301	1.783	1.801	1 834	1007	1696		2.15	
960	1007	1.79	1.807	1 84	1008	535	2 04	2.15	2 15
961	2376	1.775	1.815	1 825	1009	335	2 14	2.15	2.18
962	2582		1 816		1010	1421	2.12	2 17	2.31
963	583	1.74	1 82		1011	2374	1.77	2 18	2.35
964	1009	1 820	1.826	1 88	1012	473	2 13	2 19	2.20
965	2346	1 800	1.831	1 846	1013	1336	1 94	2 20	2 51
966	2802	1 750	1.832	1 832	1014	1327	2 10	2.20	2 31
967	1049	1 8090	1 8380	1 8593	1015	1391	2.19	2 20	2.33
968	999	1.69	1 84	1 85	1016	529	2.1992	2 2172	2 2596
969	1430	1 773	1 840	1 845	1017	1697	2.17	2 22	2 32
970	2363	1 825	1 842	1 857	1018	1671	2 09	2 24	2 26
971	2221	1.85	1.85	1.99	1019	1807	2 22	2 25	2 29
972	2220	1 85	1 85	2 02	1020	1784	2 17	2 26	2 32
973	639	1 789	1 852	1 877	1021	1781	2 18	2 27	2 35
974	2492		1 865		1022	536	2.24	2 27	2 31
975	707	1 8600	1 8671	1 8853	1023	1694	2.27	2 27	2 30
976	1010	1 73	1 870	1 91	1024	279	2.18	2 35	2 35
977	1027	1 655	1 875	1 909	1025	2331		2 38	
978	1407	1 835	1 877	1 886	1026	1335	2 26	2 39	2.40
979	1794	1 817	1 879	2 057	1027	878	2 37	2 5	2 65
980	1302	1 87	1 88	1 93	1028	446	2.583	2.586	2.741
981	553	1 8771	1 8823	1 8937	1029	917		3	
982	3010	1 527	1 903	1 952	1030	1096		3	
983	2334	1 900	1.907	2 034	1031	1101		3	
984	2361		1.91	1 91	1032	296	3 194	4 046	4 303

MISCELLANEOUS

Serial No.	Gen. index No.	α	β	γ	Serial No.	Gen. index No.	α	β	γ
1033	944	1 831	1.861 (green)	1 880	1037.1	3143.5	1.461		1 449
1034	429	1 3996		1 4102	1037.2	3017 5	1 466		1.455
1035	432	1.4057		1.4165	1038	3009	1.4676		1.620
1036	418	1.4248		1 4382	1039	1399	1.500		1 660
1037	2994	1.452		1.465	1040	2776	1.518		1 527

Serial No.	Gen. index No	Refractive index			Serial No.	Gen. index No.	Refractive index		
		α	β	γ			α	β	γ
1041	2213	1 575		1 649	1061	1412	2.38	2.39 (Li)	2.42
1042	2644	1 584		1 604	1062	1698		2.40 (Li)	
1043	2646	1 594		1 611	1063	1800		2.40 (Li)	
1044	1322	1 62		1 63	1064	1766	2.41	2.50 (Li)	2.51
1045	2348	1 6226		1 7643	1065	1661		2.55 (Li)	
1046	2323	1 611		1 650	1066	1093	2.48	2.58 (Li)	2.60
1047	2570	1 6704			1067	271	2.46	2.59 (Li)	2 61
1048	2414	1 675		1 685	1068	525	2.51	2 61 (Li)	2 71
1049	2319	1 717		1 735	1069	1411		2 62 (Li)	
1050	1075	1 729		1 788	1070	887	2 35	2 64 (Li)	2 66
1051	2549			1 789	1071	272		> 2.72 (Li)	
1052	2560	1.810		1 830	1072	723	> 2.72	> 2.72 (Li)	
1053	716	1.817			1073	298	2.74(La)		> 2 72 (Li)
1054	582	1 90		1 97	1074	2770		1.473 (red)	
1055	3081	1 553	1 555 (La)	1 571	1075	3177		1.5226 (red)	
1056	82	2 00	2 18 (Li)	2 35	1076	2524		1 532 (red)	
1057	2355	2 200	2.200 (Li)	2 290	1077	3114		1 591 (red)	
1058	1263	2.24	2.24 (Li)	2 53	1078	935		2.63 (red)	
1059	599	2.30	2 35 (La)	2 40					
1060	1631	2 31	2 37 (La)	2 66					

INDEX OF MINERAL NAMES

ℭ-TABLE

[Compounds of carbon with elements having key-numbers below 16]

Gen. index No.	Formula	Name of Table p. 250	Molecular weight (I.C.T. atomic weights, t. p. 43)	Normal melting point °C	Boiling point under 1 atm. (or mm of Hg indicated by superscript)	Specific gravity 20°/4° (or at other indicated temperature)	Refractive index finding No. r Table, p. 276
1	CBi_2O_4	Bismutospherite	510 00	d.		7.35	
1 1	$CBrClO$	Carbonyl bromochloride	143 37		25	1 82[15]	
2	$CBrCl_3$	Bromotrichloromethane	198 29	−21	172	1.959[14 6]	697
3	$CBrN$	Cyanogen bromide	105 92	52	61 6	2 015	
4	CBr_2O	Carbonyl bromide	187 83		64 5	2.44	
5	CBr_2NO_2	Bromopicrin	297 76	10 3	127[114]	2.799	826
6	CBr_4	Carbon tetrabromide	331 66	{ α48.4 β90.1	189.5	3 42	
7	$CClN$	Cyanogen chloride	61 466	−6	13.8	1.186	
8	$CCl_2N_2O_4$	Dichlorodinitromethane $Cl_2C(NO_2)$	171 93	122 5			
9	CCl_2O	Carbonyl chloride (Phosgene)	98 916	−104	8.3	1.392	
10	CCl_2S	Thiophosgene	114 98		73 5	1.509[16]	721
11	CCl_3NO_2	Chloropicrin Cl_3CNO_2	164 38	−64	112 4	1.692[0]	470
12	CCl_4	Carbon tetrachloride	153 83	−23 0	76 8	1.595	476
13	CF_4	Carbon tetrafluoride	88 00	−80	−15		
14	CIN	Cyanogen iodide	152 94	146 5			
15	CIN_3O_6	Iodotrinitromethane $CI(NO_2)_3$	276 96	56			
16	CI_4	Carbon tetraiodide	519 73	d.		4 32	
17	CN_4O_8	Tetranitromethane $C(NO_2)_4$	196 03	13	125.7	1 650[11]	364
17 1	COS	Carbonyl sulfide	61 065	−138	−48	1 24[−87]	
17 2	$CSSe$	Carbon sulfoselenide	123 265		84.5		
17 3	CS_2	Carbon disulfide	76.130	−111.6	46.3	1 261[22/20]	
17 4	$CHBrCl_2$	Bromodichloromethane	163 84		92	1 925[15]	
18	$CHBr_3$	Bromoform	252 76	7 7	150.4	2 890	772
19	$CHCl_3$	Chloroform	119 38	−63.5	61 2	1 489	417
20	CHF_3	Fluoroform	70 008		20[40 at.]	2 53	
21	CHI_3	Iodoform	393 80	119		4 1	1189
22	CHN	Hydrocyanic acid HCN	27 016	−14	26	0 699	809
23	$CHNO$	Cyanic acid HCNO	43 016	d.		1.140[0]	
24	$CHNS$	Thiocyanic acid HCNS	59.081	5	d.		
25	CHN_3O_6	Nitroform $CH(NO_2)_3$	151 032	15	> 100 d.		
26	CH_2Br_2	Methylene bromide	173 85	−52 8	97 8	2 46[15/15]	
27	CH_2ClNO	Carbamyl chloride $ClCONH_2$	79 481	50	62		
28	CH_2Cl_2	Methylene chloride	84 931	−96 7	40 1	1 336	273
29	CH_2I_2	Methylene iodide	267 88	5 2; 5 7	180 d.	3 325	870
30	CH_2N_2	Cyanamide $CN.NH_2$	42 031	44	140[19] d.	1 083	1073
31	CH_2N_2	Diazomethane $H_2C:N_2$	42 031	−145	−23		
32	$CH_2N_2O_3$	Methylnitrolic acid $O_2NCHNOH$	90 031	64			
33	$CH_2N_2O_4$	Dinitromethane $H_2C(NO_2)_2$	106 031	< −15	100 d.		
34	CH_2N_4	Tetrazole	70 047	155			
35	CH_2O	Formaldehyde HCHO	30 015	−92	−21	0 815[−20]	
36	$(CH_2O)_x$	Paraformaldehyde	(30 015)$_x$	160			
37	CH_2O_2	Formic acid HCO_2H	46 015	8 4	100 5	1 220	25
38	CH_2AsCl_2	Methylarsine dichloride	160 90	−59	136	1 838	
39	CH_3AsO	Methylarsinous oxide	105 98	95			
40	CH_3Br	Methyl bromide	94 939	−93	4 6	1 732[0]	
41	CH_3Cl	Methyl chloride	50 481	−97.6	−23 7	0.920[18]	
42	CH_3ClO	Methyl hypochlorite CH_3OCl	66 481		13 4		
43	CH_3ClO_2S	Methylsulfone chloride	114 546		160	1 510	
44	CH_3F	Methyl fluoride	34 023		−78 0		
45	CH_3I	Methyl iodide	141.96	−66 1	42.6	2 279	696
46	CH_3NO	Formamide $HCONH_2$	45 031	−5	193	1.139	995
47	CH_3NO	Formaldoxime $H_2C:NOH$	45 031		84		
48	CH_3NO_2	Nitromethane CH_3NO_2	61.031	−29.2	101.9	1 139	43
49	CH_3NO_2	Methyl nitrite CH_3ONO	61.031		−12	0.991[14]	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
50	CH₃NO₃	Methyl nitrate CH₃ONO₂	77 031		exp. 65	1 217¹⁰	
51	CH₃NS	Thioformamide HCSNH₂	61 096	29			
52	CH₃N₃	Methyl azide	57 047		21	0 869⁴₁₄	
53	CH₃N₃O₃	Nitrourea O₂NNHCONH₂	105 05	150 d.			
54	CH₄	Methane	16 0308	-184	-161 4	0 415⁻¹⁶⁴	
55	CH₄N₂O	Urea H₂NCONH₂	60 047	132 7		1 335	1167
56	CH₄N₂O₂	Methylnitramine CH₃NHNO₂	76 047	38		1 243⁴⁸ ⁰	1077
57	CH₄N₂S	Ammonium thiocyanate	76 112	149 6	d 160	1 305	
58	CH₄N₂S	Thiourea H₂NCSNH₂	76 112	182		1 405	
59	CH₄N₄O₂	Nitroguanidine H₂NC(:NH)NHNO₂	104 063	231			
60	CH₄O	Methyl alcohol CH₃OH	32 031	97 8		0 792	2
61	CH₄O₃S	Methylsulfonic acid CH₃SO₃H	96 096		167¹⁰	1 481	
62	CH₄O₄S	Methyl sulfuric acid CH₃SO₄H	112 09	< -30			
63	CH₄S	Methylmercaptan CH₃SH	48 096	121 0	7 6	0 868	
64	CH₅As	Methylarsine CH₃AsH₂	91 999		2		
64 1	CH₅AsO₃	Methyl arsenate CH₃AsO(OH)₂	139 999	161			1234
65	CH₅N	Methylamine CH₃NH₂	31 047	92 5	-6 5	0 699⁻¹¹	
66	CH₅NO	N-Methylhydroxylamine CH₃NHOH	47 017	42	62 5¹⁰	1 0003	226
67	CH₅NO₂	Ammonium formate HCO₂NH₄	63 047	116		1 266	1223
67 1	CH₅NO₃	Ammonium hydrogen carbonate	79 047	d		1 573	
68	CH₅N₃	Diazoaminomethane	59 063	-12	92 s d		
69	CH₅N₃O	Semicarbazide H₂NCONHNH₂	75 063	96			
70	CH₅N₃O₄	Urea nitrate H₂NCONH₂·HNO₃	123 06	153 d		1 664	
71	CH₅N₃S	Thiosemicarbazide H₂NCSNHNH₂	91 128	183			
72	CH₅O₂P	Methylphosphinic acid CH₃PO(OH)₂	96 063	105			
73	CH₅P	Methylphosphine CH₃PH₂	48 063		-14		
74	CH₆ClN	Methylamine hydrochloride	67 512	226	230¹⁶		1333
75	CH₆ClN₃	Guanidine hydrochloride	95 528				
76	CH₆ClN₃O	Semicarbazide hydrochloride	111 53	173 d.			
77	CH₆N₂	Methylhydrazine CH₃NHNH₂	46 062		87.5		
78	CH₆N₄	Methyltetrazine CH₃NHN·NNH₂	74 078	130			
79	CH₆N₄O₂	Guanidine nitrite (NH₂)₂C(NH)·HNO₂	106 08	78 5			
80	CH₆N₄O₃	Guanidine nitrate	122 079				1333
81	CH₆N₄O₄	Semicarbazide nitrate	138 08	123			
82	CH₇ClN₄	Aminoguanidine hydrochloride	110 54	163			
83	C₂Br₂	Dibromoacetylene BrC≡CBr	183 83		76	2	894
84	C₂Br₂Cl₂	1, 2-Dibromo-1, 2-dichloroethylene	254 75	4 4	172	2 304¹⁵	
84 1	C₂Br₂Cl₄	1, 2-Dibromo-1, 1, 2, 2-tetrachloroethane	325 66			2 713	1308
85	C₂Br₂O₂	Oxalyl bromide (COBr)₂	215 83	-19 5	104 4		
86	C₂Br₄	Tetrabromoethylene Br₂C·CBr₂	343 66	57 5	227		
87	C₂Br₆	Hexabromoethane Br₃CCBr₃	503 50		210	3 823	1316
88	C₂Cl₂	Dichloroacetylene ClC≡CCl	94 916	-50			
89	C₂Cl₂O₂	Oxalyl chloride (COCl)₂	126 916	-12	64	1 488¹⁴ ⁴	822
90	C₂Cl₄	Tetrachloroethylene Cl₂C·CCl₂	165 83	22 4	120 8	1 623	623
91	C₂Cl₄O₂	Trichloromethyl chloroformate	197 83	-57	127 5	1 653¹⁴	
92	C₂Cl₆	Hexachloroethane Cl₃CCCl₃	236 75	185	185	2 091	
93	C₂I₂	Diiodoacetylene IC·CI	277 86	82			
94	C₂I₄	Tetraiodoethylene I₂C·CI₂	531 73	187		2 983	
95	C₂N₂	Cyanogen CN·CN	52 016	-31 4	-20 5	0 866¹⁷ ²	
96	C₂N₂S	Cyanogen sulfide (CN)₂S	84 081	60			
97	C₂N₄O₄	Trinitroacetonitrile	176 03	41 5	exp. 220		
98	C₂N₆O₁₂	Hexanitroethane (O₂N)₃CC(NO₂)₃	300 05	142 d.			
99	C₂HBr	Bromoacetylene BrC≡CH	104 924		-2		
100	C₂HBrCl₂	1, 2-Dichloro-1-bromoethylene	175 84	-83 5	113 8	1 913¹⁴	867
101	C₂HBr₃	Tribromoethylene Br₂C·CHBr	264 76		164	2 708	778
102	C₂HBr₃Cl₂	1, 2, 2-Tribromo-1, 2-dichloroethane	335 67	6	112¹⁶	2 635¹⁴	781
103	C₂HBr₃O	Bromal Br₃CCHO	280 76		174	2 30¹⁴	
104	C₂HBr₃O₂	Tribromoacetic acid Br₃CCO₂H	296 76	130	245 d.		
105	C₂HBr₅	Pentabromoethane Br₃CCHBr₂	424 59	57	210⁴⁶⁰	3 312	
106	C₂HCl₃	Trichloroethylene Cl₂C·CHCl	131.38	-86 4	88	1 477	525
107	C₂HCl₃O	Chloral Cl₃CCHO	147 38	-57 5	98.1	1 512	455
108	C₂HCl₃O	Dichloroacetyl chloride Cl₂CHCOCl	147 38		108		
109	C₂HCl₃O₂	Trichloroacetic acid Cl₃CCO₂H	163.38	57.5	195 3	1.617⁴⁴₁₆	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
110	$C_2HCl_3O_2$	Dichloromethyl chloroformate	163 38		116	1.558[14]	
111	C_2HCl_5	Pentachloroethane Cl_3CCHCl_2	202 298	−29.0	162	1.709[0]	614
112	C_2HF_3	Trifluoroethylene	82 008		−51	1.26[−71]	
112 II	$C_2HF_3O_2$	Trifluoroacetic acid F_3CCO_2H	114 01	−15.6	72.5	1.535[0]	
113	C_2HI	Iodoacetylene $IC.CH$	151.94		32		
114	$C_2HI_3O_2$	Triiodoacetic acid I_3CCO_2H	437 80	150 d.			
115	C_2H_2	Acetylene $HC.CH$	26 015	−81.8	−83 6	Liq. 0 613[−80] Sol. 0.730[−84]	
116	$C_2H_2AsCl_3$	2-Chlorovinylarsine dichloride	207 35		190	1.888	
117	C_2H_2BrCl	cis-1-Bromo-2-chloroethylene	141 39		84 7	1.797[15]	863
118	C_2H_2BrCl	trans-1-Bromo-2-chloroethylene	141 39	41	75 4	1 777[15]	864
119	C_2H_2BrClO	Chloroacetyl bromide $ClCH_2COBr$	157.39		135	1.913[0]	
120	$C_2H_2BrClO_2$	Bromochloroacetic acid $BrClCHCO_2H$	183 39	23 8	211 7 s. d.	1.985[0]	
121	$C_2H_2BrCl_3$	1-Bromo-1, 2, 2-trichloroethane	212 31	−21	104 1	2.0554[0]	
122	$C_2H_2Br_2$	1, 1-Acetylene dibromide $CH_2:CBr_2$	185 85		92	2 178	
123	$C_2H_2Br_2$	1, 2-Acetylene dibromide $BrCH:CHBr$	185 85		110.2	2 256	719
124	$C_2H_2Br_2O$	Bromoacetyl bromide $BrCH_2COBr$	201 85		150	2.317[21]	
125	$C_2H_2Br_2O_2$	Dibromoacetic acid Br_2CHCO_2H	217.85	48	232		
126	$C_2H_2Br_3Cl$	1, 2, 2-Tribromo-1-chloroethane	301.22	20 6	220 d.	2 652[14]	780
127	$C_2H_2Br_4$	1, 1, 1, 2-Tetrabromoethane $BrCH_2CBr_3$	345 68	0 0	103 5[13 5]	2.875	794
128	$C_2H_2Br_4$	1, 1, 2, 2-Tetrabromoethane	345 68	0 1	151[54]	2.964	796
129	$C_2H_2ClIO_2$	Chloroiodoacetic acid $ClCHICO_2H$	220.41	90			
130	C_2H_2ClNO	Chloromethyl isocyanate $ClCH_2CNO$	91.481		81		
132	$C_2H_2Cl_2$	cis-1, 2-Acetylene dichloride	96 931	−50.0	48.4	1 265[15]	853
133	$C_2H_2Cl_2$	trans-1, 2-Acetylene dichloride	96 931	−80 5	60.3	1.291[15]	854
134	$C_2H_2Cl_2O$	Dichloroacetaldehyde Cl_2CHCHO	112 931		90.5		
135	$C_2H_2Cl_2O$	Chloroacetyl chloride $ClCH_2COCl$	112 931		105	1.495[0]	
136	$C_2H_2Cl_2O_2$	Dichloroacetic acid Cl_2CHCO_2H	128.931	10; −4	193.5	1 563	490
137	$C_2H_2Cl_2O_2$	Chloromethyl chloroformate	128.931		108	1 516	
138	$C_2H_2Cl_3NO$	Trichloroacetamide Cl_3CCONH_2	162.40	141	240		
139	$C_2H_2Cl_4$	1, 1, 1, 2-Tetrachloroethane	167.85		130.5	1.588	528
140	$C_2H_2Cl_4$	1, 1, 2, 2-Tetrachloroethane	167.85	−43 8	146.3	1.600	567
141	$C_2H_2F_2O_2$	Difluoroacetic acid F_2CHCO_2H	96 015	−0.35	134 2[766]	1.526	4
142	$C_2H_2F_3NO$	Trifluoroacetamide F_3CCONH_2	113.023	74.8	162 5		
143	$C_2H_2I_2O_2$	Diiodoacetic acid I_2CHCO_2H	311.88	110			
144	$C_2H_2N_4$	1, 2, 4, 5-Tetrazine	82 047	99			
145	C_2H_2O	Ketene $CH_2:CO$	42 015	−151	−56		
146	$C_2H_2O_2$	Glyoxal $CHO.CHO$	58 015	15	50 4	1 14	46
147	$C_2H_2O_4$	Oxalic acid HO_2CCO_2H	90 015	189		2	1194
148	C_2H_3Br	Vinyl bromide $CH_2:CHBr$	106 939	−137 8	15.8	1 517[14]	415
149	C_2H_3BrO	Acetyl bromide CH_3COBr	122 939	−96 5	76.7	1 52[9 5]	
150	$C_2H_3BrO_2$	Bromoacetic acid CH_2BrCO_2H	138 939	50	208	1.934	
151	$C_2H_3Br_3$	1, 1, 2-Tribromoethane $BrCH_2CHBr_2$	266 77	−26	188 4	2.579	773
152	$C_2H_3Br_3O$	Tribromoethyl alcohol Br_3CCH_2OH	282 77	80	94[11]		
152 I	$C_2H_3Br_3O_2$	Bromal hydrate	298 77	53			1333
153	C_2H_3Cl	Vinyl chloride $CH_2:CHCl$	62 481		−15		
154	C_2H_3ClO	Acetyl chloride CH_3COCl	78 481	−112 0	52	1 104	76
155	$C_2H_3ClO_2$	Methyl chloroformate $ClCO_2CH_3$	94 481		71 4	1.236[15]	
156	$C_2H_3ClO_2$	Chloroacetic acid CH_2ClCO_2H	94.481	$\alpha61$ 2 $\beta56$ 3 $\gamma50$ 1 $\delta43$ 8 (?)	189 5	1 370[15]	1099
157	$C_2H_3Cl_2NO$	Dichloroacetamide $Cl_2CHCONH_2$	127.947	98	234 6		
158	$C_2H_3Cl_3$	1, 1, 1-Trichloroethane CH_3CCl_3	133.397		74 1	1 334	350
159	$C_2H_3Cl_3$	1, 1, 2-Trichloroethane $ClCH_2CHCl_2$	133 397	−36 7	113 5	1.443	506
160	$C_2H_3Cl_3O$	Trichloroethyl alcohol Cl_3CCH_2OH	149 397	17 8	152 2	1 550[23 3]	
161	$C_2H_3Cl_3O_2$	Chloral hydrate $Cl_3CCH(OH)_2$	183 41	47 4	98 d.	1 908	1258
162	C_2H_3FO	Acetyl fluoride CH_3COF	62 023	> −60	20.5	0 993[20]	
163	$C_2H_3FO_2$	Fluoroacetic acid CH_2FCO_2H	78 023	33	165		
164	C_2H_3I	Vinyl iodide $CH_2:CHI$	153 96		56	2 08[9]	
165	C_2H_3IO	Iodoacetaldehyde CH_2ICHO	169 96		80 d.		
166	C_2H_3IO	Acetyl iodide CH_3COI	169.96		108	1.98[17]	
167	$C_2H_3IO_2$	Iodoacetic acid ICH_2CO_2H	185.96	82			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
168	C₂H₃N	Acetonitrile CH₃CN.	41 031	−41	82	0 783	6
169	C₂H₃N	Methyl isocyanide CH₃NC.	41 031	−45	59 6	0 756⁴	
170	C₂H₃NO	Glycollic nitrile HOCH₂CN	57 031		183	1 104	952
172	C₂H₃NO	Methyl isocyanate CH₃N:CO	57 031		43		
173	C₂H₃NO₂	Nitroethylene CH₂:CHNO₂	73 031		98 5	1 073¹²·⁸	
174	C₂H₃NO₃	Oxamic acid HO₂CCONH₂	89 031	210 d.			
175	C₂H₃NO₄	Nitroacetic acid O₂NCH₂CO₂H	105 03	89			
176	C₂H₃NS	Methyl thiocyanate CH₃CNS	73 096	−51	133	1 088	501
177	C₂H₃NS	Methyl isothiocyanate CH₃N:CS.	73 096	35	119	1 069¹⁷₄	1052
178	C₂H₃N₃	1, 2, 4-Triazole	69 047	121	260		
179	C₂H₃N₃O₆	1, 1, 1-Trinitroethane (O₂N)₃CCH₃	165 05	56			
180	C₂H₄	Ethylene H₂C:CH₂. .	28 0308	−169 4	103 8	0 566 ¹⁰²₈	
181	C₂H₄BrCl	1-Bromo-2-chloroethane ClCH₂CH₂Br	143 405	−16 6	103 7	1 79⁰	
182	C₂H₄BrNO	Acetobromoamide CH₃CONHBr	137.96	108			
183	C₂H₄Br₂	1, 1-Dibromoethane CH₃CHBr₂	187 86		110	2 056	647
184	C₂H₄Br₂	Ethylene bromide BrCH₂CH₂Br	187 86	10 0	131 7	2 182	710
185	C₂H₄Br₂O	Dibromoethyl alcohol Br₂CHCH₂OH	203 86		181	2 35⁰	
186	C₂H₄Br₂O	sym.-Dibromomethyl ether (BrCH₂)₂O	203 86	−34	155	2 201	
187	C₂H₄ClNO	Acetochloroamide CH₃CONHCl	93 497	110			
188	C₂H₄ClNO	Chloroacetamide ClCH₂CONH₂	93 497	119 5	225 6		
189	C₂H₄Cl₂	1, 1-Dichloroethane CH₃CHCl₂	98 947	−96 7	57 3	1 174	227
190	C₂H₄Cl₂	Ethylene chloride ClCH₂CH₂Cl	98 947	−35 3	83.7	1.257	400
191	C₂H₄Cl₂O	Dichloroethyl alcohol Cl₂CHCH₂OH	114 947		146	1 145¹⁵	
192	C₂H₄Cl₂O	sym.-Dichloromethyl ether (ClCH₂)₂O	114 947		106	1 315	349
193	C₂H₄Cl₂OS	Di-(chloromethyl) sulfoxide	117 01	40			
194	C₂H₄Cl₂S	sym.-Dichloromethyl sulfide.	131 012		58 5¹⁸	1 114¹⁴₄	
195	C₂H₄Cl₃NO	Chloral ammonia Cl₃CCHO.NH₃.	164 41	74	100 d.		
196	C₂H₄I₂	1, 1-Diiodoethane CH₃CHI₂	281.9		179	2 84⁰	
197	C₂H₄I₂	Ethylene iodide ICH₂CH₂I.. .	281.9	82	d.	2 132¹⁰	
199	C₂H₄N₂O₂	Oxamide H₂NOCCONH₂...	88 047	419 d.		1 667	
200	C₂H₄N₂O₂	Glyoxime NOH:CHCH:NOH	88 047	178			
201	C₂H₄N₂O₃	Ethylnitrolic acid CH₃C(NO₂):NOH	104 047	88	d.		
202	C₂H₄N₂O₄	1, 1-Dinitroethane CH₃CH(NO₂)₂	120 047		186	1 350²¹₄	
203	C₂H₄N₂O₄	Ethylene dinitrite ONOCH₂CH₂ONO..	120 047	37.5	98	1 216⁰	
204	C₂H₄N₂O₅	Ethylene nitrite nitrate..	136 047	d.		1.472	
205	C₂H₄N₂O₆	Dinitroglycol (CH₂ONO₂)₂	152 047	−20	exp. 116	1 496¹⁵	
207	C₂H₄N₄	Dicyandiamide H₂NC(:NH)NHCN	84 063	207			
208	C₂H₄O	Acetaldehyde CH₃CHO ..	44 031	−123 5	20 2	0.781	3
209	C₂H₄O	Ethylene oxide... .	44.031	−111.3	10.7	0 887⁷₄	803
210	C₂H₄OS	Thioacetic acid CH₃COSH	76 096	<−17	93	1 074¹⁰	
211	C₂H₄O₂	Glycollic aldehyde HOCH₂CHO	60 031	97			
212	C₂H₄O₂	Acetic acid CH₃CO₂H....	60 031	16.6	118 1	1 049	26
213	C₂H₄O₂	Methyl formate HCO₂CH₃ .	60.031	−99 8	31 8	0 975	5
214	C₂H₄O₃	Glycollic acid HOCH₂CO₂H	76.031	α63 0 β79			
215	C₂H₄O₃	Methyl acid carbonate CH₃HCO₃	76.031	−57			
216	C₂H₄O₃	Ethylene ozonide.	76 031		18¹⁶		
217	C₂H₄O₅S	Sulfoacetic acid HO₃SCH₂CO₂H	140.10	86			
218	C₂H₄S	Ethylene sulfide	60 096		55	1.034	
219	C₂H₅AsO₄	Arsonoacetic acid (OH)₂AsOCH₂COOH	184 00	152			
220	C₂H₅Br	Ethyl bromide	108 955	−119 0	38.0	1.430	275
221	C₂H₅BrO	2-Bromoethyl alcohol BrCH₂CH₂OH	124 955		150.3	1.685	555
222	C₂H₅BrO	Bromomethyl methyl ether .	124.955		87	1.531¹²·³	458
224	C₂H₅Cl	Ethyl chloride ...	64 497	−138 7	12 2	0 910	
225	C₂H₅ClO₃S	Chloromethyl methyl sulfate	160 56		92¹⁸	1.473	
226	C₂H₅Cl₂N	Ethyl dichloramine C₂H₅NCl₂	113 963		89		
227	C₂H₅ClO	2-Chloroethyl alcohol ClCH₂CH₂OH	80 497	−69 0	128 8	1 213	
228	C₂H₅ClO	Chloromethyl methyl ether.	80.497		59 5	1.063¹⁰	107
229	C₂H₅ClO	Ethyl hypochlorite . .	80.497		36 6		
230	C₂H₅ClO₂S	Ethylsulfone chloride CH₃CH₂SO₂Cl	128.562		177 5	1.357	
231	C₂H₅ClO₄	Ethyl perchlorate	128.497		74		
232	C₂H₅F	Ethyl fluoride .	48.039		−32	1.7	
233	C₂H₅FO	2-Fluoroethyl alcohol FCH₂CH₂OH	64.039	−26.5	103.4	1.114	21

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
234	C₂H₅I	Ethyl iodide	155 97	−108 5	72 2	1.933	644
235	C₂H₅IO	2-Iodoethyl alcohol ICH₂CH₂OH	171 97		177 s. d.	2.905	
236	C₂H₅IO	Iodomethyl methyl ether ICH₂OCH₃.	171 97		125	2.025^{16}	728
237	C₂H₅N	Vinylamine H₂C:CHNH₂	43 047		56	0.832	
238	C₂H₅NO	Acetamide CH₃CONH₂	59 047	{ 81 0 69.4	222	1.159	1107, 1173, 1198
239	C₂H₅NO	Acetaldoxime CH₃CH:NOH	59.047	**47**	115	0 966	1070
240	C₂H₅NO₂	Acetohydroxamic acid CH₃CONHOH	75 047	88		1 161	1274
241	C₂H₅NO₂	Aminoacetic acid H₂NCH₂CO₂H	75 047	233 d.		1.056$^{15}_{15}$	84
242	C₂H₅NO₂	Nitroethane CH₃CH₂NO₂ .	75 047	< −50	114 8	0.900$^{15\ 5}$	
243	C₂H₅NO₂	Ethyl nitrite CH₃CH₂ONO	75 047		17		
244	C₂H₅NO₂	Methyl carbamate CH₃CONH₂	75 047	52	177		
245	C₂H₅NO₂	Glycolleamide HOCH₂CONH₂	75 047	120			
246	C₂H₅NO₃	Nitroethyl alcohol O₂NCH₂CH₂OH	91 047	< −80	193 8	1.270^{16}	
247	C₂H₅NO₃	Ethyl nitrate CH₃CH₂ONO₂	91 047	−102 0	88 7	1 105	54
248	C₂H₅NO₄ (H₂O)	Ammonium hydrogen oxalate	107 047			1 556	
249	C₂H₅NO₄	Nitroglycol HOCH₂CH₂NO₃	107 047	d.		1 31^{11}	
250	C₂H₅NS	Thioacetamide CH₃CSNH₂	75 112	108.5			
251	C₂H₅N₃O₂	Biuret NH(CONH₂)₂	103 063	193			
252	C₂H₆	Ethane CH₃CH₃ .	30 0162	−172 0	−88.3	0 546^{-88}	
253	C₂H₆AsBr	Cacodyl bromide (CH₃)₂AsBr	184 92		130		
254	C₂H₆AsCl	Cacodyl chloride (CH₃)₂AsCl	140 464		106 5	>1	
255	C₂H₆AsCl₃	Cacodyl trichloride (CH₃)₂AsCl₃	211 38	50 d.			
256	C₂H₆AsI	Cacodyl iodide (CH₃)₂AsI	231 94		160		
257	C₂H₆N₂O	Aminoacetamide H₂NCH₂CONH₂	74 06	65			
258	C₂H₆N₂O	Dimethylnitrosamine (CH₃)₂N NO	74 062		152 5	1 003	356
259	C₂H₆N₂O	N-Methylurea CH₃NHCONH₂	74 062	101		1 204	
260	C₂H₆N₄O₂	Oxalyl dihydrazide (CONHNH₂)₂	118 08	235 d.			
261	C₂H₆N₄S	Guanidine thiocyanate	118 143	118			
262	C₂H₆O	Ethyl alcohol C₂H₅OH	46 016	−117 3	78 5	0 789	17
263	C₂H₆O	Methyl ether CH₃OCH₃	46 046	−138 0	−24 9	1 617	
264	C₂H₆O₂	Glycol HOCH₂CH₂OH	62 046	−17 4	197 5	1 115	305
265	C₂H₆O₂S	Dimethyl sulfone (CH₃)₂SO₂	94 111	193	238		
266	C₂H₆O₃S	Methyl sulfite (CH₃)₂SO₃	110 111		126 5	1.046	
267	C₂H₆O₄	Acetyl peroxide (CH₃CO)₂O₂	94 046	30	63^{21}		
268	C₂H₆O₄S	Ethylsulfuric acid C₂H₅SO₄H	126 111		d.	1 316^{17}	
269	C₂H₆O₄S	Methyl sulfate (CH₃)₂SO₄	126 111	−31 8	188 8	1 333$^{15}_{15}$	66
270	C₂H₆O₆	Oxalic acid dihydrate	126 046	101 5		1 64	1206
271	C₂H₆O₆S₂	Ethane-1, 2-disulfonic acid	190 18	104			
272	C₂H₆S	Methyl sulfide (CH₃)₂S	62 111	−83 2	36 2	0 849	323
273	C₂H₆S	Ethylmercaptan C₂H₅SH	62 111	−121 0	34 7	0 840	
274	C₂H₆S₂	Methyl disulfide CH₃SSCH₃	94 176		118	1 046	
275	C₂H₆S₂	Ethylenemercaptan HSCH₂CH₂SH	94 176		146	1 123	
276	C₂H₆Se	Ethylhydroselenide C₂H₅SeH	109 246		53 5	1 395	
277	C₂H₆Te	Methyl telluride (CH₃)₂Te	157 546		82		
278	C₂H₇As	Dimethylarsine (CH₃)₂AsH	106 014		36	1 213^{29}	
279	C₂H₇As	Ethylarsine C₂H₅AsH₂	106 014		36	1.217	
280	C₂H₇AsO₂	Cacodylic acid (CH₃)₂AsO.OH	138 014	200			
281	C₂H₇AsO₃	Ethylarsonic acid C₂H₅AsO(OH)₂	154 014	95			
282	C₂H₇N	Dimethylamine (CH₃)₂NH	45 062	−96 0	7 4	0 680$^{0}_{4}$	
283	C₂H₇N	Ethylamine C₂H₅NH₂	45 062	−80 6	16 6	0 689$^{15}_{15}$	1333
284	C₂H₇NO	Acetaldehyde ammonia CH₃CHO NH₃	61 062	97	110 s. d.		446
285	C₂H₇NO	2-Aminoethyl alcohol H₂NCH₂CH₂OH	61 062		171	1 022^{20}	
286	C₂H₇NO	Dimethylhydroxylamine (CH₃)₂NOH.	61 062		42 4		
287	C₂H₇NO	α-Ethylhydroxylamine NH₂OC₂H₅	61 062		68	0 883$^{7\ 5}$	1098
288	C₂H₇NO	β-Ethylhydroxylamine C₂H₅NHOH	61 062	59 d.		0 908	
289	C₂H₇NO₂	Ammonium acetate CH₃CO₂NH₄	77 062	114		1 073	
290	C₂H₇NO₃S	Taurine H₂NCH₂CH₂SO₃H	125 127	88			
290 1	C₂H₇N₃	Diazoaminoethane C₂H₅N.N.NH₂	73 08	−12	92 s. d.		
291	C₂H₇N₃O₄	Methylurea nitrate	137 08	128			
292	C₂H₇O₂P	Dimethylphosphinic acid (CH₃)₂PO.OH	94 08	76			
293	C₂H₇O₂P	Ethylphosphinic acid C₂H₅PO(OH)₂. ..	110.08	44			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
294	C₂H₇P	Dimethylphosphine (CH₃)₂PH	62 078		25		
295	C₂H₇P	Ethylphosphine C₂H₅PH₂	62 078		25	<1	
296	C₂H₆BrN	Ethylamine hydrobromide	125 986	159 5		1 741	
297	C₂H₆ClN	Dimethylamine hydrochloride	81 528	171			
298	C₂H₆ClN	Ethylamine hydrochloride	81 528	109		1 216	
299	C₂H₆IN	Ethylamine hydroiodide C₂H₅NH₂·HI	173 00	188 5		2.100	
300	C₂H₈N₂	Ethylenediamine H₂NCH₂CH₂NH₂	60 078	8 5	117	0 892	1032
301	C₂H₈N₂	unsym.-Dimethylhydrazine	60 078		61	0 794	987
302	C₂H₈N₂	Ethylhydrazine C₂H₅NHNH₂	60 078		101 5		
303	C₂H₈N₂O₄(H₂O)	Ammonium oxalate	124 078			1 501	1233
304	C₂H₈N₄	Ethyltetrazine	88 094	< 20	140 d.		
305	C₂H₈N₄O₃	Methylguanidine nitrate	136 09	150			
306	C₂H₁₀Cl₂N₂	Ethylenediamine hydrochloride	133 01				1284
307	C₂H₁₀N₂O	Ethylenediamine hydrate	78 093	10	118	0 963	433
308	C₂H₁₄N₆O₄S	Aminoguanidine sulfate	246 24	161			
308 1	C₃Cl₃N₃	Cyanuric trichloride	184 40	146		1 32	
309	C₃Cl₈	Octachloropropane Cl₃CCl₂CCl₃	319 66	160	269		802
310	C₃O₂	Carbon suboxide OC:C:CO	68 00	−107	6 3	1 114⁰	
311	C₃HCl₃O₂	Trichloroacrylic acid Cl₂C:CClCO₂H	175 38	72 9	223		
312	C₃HCl₇	Heptachloropropane Cl₃CHClCCl₂CCl₃	285 21	30	248	1 805	911
313	C₃HN	Cyanoacetylene HC:CCN	51 016	5	42 5	0 816	
313 1	C₃H₂Br₂N₂O	Dibromocyanoacetamide	245 86	123		2 375	
314	C₃H₂Cl₂O₂	Malonyl chloride H₂C(COCl)₂	140 93		58°	1 450	1000
315	C₃H₂Cl₃NO	2, 2, 2-Trichlorolactic nitrile	174 40	61	220		
316	C₃H₂N₂	Malonic nitrile H₂C(CN)₂	66 031	32 1	220	1 049	1042
317	C₃H₂N₂O₃	Parabanic acid CO<(NHCO)₂>	114 031	227 d			1333
318	C₃H₄O	Propargyl aldehyde HC:CCHO	54 015		61		
319	C₃H₄O₂	Propiolic acid HC:C.CO₂H	70 015	9	144 d	1 139	
320	C₃H₃BrO₂	1-Bromoacrylic acid CH₂:CBrCO₂H	150 94	70			
321	C₃H₃BrO₂	2-Bromoacrylic acid BrCH:CHCO₂H	150 94	116			
322	C₃H₃BrO₄	Bromomalonic acid BrCH(CO₂H)₂	182 94	112 d.			
323	C₃H₅Cl	3-Chloroallylene ClCH₂C:CH	74 481		65	1 045	
323 1	C₃H₃ClO	Acryl chloride H₂C:CHCOCl	90 481		76	1 14⁰	
324	C₃H₃ClO₂	1-Chloroacrylic acid CH₂:CClCO₂H	106 48	65			
325	C₃H₃ClO₂	2-Chloroacrylic acid ClCH:CHCO₂H	106 48	85			
326	C₃H₃ClO₄	Chloromalonic acid ClCH(CO₂H)₂	138 48	133			
327	C₃H₃Cl₃O	1, 1, 1-Trichloroacetone CH₃COCCl₃	161 40		149		
328	C₃H₃Cl₃O	1, 1, 1'-Trichloroacetone	161 40		172		
329	C₃H₃Cl₃O₂	Methyl trichloroacetate Cl₃CCO₂CH₃	177 40	−17 5	153 8	1 489	
330	C₃H₃Cl₃O₃	2, 2, 2-Trichlorolactic acid	193 40	124	170°		
331	C₃H₃Cl₅	Pentachloropropane	216 31		198	1 607	645
332	C₃H₃N	Acrylic nitrile CH₂:CHCN	53 031	−82 0	79		
332 1	C₃H₃NO	Pyruvic nitrile CH₃COCN	69 04		93		
333	C₃H₃NO₂	Cyanoacetic acid NCCH₂CO₂H	85 031	66	108°		
334	C₃H₃NS	Thiazole	85 096		116 8	1 198	
335	C₃H₃N₃O₃	Cyanuric acid	129 047	>360			1333
336	C₃H₃N₃O₃	Fulminuric acid (CNOH)₃	129 05	145 d.			
337	C₃H₄	Allene H₂C:C:CH₂	40 031	−146	−32	0.660⁻¹²	
338	C₃H₄	Allylene HC:CCH₃	40 031	−104 7	−27 5	2 024	924
339	C₃H₄Br₂	cis-1, 2-Dibromopropylene	199 86		135 2	2 024	925
340	C₃H₄Br₂	trans-1, 2-Dibromopropylene	199 86		126	1 934	
341	C₃H₄Br₂	2, 3-Dibromopropylene	199 86		142 3		
342	C₃H₄Br₂O₂	1, 1-Dibromopropionic acid	231 86	61	221		
343	C₃H₄Br₂O₂	1, 2-Dibromopropionic acid	231.86	64; 51	160²⁰		
344	C₃H₄Br₄	1, 1, 2, 2-Tetrabromopropane	359.69		230 s. d.	2.94⁰	
345	C₃H₄Br₄	1, 2, 2, 3-Tetrabromopropane	359 69	11	230 d.	2 653	
346	C₃H₄Cl₂O	sym.-Dichloroacetone (ClCH₂)₂CO	126.947	45	173 4	1.383	
347	C₃H₄Cl₂O	unsym.-Dichloroacetone	126 947		120	1 234	
348	C₃H₄Cl₂O₂	2, 2-Dichloropropionic acid	142.947	56	190		
349	C₃H₄Cl₃NO	Chloral formamide Cl₃CCHO.HCONH₂	192.41	116			
350	C₃H₄N₂	Imidazole	68 047	90	256		
351	C₃H₄N₂	Pyrazole	68.047	70	188		
352	C₃H₄N₂O	Cyanoacetamide NCCH₂CONH₂	84.047	120			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
353	$C_3H_4N_2O$	Pyrazolone —NHCOCH$_2$CH:N—....	84.047	165			
354	$C_3H_4N_2O_2$	Hydantoin —NHCONHCH$_2$CO—	100.047	220			
355	C_3H_4O	Propargyl alcohol HC:CCH$_2$OH.	56.031	−17	115	0.972	324
356	C_3H_4O	Acrolein H$_2$C.CH.CHO	56.031	−87.7	52.5	0.841	119
357	C_3H_4O	Allylene oxide. . .	56 031		63		
358	$C_3H_4O_2$	Acrylic acid H$_2$CCHCO$_2$H	72 031	12.3	141.9	1.051	264
359	$C_3H_4O_3$	Pyruvic acid CH$_3$COCO$_2$H	88.031	13.6	165	1.267	873
360	$C_3H_4O_4$	Malonic acid CH$_2$(CO$_2$H)$_2$	104.031	135 6			
361	$C_3H_4O_4$	Methyl hydrogen oxalate	104 031	54	163.3	1.422^{54}	1191
362	$C_3H_4O_5$	Tartronic acid HOCH(CO$_2$H)$_2$.	120 031	158 d.			1333
363	$C_3H_4O_5$	Mesoxalic acid (HO)$_2$C(CO$_2$H)$_2$.	136 03	121			
364	C_3H_5Br	1-Bromopropylene CH$_3$CH.CHBr	120 955	−116.6	60 2	1.428$^{19.6}$	452
365	C_3H_5Br	2-Bromopropylene CH$_3$CBr:CH$_2$	120 955	−124.8	48.4	1.362^{20}	
366	C_3H_5Br	3-Bromopropylene BrCH$_2$CH:CH$_2$	120.955	−119.4	71.3	1.398	489
367	C_3H_5BrO	Bromoacetone CH$_3$COCH$_2$Br	136 955	−54	127	1 603	
368	$C_3H_5BrO_2$	dl-1-Bromopropionic acid	152 955	25 7	203.5	1.700	522
369	$C_3H_5BrO_2$	2-Bromopropionic acid	152 96	61			
370	$C_3H_5Br_3$	1, 1, 2-Tribromopropane	280 79		201	2.356	
371	$C_3H_5Br_3$	1, 2, 2-Tribromopropane .	280 79		191	2.33^{12}	
372	$C_3H_5Br_3$	1, 2, 3-Tribromopropane	280.79	17	222	2.436^{23}	767
373	C_3H_5Cl	1-Chloropropylene CH$_3$CH:CHCl	76.497		36		
374	C_3H_5Cl	2-Chloropropylene CH$_3$CCl:CH$_2$	76 497	−137 4	22.7	0 931^0	
375	C_3H_5Cl	3-Chloropropylene ClCH$_2$CH:CH$_2$	76 497	−136 4	44 6	0.938	222
376	$C_3H_5ClN_2O_6$	Chlorodinitrohydrin	200 51	6 8	123^{15}	1.54^{15}	
377	C_3H_5ClO	Chloroacetone CH$_3$COCH$_2$Cl	92 497	−44 5	121	1 162^{16}	
378	C_3H_5ClO	Propionyl chloride C$_2$H$_5$COCl	92 497	−94.0	80	1 065	152
379	C_3H_5ClO	α-Epichlorohydrin .	92 497	−25 6	117	1.184	895
380	$C_3H_5ClO_2$	Chloroacetyl carbinol	108 497	74 d.			
381	$C_3H_5ClO_2$	1-Chloropropionic acid.	108.497		186	1.306^9	
382	$C_3H_5ClO_2$	2-Chloropropionic acid	108.497	61	204		
383	$C_3H_5ClO_2$	Ethyl chloroformate ClCO$_2$C$_2$H$_5$ . .	108 497	−80 6	95	1.139$^{15.2}_{15}$	
384	$C_3H_5ClO_2$	Methyl chloroacetate ClCH$_2$CO$_2$CH$_3$	108 497	−32 7	131.5	1.22	
385	$C_3H_5Cl_3$	1, 1, 2-Trichloropropane	147.413		137	1 372^{26}	
386	$C_3H_5Cl_3$	1, 1, 3-Trichloropropane. ..	147 413		148	1 362^{16}	
387	$C_3H_5Cl_3$	1, 2, 2-Trichloropropane.	147.413		123	1 318^{16}	
388	$C_3H_5Cl_3$	1, 2, 3-Trichloropropane	147.413	−14.7	156	1.417^{16}	
389	$C_3H_5Cl_3O$	1, 1, 1-Trichloroisopropyl alcohol	163 413	50	161.3		
390	C_3H_5I	2-Iodopropylene CH$_3$CI:CH$_2$ ..	167.97		103	1.835	
391	C_3H_5I	3-Iodopropylene ICH$_2$CH:CH$_2$	167 97	−99 3	103.1	1.848$^{12}_{13}$	
392	C_3H_5IO	Iodoacetone CH$_3$COCH$_2$I	183.97		58 4^{11}	2 17^{16}	
393	$C_3H_5IO_2$	1-Iodopropionic acid CH$_3$CHICO$_2$H.	199.97	45.5	105$^{0.3}$		
394	$C_3H_5IO_2$	2-Iodopropionic acid ICH$_2$CH$_2$CO$_2$H	199 97	82			
395	C_3H_5N	Propionitrile C$_2$H$_5$CN......	55.047	−91 9	97.1	0 783	22
396	C_3H_5N	Ethyl isocyanide C$_2$H$_5$NC.	55 047	< −66	79	0 742$^{21.3}$	19
397	C_3H_5NO	Ethyl isocyanate C$_2$H$_5$CNO... .	71 047		60	0 898	
398	C_3H_5NO	Acrylamide CH$_2$:CHCONH$_2$	71 047	85			
399	C_3H_5NO	2-Hydroxypropionitrile HOCH$_2$CH$_2$CN	71.047		221	1.059	
400	C_3H_5NO	Lactonitrile CH$_3$CH(OH)CN .	71.047	−40 0	184 s. d.	0 992	944
401	$C_3H_5NO_2$	Isonitrosoacetone CH$_3$COCH(:NOH).	87.407	69			
402	$C_3H_5NO_2$	Allyl nitrite C$_3$H$_5$ONO.	87.047		44	0.955^0	
403	C_3H_5NS	Ethyl thiocyanate C$_2$H$_5$CNS	87 112	−85 5	144 4	0.996	494
404	C_3H_5NS	Ethyl isothiocyanate C$_2$H$_5$CSN	87 112	−5 9	132	0 995	651
405	$C_3H_5NS_2$	μ-Mercaptothiazoline	119.177		217		
406	$C_3H_5N_3O_6$	Glycerol trinitrate	179.06		154	1 291$^{10}_{15}$,	
407	$C_3H_5N_3O_9$	Glycerol trinitrate	227.06	2.9 13 2	160^{15} exp. 260	1 601^{15}	
408	C_3H_6	Cyclopropane	42.046	−126.6	−34.4	0 720$^{−79}$	
409	C_3H_6	Propylene CH$_3$CH:CH$_2$	42.046	−185 2	−47.0	0 609$^{−47}_{−4}$	
410	C_3H_6AsN	Cacodyl cyanide (CH$_3$)$_2$AsCN..... ..	131.014		138		
411	$C_3H_6Br_2$	1, 1-Dibromopropane CH$_3$CH$_2$CHBr$_2$	201.88		130		
412	$C_3H_6Br_2$	1, 2-Dibromopropane CH$_3$CHBrCH$_2$Br	201 88	−55 5	140	1.933	664
413	$C_3H_6Br_2$	1, 3-Dibromopropane	201 88	−34.4	167.0	1 979	671
414	$C_3H_6Br_2$	2, 2-Dibromopropane CH$_3$CBr$_2$CH$_3$.	201.88		114 5	1 783	
415	$C_3H_6Br_2O$	1, 1'-Dibromoisopropyl alcohol	217.88		219	2.11^{15}	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
416	C₃H₆Br₂O	2, 3-Dibromopropyl alcohol	217 88		219	2.168⁰	
417	C₃H₆Cl₂	1, 1-Dichloropropane CH₃CH₂CHCl₂	112 962		87	1.143¹⁴	
418	C₃H₆Cl₂	1, 2-Dichloropropane CH₃CHClCH₂Cl	112 962		96 8	1 166¹⁴	
419	C₃H₆Cl₂	1, 3-Dichloropropane ClCH₂CH₂CH₂Cl	112 962		125	1 201¹⁴	177
420	C₃H₆Cl₂	2, 2-Dichloropropane CH₃CCl₂CH₃	112 962		69 7	1 093	
421	C₃H₆Cl₂O	1, 1-Dichloroisopropyl alcohol	128 96		147.8	1 333	532
422	C₃H₆Cl₂O	1, 1'-Dichloroisopropyl alcohol	128 96		174	1 367	
423	C₃H₆Cl₂O	2, 3-Dichloropropyl alcohol	128.96		183	1 355	
424	C₃H₆Cl₂O₂	Dichloromethylal H₂C(OCH₂Cl)₂	144 96		166	1 352¹¹	
425	C₃H₃Cl₃N₃	cis-Chloralimide	103 19	155			
426	C₃H₅INO	Iodoacetoxime ICH₂C(:NOH)CH₃	198 99	64 5		2 490	
427	C₃H₆I₂	1, 2-Diiodopropane CH₃CHICH₂I	295 91		d.	2 576¹⁴	797
428	C₃H₆I₂	1, 3-Diiodopropane ICH₂CH₂CH₂I	295 91	−13 0	224	2 446⁰	
429	C₃H₆I₂	2, 2-Diiodopropane (CH₃)₂CI₂	295 91		118 d		
431	C₃H₆N₂	Pyrazoline	70 062		141		
432	C₃H₆N₂O	Ethyleneurea —CH₂NHCONHCH₂	86 062	131			
433	C₃H₆N₂O	Ethylideneurea CH₃CH:NCONH₂	86 062	154	160 d		
434	C₃H₆N₂OS	Acetylthiourea CH₃CONHCSNH₂	118 13	165			
435	C₃H₆N₂O₂	Acetylurea NH(COCH₃)₂	102 062	217			
436	C₃H₆N₂O₂	Malonamide H₂C(CONH₂)₂	102 062	170			
437	C₃H₆N₂O₂	Methylglyoxime.	102 06	153			
438	C₃H₆N₂O₃	Hydantoic acid . . .	118 062	171			
439	C₃H₆N₂O₃	Propylnitrolic acid	118 06	66			
440	C₃H₆N₂O₃	Methyl allophanate	118 06	208			
441	C₃H₆N₂O₃	Propylpseudonitrole	118 06	76			
442	C₃H₆N₂O₄	Nitrourethane C₂H₅CO₂NHNO₂	134 06	64			
443	C₃H₆N₂O₇	Glycerol-1, 3-dinitrate . .	182 06	<−30	148¹⁵	1 47¹⁴	1166
444	C₃H₆N₄O₃	Ammonium fulminurate	146 078	d.		1 573¹⁵⁰	1311
445	C₃H₆N₆	Melamine (CNNH₂)₃	126 094	<250			204
446	C₃H₆O	Allyl alcohol CH₂:CHCH₂OH	58 046	−129	97 0	0 855	20
447	C₃H₆O	Propionaldehyde C₂H₅CHO	58 046	−81	48 8	0 807	14
448	C₃H₆O	Acetone CH₃COCH₃ . .	58 046	−94.3	56.1	0 7915	315
449	C₃H₆O₂	Acetyl carbinol CH₃COCH₂OH	74 046	−17	146	1 082²⁰₂₀	63
450	C₃H₆O₂	Propionic acid C₂H₅CO₂H	74 046	−22	141.1	0 992	15
451	C₃H₆O₂	Ethyl formate HCO₂C₂H₅	74 046	−80 5	54 3	0 906	18
452	C₃H₆O₂	Methyl acetate CH₃CO₂CH₃	74 046	−98 1	57 1	0.933	
453	C₃H₆O₂	Glycide C₂H₃OCH₂OH	74 046		162 d.	1.165	
454	C₃H₆O₃	Glyceric aldehyde HOCH₂CHOHCHO	90 046	138			
455	C₃H₆O₃	Dihydroxyacetone HOCH₂COCH₂OH	90 046	75			
456	C₃H₆O₃	d(l)-Lactic acid CH₃CH(OH)CO₂H.	90 046	27	122¹⁴	1 249¹⁴	381
457	C₃H₆O₃	dl-Lactic acid CH₃CH(OH)CO₂H	90 046	18	89.7	1 069²²	
458	C₃H₆O₃	Dimethyl carbonate (CH₃O)₂CO	90 046	0 5			
459	C₃H₆O₃	Ethyl acid carbonate C₂H₅HCO₃	90 046	−57	151.2	1 108¹⁴	
460	C₃H₆O₃	Methyl glycollate HOCH₂CO₂CH₃	90 046	64	s. 46		
461	C₃H₆O₃	α-Trihydroxymethylene	74 111		90		
462	C₃H₆S	Allyl mercaptan CH₂:CHCH₂SH	166 01	128	70 9	1 353	346
463	C₃H₇AsO₃	Allylarsonic acid	122 97	−110 0	59 6	1.310	289
464	C₃H₇Br	n-Propyl bromide CH₃CH₂CH₂Br	122 97	−89 0	148		
465	C₃H₇Br	Isopropyl bromide (CH₃)₂CHBr	138 97		112¹⁵⁶	1.537	
466	C₃H₇BrO	Bromoisopropyl alcohol	138 97		46 6	0.890	71
467	C₃H₇BrO	3-Bromopropyl alcohol	78 512	−122 8	36 5	0 860	
468	C₃H₇Cl	n-Propyl chloride CH₃CH₂CH₂Cl	78 512	−117 0	126	1 115²⁰₂₀	371
469	C₃H₇Cl	Isopropyl chloride (CH₃)₂CHCl	94.512		134	1 103	354
470	C₃H₇ClO	Chloroisopropyl alcohol.	94.512		124 5¹⁴⁶	1 321	
471	C₃H₇ClO	2-Chloropropyl alcohol . .	110 512		213 d.	1 322	
472	C₃H₇ClO₂	2-Chloro-1, 3-dihydroxypropane	110 512		2		
473	C₃H₇ClO₂	3-Chloro-1, 2-dihydroxypropane	62 054		102 4	1.747	621
474	C₃H₇F	n-Propyl fluoride CH₃CH₂CH₂F	169.99	−101 4	89 5	1.703	597
475	C₃H₇I	n-Propyl iodide CH₃CH₂CH₂I	169.99	−90 8	105⁶⁰		
476	C₃H₇I	Isopropyl iodide (CH₃)₂CHI	185.99		225.4	2.349¹³	
477	C₃H₇IO	Iodoisopropyl alcohol.	185 99		53.2	0.761	237
478	C₃H₇IO	3-Iodopropyl alcohol . . .	57.062				
479	C₃H₇N	Allylamine CH₂:CHCH₂NH₂					

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
480	C_3H_7NO	Aminoacetone $CH_3COCH_2NH_2$	73 062		189 d.		
481	C_3H_7NO	Acetoxime $CH_3CH:NOH$	73.062	61	136.3	0.97^{20}_{20}	1162
482	C_3H_7NO	Propionamide $C_2H_5CONH_2$	73 062	79	213	1.042	1153
483	C_3H_7NOS	Thiourethane $C_2H_5COSNH_2$	105.13	108			
484	$C_3H_7NO_2$	d-Alanine $CH_3CH(NH_2)CO_2H$	89.062				
485	$C_3H_7NO_2$	dl-Alanine	89.062	295	s. >200		1225
486	$C_3H_7NO_2$	Sarcosine $CH_3NHCH_2CO_2H$	89 062	210 d.			
487	$C_3H_7NO_2$	1-Nitropropane $C_2H_5CH_2NO_2$	89.062		131.5	1.011^{15}	136
488	$C_3H_7NO_2$	2-Nitropropane $CH_3CH(NO_2)CH_3$	89 062		120	1.024^0	
489	$C_3H_7NO_2$	Propyl nitrite C_3H_7ONO	89 062		57	0.935	16
490	$C_3H_7NO_2$	Isopropyl nitrite $(CH_3)_2CHONO$	89 062		45	0.844^{25}	
491	$C_3H_7NO_2$	Lactamide $CH_3CH(OH)CONH_2$	89 062	74		1.138^{80}_4	
492	$C_3H_7NO_2$	Urethane $C_2H_5OCONH_2$	89 062	48	180	1.11^{20}_{20}	
493	$C_3H_7NO_3$	dl-Serine $HOCH_2CH(NH_2)CO_2H$	105 062	246 d.			
493 1	$C_3H_7NO_3$	d-Serine $HOCH_2CH(NH_2)CO_2H$	105 062	228 d.			
494	$C_3H_7NO_3$	Isoserine $H_2NCH_2CH(OH)CO_2H$	105 062	242 d.			1249
495	$C_3H_7NO_3$	Propyl nitrate $C_3H_7ONO_2$	105 062		100 5	1.053^{25}	105
496	$C_3H_7NO_3$	Isopropyl nitrate $(CH_3)_2CHONO_2$.	105 062		102	1.036	
497	$C_3H_7NO_5$	Glycerol-1-nitrate	137 06	58	160	1.40	
498	$C_3H_7NO_5$	Glycerol-2-nitrate	137 06	54	160	1.40	
499	$C_3H_7N_4O$	Acetaldehyde semicarbazone	101 08	162			
500	C_3H_8	Propane $CH_3CH_2CH_3$	44.062	−189 9	−44 5	$0 585^{-44 \, 5}$	
501	$C_3H_8ClNO_2S$	Cysteine hydrochloride	157 59	175			
502	$C_3H_8N_2O$	1, 2-Dimethylurea $CO(NHCH_3)_2$	88 078	102 5	270	1.142	
503	$C_3H_8N_2O$	1, 1-Dimethylurea $(CH_3)_2NCONH_2$	88 078	182		1.255	
504	$C_3H_8N_2O$	Ethylurea $C_2H_5NHCONH_2$	88 078	92		1.213^{18}	
505	C_3H_8O	n-Propyl alcohol $C_2H_5CH_2OH$	60 062	−127	97 8	0.804	59
506	C_3H_8O	Isopropyl alcohol $(CH_3)_2CHOH.$	60 062	−85 8	82 3	0 786	37
508	C_3H_8O	Methyl ethyl ether $CH_3OC_2H_5$	60 062		7 9	0 697	
509	$C_3H_8OS_2$	1, 2-Dithioglycerol	124 192	130 d.		$1 342^{11 \, 4}$	
510	$C_3H_8O_2$	1, 2-Propyleneglycol	76 062		189	$1 038^{23}$	
511	$C_3H_8O_2$	Trimethyleneglycol $HO(CH_2)_3OH$	76 062		214 d.	1.053	
512	$C_3H_8O_2$	Glycol methyl ether $HOCH_2CH_2OCH_3$	76 062		124 6	$0 969^{15}_{15}$	
513	$C_3H_8O_2$	Methylal $HCH(OCH_3)_2$	76 062	−104 8	44	0.862	8
514	$C_3H_8O_2S$	1-Thioglycerol $HOCH_2CH_2(OH)CH_2SH$	108 127		d.	$1.295^{14 \, 4}$	
515	$C_3H_8O_3$	Glycerol $HOCH(CH_2OH)_2$	92 062	17 9	290	1.260	512
516	$C_3H_8S_3$	Trithioglycerol $HSCH(CH_2SH)_2$	110 257	d.		$1.391^{11 \, 4}$	
517	C_3H_8S	Methyl ethyl sulfide $CH_3SC_2H_5$	76 127	−104 8	66	0 837	
518	C_3H_8S	n-Propyl mercaptan C_3H_7SH	76 127	−111 5	68		
519	C_3H_8S	Isopropyl mercaptan $(CH_3)_2CHSH.$.	76 127		60		
520	C_3H_9As	Trimethylarsine $(CH_3)_3As$	120 029		52 8	$1 124^{22}$	
521	$C_3H_9AsO_3$	Propylarsonic acid $C_3H_7AsO_3H$	168 03	126			
522	C_3H_9Bi	Trimethyl bismuthine $(CH_3)_3Bi$	254 07		110	$2 300^{18}$	
523	$C_3H_9ClN_2O$	Lactamidine hydrochloride	124 54	171			
524	C_3H_9N	n-Propylamine $C_3H_7NH_2$	59 077	−83 0	48 7	0.719	72
525	C_3H_9N	Isopropylamine $(CH_3)_2CHNH_2$	59 077	−101 2	34	0 694	875
526	C_3H_9N	Trimethylamine $(CH_3)_3N$	59 077	−124 0	3 5	$0 662^{-5 \, 2}$	
527	$C_3H_9N_3O_2$	Guanidine acetate	119 09	230			
528	$C_3H_9O_4P$	Trimethyl phosphate $(CH_3)_3PO_4$	140 09		193	$1 220^{15}$	
529	C_3H_9P	Propylphosphine $C_3H_7PH_2$	76 093		53 5		
530	C_3H_9P	Trimethylphosphine $(CH_3)_3P$	76 093		42	>1	
531	C_3H_9Sb	Trimethylstibine $(CH_3)_3Sb$	166 84		80 6	$1 523^{15}$	
532	$C_3H_{10}ClN$	Trimethylamine hydrochloride	95 543	275 d.			
533	$C_3H_{10}N_2$	dl-Propylenediamine $CH_3CH(CH_2NH_2)_2$	74 093		119	0 878	
534	$C_3H_{10}N_2$	Trimethylenediamine $H_2N(CH_2)_3NH_2$	74 093		135 5		
535	$C_3H_{12}N_6O_3$	Guanidine carbonate	180 14	197			
537	C_4Br_4S	Thiophene tetrabromide	399.73	112		$1 251^4$	1169
538	$C_4Cl_{10}O$	Perchloroether $(C_2Cl_5)_2O$	418 58	69			
539	$C_4F_6O_3$	Trifluoroacetic anhydride $(F_3CCO)_2O$	210.00	−65	40 5	1.900^{14}	
540	C_4I_2	Diiododiacetylene $IC:CC:CI$	301.86	101			
541	C_4HBr_4N	Tetrabromopyrrole	382.68	250			
542	C_4HI_4N	Tetraiodopyrrole	570.74	150 d.			
543	C_4HN_3	Cyanoform $CH(CN)_3$.	91 032	93.5			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
544	C₄H₂ClN₂O₃	5, 5-Dichlorobarbituric acid .	196 95	211 d.			
545	C₄H₂Cl₂O₂	Fumaryl chloride ClOCCH:CHCOCl.	152 93		160	1.410	938
546	C₄H₂I₂S	Thiophene diiodide	335 94	40			
547	C₄H₂N₂O₄	Alloxan OC(NHCO)₂CO	142 03	256 d.			
548	C₄H₂O₃	Maleic anhydride (:CHCO)₂O	98 015	57	202	0 934	
549	C₄H₂O₄	Acetylenedicarboxylic acid	114 02	179			
550	C₄H₃BrO₄	Bromofumaric acid........ .	194 94	186			
551	C₄H₃BrO₄	Bromomaleic acid HO₂CCBr:CHCO₂H	194 94	141			
552	C₄H₃ClN₂O₃	5-Chlorobarbituric acid....	162 50	295 s. d.			
553	C₄H₃NO₂S	2-Nitrothiophene.......... .	129 096	46 5	225		
554	C₄H₃N₃O₄	Violuric acid............	157 05	224 d.			
555	C₄H₃AsCl₂	bis-2-Chlorovinyl chloroarsine	233 36		230	1 702	
556	C₄H₄BrNS	2-Bromoallyl isothiocyanate	178 02		200		
557	C₄H₄Br₂O₄	1, 2-Dibromosuccinic acid	275 86	255			
558	C₄H₄Cl₂O₂	Succinyl chloride (CH₂COCl)₂	154 95	17	192	1 395	872
559	C₄H₄Cl₂O₃	Chloroacetic anhydride (ClCH₂CO)₂O	170 95	46	163¹¹⁴		
560	C₄H₄N₂	Succinyl nitrile (CH₂CN)₂ .	80 047	54 5	267	0 985⁶¹ ¹	1097
561	C₄H₄N₂	Pyridazine (1, 2-Diazine) .	80 017	−8	208	1 107	1015
562	C₄H₄N₂	Pyrimidine (1, 3-Diazine) .	80 047	22	124		
563	C₄H₄N₂	Pyrazine (1, 4-Diazine)..	80 017	53	118	1 031⁶¹	1091
564	C₄H₄N₂O₂	Uracil —NHCONHCH:CHCO	112 05	338			
565	C₄H₄N₂O₃	Barbituric acid OC(NHCO)₂CH₂	128 017	245	260 d.		
567	C₄H₄N₄	Hydrocyanic acid (tetramer)	108 063	179 d			
568	C₄H₄O	Tetrolic aldehyde CH₃C:C:CHO	68 031	−26	107	0 927¹⁷	913
569	C₄H₄O	Furfural (Furan).... .	68 031		31	0 937	260
570	C₄H₄O₂	Tetrolic acid CH₃C:CCO₂H	84 031	76 5	203		
571	C₄H₄O₃	Succinic anhydride . .	100 031	119 6	261	1 104	
572	C₄H₄O₃	Tetronic acid —OCH₂C(OH).CHCO ·	100 03	141			
573	C₄H₄O₄	Fumaric acid (:CHCO₂H)₂	116 031	287	290	1 635	
574	C₄H₄O₄	Maleic acid (:CHCO₂H)₂	116 031	130 5	135 d.	1 590	
575	C₄H₄O₅	Hydroxymaleic acid . .	132 03	152			
576	C₄H₄S	Thiophene	84 096	−40 0	85	1 065	693
577	C₄H₅BrO₄	Bromosuccinic acid.. .	196 95	159			
578	C₄H₅ClO	Crotonyl chloride CH₃CH:CHCOCl	104 497		125	1 091	
579	C₄H₅ClO₂	1-Chloro-α-crotonic acid	120 50	99			
580	C₄H₅ClO₂	1-Chloro-β-crotonic acid	120 50	66			
581	C₄H₅ClO₂	2-Chloro-β-crotonic acid	120 50	61			
582	C₄H₅Cl₃O	1, 1, 2-Trichlorobutyraldehyde	175.41		165 4	1 396	523
583	C₄H₅Cl₃O₂	1, 1, 2-Trichlorobutyric acid	191 41	60	238		
584	C₄H₅Cl₃O₂	1, 1, 3-Trichlorobutyric acid	191 41	75			
585	C₄H₅Cl₃O₂	Ethyl trichloroacetate Cl₃CCO₂C₂H₅	191 41		168	1 383	437
586	C₄H₅F₃O₂	Ethyl trifluoroacetate F₃CCO₂C₂H₅	142 039		61.7	1 195¹⁸	1
587	C₄H₅N	Allyl cyanide CH₂:CHCH₂CN	67 017		116.1	0 832	212
588	C₄H₅N	Allyl isocyanide CH₂:CHCH₂NC.	67 017		106	0.794¹⁷	
589	C₄H₅N	Pyrrole.	67 047		131	0 948	612
590	C₄H₅NO₂	Ethyl cyanoformate NCCO₂C₂H₅	99 047		116	1 013	
591	C₄H₅NO₂	Methyl cyanoacetate NCCH₂CO₂CH₃	99 047		200	1.123¹⁴	
592	C₄H₅NO₂	Succinimide............	99 047	124	288	1.412¹⁵	1333
593	C₄H₅NS	Allyl thiocyanate CH₂:CHCH₂CNS	99 112		161	1.050	
594	C₄H₅NS	Allyl isothiocyanate CH₂:CHCH₂CSN.	99.112	−100 0	150.7	1 010²⁰	687
595	C₄H₆	1, 2-Butadiene CH₂:C:CHCH₃	54 046		19		
596	C₄H₆	1, 3-Butadiene CH₂:CHCH.CH₂	54 046		−2 6		
597	C₄H₆	Dimethylacetylene (CH₃C·)₂	54 046		28 9		
598	C₄H₆	Ethylacetylene C₂H₅C:CH	54 046	−130	18.5	0.668⁰	101
599	C₄H₆As₂O₄	Diarsenodiacetic acid . .	267 97	205 d.			
600	C₄H₆Br₂O₂	Ethyl dibromoacetate Br₂CHCO₂C₂H₅.	245 88		194	1.903	588
601	C₄H₆Br₄	1, 1, 4, 4-Tetrabromobutane	373 71		145¹⁰	2 529	782
602	C₄H₆Br₄	1, 2, 3, 4-Tetrabromobutane	373 71	19; 39	181⁶⁰		
603	C₄H₆Br₄	2, 2, 3, 3-Tetrabromobutane	373 71	39	230		
604	C₄H₆Cl₂O₂	Ethyl dichloroacetate . . .	156 96		158.2	1 282	367
604.1	C₄H₆Cl₂O₂	Methyl 1, 2-dichloropropionate	156 96		92⁵⁰	1 328	
605	C₄H₆Cl₄O	1, 2, 2, 2-Tetrachloroethyl ether . .	211 88		189.7	1 422	
606	C₄H₆N₂	1-Methylimidazole........	82.062	−6	199	1.036¹⁰	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
607	C₄H₆N₂	4-Methylimidazole	82 062	56	262.9	1.008	
608	C₄H₆N₂	1-Methylpyrazole	82 062		127	0 993¹⁴	828
608 1	C₄H₆N₂	3-Methylpyrazole	82 062			1 020	898
608 2	C₄H₆N₂	5-Methylpyrazole	82 062		204	1 022	
609	C₄H₆N₂O₂	Ethyl diazoacetate	114 062	−22	59¹²	1.085¹⁷ ⁶	927
609 1	C₄H₆N₂O₃S	3-Methylpyrazole-4-sulfonic acid	162 22	258			1267
610	C₄H₆N₄O₃	Allantoin	158 08	235			1328
611	C₄H₆N₄O₁₂	Erythritol tetranitrate	302 08	61			
612	C₄H₆O	Methyl propargyl ether	70 046		62	0 83¹² ⁵	
613	C₄H₆O	Vinyl ether (CH₂.CH)₂O	70 046		39		
614	C₄H₆O	Crotonaldehyde CH₃CH.CHCHO	70 046	−75	104	0 859¹⁴	361
615	C₄H₆O	Dimethylketene (CH₃)₂C·CO	70 046	−97 5	34 3		
616	C₄H₆O₂	Succinic dialdehyde (CH₂CHO)₂	86 046		57¹⁰	1 064	290
617	C₄H₆O₂	α-Crotonic acid CH₃CH:CHCO₂H.	86 046	72	185	0 964⁷⁹ ⁷	1112
619	C₄H₆O₂	β-Crotonic acid CH₂:C(CH₃)CO₂H	86 046	14 6	171 9 d.	1 027	411
620	C₄H₆O₂	1-Methylacrylic acid	86.046	16	163	1.015	333
621	C₄H₆O₂	Trimethylenecarboxylic acid	86 046	17	182 5	1.088	
622	C₄H₆O₂	Vinylacetic acid CH₂:CHCH₂CO₂H	86 046	−39	163	1.013¹⁵	849
623	C₄H₆O₂	Allyl formate HCO₂C₃H₅	86 046		83	0.948¹⁴	
624	C₄H₆O₂	Methyl acrylate CH₂:CHCO₂CH₃	86 046		80 5	0.956¹⁸	113
625	C₄H₆O₂	Diacetyl CH₃COCOCH₃	86 046		88	0 975	85
626	C₄H₆O₃	Acetic anhydride (CH₃CO)₂O.	102 046	−73 0	139 6	1.082	81
627	C₄H₆O₃	1-Ketobutyric acid C₂H₅COCO₂H	102 046	32	85²¹		
628	C₄H₆O₃	Methyl pyruvate CH₃COCO₂CH₃	102 046		137	1 154⁰	
629	C₄H₆O₄	Succinic acid (CH₂CO₂H)₂	118 046	185	235	1 562	1220
630	C₄H₆O₄	Isosuccinic acid CH₃CH(CO₂H)₂	118 046	135		1 455	
631	C₄H₆O₄	Dimethyl oxalate (CO₂CH₃)₂ .	118 046	54 0	163 3	1 120⁵²	1122
632	C₄H₆O₄	Ethyl hydrogen oxalate HO₂CCO₂C₂H₅.	118 046		117¹⁵	1 218	
633	C₄H₆O₅	Diglycollic acid O(CH₂CO₂H)₂	134 05	148			
634	C₄H₆O₅	Glycollic anhydride (CH₂OHCO)₂O	134 05	130			
635	C₄H₆O₅	l-Malic acid HO₂CCH₂CH(OH)CO₂H	134 05	100	110 d.	1 595	
636	C₄H₆O₅	dl-Malic acid . .	134 05	129	150 d.	1 601	
637	C₄H₆O₅	Isomalic acid CH₃C(OH)(CO₂H)₂	134 05	160 d.			
638	C₄H₆O₆	Mesotartaric acid	150 05	140		1 666	1224
639	C₄H₆O₆	d-Tartaric acid .	150 05	170		1 760	1222
640	C₄H₆O₆	dl-Tartaric acid ...	150 05	206		1.687	
641	C₄H₆O₈	Dihydroxytartaric acid	182 05	114			
642	C₄H₆S	Divinyl sulfide (CH₂:CH)₂S .	86 111		101	0 912	
643	C₄H₇Br	Vinylethyl bromide CH₂:CHCH₂CH₂Br	134 97		99 0		
644	C₄H₇BrO	Bromomethyl ethyl ketone	150 97		146		
645	C₄H₇BrO₂	1-Bromobutyric acid C₂H₅CHBrCO₂H	166 97	−4	115²⁰	1 574¹⁵	
646	C₄H₇BrO₂	2-Bromobutyric acid	166 97	18	122¹⁶		
647	C₄H₇BrO₂	3-Bromobutyric acid	166 97	32			
648	C₄H₇BrO₂	1-Bromoethyl acetate	166 97		63³⁹	1 4620	395
648.1	C₄H₇BrO₂	2-Bromoethyl acetate	166 97		70²⁷	1 5140	450
648 2	C₄H₇BrO₂	Ethyl bromoacetate BrCH₂CO₂C₂H₅	166 97		159	1 514¹³	438
648 3	C₄H₇BrO₂	Methyl 1-bromopropionate	166 97		68 5⁴⁸	1 4917	436
648 4	C₄H₇BrO₂	Methyl 2-bromopropionate	166 97		79⁴⁶	1 5192	460
649	C₄H₇Br₃	1, 2, 3-Tribromobutane	294 80		113¹⁹	2 190	752
650	C₄H₇Br₃O	1, 1, 1-Tribromo-tert -butyl alcohol	310 80	176			
651	C₄H₇ClO	Butyryl chloride C₃H₇COCl ..	106 51	−89 0	102	1.028	194
652	C₄H₇ClO	Isobutyryl chloride (CH₃)₂CHCOCl	106 51	−90 0	92	1 017	168
653	C₄H₇ClO₂	1-Chlorobutyric acid C₂H₅CHClCO₂H.	122 51		101 3¹⁵		
654	C₄H₇ClO₂	d-2-Chlorobutyric acid ..	122 51	44	100¹³		
655	C₄H₇ClO₂	dl-2-Chlorobutyric acid	122 51	16 5	116²²	1 186	386
656	C₄H₇ClO₂	3-Chlorobutyric acid .	122.51	16	196²²	1 250¹⁰	
657	C₄H₇ClO₂	1-Chloroethyl acetate . . .	122 51		46⁴⁵	1 1124	190
657.1	C₄H₇ClO₂	2-Chloroethyl acetate	122 51		145	1.178⁰	285
658	C₄H₇ClO₂	Ethyl chloroacetate ClCH₂CO₂C₂H₅.	122 51		144 2	1.159	267
659	C₄H₇ClO₂	Methyl 2-chloropropionate .	122.51		148	1.187	
660	C₄H₇ClO₂	n-Propyl chloroformate ClCO₂C₃H₇	122 51		116	1 083¹⁵	
661	C₄H₇Cl₃O	1, 2, 2-Trichloroethyl ethyl ether	177 43		170	1.330¹⁴	
662	C₄H₇Cl₃O	1, 1, 1-Trichloro-tert.-butyl alcohol... ..	177.43	97	166.4		

No	Formula	Name	Mol. wt	M. P.	B. P.	d	R. I. No.
663	C₄H₇Cl₃O₂	Chloral alcoholate Cl₃CCHO.C₂H₅OH	193 43	55	115	1 143⁴⁰	
664	C₄H₇Cl₃O₂	1, 1, 2-Trichlorobutyraldehyde hydrate	193 43	78		1 694⁴	
665	C₄H₇FO₂	Ethyl fluoroacetate FCH₂CO₂C₂H₅	106 054			1 093	33
666	C₄H₇IO₂	Ethyl iodoacetate ICH₂CO₂C₂H₅	213 99		180	1 817¹³ ⁷	618
667	C₄H₇N	n-Butyronitrile C₃H₇CN ...	69 062	−112 0	118	0 794	47
668	C₄H₇N	Isobutyronitrile (CH₃)₂CHCN	69 062		108		
669	C₄H₇N	Isopropylisocyanide (CH₃)₂CHNC	69 062		87	0 760	
670	C₄H₇N	Pyrroline	69 062		91	0 910	
671	C₄H₇NO	Acetonecyanhydrin (CH₃)₂C(OH)CN.	85 062	−19	82²³	0 932¹⁰	117
672	C₄H₇NO	α-Pyrrolidone.	85 062	25	250 8	1 116	
673	C₄H₇NO₂	Diacetamide NH(COCH₃)₂	101 062	78	223 5		
674	C₄H₇NO₂	Diacetylmonoxime CH₃COC(:NOH)CH₃	101 062	74	186		
675	C₄H₇NO₂S	Ethyl thiooxamate H₂NCSCO₂C₂H₅.	133 13	63			
676	C₄H₇NO₃	Acetylaminoacetic acid...	117 062	206			
677	C₄H₇NO₃	Diacetohydroxamic acid.	117 06	89			
678	C₄H₇NO₃	Ethyl oxamate H₂NCO.CO₂C₂H₅...	117 06	115			
679	C₄H₇NO₄	l-Aspartic acid ...	133 06	270		1 661¹³ ¹	
679 1	C₄H₇NO₇	Nitrotetronic acid dihydrate	181 06	d. 184		1 684	1190
680	C₄H₇NO₈	Ammonium tetraoxalate	197 06	130 5		1 607	
681	C₄H₇NS	Propyl isothiocyanate	101 127		153	0 991	
682	C₄H₇N₃O	Creatinine.	113 078	260 d.			
683	C₄H₈	Cyclobutane (CH₂)₄	56 062	−50	13	0 703⁰	801
684	C₄H₈	1, 1-Dimethylethylene CH₂:C(CH₃)₂	56 062		−6		
685	C₄H₈	1, 2-Dimethylethylene CH₃CH:CHCH₃	56 062		1 4		
686	C₄H₈	Ethylethylene C₂H₅CH:CH₂	56 062	−130	−18	0.668⁰	102
687	C₄H₈	Methylcyclopropane (CH₂)₂CHCH₃	56 062		5	0 691⁻²⁰	
688	C₄H₈Br₂	1, 2-Dibromobutane C₂H₅CHBrCH₂Br	215 89		166	1.820	
689	C₄H₈Br₂	1, 3-Dibromobutane.	215 89		174	1 807	632
690	C₄H₈Br₂	1, 4-Dibromobutane Br(CH₂)₄Br	215 89	−20	198 d.	1 79¹⁴	
691	C₄H₈Br₂	2, 3-Dibromobutane (CH₃CHBr)₂CH₃.	215 89		158	1 83⁰	
693	C₄H₈Br₂	1, 2-Dibromo-2-methylpropane	215 89	−70.3	149 0	1 759	639
694	C₄H₈Br₂S	Di-(1-bromoethyl) sulfide. ..	247 96		87¹⁵	1 742	
695	C₄H₈Cl₂	1, 2-Dichloro-2-methylpropane.	126 98		108		
696	C₄H₈Cl₂O	2-Chloroethyl ether (ClCH₂CH₂)₂O	142 98		178	1 213²⁰	461
697	C₄H₈Cl₂O	1, 2-Dichloroethyl ethyl ether	142.98		145	1 174²³	
697.1	C₄H₈Cl₂O₂	Dichlorobutylene glycol	158 98	126			1177
698	C₄H₈Cl₂S	Di-(1-chloroethyl) sulfide...	159 04		67 5²⁷	1 199⁴	
699	C₄H₈Cl₂S	Di-(2-chloroethyl) sulfide (CH₃CHCl)₂S	159 04	13 5	120²⁴	1 285⁵	701
700	C₄H₈Cl₂OS	Di-(2-chloroethyl) sulfoxide	175 04	110	140²⁴ d.		
701	C₄H₈Cl₂O₂S	Di-(2-chloroethyl) sulfone.	191 04	53.5	181¹⁵		
702	C₄H₈N₂	2-Methyl-4, 5-dihydroimidazole	84 078	106	198		
703	C₄H₈N₂O₂	1-Acetyl-2-methylurea	116 08	180			
704	C₄H₈N₂O₂	Dimethyloxamide (CONHCH₃)₂	116 08	210			
705	C₄H₈N₂O₂	Dimethylglyoxime ...	116 08	246			
706	C₄H₈N₂O₂	Succinamide (CH₂CONH₂)₂	116 078	243			
707	C₄H₈N₂O₃	Ethyl allophanate H₂NCONHCO₂C₂H₅	132 08	192			
708	C₄H₈N₂O₃	l-Asparagine..	132 08	226	235 d.	1 543¹⁵	1254
709	C₄H₈N₂O₄	d-Tartaramide [CH(OH)CONH₂]₂	148 08	195			
710	C₄H₈N₂S	Allylthiourea CH₂:CHCH₂NHCONH₂	116 143	78 4		1 219²⁰	
711	C₄H₈O	Crotonyl alcohol CH₃CH:CHCH₂OH.	72 062	> −30	118	0 854	276
712	C₄H₈O	Cyclobutanol (CH₂)₃CHOH	72 062		124 1	0 923¹⁴	343
713	C₄H₈O	Cyclopropyl carbinol (CH₂)₂CHCH₂OH	72 062		124 3	0 899	850
714	C₄H₈O	Vinylethyl alcohol CH₂:CHCH₂CH₂OH	72 062		114	0 856⁰	
715	C₄H₈O	Methyl allyl ether CH₂:CHCH₂OCH₃	72 062		46	0 77¹¹	
716	C₄H₈O	Vinyl ethyl ether CH₂:CHOC₂H₅	72 062		35.5	0 763¹⁴ ¹	
717	C₄H₈O	n-Butyraldehyde C₃H₇CHO	72 062	−99 0	75 7	0.817	50
718	C₄H₈O	Isobutyraldehyde (CH₃)₂CHCHO	72 062	− 65 9	61	0.794	30
719	C₄H₈O	Methyl ethyl ketone CH₃COC₂H₅	72 062	−86 4	79 6	0.805	40
720	C₄H₈O₂	Erythrol................	88 047		196 5	1 047	
721	C₄H₈O₂	Methylacetyl carbinol (Acetoin)	88.062	15	142	1.002¹⁵	303
722	C₄H₈O₂	2-Hydroxybutyraldehyde (Aldol)	88.062		83²⁰	1.103	
723	C₄H₈O₂	n-Butyric acid C₃H₇CO₂H .	88 062	− 7 9	163.5	0 959	109
724	C₄H₈O₂	Isobutyric acid (CH₃)₂CHCO₂H	88.062	−47.0	154.4	0.949	88

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
725	C₄H₈O₂	Ethyl acetate CH₃COC₂H₅	88.062	−83.6	77.1	0.899	29
726	C₄H₈O₂	Methyl propionate C₂H₅CO₂CH₃	88.062	−87.5	79.9	0.917	36
727	C₄H₈O₂	n-Propyl formate HCO₂C₃H₇	88 062	−92.9	81.3	0.901	35
728	C₄H₈O₂	Isopropyl formate HCO₂CH(CH₃)₂.	88 062		71.3	0.883⁰	
729	C₄H₈O₃	Ethoxyacetic acid C₂H₅OCH₂CO₂H	104 062		206		
730	C₄H₈O₃	1-Hydroxybutyric acid	104 062	42 5	260		
731	C₄H₈O₃	1-Hydroxyisobutyric acid	104 062	79	212		
732	C₄H₈O₃	2-Hydroxybutyric acid	104 062		130¹⁴		
733	C₄H₈O₃	Ethyl glycollate HOCH₂CO₂C₂H₅	104 062		160	1.083²²	
734	C₄H₈O₃	Glycol acetate HOCH₂CH₂OCOCH₃	104 062		182		
735	C₄H₈O₃	Methylethyl carbonate CH₃(C₂H₅)CO₃	104 062	−14 5	109.2	1.002²⁷	
736	C₄H₈O₃	Methyl hydracrylate.	104 062		79¹²	1.118	336
737	C₄H₈O₃	Methyl lactate CH₃CH(OH)CO₂CH₃	104 062		144.8	1.08¹⁶	883
738	C₄H₈O₄	1, 2-Dihydroxybutyric acid. . .	120 06	75			
739	C₄H₈O₄	d-Methyl glycerinate	120 06		120¹⁴	1.280¹⁵	
740	C₄H₈S₂	Diethylene disulfide	120 192	112	200		
741	C₄H₉Br	n-Butyl bromide C₄H₉Br	136 99	−112 4	101 6	1.275	372
742	C₄H₉Br	Isobutyl bromide (CH₃)₂CHCH₂Br	136 99	−118 5	91.5	1.264	352
743	C₄H₉Br	sec.-Butyl bromide C₂H₅CHBrCH₃.	136 99		91 3	1 251²⁵	347
744	C₄H₉Br	tert-Butyl bromide (CH₃)₃CBr	136 99	−20	73 3	1 222	309
745	C₄H₉BrO	2-Bromoethyl ethyl ether . .	152 99		128 2	1 370⁰	
746	C₄H₉Cl	n-Butyl chloride C₄H₉Cl .	92 527	−123 1	78 0	0 884	132
747	C₄H₉Cl	Isobutyl chloride (CH₃)₂CHCH₂Cl	92 527	−131 2	68 9	0 875	98
748	C₄H₉Cl	sec-Butyl chloride C₂H₅CHClCH₃	92 527		68	0 871	110
749	C₄H₉Cl	tert.-Butyl chloride (CH₃)₃CCl	92 527	−28 5	51 0	0.840	60
751	C₄H₉ClO	1-Chloroethyl ethyl ether .	108 527		98		
752	C₄H₉ClO	tert-Butyl hypochlorite (CH₃)₃CClO .	108 527		80	0.958	
753	C₄H₉ClS	2-Chloroethyl ethyl sulfide . .	124.59		157		
754	C₄H₉I	n-Butyl iodide C₄H₉I .	184 00	−103 5	127	1 617	600
755	C₄H₉I	Isobutyl iodide (CH₃)₂CHCH₂I	184 00	−93 5	120.4	1 605	578
756	C₄H₉I	sec.-Butyl iodide C₂H₅CHICH₃	184 00	−104 0	117 5	1 595	
757	C₄H₉IO	2-Iodoethyl ethyl ether C₂H₅OCH₂CH₂I	200.00		155	1.670	
758	C₄H₉N	Crotonylamine CH₃CH CHCH₂NH₂	71 077		81		
759	C₄H₉N	Tetrahydropyrrole (Pyrrolidine)	71 077		88 5	0 871¹⁰	
760	C₄H₉NO	n-Butyramide C₃H₇CONH₂	87 077	116	216	1 032	
761	C₄H₉NO	Isobutyramide (CH₃)₂CHCONH₂	87 077	129	220	1 013	
762	C₄H₉NO	N-Dimethylacetamide CH₃CON(CH₃)₂.	87 077		165 7	0 943	365
763	C₄H₉NO	N-Ethylacetamide CH₃CONHC₂H₅	87 077		205	0 942	
764	C₄H₉NO	Methyl ethyl ketoxime	87 077		152	0.923	393
765	C₄H₉NO₂	Iminoethyl alcohol HN(CHCH₂O₂H)₂	103 077	28	270		
766	C₄H₉NO₂	1-Aminobutyric acid	103 077	285			
767	C₄H₉NO₂	2-Aminobutyric acid	103 077	184			
768	C₄H₉NO₂	3-Aminobutyric acid	103 08	193			
769	C₄H₉NO₂	1-Aminoisobutyric acid..........	103 077		280		
770	C₄H₉NO₂	Ethylaminoacetic acid	103 08	>160			
771	C₄H₉NO₂	Propyl carbamate C₃H₇OCONH₂	103 077	53	200		
772	C₄H₉NO₂	n-Butyl nitrite C₄H₉ONO	103 077		75	0 911⁰	
773	C₄H₉NO₂	Isobutyl nitrite (CH₃)₂CHCH₂ONO	103 077		67	0 877¹⁶	28
773 1	C₄H₉NO₂	Methy urethane CH₃NHCO₂C₂H₅.	103 077		170	1 009¹⁸·⁹	950
774	C₄H₉NO₃	n-Butyl nitrate C₄H₉ONO₂	119 077		136	1 048⁰	
775	C₄H₉NO₃	Isobutyl nitrate (CH₃)₂CHCH₂ONO₂	119 077		122 9	1.014²⁵	137
776	C₄H₉NO₅	d-Ammonium hydrogen malate	151 077	170			1205
777	C₄H₉NO₅	l-Ammonium hydrogen malate	151.077	161		1 509	
778	C₄H₉NO₆	Ammonium hydrogen tartrate	167.077	d.		1 680	1241
779	C₄H₉NS	1, 4-Thiazan	103.142		169		
780	C₄H₉N₃O₂	Creatine . .	131.093	295			
781	C₄H₁₀ClNO₂	Ethylaminoacetic acid hydrochloride.	139.54	144			
781 1	C₄H₁₀	n-Butane CH₃CH₂CH₂CH₃	58 077	−135 0	0 6	0 601⁰ (liq.)	
781 2	C₄H₁₀	Trimethylmethane (Isobutane) .	58 077	−145 0	−10.2		
782	C₄H₁₀N₂	Diethylenediamine (Piperazine)	86 093	105 6	146		1156
783	C₄H₁₀N₂O	Nitrosodiethylamine (C₂H₅)₂NNO.	102 093		175 4	0.951¹⁷·⁵	
784	C₄H₁₀N₂O	Trimethylurea (CH₃)₃NCONHCH₃. .	102 093	75 5	232.5		
785	C₄H₁₀N₂S	Propylthiourea C₃H₇NHCSNH₂. ..	118 16	110			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
786	C₄H₁₀N₄O₃	Guanidine lactate	132.10	d.			1236
788	C₄H₁₀N₄S₂	Ethylenediamine thiocyanate	178.24				1285
789	C₄H₁₀O	n-Butyl alcohol C₄H₉OH	74.077	−89.8	117.7	0.810	116
790	C₄H₁₀O	Isobutyl alcohol (CH₃)₂CHCH₂OH	74.077	−108	107.3	0.802	99
791	C₄H₁₀O	sec.-Butyl alcohol C₂H₅CH(OH)CH₃	74.077		99.5	0.808	104
792	C₄H₁₀O	tert.-Butyl alcohol (CH₃)₃COH..	74.077	25.5	82.8	0.789	64
793	C₄H₁₀O	Ether (C₂H₅)₂O	74.077	α −116.3 β −123.3	34.5	0.714	7
794	C₄H₁₀O	Methyl propyl ether CH₃OC₃H₇..	74.077		38.9	0.738	13
794.1	C₄H₁₀O	Methyl isopropyl ether	74.077		32.5²⁷	0.735²⁰₂₀	12
795	C₄H₁₀O₂	1, 4-Dihydroxybutane (CH₂CH₂OH)₂	90.077	16	230	1.020	
796	C₄H₁₀O₂	2, 3-Dihydroxybutane (CH₃CHOH)₂	90.077		184	1.018⁰	
797	C₄H₁₀O₂	1, 2-Dihydroxy-2-methylpropane	90.077		177	1.003	
798	C₄H₁₀O₂	Glycol dimethyl ether (CH₃OCH₂)₂	90.077		84.5	0.873	
799	C₄H₁₀O₂	Glycol ethyl ether HOCH₂CH₂OC₂H₅	90.077		135.3	0.935	
800	C₄H₁₀O₂	Diethyl peroxide (C₂H₅O)₂	90.077		65	0.827	
801	C₄H₁₀O₂	Dimethyl acetal CH₃CH(OCH₃)₂	90.077		64.4	0.866	
802	C₄H₁₀O₂S	Ethyl sulfone (C₂H₅)₂SO₂....	122.142	70	248	1.357	
803	C₄H₁₀O₂S₂	Diethyl disulfoxide C₂H₅(SO)₂C₂H₅	154.21		110 d.	1.24	
804	C₄H₁₀O₃	1, 2, 3-Trihydroxybutane.....	106.077		136²⁸	1.232¹⁷	
805	C₄H₁₀O₃	Di-(2-hydroxyethyl) ether	106.077		250	1.132	
806	C₄H₁₀O₃	Glycerol 1-methyl ether.	106.077		197	1.270²⁵₄	
807	C₄H₁₀O₃S	Diethyl sulfite (C₂H₅)₂SO₃	138.11		161.3	1.077	811
808	C₄H₁₀O₄	dl-Erythritol HOCH₂(CHOH)₂CH₂OH	122.08	126	331	1.451	1174
809	C₄H₁₀O₄S	Diethyl sulfate (C₂H₅O)₂SO₂	154.14	−26.0	208 s. d	1.172²⁵	78
810	C₄H₁₀S	n-Butyl mercaptan C₄H₉SH	90.142		98	0.836²⁰	
811	C₄H₁₀S	Isobutyl mercaptan (CH₃)₂CHCH₂SH	90.142	< −79	88	0.836	368
812	C₄H₁₀S	sec.-Butyl mercaptan C₂H₅CH(SH)CH₃	90.142		85	0.830¹⁷	
813	C₄H₁₀S	tert.-Butyl mercaptan (CH₃)₃CSH	90.142		67		
814	C₄H₁₀S	Ethyl sulfide (C₂H₅)₂S	90.142	−102.1	91.6	0.837	390
815	C₄H₁₀S₂	Ethyl disulfide (C₂H₅S)₂	122.21		153.5	0.993	630
816	C₄H₁₀Se	Ethyl selenide (C₂H₅)₂Se	137.28		108	1.230²⁷₄	1035
817	C₄H₁₀Te	Ethyl telluride (C₂H₅)₂Te..	185.58		138		
818	C₄H₁₁AsO₂	Diethylarsonic acid (C₂H₅)₂AsO(OH).	166.05	190			
819	C₄H₁₁AsO₃	N-Butylarsonic acid C₄H₉AsO(OH)₂	182.05	159			
820	C₄H₁₁N	n-Butylamine C₄H₉NH₂.....	73.093	−50.5	76	0.740²⁰	131
821	C₄H₁₁N	Isobutylamine (CH₃)₂CHCH₂NH₂	73.093	−85.5	68	0.736	111
822	C₄H₁₁N	sec.-Butylamine C₂H₅CH(NH₂)CH₃	73.093	−104.5	63	0.718²⁰	93
823	C₄H₁₁N	tert.-Butylamine (CH₃)₃CNH₂	73.093	−67.5	43.8	0.696	39
824	C₄H₁₁N	Diethylamine (C₂H₅)₂NH	73.093	−50.0	56.0	0.711	65
825	C₄H₁₁P	Diethylphosphine (C₂H₅)₂PH	90.109		85		
826	C₄H₁₂As₂	Cacodyl (CH₃)₂As.As(CH₃)₂	210.01	−6	170	> 1	
827	C₄H₁₂As₂O	Cacodyl oxide [(CH₃)₂As]₂O	226.01	−25	120	1.462¹⁵	
828	C₄H₁₂As₂S	Cacodyl sulfide [(CH₃)₂As]₂S	242.08		211		
829	C₄H₁₂BrN	Tetramethylammonium bromide	154.02			1.56	
830	C₄H₁₂BrNO	Diethylbromoacetamide...	170.02	67		1.048	
831	C₄H₁₂ClN	Diethylamine hydrochloride	109.56	217	330	1.048	
832	C₄H₁₂ClN	Tetramethylammonium chloride	109.56			1.169	
833	C₄H₁₂N₂	Tetramethylenediamine....	88.108	27	158		
834	C₄H₁₂N₂O₄	Ammonium succinate	152.11			1.367¹⁰	
835	C₄H₁₂N₂O₆	Ammonium d-tartrate	184.11	d.		1.608	1253
835.1	C₄H₁₂N₂O₆	Ammonium dl-tartrate.......	184.11			1.601	1323
836	C₄H₁₂N₄	Tetramethylammonium trinitride	116.124	125 d.			
837	C₄H₁₂OS	Dimethylethylsulfonium hydroxide	108.15	−99.5	93	0.837	
838	C₄H₁₃NO	Tetramethylammonium hydroxide	91.108	63	d.		
839	C₄H₁₆N₆O₄S	Methylguanidine sulfate	214.24	240			
840	C₅HCl₃N₄	2, 6, 8-Trichloropurine.........	223.41	187			
841	C₅HCl₄N	2, 3, 4, 5-Tetrachloropyridine	216.85	21	137²⁴		
842	C₅HCl₄N	2, 3, 4, 6-Tetrachloropyridine	216.85	75	135²⁰		
843	C₅HCl₄N	2, 3, 5, 6-Tetrachloropyridine	216.85	91	130²⁰		
844	C₅H₂Cl₃N	2, 3, 5-Trichloropyridine.....	182.40	50	120¹⁶		
845	C₅H₃Cl₂N	3, 5-Dichloropyridine..........	147.95	67			
846	C₅H₃N₃	1, 1, 1-Tricyanoethane CH₃C(CN)₃..	105.05	93.5		0.760	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
847	C₅H₄BrN	3-Bromopyridine	157.96		173	1.632¹⁰	
848	C₅H₄ClN	2-Chloropyridine	113 50		167.5	1.205¹⁵	
849	C₅H₄ClN	3-Chloropyridine	113 50		148.5		
850	C₅H₄ClN	4-Chloropyridine	113 50		148		
851	C₅H₄N₂	Glutaconic nitrile NCCH₂CH:CHCN	92 047	31.5	130¹²		
852	C₅H₄N₂O₂	3-Nitropyridine	124 05	41	216		
853	C₅H₄N₂O₄	Methylalloxan	156 05	156 d.			
853.1	C₅H₄N₂O₄ (H₂O)	3, 5-Pyrazoledicarboxylic acid	156 05			1.626	1239
854	C₅H₄N₄	Purine	120 06	217			
855	C₅H₄N₄O	Hypoxanthine	136 06	>150			
857	C₅H₄N₄O₃	Uric acid	168 06	d.		1 893	
858	C₅H₄OS	Thiophene-2-aldehyde	112 10		198	1 215	
859	C₅H₄O₂	Furfural	96 031	−38 7	161 7	1 159	685
860	C₅H₄O₂	1, 4-Pyrone	96 031	32 5	217 7	1 190⁴⁰ ⁵	1063
861	C₅H₄O₂S	Thiophene-2-carboxylic acid	128 10	126 5	260 d.		
862	C₅H₄O₂S	Thiophene-3-carboxylic acid	128 10	136			
863	C₅H₄O₃	Citraconic anhydride	112 03	7	228	1.245	508
864	C₅H₄O₃	Glutaconic anhydride	112 03	87	152¹⁵		
865	C₅H₄O₃	Itaconic anhydride	112 03	68			
866	C₅H₄O₃	Pyromeconic acid	112 03	117	228		
867	C₅H₄O₃	Pyromucic acid	112 03	133			
868	C₅H₄O₄	Aconic acid	128 03	164			1324
869	C₅H₄O₄	Glutinic acid HO₂CC:CCH₂CO₂H	128 03	146			
870	C₅H₅N	Pyridine	79 017	−42	115 3	0 982	641
871	C₅H₅NO	2-Hydroxypyridine	95 047	107	281		
872	C₅H₅NO	3-Hydroxypyridine HOC₅H₄N	95 047	129			
873	C₅H₅NO	4-Hydroxypyridine	95 047	148 5			
874	C₅H₅NO	Pyrrole-2-aldehyde CHOC₄H₄N	95 047	47			
875	C₅H₅NO₂	2, 4-Dihydroxypyridine (HO)₂C₅H₃N.	111 05	265			
876	C₅H₅NO₂	2, 6-Dihydroxypyridine (HO)₂C₅H₃N.	111.05	195			
877	C₅H₅NO₂	Pyrrole-2-carboxylic acid HO₂C.C₄H₃N.	111.05	191 5			
878	C₅H₅NO₃	2, 4, 6-Trihydroxypyridine	127 05	230 d.			
879	C₅H₅N₅	Adenine	135.08	365			
880	C₅H₆	Cyclopentadiene	66 046		42.5	0.805	903
881	C₅H₆	2-Methyl-1, 3-butenine (Valylene)	66 046		50		
882	C₅H₆N₂	2-Aminopyridine	94 062	56	204		
883	C₅H₆N₂	3-Aminopyridine	94 062	64	252		
884	C₅H₆N₂	4-Aminopyridine H₂NC₅H₄N	94 062	157			
886	C₅H₆N₂	Glutaric nitrile NC(CH₂)₃NC	94 062	−29	287 4	0 995¹⁴	1007
887	C₅H₆N₂O	2-Hydroxyglutaric nitrile	110 06		203¹¹	1 181	534
888	C₅H₆N₂O₂	Thymine	126 06	335 d.			
889	C₅H₆N₂O₃	Dimethylparabanic acid	142 06	145	277		
890	C₅H₆N₂O₃	Pyridine nitrate	142.06				1333
891	C₅H₆O	2-Methylfurfuran	82 046		64 3	0.916	
892	C₅H₆OS	Thiophene-2-alcohol	114 11		207		
893	C₅H₆O₂	Furfuryl alcohol	98.046		170 2	1 136	996
894	C₅H₆O₂	Pentinoic acid	98.046	103			
895	C₅H₆O₂	Ethyl propiolate CH·CCO₂C₂H₅	98 046		119 5	0 968¹⁴₁₅	
896	C₅H₆O₂	Propargyl acetate CH·CCH₂O₂CCH₃	98 046		125	1.005	252
897	C₅H₆O₃	Glutaric anhydride	114 05	57	287		
898	C₅H₆O₄	Citraconic acid CH₃C(CO₂H):CHCO₂H	130 05	91		1 617	
899	C₅H₆O₄	Glutaconic acid	130 05	134			
900	C₅H₆O₄	Itaconic acid CH₂:C(CO₂H)CH₂CO₂H.	130 05	161 d.		1 632	
901	C₅H₆O₄	Mesaconic acid CH₃(CO₂H)C:CHCO₂H	130 05	202	250		
902	C₅H₆O₄	Paraconic acid	130 05	58			
903	C₅H₆O₄	Trimethylene-1, 1-dicarboxylic acid	130 05	175	210³⁰		
904	C₅H₆O₅	Acetone-1-1'-dicarboxylic acid	146 05	135 d.			
905	C₅H₆O₅	1-Ketoglutaric acid	146 05	113			
906	C₅H₆N₂O₃	1-Methylbarbituric acid	142 06	132			
907	C₅H₇Cl₃O₂	Chloral acetone	205 43	76			
908	C₅H₇N	1-Methylpyrrole	81.062		115.4	0 911	892
909	C₅H₇N	2-Methylpyrrole	81 062		148	0 945	
910	C₅H₇N	3-Methylpyrrole	81.062		143		

No	Formula	Name	Mol. wt	M. P.	B. P.	d	R. I. No.
911	C₅H₇NO₂	Ethyl cyanoacetate NCCH₂CO₂C₂H₅	113 06	−22 5	206	1 063	232
912	C₅H₇NS	Crotonyl isothiocyanate	113 13		85⁵⁰	0 993⁰	
913	C₅H₈	Cyclopentene	68 062		43.6	0 776	
914	C₅H₈	2, 3-Pentadiene CH₃CH:C:CHCH₃	68 082		51	0 702	
915	C₅H₈	unsym.-Dimethylallene (CH₃)₂C⁚C⁚CH₂	68 062	−120	40 5	0 678	
916	C₅H₈	Isoprene CH₂:C(CH₃)CH:CH₂	68 062	·120	34	0 679	043
917	C₅H₈	Methylethylacetylene CH₃C:CC₂H₅	68 062		56	0 687	121
918	C₅H₈	1, 3-Pentadiene CH₃CH:CHCH:CH₂	68 062		44	0 696	901
920	C₅H₈	Propylacetylene C₃H₇C:CH	68 062	−95	40	0 722⁰	932
921	C₅H₈	Isopropylacetylene (CH₃)₂CHC:CH	68 062		29 3	0 685⁰	
921 1	C₅H₈Cl₂O₂	Ethyl 1, 2-dichloropropionate	170 98		184	1 246	424
921 2	C₅H₈N₂	3, 4-Dimethylpyrazole	96 078	58		0 933⁹⁷ ³	1131
922	C₅H₈N₂	3, 5-Dimethylpyrazole	96 078	107	220		
923	C₅H₈N₄O₆	Uroxanic acid	220 09	162 d			
924	C₅H₈O	Cyclopentanone	84 062		130 6	0 951	353
925	C₅H₈O	Ethyl propargyl ether CH:CCH₂OC₂H₅	84 062		80	0 833	325
926	C₅H₈O	Tiglic aldehyde CH₃CH:C(CH₃)CHO	84 062		116 5	0 870	430
927	C₅H₈O	Ethylideneacetone CH₃CH:CHCOCH₃	84 062		121	0 856	370
928	C₅H₈O₂	Levulinic aldehyde	100 062		188	1 018	295
929	C₅H₈O₂	Acetylacetone CH₃COCH₂COCH₃	100 062	−23 2	137	0 976	439
930	C₅H₈O₂	Allylacetic acid CH₂:CH(CH₂)₂CO₂H	100 062	< −18	189	0 984	805
931	C₅H₈O₂	Angelic acid	100 062	45	185	0 983⁴⁴ ⁷	1069
932	C₅H₈O₂	2, 2-Dimethylacrylic acid	100 062	70	195		
933	C₅H₈O₂	1-Ethylacrylic acid CH₂:C(C₂H₅)CO₂H	100 062	45	180		
934	C₅H₈O₂	1, 2-Pentenic acid C₂H₅CH:CHCO₂H	100 062	10	108¹⁷	0 990	904
935	C₅H₈O₂	2, 3-Pentenic acid	100 062		95¹⁴	0.987	949
936	C₅H₈O₂	Tiglic acid CH₃CH:C(CH₃)CO₂H	100 062	64	198 5	0 872	1121
937	C₅H₈O₂	Allyl acetate CH₃CO₂C₃H₅	100 062		105	0 928	146
938	C₅H₈O₂	Ethyl acrylate C₂H₃COC₂H₅	100 062		99 8	0 924	
939	C₅H₈O₂	Methyl α-crotonate	100 062		120 7	0 981⁴	
941	C₅H₈O₃	Levulinic acid CH₃COCH₂CH₂CO₂H	116 06	33 1	246	1 143¹⁷	383
942	C₅H₈O₃	Ethyl pyruvate CH₃COCO₂C₂H₅	116 06		144	1 060¹⁶	882
943	C₅H₈O₃	Methyl acetoacetate	116 06		170	1 077	241
944	C₅H₈O₄	Dimethylmalonic acid (CH₃)₂C(CO₂H)₂	132 06	193			
945	C₅H₈O₄	Ethylmalonic acid C₂H₅CH(CO₂H)₂	132 06	111 5	160 d		
946	C₅H₈O₄	Glutaric acid CH₂(CH₂CO₂H)₂	132 06	97 5	304	1 192¹⁰⁶	1151
947	C₅H₈O₄	Pyrotartaric acid	132 06	111		1 411	1333
947 1	C₅H₈O₄	Methyltetronic lactone	132 06	123			1213
948	C₅H₈O₄	Dimethyl malonate H₂C(CO₂CH₃)₂	132 06	−62	181 5	1 154	206
949	C₅H₈O₄	Ethyl hydrogen malonate	132 06		117²¹	1 176	301
950	C₅H₈O₄	Methyl ethyl oxalate	132 06		173.7	1 156⁰	
951	C₅H₈O₄	Methylene diacetate CH₂(CO₂CH₃)₂	132 06		170		
952	C₅H₈O₅	α-Citramalic acid	148 06	95			
953	C₅H₈O₅	dl-Citramalic acid	148 06	117			
954	C₅H₈O₅	β-Methylmalic acid	148 06	123			
955	C₅H₈O₅	Arabonic lactone	148 06	98			
956	C₅H₈O₅	Dimethyl tartronate	148 06	53 3			
957	C₅H₈O₆ (H₂O)	d-Methyl hydrogen tartrate	164 06	76			
958	C₅H₈O₇	Aposorbinic acid	180 06	110			
959	C₅H₉BrO₂	1-Bromovaleric acid C₃H₇CHBrCO₂H	180 99		105¹⁰		
960	C₅H₉BrO₂	2-Bromovaleric acid	180 99	60			
961	C₅H₉BrO₂	3-Bromovaleric acid	180 99	40			
962	C₅H₉BrO₂	2-Bromoisovaleric acid	180 99	73 5			
963	C₅H₉BrO₂	Ethyl 1-bromopropionate	180 99		160	1 393	419
964	C₅H₉Br₃	1, 2, 3-Tribromopentane	308 82		128²¹	2 095¹⁴	743
965	C₅H₉Cl	Isoprene hydrochloride	104 53		109	0 933	
966	C₅H₉ClO	n-Valeryl chloride C₄H₉COCl	120 53		128	1 016¹⁵	223
967	C₅H₉ClO	Isovaleryl chloride (CH₃)₂CHCH₂COCl	120 53		113		
968	C₅H₉ClO₂	Ethyl 1-chloropropionate	136 53		146	1 087	235
969	C₅H₉ClO₂	Ethyl 2-chloropropionate	136 53		162 5	1.114	236
969.1	C₅H₉ClO₂	n-Butyl chloroformate ClCO₂C₄H₉	136 53		138 9	1.078	807
970	C₅H₉ClO₂	Isobutyl chloroformate	136 53		130	1.040²⁵	
971	C₅H₉IO₂	Ethyl 2-iodopropionate	228 00		202	1.679¹⁴	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
972	C₅H₉N	n-Valeryl nitrile C₄H₉CN .	83.077		141	0.801	82
973	C₅H₉N	Isovaleryl nitrile (CH₃)₂CHCH₂CN...	83.077		129.3	0.802	
974	C₅H₉NO	Piperidone	99.077	40	256		
975	C₅H₉NO₃	Acetylurethane CH₃CONHCO₂C₂H₅..	131.08	78	215		
975.1	C₅H₉NO₃	α-Acetylaminopropionic acid	131.08	133			1215
976	C₅H₉NO₄	dl-Glutaminic acid	147.08	198		1.460	1261
977	C₅H₉NO₄	d-Glutaminic acid	147.08	208 d.		1.538	1266
978	C₅H₉NS	Isobutyl isothiocyanate	115.14		162	0.943	
979	C₅H₁₀	Cyclopentane CH₂<(CH₂CH₂)₂>	70.077	−93 3	49.5	0.754	843
980	C₅H₁₀	1, 1-Dimethyltrimethylene	70.077		21	0.660	
981	C₅H₁₀	Methylcyclobutane	70.077		42		
982	C₅H₁₀	β-Amylene CH₃CH CHC₂H₅.	70.077	−139	36 4	0 651	921
983	C₅H₁₀	α-Amylene C₂H₅C(CH₃):CH₂	70.077		32	0.667⁰₀	880
984	C₅H₁₀	n-Propylethylene C₃H₇CH:CH₂ ..	70.077		40		31
985	C₅H₁₀	2-Methyl-3-butene CH₂:CHCH(CH₃)₂.	70.077	−135	20 1	0 632¹⁵	
986	C₅H₁₀	2-Methyl-2-butene CH₃CH:C(CH₃)₂ .	70.077	−124	38 4	0 668¹³	
987	C₅H₁₀Br₂	1, 5-Dibromopentane CH₂(CH₂CH₂Br)₂	229.91	−35	224	1.706¹⁸	
988	C₅H₁₀Br₂	2, 3-Dibromopentane C₂H₅(CHBr)₂CH₃	229.91		175	1.7087⁰	866
988.1	C₅H₁₀ClNO₄	d(l)-Glutaminic acid hydrochloride	183.54	193			1240
989	C₅H₁₀Cl₂	3, 3-Dichloro-2-methylbutane	140.99		145	1 065	
990	C₅H₁₀Cl₂	1, 4-Dichloropentane	140.99		61¹⁷		
991	C₅H₁₀Cl₂	1, 5-Dichloropentane CH₂(CH₂CH₂Cl)₂.	140.99		178		
992	C₅H₁₀Cl₂	2, 3-Dichloropentane C₂H₅(CHCl)₂CH₃	140.99		139		
993	C₅H₁₀N₂	Diethyleyanamide NCN(C₂H₅)₂	98.093		187 d.	0 854	1072
994	C₅H₁₀N₂O₂	1-Nitropiperidine	130.09	−5.5	245	1 158	1033
994.1	C₅H₁₀N₂O₂	Dimethylmalonamide	130.09	198			1208
995	C₅H₁₀N₂O₃	dl-Glutamine .	146.09	256			
996	C₅H₁₀N₂O₄	Amylene nitrosate	162.09	99			1207
997	C₅H₁₀O	Cyclopentanol	86.077		141	0 946	
998	C₅H₁₀O	Methylallyl carbinol	86.077		116 4	0 834	
999	C₅H₁₀O	Vinylethyl carbinol	86.077		114 7	0 837	277
1000	C₅H₁₀O	2-Pentene-1-ol	86.077		64⁶²	0.838	933
1001	C₅H₁₀O	Ethyl allyl ether C₂H₅OCH₂CH:CH₂	86.077		67 6	0 765	69
1002	C₅H₁₀O	Isovaleraldehyde (CH₃)₂CHCH₂CHO	86.077	−51	92 5	0 803¹⁷	79
1003	C₅H₁₀O	Trimethylacetaldehyde (CH₃)₃CCHO	86.077	3	75	0.793	
1004	C₅H₁₀O	n-Valeric aldehyde C₄H₉CHO	86.077		103 4	0 819¹¹	70
1005	C₅H₁₀O	Diethyl ketone (C₂H₅)₂CO .	86.077	−42.0	101 7	0 814	86
1006	C₅H₁₀O	Methyl propyl ketone CH₃COC₃H₇.	86.077	−77.8	101.7	0 812¹⁵₁₅	75
1007	C₅H₁₀O	Methyl isopropyl ketone	86.077	−92.0	93	0 815¹⁵	62
1008	C₅H₁₀O	Pentamethylene oxide	86.077		87	0 880⁰	
1009	C₅H₁₀O₂	3-Acetylpropyl alcohol	102.08		209	1 016⁰	
1010	C₅H₁₀O₂	dl-Methylethylacetic acid .	102.08	< −80	174	0 941	153
1011	C₅H₁₀O₂	Trimethylacetic acid (CH₃)₃CCO₂H...	102.08	35 5	163 8	0 905⁵⁰	1050
1012	C₅H₁₀O₂	n-Valeric acid C₄H₉CO₂H	102.08	−59; −31 5	187 0	0 942	175
1013	C₅H₁₀O₂	Isovaleric acid (CH₃)₂CHCH₂CO₂H	102.08	−37 6	176 7	0 937¹⁵₅	145
1014	C₅H₁₀O₂	n-Butyl formate HCO₂C₄H₉ ..	102.08	−90 0	106 8	0 911⁰	74
1015	C₅H₁₀O₂	d-sec-Butyl formate..	102.08		97	0 882	48
1016	C₅H₁₀O₂	Isobutyl formate (CH₃)₂CHCH₂CO₂H	102.08	−95 3	98 2	0 875	58
1017	C₅H₁₀O₂	Ethyl propionate C₂H₅CO₂C₂H₅...	102.08	−72 6	99 1	0 891	51
1018	C₅H₁₀O₂	Methyl n-butyrate C₃H₇CO₂CH₃	102.08	< −95	102 3	0.898	68
1019	C₅H₁₀O₂	Methyl isobutyrate (CH₃)₂CHCO₂CH₃	102.08	−84 7	92 6	0 891	49
1020	C₅H₁₀O₂	n-Propyl acetate CH₃CO₂C₃H₇......	102.08	−92 5	101.6	0 887	52
1021	C₅H₁₀O₂	Isopropyl acetate CH₃COCH₂(CH₃)₂	102.08	−73.4	89	0 877¹⁵ ⁶	
1022	C₅H₁₀O₂S	Ethyl thiocarbonate CS(OC₂H₅)₂.	134.14		162	1 028	939
1023	C₅H₁₀O₃	1-Hydroxyvaleric acid..........	118.08	31			
1024	C₅H₁₀O₃	1-Hydroxyisovaleric acid..... .	118.08	86			
1025	C₅H₁₀O₃	2-Hydroxyvaleric acid...... .	118.08	< −32			
1026	C₅H₁₀O₃	Diethyl carbonate (C₂H₅O)₂CO .. .	118.08	−43.0	125.8	0 979	57
1027	C₅H₁₀O₃	Ethyl hydracrylate	118.08		84¹²	1.064¹⁵₅	2⁷
1028	C₅H₁₀O₃	Ethyl lactate CH₃CH(OH)CO₂C₂H₅...	118.08		154	1 031	
1028.1	C₅H₁₀O₃	Methyl l-1-methoxypropionate	118.08		131	0.9986¹⁴ ⁴	892
1029	C₅H₁₀O₃	Propyl glycollate HOCH₂CO₂C₃H₇ .	118.08		170.5	1.062¹⁸	
1030	C₅H₁₀O₄	Ethyl glycerate................. .	134.08		121¹⁴	1.191¹⁵₁₅	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1031	C₅H₁₀O₄	Glycerol acetate (Monoacetin)	134.08		158¹⁴⁴	1 20	
1032	C₅H₁₀O₅	d(l)-α-Arabinose	150 08	159 5		1 585	1243
1033	C₅H₁₀O₅	d(l)-β-Arabinose	150 08			1 605	1248
1034	C₅H₁₀O₅	dl-Arabinose	150 08	164 5			
1035	C₅H₁₀O₅	d-Lyxose	150 08	105		1.545	1228
1036	C₅H₁₀O₅	d-Ribose	150 08	87			
1037	C₅H₁₀O₅	l-Xylose	150 08	153		1.525	1231
1038	C₅H₁₀O₅	dl-Xylose	150 08	131			
1039	C₅H₁₀O₆	Arabonic acid HO₂C(CHOH)₃CH₂OH	166 08	89			
1040	C₅H₁₁Br	n-Amyl bromide CH₃(CH₂)₄Br	151 00		127.9	1 223	401
1041	C₅H₁₁Br	Isoamyl bromide (CH₃)₂CHCH₂CH₂Br	151 00		121	1 215	378
1042	C₅H₁₁Br	tert.-Amyl bromide (CH₃)₂(C₂H₅)CBr	151 00		109 2	1 190	389
1043	C₅H₁₁Cl	n-Amyl chloride CH₃(CH₂)₄Cl	106 54		105 7	0 883	191
1044	C₅H₁₁Cl	Isoamyl chloride (CH₃)₂CHCH₂CH₂Cl	106 54		99 1	0 893	181
1045	C₅H₁₁Cl	tert.-Amyl chloride (CH₃)₂(C₂H₅)CCl	106 54	−72 9	85 7	0 870¹⁵	155
1046	C₅H₁₁Cl	sec.-Amyl chloride C₃H₇(CH₃)CHCl	106 54		105	0.870	157
1047	C₅H₁₁Cl	3-Chloropentane (C₂H₅)₂CHCl	106 54		105	0 895	
1048	C₅H₁₁ClO	tert.-Amyl hypochlorite	122 54		76 3	0 855	
1049	C₅H₁₁F	n-Amyl fluoride CH₃(CH₂)₄F	90 085	> −80	62 8	0 788	11
1050	C₅H₁₁F	Isoamyl fluoride (CH₃)₂CHCH₂CH₂F	90 085	< −11	53 5		
1051	C₅H₁₁I	n-Amyl iodide CH₃(CH₂)₄I	198 02		156	1 517	572
1052	C₅H₁₁I	Isoamyl iodide (CH₃)₂CHCH₂CH₂I	198 02		148	1 510	
1053	C₅H₁₁I	tert.-Amyl iodide (CH₃)₂(C₂H₅)CHI	198 02		125	1 497¹⁹	
1054	C₅H₁₁N	Piperidine	85 093	−9	105 8	0 860	444
1055	C₅H₁₁NO	Diethylketoxime (C₂H₅)₂C:NOH	101 09		168 3	0 914	407
1056	C₅H₁₁NO	Methylpropylketoxime	101 09		168	0 900	403
1057	C₅H₁₁NO	Valeramide C₄H₉CONH₂	101 09	106		1.023	
1058	C₅H₁₁NO	Isovaleramide (CH₃)₂CHCH₂CONH₂	101 09	137	232	0 965	
1059	C₅H₁₁NO₂	1-Aminovaleric acid	117.09	291.5			
1060	C₅H₁₁NO₂	3-Aminovaleric acid	117.09	193			
1061	C₅H₁₁NO₂	4-Aminovaleric acid	117.09	157			
1062	C₅H₁₁NO₂	2-Aminoisovaleric acid	117.09	217			
1063	C₅H₁₁NO₂	n-Amyl nitrite CH₃(CH₂)₄ONO	117.09		104⁷⁶	0 853	56
1064	C₅H₁₁NO₂	Isoamyl nitrite (CH₃)₂CH(CH₂)₂ONO	117.09		99	0 872	67
1065	C₅H₁₁NO₂	tert.-Amyl nitrite (CH₃)₂(C₂H₅)CONO	117 09		93	0.903⁶	
1066	C₅H₁₁NO₂	n-Butyl carbamate C₄H₉CO₂NH₂	117 09	51			
1067	C₅H₁₁NO₂	Isobutyl carbamate H₂NCO₂C₄H₉	117 09	67	206		
1067 1	C₅H₁₁NO₂	Ethylurethane C₂H₅NHCO₂C₂H₅	117.09		176	0 981	262
1068	C₅H₁₁NO₂	Betaine	117 09	273 d			
1069	C₅H₁₁NO₂	dl-Valine (CH₃)₂CHCH(NH₂)CO₂H	117.09	298 d.			
1069 1	C₅H₁₁NO₂	d-Valine	117.09	315			1327
1070	C₅H₁₁NO₃	Isoamyl nitrate	133 09		148	0 996²¹⁷	200
1070.1	C₅H₁₁NO₃	Bios	133 09	223			1163
1070 2	C₅H₁₁NO₄	Methyltetronic amide	149 09	135 d.			1218
1071	C₅H₁₁NO₅	l-Arabinose oxime	165 09	139			
1072	C₅H₁₂	2-Methylbutane (Isopentane)	72 092	−159 7	28 0	0 621¹⁹⁴	9
1073	C₅H₁₂	n-Pentane CH₃(CH₂)₃CH₃	72 092	−131 5	36 2	0.631	10
1074	C₅H₁₂	2, 2-Dimethypropane (CH₃)₄C	72 092	−20	9.5		
1075	C₅H₁₂ClN	Piperidine hydrochloride	121 56	237			
1076	C₅H₁₂ClNO₂	Betaine hydrochloride	153 56	235			
1077	C₅H₁₂N₂O	1, 2-Diethylurea CO(NHC₂H₅)₂	116 11	106	263	1.042	
1078	C₅H₁₂O	n-Amyl alcohol CH₃(CH₂)₃CH₂OH	88 092	−78 5	137 9	0.817²⁰	823
1079	C₅H₁₂O	Isoamyl alcohol* (CH₃)₂CHCH₂CH₂OH	88 092	−117 2	130 5	0 812	166
1080	C₅H₁₂O	Diethyl carbinol (C₂H₅)₂CHOH	88 092		115 6	0 815²⁵	179
1081	C₅H₁₂O	tert.-Amyl alcohol (CH₃)₂(C₂H₅)COH	88 092	−11 9	101.8	0 809	158
1082	C₅H₁₂O	tert.-Butyl carbinol	88 092	53	114		
1083	C₅H₁₂O	d-Amyl alcohol CH₃(C₂H₅)CHCH₂OH	88.092		128	0.816	
1084	C₅H₁₂O	sec.-Amyl alcohol CH₃(C₃H₇)CH₂OH	88 092		119.5	0.809	165
1084.1	C₅H₁₂O	d-sec.-Amyl alcohol	88 092		118	0.8103	154
1085	C₅H₁₂O	Methyl isopropyl carbinol	88.092		114	0.819	
1085.1	C₅H₁₂O	d-Methyl isopropyl carbinol	88.092			0.818	106
1086	C₅H₁₂O	Ethyl propyl ether C₂H₅OC₃H₇	88.092	< −79	61.4	0.732	24
1087	C₅H₁₂O	Ethyl isopropyl ether C₂H₅OCH(CH₃)₂	88.092		54	0.745⁰	

* Commercially known as "Amyl alcohol."

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1088	$C_5H_{12}O$	Methyl n-butyl ether $CH_3OC_4H_9$	88 092		70 3	0.764^0	
1089	$C_5H_{12}O_2$	Pentane-1, 2-diol $C_3H_7CHOHCH_2OH$	104 09		211 8	0.980^{20}_{20}	376
1090	$C_5H_{12}O_2$	Pentane-1, 5-diol $CH_2(CH_2CH_2OH)_2$	104 09		239.4	0.994^{20}_{20}	432
1091	$C_5H_{12}O_2$	Methylene diethyl ether $CH_2(OC_2H_5)_2$	104 09		89	0.851^0	
1092	$C_5H_{12}O_3$	Glycerol 1-ethyl ether	120 09		230	1 091	
1093	$C_5H_{12}O_4$	Pentaerythritol	136 09	253			1178
1094	$C_5H_{12}O_5$	Adonitol	152 09	102			1333
1095	$C_5H_{12}O_5$	d-Arabitol	152 09	103			
1096	$C_5H_{12}S$	n-Amyl mercaptan $C_5H_{11}SH$	104.16		126	0.857^{20}	396
1097	$C_5H_{12}S$	act-Amyl mercaptan	104.16		118	0.848^{13}	
1098	$C_5H_{12}S$	Isoamyl mercaptan	104.16		129 5	0 835	379
1099	$C_5H_{13}N$	n-Amylamine $C_5H_{11}NH_2$	87.108	$-55\ 0$	104	0.766^{18}	
1100	$C_5H_{13}N$	Isoamylamine $(CH_3)_2CHCH_2CH_2NH_2$	87.108		95	0 751	176
1101	$C_5H_{13}N$	sec-Amylamine $CH_3(C_3H_7)CH_2NH_2$	87.108		91	0 749	
1102	$C_5H_{13}N$	$tert$-Amylamine $(CH_3)_2(C_2H_5)CNH_2$	87.108	$-105\ 0$	78		
1103	$C_5H_{13}NO_2$	Ammonium valerate	119 11				1333
1105	$C_5H_{14}N_2$	Pentamethylenediamine	102 12	9	178	0.885^{15}_{14}	482
1106	$C_6Br_4O_2$	Bromanil $OC:(CBr(CBr)_2CO$	423 66	300			
1107	C_6Br_6	Hexabromobenzene	551.50	306			
1108	C_6Br_6O	"Hexabromophenol".	367 50	128			
1109	$C_6Cl_4O_2$	Chloranil $OC.(CClCCl)_2:CO$	245 83	290			
1110	C_6Cl_6	Hexachlorobenzene	284 75	226	326 *	1.569^{234}	
1111	C_6Cl_6O	"Hexachlorophenol"	300.75	46			
1111.1	C_6Cl_8O	β-Octachlorocyclohexenone	371.67	90		2 016	1292
1111.2	C_6Cl_8O	γ-Octachlorocyclohexenone	371 67	89		2 058	1305
1112	C_6I_6	Hexaiodobenzene	833 59	350 d.			
1113	C_6HBr_5	Pentabromobenzene	472.59	293			
1114	C_6HBr_5O	Pentabromophenol $C(Br_5)OH$	488 59	225			
1115	$C_6HCl_3O_2$	Trichloroquinone	211.38	168			
1116	$C_6HCl_4NO_2$	2, 3, 4, 5-Tetrachloronitrobenzene	260 85	64 5			
1117	$C_6HCl_4NO_2$	2, 3, 4, 6-Tetrachloronitrobenzene	260 85	22			
1118	$C_6HCl_4NO_2$	2, 3, 5, 6-Tetrachloronitrobenzene	260 85	99	304 d.		
1119	C_6HCl_5	Pentachlorobenzene	250 30	86	277	$1 842^{10}$	
1120	C_6HCl_5O	Pentachlorophenol HOC_6Cl_5	266 30	188	310 2	1.978	
1121	$C_6HN_5O_{11}$	Pentanitrophenol $C_6(NO_2)_5OH$	319 05	190 d.			
1122	$C_6H_2BrN_3O_6$	Picryl bromide 2, 4, 6$(NO_2)_3C_6H_2Br$.	291 96	123			
1122.1	$C_6H_2Br_2N_2O_4$	1, 2-Dinitro-4, 5-dibromobenzene	325.86	115		2.313	
1122.2	$C_6H_2Br_2N_2O_4$	1, 3-Dinitro-4, 6-dibromobenzene	325 86	117		2.295	
1123	$C_6H_2Br_4$	1, 2, 3, 5-Tetrabromobenzene	393 68	98.5	329		
1124	$C_6H_2Br_4$	1, 2, 4, 5-Tetrabromobenzene	393.68	178		3.027	
1125	$C_6H_2Br_4O$	2, 3, 4, 6-Tetrabromophenol	409 68	120			
1126	$C_6H_2Br_5N$	Pentabromoaniline $C_6(Br_5)NH_2$	487.60	222			
1127	$C_6H_2ClN_3O_6$	Picryl chloride $(NO_2)_3C_6H_2Cl$	247.50	83		1.797	
1128	$C_6H_2ClN_3O_6$	5-Chloro-1, 2, 4-trinitrobenzene	247.50	116			
1129	$C_6H_2Cl_2O_2$	2, 5-Dichloroquinone	176.93	161			
1130	$C_6H_2Cl_2O_2$	2, 6-Dichloroquinone	176.93	121			
1131	$C_6H_2Cl_3NO_2$	2, 3, 4-Trichloronitrobenzene	226.40	56			
1132	$C_6H_2Cl_3NO_2$	2, 3, 6-Trichloronitrobenzene	226.40	89			
1133	$C_6H_2Cl_3NO_2$	2, 4, 5-Trichloronitrobenzene	226.40	57	288	1 790	
1134	$C_6H_2Cl_3NO_2$	2, 4, 6-Trichloronitrobenzene	226 40	68			
1135	$C_6H_2Cl_4$	1, 2, 3, 4-Tetrachlorobenzene	215.85	47 5	254		
1136	$C_6H_2Cl_4$	1, 2, 3, 5-Tetrachlorobenzene	215.85	51	246		
1137	$C_6H_2Cl_4$	1, 2, 4, 5-Tetrachlorobenzene	215.85	138	246	1.734^{10}	
1138	$C_6H_2Cl_4O$	2, 3, 4, 6-Tetrachlorophenol	231.85	69	164^{21}		
1139	$C_6H_2Cl_4O_2$	Tetrachlorohydroquinone	247.85	232			
1140	$C_6H_2Cl_5N$	Pentachloroaniline $C_6(Cl_5)NH_2$	265.31	232			
1141	$C_6H_2IN_3O_6$	Picryl iodide $(NO_2)_3C_6H_2I$	338.97	165		$2 285^{22\ 5}$	
1142	$C_6H_2I_2N_2O_4$	2, 4-Diiodo-1, 3-dinitrobenzene	419.90	162			1315
1143	$C_6H_2I_2N_2O_4$	4, 6-Diiodo-1, 3-dinitrobenzene	419.90	168.4		2 744	
1144	$C_6H_2I_4$	1, 2, 3, 4-Tetraiodobenzene	581.74	136			
1145	$C_6H_2I_4$	1, 2, 3, 5-Tetraiodobenzene	581.74	148			
1146	$C_6H_2I_4$	1, 2, 4, 5-Tetraiodobenzene	581.74	254			
1147	$C_6H_2N_4O_9$	2, 3, 4, 6-Tetranitrophenol	274.05	140	d.		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1148	C₆H₂O₄	Diacetylenedicarboxylic acid	138.02	178 exp.			
1149	C₆H₃BrN₂O₄	3-Bromo-1, 2-dinitrobenzene	246.96	101.5	320		
1150	C₆H₃BrN₂O₄	4-Bromo-1, 2-dinitrobenzene	246.96	59.4			1302
1151	C₆H₃BrN₂O₄	4-Bromo-1, 3-dinitrobenzene	246.96	75.3			
1152	C₆H₃Br₂NO₂	2, 4-Dibromonitrobenzene	280.86	62			
1153	C₆H₃Br₂NO₂	2, 5-Dibromonitrobenzene	280.86	85		2.356	
1154	C₆H₃Br₂NO₂	3, 4-Dibromonitrobenzene	280.86	58	296	2.368	
1155	C₆H₃Br₂NO₂	3, 5-Dibromonitrobenzene	280.86	106		2.354	
1155.1	C₆H₃Br₂NO₃	4, 6-Dibromo-2-nitrophenol	296.86	117.5			
1156	C₆H₃Br₃	1, 2, 3-Tribromobenzene	314.77	87.4		2.434	
1157	C₆H₃Br₃	1, 2, 4-Tribromobenzene	314.77	44	276	2.658	
1158	C₆H₃Br₃	1, 3, 5-Tribromobenzene	314.77	119.6	278		
1159	C₆H₃Br₃O	2, 3, 5-Tribromophenol Br₃C₆H₂OH	330.77	92.5			
1160	C₆H₃Br₃O	2, 4, 6-Tribromophenol Br₃C₆H₂OH	330.77	96		2.55	
1161	C₆H₃Br₃O₂	2, 4, 6-Tribromoresorcinol	346.77	111			
1162	C₆H₃ClN₂O₄	3-Chloro-1, 2-dinitrobenzene	202.50	86.8			
1163	C₆H₃ClN₂O₄	4-Chloro-1, 2-dinitrobenzene	202.50	α 36.3 β 37.1 γ 38.8 δ 28	315 d		
1164	C₆H₃ClN₂O₄	2-Chloro-1, 3-dinitrobenzene	202.50	87			
1165	C₆H₃ClN₂O₄	α-4-Chloro-1, 3-dinitrobenzene	202.50	53.4	315	1.697	
1166	C₆H₃ClN₂O₄	β-4-Chloro-1, 3-dinitrobenzene	202.50	43	315	1.680	
1167	C₆H₃ClN₂O₄	5-Chloro-1, 3-dinitrobenzene	202.50	59			
1168	C₆H₃ClN₂O₄	2-Chloro-1, 4-dinitrobenzene	202.50	60			
1169	C₆H₃Cl₂NO₂	2, 3-Dichloronitrobenzene	191.95	62	258	1.721¹⁴	
1170	C₆H₃Cl₂NO₂	2, 4-Dichloronitrobenzene	191.95	33		1.439⁹⁰	
1171	C₆H₃Cl₂NO₂	2, 5-Dichloronitrobenzene	191.95	54.5	266	1.069²²	
1172	C₆H₃Cl₂NO₂	2, 6-Dichloronitrobenzene	191.95	72.5	130⁸	1.603¹⁷	
1173	C₆H₃Cl₂NO₂	3, 4-Dichloronitrobenzene	191.95	43	256	1.451⁸⁰	
1174	C₆H₃Cl₂NO₂	3, 5-Dichloronitrobenzene	191.95	65.4		1.692¹⁴	
1174.1	C₆H₃Cl₂NO₃	4, 6-Dichloro-2-nitrophenol	207.95	122		1.822	
1175	C₆H₃Cl₃	1, 2, 3-Trichlorobenzene	181.40	52	219		
1176	C₆H₃Cl₃	1, 2, 4-Trichlorobenzene	181.40	17	213	1.574¹⁰	754
1177	C₆H₃Cl₃	1, 3, 5-Trichlorobenzene	181.40	63	208.5		
1178	C₆H₃Cl₃O	2, 3, 5-Trichlorophenol	197.40	53.4	253		
1179	C₆H₃Cl₃O	2, 4, 6-Trichlorophenol	197.40	68	244.5		
1180	C₆H₃Cl₃O₂	2, 3, 5-Trichlorohydroquinone	213.40	134			
1181	C₆H₃Cl₃O₂	2, 4, 6-Trichlororesorcinol	213.40	83			
1182	C₆H₃Cl₃O₆S₃	Benzene-1, 3, 5-trisulfonyl chloride	373.59	184			
1183	C₆H₃Cl₄N	2, 3, 4, 5-Tetrachloroaniline	230.86	118			
1184	C₆H₃Cl₄N	2, 3, 4, 6-Tetrachloroaniline	230.86	88			
1185	C₆H₃Cl₄N	2, 3, 5, 6-Tetrachloroaniline	230.86	90			
1186	C₆H₃I₃	1, 2, 3-Triiodobenzene	455.82	116			
1187	C₆H₃I₃	1, 2, 4-Triiodobenzene	455.82	84			
1188	C₆H₃I₃	1, 3, 5-Triiodobenzene	455.82	181			
1189	C₆H₃I₃O	2, 4, 6-Triiodophenol I₃C₆H₂(OH)	471.82	156			
1190	C₆H₃N₃O₆	1, 2, 3-Trinitrobenzene	213.05	127.5			
1191	C₆H₃N₃O₆	1, 2, 4-Trinitrobenzene	213.05	61		1.73¹⁵·⁵	
1192	C₆H₃N₃O₆	1, 3, 5-Trinitrobenzene	213.05	121; 61	d.	1.688	
1193	C₆H₃N₃O₆S	Thiopicric acid	245.11	114	exp. 115		
1194	C₆H₃N₃O₇	2, 3, 5-Trinitrophenol C₆H₂(NO₂)₃OH	229.05	120			
1195	C₆H₃N₃O₇	2, 3, 6-Trinitrophenol C₆H₂(NO₂)₃OH	229.05	118			
1196	C₆H₃N₃O₇	2, 4, 5-Trinitrophenol C₆H₂(NO₂)₃OH	229.05	96			
1197	C₆H₃N₃O₇	Picric acid (NO₂)₃C₆H₂OH	229.05	121.8	exp. > 300	1.763	1313
1198	C₆H₃N₃O₈	Styphnic acid	245.05	180		1.829	
1199	C₆H₃N₃O₉S	Picrylsulfonic acid	293.11	100			
1200	C₆H₃N₄O₈	2, 3, 4, 6-Tetranitroaniline	273.06	170	exp 237	1.89	1314
1200.1	C₆H₄BrCl	o-Bromochlorobenzene	191.40	−12.6	204⁷⁵⁵	1.656¹²·⁵	765
1200.2	C₆H₄BrCl	m-Bromochlorobenzene	191.40	−21.2	196	1.627¹⁴	764
1200.3	C₆H₄BrCl	p-Bromochlorobenzene	191.40	67.4	196.3		
1200.4	C₆H₄BrI	o-Bromoiodobenzene	282.88	2.1	257.4⁷⁴⁴		
1200.5	C₆H₄BrI	m-Bromoiodobenzene	282.88	−9.3	252⁷⁴⁴		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1200 6	C_6H_4BrI	p-Bromoiodobenzene	282 88	92	251.6[7.64]		
1201	$C_6H_4BrNO_2$	o-Bromonitrobenzene	201 96	43.0	261	1.623[40]	
1202	$C_6H_4BrNO_2$	m-Bromonitrobenzene	201 96	56 0	256 5	1.704	777
1203	$C_6H_4BrNO_2$	p-Bromonitrobenzene	201 96	127	256		
1204	$C_6H_4Br_2$	o-Dibromobenzene	235 86	1 8	221	1 966[15]	787
1205	$C_6H_4Br_2$	m-Dibromobenzene	235 86	−6 9	217	1.955	783
1206	$C_6H_4Br_2$	p-Dibromobenzene	235 86	86 8	219	1.954	1132
1207	$C_6H_4Br_2O$	2, 4-Dibromophenol	251 86	36	239		
1208	$C_6H_4Br_2O$	2, 6-Dibromophenol	251 86	56			
1209	$C_6H_4Br_2O$	3, 4-Dibromophenol	251 86	80			
1210	$C_6H_4Br_2O$	3, 5-Dibromophenol	251 83	76 5			
1211	$C_6H_4Br_2O_2$	2, 4-Dibromoresorcinol	267 86	92 5			
1212	$C_6H_4Br_2O_2$	4, 6-Dibromoresorcinol	267 86	112	130 (in CO_2)		
1213	$C_6H_4Br_3N$	2, 4, 6-Tribromoaniline	329 79	119	300		
1214	$C_6H_4Br_3N$	3, 4, 5-Tribromoaniline	329 79	118			
1214 1	C_6H_4ClI	p-Chloroiodobenzene	238 42	57	227 6[7.61]		
1215	$C_6H_4ClNO_2$	o-Chloronitrobenzene	157 50	32 5	245 7	1.365	
1216	$C_6H_4ClNO_2$	m-Chloronitrobenzene	157 50	44 4; 23 7	235 6	1 534	
1217	$C_6H_4ClNO_2$	p-Chloronitrobenzene	157 50	83 5	242	1.520	
1218	$C_6H_4ClNO_3$	4-Chloro-2-nitrophenol	173 50	87			
1219	$C_6H_4ClNO_3$	5-Chloro-2-nitrophenol	173 50	38 9			
1220	$C_6H_4ClNO_3$	6-Chloro-2-nitrophenol	173 50	70			
1221	$C_6H_4ClNO_3$	2-Chloro-3-nitrophenol	173 50	120			
1222	$C_6H_4ClNO_4$	4-Chloro-3-nitrophenol	173 50	127			
1223	$C_6H_4ClNO_3$	5-Chloro-3-nitrophenol	173 50	147			
1224	$C_6H_4ClNO_3$	6-Chloro-3-nitrophenol	173 50	118			
1225	$C_6H_4ClNO_3$	2-Chloro-4-nitrophenol	173 50	111			
1226	$C_6H_4ClNO_3$	3-Chloro-4-nitrophenol	173 50	133			
1227	$C_6H_4ClNO_5S$	2-Chloronitrobenzene-5-sulfonic acid	237 56	>200 d.			
1228	$C_6H_4ClNO_5S$	5-Chloronitrobenzene-3-sulfonic acid	237 56	200 d.			
1229	$C_6H_4Cl_2$	o-Dichlorobenzene	146 95	−17 6	179	1.298	731
1230	$C_6H_4Cl_2$	m-Dichlorobenzene	146 95	−24 8	173	1.288	723
1231	$C_6H_4Cl_2$	p-Dichlorobenzene	146 95	52 9	173	1.458	1101
1232	$C_6H_4Cl_2O$	2, 3-Dichlorophenol	162 95	57			
1233	$C_6H_4Cl_2O$	2, 4-Dichlorophenol	162 95	45	210		
1234	$C_6H_4Cl_2O$	2, 5-Dichlorophenol	162 95	58	211 7		
1235	$C_6H_4Cl_2O$	2, 6-Dichlorophenol	162 95	67	220		
1236	$C_6H_4Cl_2O$	3, 4-Dichlorophenol	162 95	68	253 5		
1237	$C_6H_4Cl_2O$	3, 5-Dichlorophenol	162 95	68	233 1		
1238	$C_6H_4Cl_2O_2$	2, 3-Dichlorohydroquinone	178 95	145			
1239	$C_6H_4Cl_2O_2$	2, 5-Dichlorohydroquinone	178 95	170		1.824	
1240	$C_6H_4Cl_2O_2$	2, 6-Dichlorohydroquinone	178 95	164			
1241	$C_6H_4Cl_2O_3S$	2, 5-Dichlorobenzenesulfonic acid	227 01	97			
1242	$C_6H_4Cl_2O_4S_2$	o-Benzenedisulfonyl chloride	275 08	105			
1243	$C_6H_4Cl_2O_4S_2$	m-Benzenedisulfonyl chloride	275 08	63			
1244	$C_6H_4Cl_2O_4S_2$	p-Benzenedisulfonyl chloride	275 08	131			
1245	$C_6H_4Cl_3N$	2, 3, 4-Trichloroaniline	196 41	67.5	291.5		
1246	$C_6H_4Cl_3N$	2, 4, 5-Trichloroaniline	196.41	96	270		
1247	$C_6H_4Cl_3N$	2, 4, 6-Trichloroaniline	196 41	77 5	262 4		
1248	$C_6H_4Cl_3N$	3, 4, 5-Trichloroaniline $Cl_3C_6H_2NH_2$.	196 41	100			
1249	$C_6H_4FNO_2$	o-Fluoronitrobenzene	141.04	−5.9	214.6	1.338	700
1250	$C_6H_4FNO_2$	m-Fluoronitrobenzene	141.04	1 7	205	1.327	688
1251	$C_6H_4FNO_2$	p-Fluoronitrobenzene .	141.04	26 5; 21 5	205	1 326	1084
1252	$C_6H_4F_2$	m-Difluorobenzene	114.03		83	1.172	384
1253	$C_6H_4F_2$	p-Difluorobenzene	114.03	−23.7	88.9	1 164	362
1254	$C_6H_4INO_2$	o-Iodonitrobenzene	248.97	49 4	290	1 810[15.5]	
1255	$C_6H_4INO_2$	m-Iodonitrobenzene	248.97	36	280	1 804[15.5]	
1256	$C_6H_4INO_2$	p-Iodonitrobenzene	248.97	171.5	288.1	1 809[15.5]	
1257	$C_6H_4INO_3$	4-Iodo-6-nitrophenol $IC_6H_3(NO_2)OH$.	264.97	81			
1258	$C_6H_4I_2$	o-Diiodobenzene	329.90	23.4	286.8		
1259	$C_6H_4I_2$	m-Diiodobenzene	329.90	34.2	284.8		
1260	$C_6H_4I_2$	p-Diiodobenzene....	329.90	129.4	285		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1261	C₆H₄I₂O	2, 4-Diiodophenol..	345 90	72	100		
1262	C₆H₄I₂O	2, 6-Diiodophenol I₂C₆H₃OH	345 90	68			
1263	C₆H₄I₂O	3, 4-Diiodophenol I₂C₆H₃OH	345 90	83			
1264	C₆H₄I₂O	3, 5-Diiodophenol I₂C₆H₃OH	345 90	104			
1265	C₆H₄I₂O₄S	2, 6-Diiodophenol-4-sulfonic acid	425 96	120	190 d.		
1266	C₆H₄I₃N	2, 4, 6-Triiodoaniline I₃C₆H₂NH₂	470 84	185 5			
1267	C₆H₄N₂	Pyridyl-2-cyanide CN.C₅H₄N	104 05	29			
1268	C₆H₄N₂	Pyridyl-3-cyanide CN.C₅H₄N	104 05	50			
1269	C₆H₄N₂	Pyridyl-4-cyanide CN.C₅H₄N	104 05	79			
1270	C₆H₄N₂O	p-Diazophenol . .	120 05	exp. 38			
1271	C₆H₄N₂O₄	o-Dinitrobenzene	168 05	116 5	319	1 59	
1272	C₆H₄N₂O₄	m-Dinitrobenzene	168 05	89 7	302	1 575	
1273	C₆H₄N₂O₄	p-Dinitrobenzene... .	168 05	172 1	299	1 625	
1274	C₆H₄N₂O₅	2, 3-Dinitrophenol (NO₂)₂C₆H₃OH	184 05	144			
1275	C₆H₄N₂O₅	2, 4-Dinitrophenol	184 05	111 6		1 683	
1276	C₆H₄N₂O₅	2, 5-Dinitrophenol (NO₂)₂C₆H₃OH	184 05	104			
1277	C₆H₄N₂O₅	2, 6-Dinitrophenol (NO₂)₂C₆H₃OH	184 05	61 8			
1278	C₆H₄N₂O₅	3, 4-Dinitrophenol (NO₂)₂C₆H₃OH	184 05	134			
1279	C₆H₄N₂O₅	3, 5-Dinitrophenol . .	184 05	126 1			
1280	C₆H₄N₂O₆	2, 4-Dinitroresorcinol ..	200 05	148	d.		
1281	C₆H₄N₂O₆	4, 6-Dinitroresorcinol	200 05	215			
1282	C₆H₄N₂O₇S	2, 4-Dinitrobenzenesulfonic acid	248 11	108			
1283	C₆H₄N₂S	Benzisothiodiazole	136 11	44	206		
1284	C₆H₄N₄O₆	Picramide 2, 4, 6-(NO₂)₃C₆H₂NH₂	228 06	188			
1285	C₆H₄N₄O₇	2, 4, 6-Trinitroaminophenol	244 06	178			
1286	C₆H₄N₆	Hexaazobenzene	160 08	83			
1287	C₆H₄O₂	Quinone	108 03	115 7		1 318	
1288	C₆H₄O₄	2, 5-Dihydroxyquinone	140 03	220			
1289	C₆H₄O₆	Sarsapic acid. . .	172 03	305			
1290	C₆H₄O₈	Ethanetetracarboxylic acid	204 03	169 d.			
1291	C₆H₅AsCl₂	Phenyl dichloroarsine	222 92		253		
1292	C₆H₅AsO	Phenylarsine oxide	168 00	120			
1294	C₆H₅Br	Bromobenzene	156 96	−30 6	156 2	1 497	747
1295	C₆H₅BrN₂O₂	4-Bromo-2-nitroaniline	216 97	111			
1296	C₆H₅BrO	o-Bromophenol .	172 96	5.6	195	1.553⁸⁰	
1297	C₆H₅BrO	m-Bromophenol .	172 96	33	236 5		
1298	C₆H₅BrO	p-Bromophenol . ..	172 96	63.5	238	1 588ⁿ⁰	
1299	C₆H₅BrO₂	Bromohydroquinone ..	188 96	115			
1300	C₆H₅BrO₂	2(4)-Bromoresorcinol	188 96	91			
1301	C₆H₅BrO₃S	p-Bromobenzenesulfonic acid	237 02	88			
1302	C₆H₅Br₂N	2, 4-Dibromoaniline .	250 88	79.5			
1303	C₆H₅Br₂N	2, 5-Dibromoaniline	250 88	52			
1304	C₆H₅Br₂N	2, 6-Dibromoaniline	250 88	84	264		
1305	C₆H₅Br₂N	3, 4-Dibromoaniline	250 88	80 4			
1306	C₆H₅Br₂N	3, 5-Dibromoaniline	250 88	56 5			
1307	C₆H₅Cl	Chlorobenzene ..	112 50	−45 2	132 1	1.107	681
1308	C₆H₅ClN₂O₂	2-Chloro-4-nitroaniline	172 51	105			
1309	C₆H₅ClN₂O₂	2-Chloro-5-nitroaniline	172 51	118			
1310	C₆H₅ClN₂O₂	3-Chloro-4-nitroaniline	172 51	157			
1311	C₆H₅ClN₂O₂	3-Chloro-6-nitroaniline	172 51	125			
1312	C₆H₅ClN₂O₂	4-Chloro-2-nitroaniline	172 51	115			
1313	C₆H₅ClN₂O₂	4-Chloro-3-nitroaniline	172 51	103			
1314	C₆H₅ClO	o-Chlorophenol. ...	128.50	α 7; β 0; γ −4 1	173	1 241¹⁸·²	1058
1315	C₆H₅ClO	m-Chlorophenol .	128.50	32.8	214		1059
1316	C₆H₅ClO	p-Chlorophenol.... .	128 50	37	217	1.306	1060
1317	C₆H₅ClO₂	Chlorohydroquinone	144 50	106	263		
1318	C₆H₅ClO₂S	Benzenesulfone chloride	176 56	14.5	247	1.383¹⁵	
1319	C₆H₅ClO₃S	p-Chlorobenzenesulfonic acid	192 56	67	146²⁵		
1320	C₆H₅Cl₂N	2, 3-Dichloroaniline. .	161 96	24	252		
1321	C₆H₅Cl₂N	2, 4-Dichloroaniline	161 96	63	245	1.567	
1322	C₆H₅Cl₂N	2, 5-Dichloroaniline	161.96	50	251		
1323	C₆H₅Cl₂N	2, 6-Dichloroaniline Cl₂C₆H₃NH₂.	161.96	39			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1324	C₆H₄Cl₂N	3, 4-Dichloroaniline	161.96	71 5	272		
1325	C₆H₄Cl₂N	3, 5-Dichloroaniline	161 96	50 5	260		
1326	C₆H₄Cl₂OP	Phosphenyl oxychloride	194.98		258	1.375	
1327	C₆H₄Cl₂P	Phosphenyl chloride	178 98		224 6	1 319	804
1328	C₆H₅F	Fluorobenzene	96 039	−41 2	86	1 024	487
1329	C₆H₅FO	o-Fluorophenol FC₆H₄OH	112 04	16 1			
1330	C₆H₅FO	m-Fluorophenol	112 04	13 8	183⁴⁹	1 222	652
1331	C₆H₅FO	p-Fluorophenol	112 04	28 5; 48 2	188	1 189⁵⁶	1083
1332	C₆H₅F₂N	2, 5-Difluoroaniline	129 05	13 5	85 8³⁰	1 288¹⁷ ²	
1333	C₆H₅I	Iodobenzene	203 97	−31 4	188 6	1 832	792
1334	C₆H₅IO	o-Iodophenol	219 97	40 4	187¹⁶⁰	1 876⁸⁰	
1335	C₆H₅IO	m-Iodophenol IC₆H₄OH	219 97	40			
1336	C₆H₅IO	p-Iodophenol IC₆H₄OH	219 97	94			
1337	C₆H₅IO	Iodosobenzene	219 97	exp. 210			
1338	C₆H₅IO₂	Iodoxybenzene	235 97	exp. 238			
1339	C₆H₅IO₂S	Benzenesulfone iodide C₆H₅SO₂I	268 04	45			
1340	C₆H₅I₂N	2, 4-Diiodoaniline I₂C₆H₃NH₂	344 91	96			
1341	C₆H₅NO	Pyridyl-α-aldehyde	107 05		181	1 126	947
1342	C₆H₅NO	Pyridyl-β-aldehyde	107 05		97¹⁵		
1343	C₆H₅NO	Nitrosobenzene	107 05	68	59¹⁸		
1344	C₆H₅NO₂	Picolinic acid	123 05	137			
1345	C₆H₅NO₂	Nicotinic acid	123 05	232			
1346	C₆H₅NO₂	Isonicotinic acid	123 05	317			
1347	C₆H₅NO₂	Nitrobenzene	123 05	5 7	210 9	1.207	736
1348	C₆H₅NO₂	p-Nitrosophenol ONC₆H₄OH	123 05	126			
1349	C₆H₅NO₃	o-Nitrophenol	139 05	45	214 5	1 447	
1350	C₆H₅NO₃	m-Nitrophenol	139 05	96	194⁷⁰	1 485	
1351	C₆H₅NO₃	p-Nitrophenol	139 05	113		1 468	
1352	C₆H₅NO₄	2-Nitroresorcinol m-(OH)₂C₆H₃NO₂	155 05	85			
1353	C₆H₅NO₄	4-Nitroresorcinol m-(OH)₂C₆H₃NO₂	155 05	115			
1354	C₆H₅NO₄	Nitrohydroquinone	155 05	134			
1355	C₆H₅NO₆S	2-Nitrophenol-4-sulfonic acid	219 11	141			
1356	C₆H₆N₂	Aziminobenzene	119.06	99			
1357	C₆H₅N₃	Triazobenzene	119 06		73.5²⁴	1 098¹⁰	991
1358	C₆H₅N₃O₄	2, 3-Dinitroaniline (NO₂)₂C₆H₃NH₂	183 06	127			
1359	C₆H₅N₃O₄	2, 4-Dinitroaniline	183 06	188			
1360	C₆H₅N₃O₄	2, 5-Dinitroaniline (NO₂)₂C₆H₃NH₂	183 06	137			
1361	C₆H₅N₃O₄	2, 6-Dinitroaniline	183 06	138			
1362	C₆H₅N₃O₄	3, 4-Dinitroaniline (NO₂)₂C₆H₃NH₂	183 06	154			
1363	C₆H₅N₃O₄	3, 5-Dinitroaniline (NO₂)₂C₆H₃NH₂	183 06	159			
1364	C₆H₅N₃O₆	Picramic acid	199 06	165			1320
1365	C₆H₆	Benzene	78 046	5.5	79 6	0 878	606
1366	C₆H₆	Dipropargyl	78 046	−6	85.4	0 805	380
1367	C₆H₆AsCl₃	Tri- (2-chlorovinyl)arsine	259 38		260	1 572	
1368	C₆H₆BrN	o-Bromoaniline	171 97	31.5	251		
1369	C₆H₆BrN	m-Bromoaniline	171 97	18.5	251	1.587¹⁵ ³	793
1370	C₆H₆BrN	p-Bromoaniline BrC₆H₄NH₂	171 97	66 4			
1371	C₆H₆Br₂N₂	3, 4-Dibromophenylhydrazine	265 89	75			
1372	C₆H₆Br₂N₂	3, 5-Dibromophenylhydrazine	265 89	95 5			
1373	C₆H₆Br₆	α-trans-Benzenehexabromide	557.54	212			
1374	C₆H₆Br₆	β-cis-Benzenehexabromide	557.54	253			
1375	C₆H₆ClN	o-Chloroaniline ClC₆H₄NH₂	127.51	0	210.5	1 213	774
1376	C₆H₆ClN	m-Chloroaniline	127.51	−10.4	229.8	1.215	776
1377	C₆H₆ClN	p-Chloroaniline	127.51	71	231	1.170⁷⁰	
1378	C₆H₆ClNO	2-Chloro-3-aminophenol	143.51	87			
1379	C₆H₆ClNO	2-Chloro-4-aminophenol	143.51	153			
1380	C₆H₆ClNO₃S	p-Chlorometanilic acid	207.58	280 d.			
1381	C₆H₆Cl₂N₂	2, 4-Dichlorophenylhydrazine	176.98	94			
1382	C₆H₆Cl₂N₂	2, 5-Dichlorophenylhydrazine	176.98	105			
1383	C₆H₆Cl₂N₂	3, 5-Dichlorophenylhydrazine	176.98	118			
1384	C₆H₆Cl₆	α-trans-Benzenehexachloride	290 79	157	288	1.87	
1385	C₆H₆Cl₆	β-cis-Benzenehexachloride	290.79	310		1.89¹⁹	

No.	Formula	Name	Mol wt	M. P	B. P.	d	R. I. No.
1386	C₆H₆Cl₆	γ-Benzenehexachloride	290 79	112			
1387	C₆H₆Cl₆	δ-Benzenehexachloride	290 79	129			
1388	C₆H₆FN	o-Fluoroaniline	111 05	−34 6	68 5¹⁴	1 151	716
1389	C₆H₆FN	m-Fluoroaniline	111 05		186 3	1 160	722
1390	C₆H₆FN	p-Fluoroaniline	111 05	−1 9	189	1 152	707
1391	C₆H₆IN	o-Iodoaniline	218 99	56 5			
1392	C₆H₆IN	m-Iodoaniline	218 99	27			
1393	C₆H₆IN	p-Iodoaniline	218 99	62			
1394	C₆H₆N₂O	p-Nitrosoaniline	122 06	174			
1395	C₆H₆N₂O₂	Phenylnitroamine	138 06	46			
1396	C₆H₆N₂O₂	o-Nitroaniline .	138 06	71 5			
1397	C₆H₆N₂O₂	m-Nitroaniline O₂NC₆H₄NH₂	138 06	111 8	286	1 430	
1398	C₆H₆N₂O₂	p-Nitroaniline	138 06	148		1 424	
1399	C₆H₆N₂O₂	Quinonedioxime p-C₆H₄(NOH)₂	138 06	240			
1400	C₆H₆N₂O₃	3-Nitro-2-aminophenol	154 06	136			
1401	C₆H₆N₂O₃	4-Nitro-2-aminophenol	154 06	143			
1402	C₆H₆N₂O₃	5-Nitro-2-aminophenol	154 06	202			
1403	C₆H₆N₂O₃	6-Nitro-2-aminophenol	154 06	111			
1404	C₆H₆N₂O₃	5-Nitro-3-aminophenol	154 06	165			
1405	C₆H₆N₂O₃	2-Nitro-4-aminophenol	154 06	206			
1406	C₆H₆N₂O₃	3-Nitro-4-aminophenol	154 06	148			
1407	C₆H₆N₂O₄	5-Acetylbarbituric acid	170 08	300			
1408	C₆H₆N₂O₄	Dimethylalloxan	170 06	255 d.			
1409	C₆H₆N₄O₃	1-Methyluric acid	182 08	400 d.			
1410	C₆H₆N₄O₃	3-Methyluric acid	182 08	>360 d.			
1411	C₆H₆N₄O₃	7-Methyluric acid	182 08	370 d.			
1412	C₆H₆N₄O₇	Ammonium picrate	246 08	d.		1.719	1318
1413	C₆H₆O	Phenol	94 046	41	182	1.071¹⁸	1064
1414	C₆H₆O₂	o-Dihydroxybenzene 1, 2-C₆H₄(OH)₂*	110 05	105	245	1.344	1272
1415	C₆H₆O₂	Resorcinol 1, 3-C₆H₄(OH), ...	110 05	110	276 5	1 285¹⁸	1275
1416	C₆H₆O₂	Hydroquinol 1, 4-C₆H₄(OH)₂ .	110 05	170 5	286 2	1 332¹⁶	1184
1417	C₆H₆O₂	5-Methylfurfural .	110 05		187	1 109¹⁶	
1418	C₆H₆O₃S	Benzenesulfinic acid	142 11	84	100 d.		
1419	C₆H₆O₃	Pyrogallol 1, 2, 3-C₆H₃(OH)₃	126 05	134	309	1.453	1333
1420	C₆H₆O₃	Hydroxyhydroquinone. .. .	126 05	140 5			
1421	C₆H₆O₃	Phloroglucinol	126 05	219			
1422	C₆H₆O₃	Acrylic anhydride	126 05		97²⁴	1 094⁰	
1423	C₆H₆O₃S	Benzenesulfonic acid ..	158 11	46	d		
1424	C₆H₆O₄	Apionol 1, 2, 3, 4-C₆H₂(OH)₄.	142 05	161			
1425	C₆H₆O₄	1, 2, 3, 5-Tetrahydroxybenzene	142.05	165			
1426	C₆H₆O₄	1, 2, 4, 5-Tetrahydroxybenzene	142 05	220			
1427	C₆H₆O₄	Muconic acid (CH.CHCO₂H)₂	142 05	320 d.			
1428	C₆H₆O₄S	o-Phenolsulfonic acid.	174 11	50			
1429	C₆H₆O₆	Aconitic acid.	174 05	191			
1430	C₆H₆S	Thiophenol C₆H₅SH .	110 11		169 5	1 074	1002
1431	C₆H₆Se	Selenophenol C₆H₅SeH.	157 25		183.6	1 487¹⁶	
1432	C₆H₆S₂	Dithioresorcinol 1, 3-C₆H₄(SH)₂	142 18	27	243		
1433	C₆H₆S₂	Dithiohydroquinone 1, 4-C₆H₄(SH)₂...	142 18	98			
1434	C₆H₇As	Phenylarsine C₆H₅AsH₂	154 01		148		
1435	C₆H₇AsO₃	Phenylarsonic acid..	202.01	158 d.		1 840	
1436	C₆H₇BrN₂	p-Bromophenylhydrazine	186 99	107			
1437	C₆H₇ClN₂	4-Chloro-o-phenylenediamine	142.53	72			
1438	C₆H₇ClN₂	4-Chloro-m-phenylenediamine	142.53	86			
1439	C₆H₇ClN₂	o-Chlorophenylhydrazine	142 53	47			
1440	C₆H₇ClN₂	p-Chlorophenylhydrazine	142.53	90			
1441	C₆H₇ClO	Sorbic chloride....	130.51		78¹⁶	1.065	741
1441 1	C₆H₇ClO₄	Methyl chloromaleate..	178.51		106.5¹⁸	1.278²¹	
1441.2	C₆H₇ClO₄	Methyl chlorofumarate . . .	178.51		115.5¹⁸	1.290²⁴	
1442	C₆H₇N	Aniline.....	93.062	−6 2	184.4	1.022	769
1443	C₆H₇N	α-Picoline....	93.062	−69.9	128.0	0.950	604
1444	C₆H₇N	β-Picoline....	93.062		143.5	0.952	1018
1445	C₆H₇N	γ-Picoline	93.062		143 1	0.957	
1446	C₆H₇NO	o-Aminophenol	109.06	170			

* Commonly known as catechol, pyrocatechol, catechin, pyrocatechin.

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1447	C₆H₇NO	m-Aminophenol . . .	109 06	123			
1448	C₆H₇NO	p-Aminophenol	109 06	184			1333
1449	C₆H₇NO	Methyl 2-pyrryl ketone	109 06	90	220		
1450	C₆H₇NO	β-Phenylhydroxylamine	109 06	82			
1451	C₆H₇NO₂	Phloramine 3, 5-(OH)₂C₆H₃NH₂	125 06	152			
1452	C₆H₇NO₂S	Benzenesulfoneamide ...	157 13	156			
1455	C₆H₇NO₃S	p-Anilinesulfonic acid .	173 13	288			
1458	C₆H₇NS	2-Aminothiophenol	125 13	26	234		
1459	C₆H₇N₃O₂	4-Nitro-o-phenylenediamine	153 08	198			
1460	C₆H₇N₃O₂	4-Nitro-m-phenylenediamine	153 08	161			
1461	C₆H₇N₃O₂	2-Nitro-p-phenylenediamine	153 08	135			
1462	C₆H₇N₅O₁₄	d-Glucose pentanitrate	405 09	135 d.			
1463	C₆H₇O₂P	Phenylphosphenous acid	142 08	70			
1464	C₆H₇O₃P	Phenylphosphenic acid	158 08	158	250 d.	1.475	
1465	C₆H₇P	Phenyl phosphine C₆H₅PH₂	110 08		160	1.001¹⁵	
1466	C₆H₈	1, 3-Cyclohexadiene . . .	80 062	−98	80.5	0.842	519
1467	C₆H₈	Diallylene (CH₂C:CH)₂.	80 062		70	0.858¹⁸ ²	
1468	C₆H₈	o-Dihydrobenzene	80 062		78 5	0.848	
1469	C₆H₈	m-Dihydrobenzene .	80 062		80 5	0.830	
1470	C₆H₈	p-Dihydrobenzene	80 062		85 5	0 848	
1471	C₆H₈AsNO₃	Arsanilic acid p-NH₂C₆H₄AsO(OH)₂.	217 03	<200			
1471.1	C₆H₈BrN	Aniline hydrobromide	173 99	286			
1472	C₆H₈ClN	Aniline hydrochloride	129 53	198	245	1 222⁴	1245
1474	C₆H₈ClNO	m-Aminophenol hydrochloride	145 53	229			
1475	C₆H₈ClNO	p-Aminophenol hydrochloride	145 53	306 d.			1333
1476	C₆H₈Cl₂O₂	Adipyl dichloride . .	182 98		132¹⁸ s. d.		
1477	C₆H₈N	Piturine	94 070		244		
1478	C₆H₈N₂	Adipyldinitrile	108 08	1	295	0 951¹⁹	471
1479	C₆H₈N₂	o-Phenylenediamine... ...	108 08	103 8	252		
1480	C₆H₈N₂	m-Phenylenediamine	108 08	62 8	287	1 107⁵⁷ ⁷	1086
1481	C₆H₈N₂	p-Phenylenediamine	108 08	139 7	267		
1482	C₆H₈N₂	2, 5-Dimethylpyrazine (Ketine) .	108 08	15	155	0.990	1017
1483	C₆H₈N₂	Phenylhydrazine C₆H₅NHNH₂	108 08	19.6	243.5	1 098	784
1484	C₆H₈N₂O	2, 5-Diaminophenol . .	124 08	68			
1485	C₆H₈N₂O	3, 4-Diaminophenol .	124 08	168			
1486	C₆H₈N₂O	3, 5-Diaminophenol	124 08	170			
1487	C₆H₈N₂O₂	1, 3-Dimethylbarbituric acid	156 08	123			
1488	C₆H₈N₂O₂	1-Ethylbarbituric acid . .	156.08	120			
1489	C₆H₈N₂O₃	Aniline nitrate . .	156.08		190 d.	1 358⁴	
1490	C₆H₈N₂O₃S	o-Phenylenediamine-3-sulfonic acid	188 14	d.			
1491	C₆H₈N₂O₃S	p-Phenylhydrazinesulfonic acid	188 14	286			
1492	C₆H₈N₂O₄S₂	o-Benzenedisulfoneamide	236 21	233			
1493	C₆H₈N₂O₄S₂	m-Benzenedisulfoneamide	236 21	229			
1494	C₆H₈N₂O₄S₂	p-Benzenedisulfoneamide	236 21	188			
1495	C₆H₈N₆O₁₄	Mannitol hexanitrate	452 11	113		1 8	
1496	C₆H₈O	2, 5-Dimethylfuran	96 062		94	0 888	974
1497	C₆H₈O₂	Dihydroresorcinol m-(OH)₂C₆H₆	112 06	104			
1498	C₆H₈O₂	Sorbic acid CH₃(CH:CH)₂CO₂H	112 06	134 5	228 d.		1333
1499	C₆H₈O₄	Dimethyl fumarate .	144 06	102	192		
1500	C₆H₈O₄	Dimethyl maleate . . .	144.06		203	1 153¹⁴	382
1501	C₆H₈O₄	Ethyl fumarate CO₂HCH.CHCO₂C₂H₅.	144.06	70			
1502	C₆H₈O₄	Lactide	144.06	125	255	0 862	
1503	C₆H₈O₅	Acetonylmalonic acid . .	160.06	150			
1504	C₆H₈O₅	Acetylmalic acid........	160.06	134			
1504 1	C₆H₈O₅	1-Ketoadipic acid........ .	160.06	124			
1505	C₆H₈O₆	Tricarballylic acid...	176.06	166	d.		
1506	C₆H₈O₆	Glycerol triformate (Triformin).......	176.06	18	266	1 320	373
1507	C₆H₈O₇	Citric acid (HO₂CCH₂)₂C(OH)CO₂H	192.06	153		1 542	1202
1508	C₆H₈O₈	Hydroxycitric acid..............	208.06	160			
1509	C₆H₈S	2, 3-Dimethylthiophene....	112.13		137	0.994	
1510	C₆H₈S	2, 4-Dimethylthiophene . .	112.13		138	0.996	
1511	C₆H₈S	2, 5-Dimethylthiophene . .	112.13		137.5	0 976¹⁷ ⁵	
1512	C₆H₈S	3, 4-Dimethylthiophene..	112.13		146	1 008²⁰₁₁.₆	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1513	$C_6H_9AsO_6$	Arsenic acetate	252 03	82	170[11]		
1514	$C_6H_9ClN_2$	Phenylhydrazine hydrochloride	144 54	243			
1515	$C_6H_9ClO_3$	Ethyl chloroacetoacetate	164 53		200	1 179[22]	
1516	C_6H_9N	1, 2-Dimethylpyrrol	95 077		65[14]		
1517	C_6H_9N	2, 3-Dimethylpyrrol	95 077		165		
1518	C_6H_9N	2, 4-Dimethylpyrrol	95 077		171	0.927[14]	829
1519	C_6H_9N	2, 5-Dimethylpyrrol	95 077		169	0 935	909
1520	C_6H_9N	1-Ethylpyrrol	95 077		131	0 888[16]	
1521	$C_6H_9NO_2$	Guavacine	127 08	285 d.			
1522	$C_6H_9NO_3$	Triacetamide $(CH_3CO)_3N$	143 08	79			
1523	$C_6H_9NO_3S$	Ammonium benzenesulfonate	175 14	256			
1524	$C_6H_9NO_4S$	m-Aminophenol sulfate	207 14	152			
1525	$C_6H_9N_3$	1, 2, 3-Triaminobenzene	123 09	103	336		
1526	$C_6H_9N_3$	1, 2, 4-Triaminobenzene	123 09	100	340		
1527	$C_6H_9N_3O$	2, 4, 6-Triaminophenol	139 09		257		
1528	$C_6H_9N_3O_2$	Cupferron	155 09	164			
1529	$C_6H_9N_3O_2$	Histidine	155 09	253 d.			
1530	$C_6H_9N_3O_3$	Phloroglucinol trioxime	171 09	155 exp.			
1531	$C_6H_9N_3O_4$	Caffuric acid	187 09	220			
1532	C_6H_{10}	n-Butylacetylene $C_4H_9C:CH$	82 077	−150	71 5		
1533	C_6H_{10}	Diisopropenyl $(CH_3C:CH_2)_2$	82 077		69 6	0 731[16]	852
1534	C_6H_{10}	1, 5-Hexadiene $(CH_2CH:CH_2)_2$	82 077		60	0 688	127
1535	C_6H_{10}	2, 4-Hexadiene $(CH:CHCH_3)_2$	82 077		82	0 718	810
1536	C_6H_{10}	Methylpropylacetylene $CH_3CC:C_3H_7$	82 077		81	0 749[0]	
1537	C_6H_{10}	1, 2, 3, 4-Tetrahydrobenzene	82 077	−103 7	83	0 810	404
1539	$C_6H_{10}ClN_3O_2$	Histidine hydrochloride	191 56	251 d.			
1540	$C_6H_{10}N_4O_{11}$	Tetranitrodiglycerol	346 11		250[8]	1 33	
1541	$C_6H_{10}O$	Cyclohexanone	98 077		156 7	0.949	874
1542	$C_6H_{10}O$	1, 2, 3, 4-Tetrahydrophenol	98 077		166 d.		
1543	$C_6H_{10}O$	1, 2, 3, 6-Tetrahydrophenol	98 077		166		
1544	$C_6H_{10}O$	Allyl ether $(CH_2:CHCH_2)_2O$	98 077		94 3	0 805	
1545	$C_6H_{10}O$	1-Ethyl-2-methylacrolein	98 077		137 3	0 858	
1546	$C_6H_{10}O$	Allylacetone $CH_2:CH(CH_2)_2COCH_3$	98 077		129 5	0 846	876
1547	$C_6H_{10}O$	Diethylketene $(C_2H_5)_2C:CO$	98 077		89 5	0 831	
1548	$C_6H_{10}O$	Mesityl oxide $(CH_3)_2C:CHCOCH_3$	98 077	−59.0	135	0 863	899
1549	$C_6H_{10}O_2$	Adipyl dialdehyde $OCH(CH_2)_4CHO$	114 08		94[9]		
1550	$C_6H_{10}O_2$	Propionylpropionic aldehyde	114 08	40	166		
1551	$C_6H_{10}O_2$	Acetonylacetone $(CH_3COCH_2)_2$	114 08	−9	194	0 970	428
1552	$C_6H_{10}O_2$	α-Ethylcrotonic acid	114 08	45	209		
1553	$C_6H_{10}O_2$	1, 2-Hexenic acid $C_3H_7CH:CHCO_2H$	114 08	32	217	0 965	1055
1554	$C_6H_{10}O_2$	2, 3-Hexenic acid	114 08		208	0 962	953
1555	$C_6H_{10}O_2$	1, 2-Isohexenic acid	114 08		108[17]	0 959	885
1556	$C_6H_{10}O_2$	Crotonyl acetate	114 08		129	0 934[0]	
1557	$C_6H_{10}O_2$	Ethyl α-crotonate	114 08		139	0 919	283
1558	$C_6H_{10}O_2$	Ethyl isocrotonate	114 08		131 2	0 925	
1559	$C_6H_{10}O_3$	Glyceryl ether	130 08		173	1 091	
1560	$C_6H_{10}O_3$	Propionic anhydride $(CH_3CH_2CO)_2O$	130 08	−45 0	196 0	1 012	142
1561	$C_6H_{10}O_3$	Ethyl acetoacetate	130 08	< −80	180	1 025	243
1562	$C_6H_{10}O_4$	Adipic acid $HO_2C(CH_2)_4CO_2H$	146 08	151	265[100]		
1563	$C_6H_{10}O_4$	1, 1-Dimethylsuccinic acid	146 08	142	165 d.		
1564	$C_6H_{10}O_4$	Ethylsuccinic acid	146.08	98			
1565	$C_6H_{10}O_4$	Methylethylmalonic acid	146 08	117 5			
1566	$C_6H_{10}O_4$	Propylmalonic acid $C_3H_7CH(CO_2H)_2$	146 08	96			
1567	$C_6H_{10}O_4$	Isopropylmalonic acid	146.08	87			
1568	$C_6H_{10}O_4$	Dimethyl succinate $(CH_2CO_2CH_3)_2$	146 08	19 5	192 8	1 121	942
1569	$C_6H_{10}O_4$	Dimethyl isosuccinate	146 08		179	1.028[25]	
1570	$C_6H_{10}O_4$	Diethyl oxalate $(CO_2C_2H_5)_2$	146 08	−40 6	186 1	1 080	182
1571	$C_6H_{10}O_4$	Glycol diacetate $(CH_2OCOCH_3)_2$	146.08	−31	190 5	1.104	216
1572	$C_6H_{10}O_4$	Ethylidene diacetate	146.08		169	0.852	
1572.1	$C_6H_{10}O_4$	Methyl l-1-acetoxypropionate	146.08		172	1.089	
1573	$C_6H_{10}O_4$	Mannide	146 08		317		
1574	$C_6H_{10}O_4$	Isomannide	146.08	87	274		
1575	$C_6H_{10}O_5$	Lactic anhydride $(CH_3CHOHCO)_2$	162.08	260 d.			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1576	$C_4H_{10}O_5$	Dimethyl malate	162 08		242	1.233	391
1577	$C_6H_{10}O_5$	β-Glucosan	162 08	178			
1578	$(C_6H_{10}O_5)_x$	Glycogen	(162.08)$_x$	240			
1578 1	$(C_6H_{10}O_5)_x$	Starch	(162 08)$_x$	d.		1.50^{11}	1164
1579	$C_6H_{10}O_5$	d-Saccharine	162 08	161			
1580	$C_6H_{10}O_6$	Dimethyl dl-tartrate (CH(OH)CO$_2$CH$_3$)$_2$	178 08	85	282		
1581	$C_6H_{10}O_6$	Dimethyl d-tartrate	178 08	48; 61 5	280	1 328	
1582	$C_6H_{10}O_6$	Ethyl d-tartrate	178 08	90			
1583	$C_6H_{10}O_8$	Allomucic acid	210 08	171 d.			
1584	$C_6H_{10}O_8$	Mucic acid HO$_2$C(CHOH)$_4$CO$_2$H	210 08	206 d.			
1585	$C_6H_{10}O_8$	d(l)-Talomucic acid	210 08	158 d.			
1586	$C_6H_{10}O_8$	Isosaccharic acid	210 08	185			
1587	$C_6H_{10}S$	Diallyl sulfide (CH$_2$.CHCH$_2$)$_2$S	114 14	−83 0	138 7	0 888$^{26.4}_{4}$	1034
1588	$C_6H_{11}Br$	Cyclohexyl bromide	163 00		165 5	1.333	575
1589	$C_6H_{11}BrN_2O_2$	Bromural	223 02	154			
1590	$C_6H_{11}BrO_2$	1-Bromocaproic acid C$_4$H$_9$CHBrCO$_2$H.	195 00		131^{10}		
1591	$C_6H_{11}BrO_2$	2-Bromocaproic acid	195 00	35			
1592	$C_6H_{11}BrO_2$	Ethyl 1-bromobutyrate	195 00		179 d.	1 325$^{25}_{25}$	
1593	$C_6H_{11}BrO_2$	Ethyl 1-bromoisobutyrate	195 00		164 d.	1 315$^{24}_{25}$	
1595	$C_6H_{11}Cl$	Cyclohexyl chloride	118 54		142 5	0 973	451
1596	$C_6H_{11}ClO$	n-Caproyl chloride C$_5$H$_{11}$COCl	134 54		153		543
1597	$C_6H_{11}ClO_2$	Isoamyl chloroformate	150 54		156	1 024$^{25}_{25}$	
1598	$C_6H_{11}Cl_2N_3O_2$	Histidine dihydrochloride	228 03	235 d.			
1599	$C_6H_{11}Cl_3O_2$	Trichloroacetal Cl$_3$CCH(OC$_2$H$_5$)$_2$	221 46		197	1.266^{15}	
1600	$C_6H_{11}Cl_3O_2$	Trichloroacetal (solid)	221 46	83	230 d.		
1601	$C_6H_{11}I$	Cyclohexyl iodide	210 02		192	1.626	
1602	$C_6H_{11}N$	Capronitrile C$_5$H$_{11}$CN	97 09		163	0.809	188
1603	$C_6H_{11}N$	Isocapronitrile (CH$_3$)$_2$CH(CH$_2$)$_2$CN	97 09	−51 1	155 5	0.806	159
1604	$C_6H_{11}N$	Isocaproisonitrile (CH$_3$)$_2$CH(CH$_2$)$_2$NC	97.09		137		
1605	$C_6H_{11}NO_2$	Hygric acid	129 09	169			
1606	$C_6H_{11}NO_2$	Nitrocyclohexane	129 09	−34	205 5	1 068	
1607	$C_6H_{11}NO_2$	Adipyl amide HO$_2$C(CH$_2$)$_4$CONH$_2$	145 09	130			
1608	$C_6H_{11}NS$	Isoamyl isothiocyanate	129 16		182		
1609	$C_6H_{11}N_3O_4$	Citramide (H$_2$NOCCH$_2$)$_2$C(OH)CONH$_2$	189 11	215			
1610	C_6H_{12}	Butylethylene C$_4$H$_9$CH.CH$_2$	84 092	−98 5	64 1	0.683	44
1611	C_6H_{12}	2, 2-Dimethyl-4-butene	84 092		42 3		
1612	C_6H_{12}	Cyclohexane	84 092	6 5	81 4	0.779	304
1613	C_6H_{12}	2-Methyl-2-pentene (CH$_3$)$_2$C:CHC$_2$H$_5$.	84 092		67 1	0.692	881
1615	C_6H_{12}	Methylcyclopentane	84 092	−140 5	73	0.750	
1616	C_6H_{12}	3-Methyl-2-pentene (isomer 1)	84 092		65 7	0.722^{16}	848
1617	C_6H_{12}	3-Methyl-2-pentene (isomer 2)	84 092		70.2	0 698	128
1618	C_6H_{12}	2, 3-Dimethyl-1-butene	84 092		59	0.680^0	
1619	C_6H_{12}	Tetramethylethylene	84 092		73	0 712	199
1620	$C_6H_{12}As_2$	Cacodyl carbide	234 01		81 5^{15}		
1621	$C_6H_{12}As_3BiO_6$	Bismuth cacodylate (8H$_2$O).	613 97	82			
1622	$C_6H_{12}Cl_2O_2$	Dichloroacetal Cl$_2$CHCH(OC$_2$H$_5$)$_2$	187 01		184	1 138^{14}	
1623	$C_6H_{12}N_2O_2$	Adipic diamide H$_2$NOC(CH$_2$)$_4$CONH$_2$.	144 11	220			
1624	$C_6H_{12}N_2O_2$	sym -Diethyloxamide	144 11	190			
1625	$C_6H_{12}N_2O_4S_2$	l-Cystine	240 24	258 d.			1187
1626	$C_6H_{12}N_4$	Hexamethylenetetramine	140 12		263		
1627	$C_6H_{12}O$	Cyclohexanol	100.09	23 9	161 5	0 962	1051
1628	$C_6H_{12}O$	2-Hexene-1-ol	100 09		59^{27}	0 837	1008
1629	$C_6H_{12}O$	Dimethyl propenyl carbinol	100.09		112	0 835	321
1630	$C_6H_{12}O$	Pinacolin (CH$_3$)$_3$CCOCH$_3$	100.09	−52 5	106.2	0 811	
1631	$C_6H_{12}O$	Ethyl isocrotonyl ether	100.09		94		
1632	$C_6H_{12}O$	Isopropyl allyl ether	100.09		84 2	0 776	
1633	$C_6H_{12}O$	n-Caproic aldehyde C$_5$H$_{11}$CHO	100.09		129	0 834	
1634	$C_6H_{12}O$	Isobutylacetaldehyde	100.09		121.7		
1635	$C_6H_{12}O$	Methylpropylacetaldehyde	100 09		121		
1636	$C_6H_{12}O$	Ethyl propyl ketone C$_2$H$_5$COC$_3$H$_7$	100 09		124	0 818$^{17.5}$	124
1637	$C_6H_{12}O$	Ethyl isopropyl ketone	100 09		114.5	0 830$^0_0$	
1638	$C_6H_{12}O$	Methyl n-butyl ketone CH$_3$COC$_4$H$_9$.	100 09	−56 9	127.2	0 830^0	
1639	$C_6H_{12}O$	Methyl isobutyl ketone	100.09	−84.7	119	0.803	96

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1640	$C_6H_{12}O$	Methyl sec.-butyl ketone	100 09		117.8	0.815	115
1641	$C_6H_{12}O_2$	Diacetone alcohol	116 09		166	0.931[16]	
1642	$C_6H_{12}O_2$	tert.-Butylacetic acid	116 09	−11	190		
1643	$C_6H_{12}O_2$	Caproic acid $C_5H_{11}CO_2H$	116 09	−9 5	202	0 929	207
1644	$C_6H_{12}O_2$	Isocaproic acid	116 09	−35	207 7	0.925	217
1645	$C_6H_{12}O_2$	Diethylacetic acid $(C_2H_5)_2CHCO_2H$	116 09	< −15	197	0 933[10 2]	201
1646	$C_6H_{12}O_2$	Dimethylethylacetic acid	116 09	−14	187		
1647	$C_6H_{12}O_2$	Methylpropylacetic acid	116 09		193 5	0 928	
1648	$C_6H_{12}O_2$	n-Amyl formate $HCO_2C_5H_{11}$	116 09		130 4	0 902[0]	
1649	$C_6H_{12}O_2$	Isoamyl formate	116 09		123 5	0 871	83
1650	$C_6H_{12}O_2$	tert.-Amyl formate	116 09		113	0 896[16]	
1651	$C_6H_{12}O_2$	n-Butyl acetate $CH_3CO_2C_4H_9$	116 09	−76 8	126 5	0 882	95
1652	$C_6H_{12}O_2$	Isobutyl acetate $CH_3CO_2CH_2CH(CH_3)_2$	116 09	−98 9	118 3	0 871	118
1653	$C_6H_{12}O_2$	sec.-Butyl acetate	116 09		112 2	0.870	73
1654	$C_6H_{12}O_2$	Ethyl n-butyrate $C_3H_7CO_2C_2H_5$	116 09	−93 3	121 3	0 879	91
1655	$C_6H_{12}O_2$	Ethyl isobutyrate	116 09	−88 2	111 7	0 871	80
1656	$C_6H_{12}O_2$	Methyl trimethylacetate	116 09		102	1 044[0]	
1657	$C_6H_{12}O_2$	Methyl n-valerate $C_4H_9CO_2CH_3$	116 09		127 3	0 910[0]	
1658	$C_6H_{12}O_2$	Methyl isovalerate	116 09		116 7	0 881	
1659	$C_6H_{12}O_2$	n-Propyl propionate $C_2H_5CO_2C_3H_7$	116 09	−75 9	123 4	0 883	92
1660	$C_6H_{12}O_2$	Isopropyl propionate	116 09		111 3	0 803[0]	
1661	$C_6H_{12}O_3$	Phloroglucite	132 09	185			
1662	$C_6H_{12}O_3$	Paraldehyde $(CH_3CHO)_3$	132 09	10 5	124	0.994	244
1663	$C_6H_{12}O_3$	1-Hydroxy-n-caproic acid	132 09	62			
1664	$C_6H_{12}O_3$	1-Hydroxyisocaproic acid	132 09	81			
1665	$C_6H_{12}O_3$	dl-1-Hydroxyisocaproic acid	132 09	76			
1666	$C_6H_{12}O_3$	1-Hydroxy-1, 1-diethylacetic acid	132 09	74 5			
1667	$C_6H_{12}O_3$	Methyl n-butyl carbonate	132 09		151		
1668	$C_6H_{12}O_5$	Fucose	164 09	145			
1669	$C_6H_{12}O_5$	Mannitan	164 09	137			
1670	$C_6H_{12}O_5$	d-Quercitol	164.09	234		1.585[16]	
1671	$C_6H_{12}O_5$	l-Quercitol	164 09	174			
1672	$C_6H_{12}O_5$ (H_2O)	β-Rhamnose	164 09	126		1.471	1219
1673	$C_6H_{12}O_5$	Rhodeose	164 09	144			
1674	$C_6H_{12}O_6$	d-Fructose (Levulose)	180 09	104		1.669[17 4]	
1675	$C_6H_{12}O_6$	d, α-Galactose	180 09	168			
1675 1	$C_6H_{12}O_6$	d, β-Galactose	180 09	168			
1676	$C_6H_{12}O_6$	dl-Galactose	180 09	144			
1677	$C_6H_{12}O_6$	d, α-Glucose	180 09	146		1.544[26]	
1678	$C_6H_{12}O_6$	d, β-Glucose	180.09	150			
1679	$C_6H_{12}O_6$	d(l)-Inosite	180 09	247	250 vac.		
1680	$C_6H_{12}O_6$	Dambose	180 09	224	d.	1.752	
1681	$C_6H_{12}O_6$	α-Mannose	180 09	133	205 d.		
1682	$C_6H_{12}O_6$	d-Mannose	180 09	132		1.539	
1683	$C_6H_{12}O_6$	dl-Mannose	180 09	133			
1684	$C_6H_{12}O_6$	d(l)-Sorbose	180 09	154		1.612	
1685	$C_6H_{12}O_6$	dl-Sorbose	180 09	154		1.638	
1686	$C_6H_{12}O_6$	d-Tagatose	180 09	124			
1687	$C_6H_{12}S$	Cyclohexyl mercaptan	116.16		160		
1688	$C_6H_{12}S_3$	α-Trithioacetaldehyde	180 29	101	247		
1689	$C_6H_{12}S_3$	β-Trithioacetaldehyde $(C_2H_4S)_3$	180 29	126			
1690	$C_6H_{12}S_3$	γ-Trithioacetaldehyde	180 29	81	100		
1690.1	$C_6H_{12}Se$	Hexamethyl selenide	163 29		172	1.122	
1691	$C_6H_{13}Br$	2-Bromo-2, 3-dimethylbutane	165.02	13	132		
1692	$C_6H_{13}Br$	n-Hexyl bromide $C_5H_{11}CH_2Br$	165 02		156	1.173	422
1693	$C_6H_{13}BrO_2$	Bromoacetal $BrCH_2CH(OC_2H_5)_2$	197.02		170		
1694	$C_6H_{13}Cl$	2-Chloro-2, 3-dimethylbutane	120.56	−10.4	112.1	0.875[26]	
1695	$C_6H_{13}Cl$	n-Hexyl chloride $C_5H_{11}CH_2Cl$	120.56		134	0.872	238
1696	$C_6H_{13}ClN_4O_4$	Hexamethylenetetramine perchlorate	240.59	158			
1697	$C_6H_{13}I$	n-Hexyl iodide $C_5H_{11}CH_2I$	212.03		180	1.441	560
1698	$C_6H_{13}IO_2$	Iodoacetal $ICH_2CH(OC_2H_5)_2$	244 03		132[00]	1.494[16]	416
1699	$C_6H_{13}N$	1-Methylpiperidine	99 108		107	0.818	416
1700	$C_6H_{13}N$	2-Methylpiperidine (α-Pipecoline)	99.108		119	0.844[23 6]	1016

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. L. No.
1701	C₆H₁₃N	3-Methylpiperidine (β-Pipecoline).	99.108		126	0.845¹⁴·⁸	1020
1702	C₆H₁₃N	4-Methylpiperidine (γ-Pipecoline).....	99 108		129	0.867⁰	
1703	C₆H₁₃NO₂	Hedonal H₂NCO₂CH(CH₃)C₃H₇,...	131 11	74	215		
1704	C₆H₁₃NO₂	Isoamyl carbamate	131.11	63 5	220		
1704.1	C₆H₁₃NO₂	Propyl urethane C₃H₇NHCO₂C₂H₅	131 11		186	0 992¹⁴	
1705	C₆H₁₃NO₂	l-Leucine (CH₃)₂CHCH(NH₂)CO₂H	131 11	295		1.293	1221
1706	C₆H₁₃NO₂	dl-Leucine	131 11	290			
1707	C₆H₁₃NO₂	d(l)-Isoleucine ...	131 11	280 d.			
1708	C₆H₁₃NO₂	dl-Isoleucine .	131 11	275			
1709	C₆H₁₃NO₅	d-Glucosamine ...	179 11	110 d.			
1710	C₆H₁₃NO₅	d-Glucosamine. .	179 11	128			
1711	C₆H₁₃NO₆	d-Glucosoxime.	195 11	138			
1712	C₆H₁₄	Diisopropyl (CH₃)₂CHCH(CH₃)₂	86 108	−135 1	58.1	0.666¹⁴	38
1713	C₆H₁₄	n-Hexane CH₃(CH₂)₄CH₃	86 108	−94 3	69 0	0 660	32
1714	C₆H₁₄	3-Methylpentane (C₂H₅)₂CHCH₃	86 108		64	0 668	34
1715	C₆H₁₄	2-Methylpentane (CH₃)₂CHC₃H₇	86 108		60 0	0 654	27
1716	C₆H₁₄	2, 2-Dimethylbutane (CH₃)₃CC₂H₅.	86 108	−98 2	49.7	0 649	23
1717	C₆H₁₄INO₅	d-Glucosamine hydroiodide	307 05	165 d.			
1718	C₆H₁₄N₂	α, 2, 5-Dimethylpiperazine	114 12	119	162		
1719	C₆H₁₄N₂O	Diacetoneamineoxime	130 12	58	135¹⁷		
1720	C₆H₁₄N₂O	Dipropylnitrosamine (C₃H₇)₂NNO	130 12		205		
1721	C₆H₁₄N₂O₇	Ammonium citrate	226 12			1 483	
1722	C₆H₁₄N₄O₂	Arginine	174 14	207 5 d.			
1723	C₆H₁₄O	tert-Amyl carbinol	102 11		135	0.844⁰₄	
1724	C₆H₁₄O	Isohexyl alcohol .	102 11		165	0 840⁰₄	429
1725	C₆H₁₄O	Dimethylisopropyl carbinol ..	102 11	−14	122	0 823	
1726	C₆H₁₄O	Ethylpropyl carbinol . ..	102 11		135	0 819	
1726.1	C₆H₁₄O	l(d)-Ethylpropyl carbinol .	102 11		134⁷³³	0 825¹³·⁵	211
1727	C₆H₁₄O	Ethylisopropyl carbinol	102 11		128	0 824	
1728	C₆H₁₄O	n-Hexyl alcohol C₆H₁₃OH	102 11	−51 6	155 8	0 820	
1730	C₆H₁₄O	Methylbutyl carbinol .	102 11		131.9	0 803²⁵	183
1730.1	C₆H₁₄O	d-Methylbutyl carbinol ...	102 11		138	0.815	205
1732	C₆H₁₄O	Methyl-sec.-butyl carbinol .	102 11		134	0.831¹⁸	245
1733	C₆H₁₄O	Pinacolyl alcohol (CH₃)₃CH(OH)CH₃	102 11	5 5	121	0 812²⁵	
1733 1	C₆H₁₄O	d-Pinacolyl alcohol	102 11		120	0 820	214
1734	C₆H₁₄O	Methyldiethyl carbinol ...	102 11	−22	122 6	0 824	242
1735	C₆H₁₄O	3-Methyl-3-ethylpropyl alcohol ...	102 11		152 1	0 830¹⁶	
1736	C₆H₁₄O	2-Methyl-2-propylethyl alcohol	102 11		147 9	0 829	231
1737	C₆H₁₄O	Ethyl n-butyl ether C₄H₉OC₂H₅	102 11		91 4	0 752	
1738	C₆H₁₄O	Ethyl isobutyl ether.. .	102 11		80	0 751	
1739	C₆H₁₄O	Methyl n-amyl ether C₅H₁₁OCH₃	102 11		88 5	0 754	53
1740	C₆H₁₄O	Methyl isoamyl ether	102 11		91	0 687²¹	
1741	C₆H₁₄O	Propyl ether (C₃H₇)₂O .	102 11	−122 0	89	0 747	41
1742	C₆H₁₄O	Isopropyl ether [(CH₃)₂CH]₂O	102 11		68 7	0 735¹⁶·²	
1743	C₆H₁₄O₂	Pinacone [(CH₃)₂COH]₂	118 11	38	172 8		
1744	C₆H₁₄O₂	Hexane-1, 5-diol .	118 11		233	0 981⁰	
1745	C₆H₁₄O₂	Hexane-1, 6-diol HOCH₂(CH₂)₄CH₂OH	118 11	42	250		
1746	C₆H₁₄O₂	Acetal CH₃CH(OC₂H₅)₂	118 11		102.2	0 831	42
1747	C₆H₁₄O₅	Diglycerol [(HO)₂C₃H₅]₂O .	166 11		230¹⁰		
1748	C₆H₁₄O₅	Fucitol . .	166 11	153			
1749	C₆H₁₄O₅	Rhamnitol	166 11	121			
1750	C₆H₁₄O₆	Dulcitol	182 11	188	205³·⁵	1.466¹⁵	1333
1751	C₆H₁₄O₆	d-Mannitol	182.11	166 1	295³·⁵	1 489	1333
1752	C₆H₁₄O₆	d-Sorbitol	182 11	110			1333
1753	C₆H₁₄O₆	d-Talitol	182 11	86			
1754	C₆H₁₄S	Dipropyl sulfide (C₃H₇)₂S	118 17		142	0.814	
1755	C₆H₁₄S	Diisopropyl sulfide [(CH₃)₂CH]₂S. . .	118.17		120 4		
1756	C₆H₁₅As	Triethyl arsine (C₂H₅)₃As............	162 08		141 d.	1.150	495
1757	C₆H₁₅AsO₃	Triethyl arsenite (C₂H₅O)₃As...	210 08		166	1.224⁰₄	
1758	C₆H₁₅AsO₄	Triethyl arsenate (C₂H₅O)₃AsO......	226 08		238	1.326⁰	
1759	C₆H₁₅Bi	Triethyl bismuthine (C₂H₅)₃Bi.	296.12		107¹⁹	1.82	
1760	C₆H₁₅N	Di-n-propylamine (C₃H₇)₂NH	101.12	−39.6	110.7	0 738	149
1761	C₆H₁₅N	Diisopropylamine [(CH₃)₂CH]₂NH.....	101.12		84	0.722²¹	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1762	$C_6H_{15}N$	n-Hexylamine $C_6H_{13}NH_2$	101.12		128		
1762.1	$C_6H_{15}N$	2-Hexylamine $C_4H_9CH(NH_2)CH_3$	101 12	– 19	130[14]	0.767[20.4]	
1763	$C_6H_{15}N$	Isohexylamine $(CH_3)_2CH(CH_2)_3NH_2$	101 12	–94 4	123 9		
1764	$C_6H_{15}N$	Triethylamine $(C_2H_5)_3N$	101 12	–114 8	89 5	0 728	129
1765	$C_6H_{15}NO_2$	Aminoacetal $H_2NCH_2CH(OC_2H_5)_2$	133 12		163		
1766	$C_6H_{15}N_3$	Acetaldehydeammonia (trimeric)	129 14	85			
1767	$C_6H_{15}O_3P$	Triethyl phosphite $(C_2H_5O)_3P$	166 14		156 5	1.076[11.4]	169
1768	$C_6H_{15}O_4P$	Triethyl phosphate $(C_2H_5O)_3PO$	182 14		216	1 072[15]	150
1769	$C_6H_{15}P$	Triethylphosphine $(C_2H_5)_3P$	118 14		128	0.800	413
1769 1	$C_6H_{15}PS$	Triethyl phosphinesulfide....	150 20	94			1182
1770	$C_6H_{15}Sb$	Triethyl stibine $(C_2H_5)_3Sb$	208 89		159 5	1 324[18]	
1771	$C_6H_{16}ClN$	Triethylamine hydrochloride	137 59	254		1 069	
1772	$C_6H_{16}N_2$	Hexamethylenediamine $H_2N(CH_2)_6NH_2$	116 14	39	196		
1773	$C_6H_{18}N_6O_4S$	1, 1-Dimethylguanidine sulfate	270 25	288 d.			
1775	$C_7HCl_5O_2$	Pentachlorobenzoic acid $C_6Cl_5CO_2H$	294 30	201			
1776	$C_7H_2Br_4O_2$	2, 3, 4, 6-Tetrabromobenzoic acid	437 68	171			
1777	$C_7H_2Cl_4O_2$	2, 3, 4, 5-Tetrachlorobenzoic acid	259 85	186			
1778	$C_7H_3Br_3O_2$	2, 3, 4-Tribromobenzoic acid	358 77	198			
1779	$C_7H_3Br_3O_2$	2, 3, 5-Tribromobenzoic acid	358 77	194			
1780	$C_7H_3Br_3O_2$	2, 4, 5-Tribromobenzoic acid	358 77	196			
1781	$C_7H_3Br_3O_2$	2, 4, 6-Tribromobenzoic acid	358 77	187			
1782	$C_7H_3Br_3O_2$	3, 4, 5-Tribromobenzoic acid	358 77	235			
1783	$C_7H_3Cl_3O_2$	2, 3, 4-Trichlorobenzoic acid	225 40	129			
1784	$C_7H_3Cl_3O_2$	2, 3, 5-Trichlorobenzoic acid	225 40	163			
1785	$C_7H_3Cl_3O_2$	2, 4, 5-Trichlorobenzoic acid	225 40	163			
1786	$C_7H_3Cl_3O_2$	2, 4, 6-Trichlorobenzoic acid	225 40	160			
1787	$C_7H_3Cl_3O_2$	3, 4, 5-Trichlorobenzoic acid	225 40	203			
1788	$C_7H_3N_3O_7$	2, 4, 6-Trinitrobenzaldehyde	241 05	119			
1789	$C_7H_3N_3O_8$	2, 4, 6-Trinitrobenzoic acid	257 05	190			
1790	C_7H_4BrClO	o-Bromobenzoyl chloride	219 41		243		
1791	C_7H_4BrClO	m-Bromobenzoyl chloride	219 41		239		
1792	C_7H_4BrClO	p-Bromobenzoyl chloride	219 41	42	247 s d		
1793	C_7H_4BrN	o-Bromobenzonitrile ...	181 96	51	253		
1794	C_7H_4BrN	m-Bromobenzonitrile..	181 96	38	225		
1795	C_7H_4BrN	p-Bromobenzonitrile	181 96	113	237		
1796	$C_7H_4Br_2O_2$	2, 3-Dibromobenzoic acid	279 86	150			
1797	$C_7H_4Br_2O_2$	2, 4-Dibromobenzoic acid	279 86	169			
1798	$C_7H_4Br_2O_2$	2, 5-Dibromobenzoic acid	279 86	153			
1799	$C_7H_4Br_2O_2$	2, 6-Dibromobenzoic acid	279 86	147			
1800	$C_7H_4Br_2O_2$	3, 4-Dibromobenzoic acid	279 86	230			
1801	$C_7H_4Br_2O_2$	3, 5-Dibromobenzoic acid	279 86	211			
1802	$C_7H_4Br_2O_5$	2, 6-Dibromo-3, 4, 5-trihydroxybenzoic acid............	327 86	150			
1803	C_7H_4ClFO	o-Fluorobenzoyl chloride	158 49		206		
1804	C_7H_4ClFO	m-Fluorobenzoyl chloride	158 49		189		
1805	C_7H_4ClFO	p-Fluorobenzoyl chloride p-FC_6H_4COCl	158 49		193		
1806	$C_7H_4ClNO_3$	o-Nitrobenzoyl chloride	185 50	75	205[105]		
1807	$C_7H_4ClNO_3$	m-Nitrobenzoyl chloride	185 50	34	278		
1808	$C_7H_4ClNO_3$	p-Nitrobenzoyl chloride	185.50	72	154[16]		
1809	$C_7H_4Cl_2O$	2, 4-Dichlorobenzaldehyde	174 95	71			
1810	$C_7H_4Cl_2O$	2, 5-Dichlorobenzaldehyde	174 95	58	233	1 231[70]	
1811	$C_7H_4Cl_2O$	3, 4-Dichlorobenzaldehyde	174 95	44	248		
1812	$C_7H_4Cl_2O$	o-Chlorobenzoyl chloride	174 95	–4	238		
1813	$C_7H_4Cl_2O$	m-Chlorobenzoyl chloride	174 95		117 5[26]		
1814	$C_7H_4Cl_2O$	p-Chlorobenzoyl chloride	174 95		119[27 5]		
1815	$C_7H_4Cl_2O_2$	2, 3-Dichlorobenzoic acid	190 95	166			
1816	$C_7H_4Cl_2O_2$	2, 4-Dichlorobenzoic acid	190.95	164 2			
1817	$C_7H_4Cl_2O_2$	2, 5-Dichlorobenzoic acid	190 95	154 4	301		
1818	$C_7H_4Cl_2O_2$	2, 6-Dichlorobenzoic acid.	190.95	143.7			
1819	$C_7H_4Cl_2O_2$	3, 4-Dichlorobenzoic acid.	190.95	204 1			
1820	$C_7H_4Cl_2O_2$	3, 5-Dichlorobenzoic acid	190.95	188.1			
1821	$C_7H_4Cl_3NO_2$	2, 3, 4-Trichloronitrotoluene	240.41	60			
1822	$C_7H_4Cl_4$	2-Chloro-1-trichloromethylbenzene	229.86	30	260	1.51	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I No.
1823	C₇H₄FNO₄	2-Fluoro-5-nitrobenzoic acid	185 04	139			
1824	C₇H₄FNO₄	3-Fluoro-4-nitrobenzoic acid	185 04	122			
1825	C₇H₄FNO₄	3-Fluoro-6-nitrobenzoic acid	185 04	134 5			
1826	C₇H₄FNO₄	4-Fluoro-2-nitrobenzoic acid	185 04	130			
1827	C₇H₄FNO₄	4-Fluoro-3-nitrobenzoic acid	185 04	121 5			
1828	C₇H₄J₂O₃	3, 5-Duodosalicylic acid	389 90	230 d.			
1829	C₇H₄N₂O₂	o-Nitrobenzonitrile	148 05	109			
1830	C₇H₄N₂O₂	m-Nitrobenzonitrile	148 05	118			
1831	C₇H₄N₂O₂	p-Nitrobenzonitrile	148 05	147			
1832	C₇H₄N₂O₄	2, 4-Dinitrobenzaldehyde	196 05	72			
1833	C₇H₄N₂O₄	2, 6-Dinitrobenzaldehyde	196 05	123			
1834	C₇H₄N₂O₆	2, 3-Dinitrobenzoic acid	212 05	201			
1835	C₇H₄N₂O₆	2, 4-Dinitrobenzoic acid	212 05	179			
1836	C₇H₄N₂O₆	2, 5-Dinitrobenzoic acid	212 05	177			
1837	C₇H₄N₂O₆	2, 6-Dinitrobenzoic acid	212 05	202 d.			
1838	C₇H₄N₂O₆	3, 4-Dinitrobenzoic acid	212 05	163			
1839	C₇H₄N₂O₆	3, 5-Dinitrobenzoic acid	212.05	205			
1840	C₇H₄N₂O₇	3, 5-Dinitro-2-hydroxybenzoic acid	228 05	174			
1841	C₇H₄N₄O₉	2, 3, 5, 6-Tetranitroanisol	288.06	154; 112			
1842	C₇H₄O₄S	o-Sulfobenzoic anhydride	184.10	130			
1843	C₇H₄O₇	Meconic acid	200.03	d.			1333
1844	C₇H₅BrO	Benzoyl bromide C₆H₅COBr	184 96	0	219	1.570	
1845	C₇H₅BrO₂	o-Bromobenzoic acid	200 96	148			
1846	C₇H₅BrO₂	m-Bromobenzoic acid	200 96	152			
1847	C₇H₅BrO₂	p-Bromobenzoic acid	200 96	251			
1848	C₇H₅BrO₃	3-Bromo-2-hydroxybenzoic acid	216.96	220			
1849	C₇H₅BrO₃	5-Bromo-2-hydroxybenzoic acid	216.96	165			
1850	C₇H₅Br₃	2, 3, 4-Tribromotoluene	328.79	45			
1851	C₇H₅Br₃	2, 3, 5-Tribromotoluene	328.79	54			
1852	C₇H₅Br₃	2, 3, 6-Tribromotoluene	328 79	59			
1853	C₇H₅Br₃	2, 4, 5-Tribromotoluene	328.79	113			
1854	C₇H₅Br₃	2, 4, 6-Tribromotoluene	328.79	66			
1855	C₇H₅Br₃	3, 4, 5-Tribromotoluene	328.79	89			
1856	C₇H₅ClO	o-Chlorobenzaldehyde	140.50	−3	205	1 252	753
1857	C₇H₅ClO	m-Chlorobenzaldehyde	140.50	18	204	1 241	751
1858	C₇H₅ClO	p-Chlorobenzaldehyde	140.50	47.5	214	1.196⁶¹	1092
1859	C₇H₅ClO	Benzoyl chloride C₆H₅COCl	140.50	−0.8	197.2	1.211	737
1860	C₇H₅ClO₂	o-Chlorobenzoic acid	156.50	140.7			
1861	C₇H₅ClO₂	m-Chlorobenzoic acid	156.50	154.9			
1862	C₇H₅ClO₂	p-Chlorobenzoic acid	156 50	241 5			
1863	C₇H₅ClO₂	Salicyl chloride o-HOC₆H₄COCl	156.50	18.0	59¹⁰ s. d.		
1864	C₇H₅ClO₃	5-Chloro-2-hydroxybenzoic acid	172.50	167.5			
1865	C₇H₅Cl₂NO₂	m-Nitrobenzal chloride	205.96	65			
1866	C₇H₅Cl₃NO₄S	Halazone	270.03	213			
1868	C₇H₆Cl₂	o-Chlorobenzal chloride	195.41		228.5	1 399¹⁵	
1869	C₇H₆Cl₂	p-Chlorobenzal chloride	195.41		234		
1870	C₇H₅Cl₃	Benzotrichloride C₆H₅CCl₃	195.41	−4.8	220.7	1 378¹⁵	
1871	C₇H₅Cl₃	2, 3, 4-Trichlorotoluene	195.41	41	234		
1872	C₇H₅Cl₃	2, 4, 5-Trichlorotoluene	195.41	82	232		
1873	C₇H₅Cl₃	3, 4, 5-Trichlorotoluene	195.41	42.5	247		
1874	C₇H₅Cl₃O	2, 4, 6-Trichloro-3-hydroxytoluene	211.41	46			
1875	C₇H₅Cl₃O	2, 4, 6-Trichloroanisol	211.41	60.5	240.7		
1876	C₇H₅FO	Benzoyl fluoride C₆H₅COF	124.04		162		
1877	C₇H₅FO₂	o-Fluorobenzoic acid	140.04	122			
1878	C₇H₅FO₂	m-Fluorobenzoic acid	140.04	124			
1879	C₇H₅FO₂	p-Fluorobenzoic acid	140.04	182			
1880	C₇H₅IO	Benzoyl iodide C₆H₅COI	231.97	3	135¹⁴		
1881	C₇H₅IO₂	o-Iodobenzoic acid	247.97	162			
1882	C₇H₅IO₂	m-Iodobenzoic acid	247.97	185			
1883	C₇H₅IO₂	p-Iodobenzoic acid	247.97	266			
1884	C₇H₅IO₃	3-Iodo-2-hydroxybenzoic acid	263.97	198			
1885	C₇H₅N	Benzonitrile C₆H₅CN	103.05	−13.1	190.7	1.008¹⁴·⁸	1028
1886	C₇H₅N	Phenyl isocyanide C₆H₅NC	103.05		166 d.	0.978¹⁵	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
1887	C₇H₅NO	Anthranil............	119 05	> −18	215	1.187¹⁴	768
1888	C₇H₅NO	Benzoxazol ..	119 05	30 5	182 5		
1889	C₇H₅NO	Phenyl isocyanate C₆H₅N:CO	119 05		165 6	1 095	
1890	C₇H₅NO	Salicylic nitrile o-OHC₆H₄CN	119 05	98			
1891	C₇H₅NOS	1-Hydroxybenzothiazole	151 11	136			
1892	C₇H₅NOS	1-Mercaptobenzoxazole	151 11	193			
1893	C₇H₅NO₂	o-Nitrobenzaldehyde	151 05	α40.9; β37 9	156¹⁴		
1894	C₆H₅NO₂	m-Nitrobenzaldehyde	151 05	58 0	164²²		
1895	C₇H₅NO₂	p-Nitrobenzaldehyde	151 05	106 5			
1896	C₇H₅NO₃S	o-Benzoicsulfimide (Saccharin)	183 11	228 d.			
1897	C₇H₅NO₄	o-Nitrobenzoic acid	167 05	147 5		1 575⁴	
1898	C₇H₅NO₄	m-Nitrobenzoic acid	167 05	141 4		1 494⁴	
1899	C₇H₅NO₄	p-Nitrobenzoic acid	167 05	242 4		1 550¹²	
1900	C₇H₅NO₄	Quinolinic acid	167 05	190 d.			
1901	C₇H₅NO₄	Lutidinic acid	167 05	248			
1902	C₇H₅NO₄	Isocinchomeronic acid	167 05	237			
1903	C₇H₅NO₄	Dipicolinic acid	167 05	226 d			
1904	C₇H₅NO₄	Cinchomeronic acid	167 05	258 d			
1905	C₇H₅NO₄	Dinicotinic acid	167 05	323			
1906	C₇H₅NO₅	Ammonchelidonic acid	183 05	220 d.			
1907	C₇H₅NO₅	3-Nitro-2-hydroxybenzoic acid	183 05	144			
1908	C₇H₅NO₅	4-Nitro-2-hydroxybenzoic acid	183 05	235			
1909	C₇H₅NO₅	5-Nitro-2-hydroxybenzoic acid	183 05	228			
1910	C₇H₅NO₅	6-Nitro-2-hydroxybenzoic acid	183 05	130			
1911	C₇H₅NO₅	2-Nitro-3-hydroxybenzoic acid	183 05	178			
1912	C₇H₅NO₅	4-Nitro-3-hydroxybenzoic acid	183 05	230			
1913	C₇H₅NO₅	5-Nitro-3-hydroxybenzoic acid	183 05	167			
1914	C₇H₅NO₅	6-Nitro-3-hydroxybenzoic acid	183 05	169			
1915	C₇H₅NO₅	3-Nitro-4-hydroxybenzoic acid	183 05	185			
1916	C₇H₅NS	Benzothiazol.........	135 11		230	1.248	
1917	C₇H₅NS	Phenyl thiocyanate C₆H₅CNS	135 11		232	1.155	
1918	C₇H₅NS	Phenyl isothiocyanate C₆H₅N:CS	135.11	−21	218 5	1.135¹⁴ ⁵	798
1919	C₇H₅N₃	1, 2, 3-Benzotriazin ..	131 06	75	240		
1920	C₇H₅N₃O₆	Chrysanisic acid.	227 06	259			
1921	C₇H₅N₃O₆	2, 3, 4-Trinitrotoluene	227 06	112	302 d.	1 620	
1922	C₇H₅N₃O₆	2, 3, 5-Trinitrotoluene	227.06	97	335 d.		
1923	C₇H₅N₃O₆	2, 3, 6-Trinitrotoluene	227 06	111	333 d.		
1924	C₇H₅N₃O₆	2, 4, 6-Trinitrotoluene (T. N. T)	227 06	80 7	240 exp	1.654	
1925	C₇H₅N₃O₆	3, 4, 5-Trinitrotoluene..	227.06	137 5	313 d.		
1926	C₇H₅N₃O₆	3, 4, 6-Trinitrotoluene	227 06	104	291 d.	1.620	
1927	C₇H₅N₃O₇	2, 3, 4-Trinitroanisol	243 06	155	exp.		
1928	C₇H₅N₃O₇	2, 3, 5-Trinitroanisol	243 06	104		1.618¹⁴	
1929	C₇H₅N₃O₇	2, 4, 6-Trinitroanisol.	243 06	68.4		1 408	
1930	C₇H₅N₃O₇	3, 4, 5-Trinitroanisol	243.06	120			
1931	C₇H₅N₃O₇	3, 4, 6-Trinitroanisol	243.06	107			
1932	C₇H₅N₃O₇	2, 4, 6-Trinitro-3-hydroxytoluene	243.06	106			
1933	C₇H₅N₅O₈	2, 4, 6-Trinitrophenylmethylnitramine (Tetryl)............	287 08	130	exp. 187		
1934	C₇H₆BrCl	o-Bromobenzyl chloride	205 42		115¹⁵		
1935	C₇H₆BrCl	p-Bromobenzyl chloride	205.42	51			
1936	C₇H₆BrCl	o-Chlorobenzyl bromide	205 42		120¹⁰		
1937	C₇H₆BrCl	p-Chlorobenzyl bromide	205.42	48			
1938	C₇H₆BrNO	o-Bromobenzamide......	199 97	156			
1939	C₇H₆BrNO	m-Bromobenzamide...	199.97	150			
1940	C₇H₆BrNO	p-Bromobenzamide.......	199.97	190			
1941	C₇H₆BrNO₂	o-Nitrobenzyl bromide..	215 97	46			
1942	C₇H₆BrNO₂	m-Nitrobenzyl bromide.	215.97	58			
1943	C₇H₆BrNO₂	p-Nitrobenzyl bromide. ..	215 97	100			
1944	C₇H₆Br₂	Benzal bromide C₆H₅CHBr₂	249 88		140²⁰	1.51¹⁶	716.1
1945	C₇H₆Br₂	o-Bromobenzyl bromide.. ..	249.88	30			
1946	C₇H₆Br₂	m-Bromobenzyl bromide	249.88	41			
1947	C₇H₆Br₂	p-Bromobenzyl bromide	249.88	61			
1948	C₇H₆Br₂	2, 3-Dibromotoluene............	249.88	31			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
1949	$C_7H_6Br_2$	2, 6-Dibromotoluene.	249.88	5.5	246	1.812²²	
1950	$C_7H_6Br_2$	3, 5-Dibromotoluene	249.88	39			
1951	C_7H_6ClNO	o-Chlorobenzamide.	155.51	141			
1952	C_7H_6ClNO	m-Chlorobenzamide...	155.51	134.5			
1953	C_7H_6ClNO	p-Chlorobenzamide.	155.51	178.3			
1954	$C_7H_6ClNO_2$	3-Chloro-2-nitrotoluene	171.51	23			
1955	$C_7H_6ClNO_2$	4-Chloro-2-nitrotoluene.	171.51	38.2	242	1 256⁶⁰	
1956	$C_7H_6ClNO_2$	5-Chloro-2-nitrotoluene	171.51	44	250		
1957	$C_7H_6ClNO_2$	6-Chloro-2-nitrotoluene	171.51	37	238		
1958	$C_7H_6ClNO_2$	2-Chloro-3-nitrotoluene	171.51	21.5	263		
1959	$C_7H_6ClNO_2$	4-Chloro-3-nitrotoluene	171.51	7	260.5	1.297²²	
1960	$C_7H_6ClNO_2$	5-Chloro-3-nitrotoluene.	171.51	61			
1961	$C_7H_6ClNO_2$	o-Nitrobenzyl chloride ...	171.51	49			1093
1962	$C_7H_6ClNO_2$	m-Nitrobenzyl chloride ..	171.51	44 5	183³⁶		1094
1963	$C_7H_6ClNO_2$	p-Nitrobenzyl chloride ...	171.51	71			1095
1964	$C_7H_6Cl_2$	Benzal chloride $C_6H_5CHCl_2$	160.96	−17 4	214	1 295¹⁶	
1965	$C_7H_6Cl_2$	o-Chlorobenzyl chloride .	160.96		214		
1966	$C_7H_6Cl_2$	p-Chlorobenzyl chloride	160.96	29	214		
1967	$C_7H_6Cl_2O$	1, 1-Dichloro-2-hydroxytoluene	176 96	82			
1968	$C_7H_6Cl_2O$	3, 5-Dichloro-2-hydroxytoluene	176.96	55			
1969	$C_7H_6Cl_2O$	4, 6-Dichloro-3-hydroxytoluene	176.96	46			
1970	$C_7H_6Cl_2O_2$	4, 5-Dichloro-2-methoxyphenol	192 96	72	270		
1971	C_7H_6FNO	o-Fluorobenzamide ...	139.05	116			
1972	C_7H_6FNO	m-Fluorobenzamide.	139.05	130			
1973	C_7H_6FNO	p-Fluorobenzamide	139.05	154 5			
1974	C_7H_6INO	o-Iodobenzamide .	246 99	183.6			
1975	C_6H_6INO	m-Iodobenzamide	246 99	186.5			
1976	C_7H_6INO	p-Iodobenzamide	246 99	217.6			
1977	$C_7H_6N_2$	Benzimidazol	118.06	170	<360		1270
1978	$C_7H_6N_2$	Cyanilide $CNNHC_6H_5$	118.06	47			
1979	$C_7H_6N_2$	Indazole...	118.06	146.5	270.6		
1980	$C_7H_6N_2O_2$	Ricinine acid	150.06	298			
1981	$C_7H_6N_2O_3$	o-Nitrobenzamide	166.06	176 6	317	1 462²²	
1982	$C_7H_6N_2O_3$	m-Nitrobenzamide	166.06	142.7	315		
1983	$C_7H_6N_2O_3$	p-Nitrobenzamide	166.06	201 4			
1984	$C_7H_6N_2O_4$	2, 3-Dinitrotoluene	182.06	59.3		1 263¹¹¹	
1985	$C_7H_6N_2O_4$	2, 4-Dinitrotoluene	182 06	69 6	300 s. d.	1.521¹⁵	1297
1986	$C_7H_6N_2O_4$	2, 5-Dinitrotoluene	182.06	50 5		1.282¹¹¹	
1987	$C_7H_6N_2O_4$	2, 6-Dinitrotoluene	182 06	61		1 283¹¹¹	1300
1988	$C_7H_6N_2O_4$	3, 4-Dinitrotoluene	182.06	59 8		1 259¹¹¹	
1989	$C_7H_6N_2O_4$	3, 5-Dinitrotoluene	182.06	93		1.277¹¹¹	
1990	$C_7H_6N_2O_5$	2, 4-Dinitroanisol	198.06	95 2		1.341	
1991	$C_7H_6N_2O_5$	2, 5-Dinitroanisol	198.06	97.0	360	1 476	
1992	$C_7H_6N_2O_5$	2, 6-Dinitroanisol	198 06	117 5		1.319	
1993	$C_7H_6N_2O_5$	3, 4-Dinitroanisol ..	198 06	69 3		1.334¹¹⁰	
1994	$C_7H_6N_2O_5$	3, 5-Dinitroanisol	198 06	105.8		1.558¹²	
1995	$C_7H_6N_2O_5$	2, 4-Dinitro-3-hydroxytoluene	198.06	99			
1996	$C_7H_6N_2O_5$	3, 5-Dinitro-4-hydroxytoluene.	198 06	85.8			
1997	$C_7H_6N_2O_5$	4, 6-Dinitro-2-methoxyphenol...	214.06	123			
1998	$C_7H_6N_2O_7S$	2, 6-Dinitrotoluene-4-sulfonic acid	262.13	165			
1999	$C_7H_6N_2S$	1-Aminobenzothiazole...........	150.13	127			
2000	$C_7H_6N_4O_7$	2, 4, 6-Trinitro-3-aminoanisol.	258.08	131			
2001	C_7H_6O	Benzaldehyde C_6H_5CHO....	106.05	−56.0	179 5	1.046	725
2002	C_7H_6OS	Thiobenzoic acid C_6H_5COSH ..	138.11	24			
2003	$C_7H_6O_2$	Furfuracrolein..................	122.05	51	200		
2004	$C_7H_6O_2$	Salicyl aldehyde o-HOC_6H_4CHO.	122.05	−7	196.5	1.167	759
2005	$C_7H_6O_2$	m-Hydroxybenzaldehyde.	122.05	106.0	240		
2006	$C_7H_6O_2$	p-Hydroxybenzaldehyde	122.05	116.0		1.129¹³⁰	
2007	$C_7H_6O_2$	Benzoic acid $C_6H_5CO_2H$	122.05	121.7	249 2	1.266¹⁵	1160, 1333
2008	$C_7H_6O_2$	Phenyl formate $HCO_2C_6H_5$.	122 05		173	1.088	
2009	$C_7H_6O_2$	Toluquinone $CH_3C_6H_3O_2$	122.05	69			
2010	$C_7H_6O_2S$	Thiosalicylic acid o-$SHC_6H_4CO_2H$	154.11	164			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2011	C₇H₆O₃	2, 3-Dihydroxybenzaldehyde	138 05	108	235		
2012	C₇H₆O₃	3, 4-Dihydroxybenzaldehyde	138 05	154			
2013	C₇H₆O₃	Salicylic acid o-HOC₆H₄CO₂H	138 05	159	s. 76	1.443	1333
2014	C₇H₆O₃	m-Hydroxybenzoic acid	138 05	201 3		1.473⁴	
2015	C₇H₆O₃	p-Hydroxybenzoic acid .	138 05	213		1.468⁴	
2016	C₇H₆O₄	2, 3-Dihydroxybenzoic acid	154 05	204			
2017	C₇H₆O₄	2, 4-Dihydroxybenzoic acid	154 05	206			
2018	C₇H₆O₄	2, 5-Dihydroxybenzoic acid	154 05	200			
2019	C₇H₆O₄	2, 6-Dihydroxybenzoic acid	154 05	167 d.			
2020	C₇H₆O₄	3, 4-Dihydroxybenzoic acid	154 05	199		1 542⁴	
2021	C₇H₆O₄	3, 5-Dihydroxybenzoic acid	154 05	227			
2022	C₇H₆O₅	Pyrogallolcarboxylic acid	170 05	200 d.			
2023	C₇H₆O₅	Gallic acid 3, 4, 5-(HO)₃C₆H₂CO₂H	170 05	220 d	d	1 694⁴	1333
2024	C₇H₆O₅S	o-Sulfobenzoic acid	202 11	141			
2025	C₇H₆O₅S	m-Sulfobenzoic acid HO₃SC₆H₄CO₂H	202 11	141			
2026	C₇H₆O₅S	p-Sulfobenzoic acid HO₃SC₆H₄CO₂H	202 11	200			
2027	C₇H₆O₆S	Salicylsulfonic acid	218 11	120			
2028	C₇H₇AsCl₂	Benzyl arsine dichloride	236 93		175⁵⁰		
2029	C₇H₇Br	Benzyl bromide	170 97	− 4 0	199	1 438²²	
2030	C₇H₇Br	o-Bromotoluene	170 97	−28 1	181 8	1 422	738
2031	C₇H₇Br	m-Bromotoluene	170 97	−39 8	183 7	1 410	734
2032	C₇H₇Br	p-Bromotoluene	170 97	28	183 6	1 310	732
2033	C₇H₇BrO	5-Bromo-2-hydroxytoluene	186 97	64	235		
2034	C₇H₇BrO	5-Bromo-3-hydroxytoluene	186 97	62			
2035	C₇H₇BrO	3-Bromo-4-hydroxytoluene	186 97		214	1 547²⁴ ᵇ	
2036	C₇H₇BrO₂	6-Bromo-2-methoxyphenol	202 97	63			
2037	C₇H₇BrO₂	4-Bromo-2-methoxyphenol	202 97	46	182⁶⁰		
2038	C₇H₇Cl	Benzyl chloride	126 51	−39	179 4	1 103¹⁸	711
2039	C₇H₇Cl	o-Chlorotoluene .	126 51	−35 1	159 4	1 080	601
2040	C₇H₇Cl	m-Chlorotoluene . . .	126 51	−47 8	162 4	1 072	672
2041	C₇H₇Cl	p-Chlorotoluene	126 51	7 8	162 5	1 071¹⁸	666
2042	C₇H₇ClO	o-Chlorobenzyl alcohol .	142 51	72	230		
2043	C₇H₇ClO	m-Chlorobenzyl alcohol .	142 51		234		
2044	C₇H₇ClO	p-Chlorobenzyl alcohol	142 51	70 5	235		
2045	C₇H₇ClO	3-Chloro-2-hydroxytoluene	142 51	86	225		
2046	C₇H₇ClO	4-Chloro-2-hydroxytoluene	142.51	49	225		
2047	C₇H₇ClO	5-Chloro-2-hydroxytoluene	142 51	49	220		
2048	C₇H₇ClO	4-Chloro-3-hydroxytoluene .	142 51	66	235		
2049	C₇H₇ClO	6-Chloro-3-hydroxytoluene	142 51	53	235		
2050	C₇H₇ClO	2-Chloro-4-hydroxytoluene	142 51		196	1 211²¹	
2051	C₇H₇ClO	3-Chloro-4-hydroxytoluene . . .	142 51	55	228		
2052	C₇H₇ClO₂	4(5)-Chloro-2-methoxyphenol	158 51	< −18	241 5		
2053	C₇H₇ClO₂S	Toluene-o-sulfonchloride	190 58	10	126²¹	1 339	
2054	C₇H₇ClO₂S	Toluene-p-sulfonchloride	190 58	69	146¹⁵		
2055	C₇H₇ClO₃S	2-Chlorotoluene-5-sulfonic acid	206 58	78			
2056	C₇H₇Cl₂NO₂S	Toluene-p-sulfonedichloroamine . . .	240 04	83			
2057	C₇H₇F	o-Fluorotoluene	110 05	< −80	114	1 001	505
2058	C₇H₇F	m-Fluorotoluene .	110 05	−110 8	116	0 999	500
2059	C₇H₇F	p-Fluorotoluene . .	110 05		117	1 001¹⁵ ¹	502
2060	C₇H₇I	Benzyl iodide .	217 99	24 1	d.	1 733²⁵	
2061	C₇H₇I	o-Iodotoluene . .	217 99		211	1 697	785
2062	C₇H₇I	m-Iodotoluene . . .	217 99		204	1 698	
2063	C₇H₇I	p-Iodotoluene .	217 99	35	211 5		
2064	C₇H₇IO	o-Iodoanisol o-CH₃OC₆H₄I	233 99		240	1.800	
2065	C₇H₇IO₂	5-Iodo-2-methoxyphenol	249 99	88			
2066	C₇H₇IO₂	4-Iodo-2-methoxyphenol	249 99	43	180 d.	1 5	
2067	C₇H₇NO	o-Aminobenzaldehyde	121.06	40			
2068	C₇H₇NO	m-Aminobenzaldehyde.	121 06	71 5			
2069	C₇H₇NO	p-Aminobenzaldehyde	121.06	71			
2070	C₇H₇NO	syn -Benzaldoxime C₆H₅C:NOH	121.06	130			
2071	C₇H₇NO	anti-Benzaldoxime C₆H₅C:NOH ..	121.06	35	153⁴³	1.111	972
2072	C₇H₇NO	Benzamide C₆H₅CONH₂	121.06	130	290	1.341⁴	
2073	C₇H₇NO	Formanilide HCONHC₆H₅	121.06	47.5	271	1.112⁶⁰	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2074	C₇H₇NO₂	Anthranilic acid o-H₂NC₆H₄CO₂H ..	137.06	145			
2075	C₇H₇NO₂	m-Aminobenzoic acid	137.06	174		1.511⁴	
2076	C₇H₇NO₂	p-Aminobenzoic acid . .	137.06	187			
2077	C₇H₇NO₂	Benzohydroxamic acid . ..	137.06	125			
2078	C₇H₇NO₂	o-Hydroxybenzamide	137.06	140	270 d.		
2079	C₇H₇NO₂	m-Hydroxybenzamide	137.06	170.5			
2080	C₇H₇NO₂	p-Hydroxybenzamide	137.06	162			
2081	C₇H₇NO₂	o-Nitrotoluene	137.06	α −10.6; β −4.1	222 3	1.168¹⁵	724
2082	C₇H₇NO₂	m-Nitrotoluene . ..	137.06	15.5	231	1.164¹⁵	729
2083	C₇H₇NO₂	p-Nitrotoluene	137.06	51.3	238	1.098⁶⁰	1096
2084	C₇H₇NO₂	Phenylnitromethane	137.06		227	1.160	702
2085	C₇H₇NO₃	o-Nitrobenzyl alcohol	153.06	74	168²⁰		
2086	C₇H₇NO₃	m-Nitrobenzyl alcohol ...	153.06	27	180³		
2087	C₇H₇NO₃	p-Nitrobenzyl alcohol	153.06	93	185¹²		
2088	C₇H₇NO₃	3-Nitro-o-cresol	153.06	145			
2089	C₇H₇NO₃	4-Nitro-o-cresol	153.06	94.6			
2090	C₇H₇NO₃	5-Nitro-o-cresol	153.06	118			
2091	C₇H₇NO₃	6-Nitro-o-cresol	153.06	69 5			
2093	C₇H₇NO₃	4-Nitro-m-cresol	153.06	129			
2094	C₇H₇NO₃	5-Nitro-m-cresol	153.06	91			
2095	C₇H₇NO₃	6-Nitro-m-cresol	153.06	56			
2096	C₇H₇NO₃	3-Nitro-4-hydroxytoluene .	153.06	36 5	125²¹	1 240¹⁹	1053
2098	C₇H₇NO₂	o-Nitroanisol .	153.06	9.4	277	1.268	749
2099	C₇H₇NO₂	m-Nitroanisol .	153.06	38	258	1 373	
2100	C₇H₇NO₃	p-Nitroanisol	153.06	54	260	1.233	
2101	C₇H₇NO₃	4-Amino-2-hydroxybenzoic acid	153.06	220			
2102	C₇H₇NO₃	5-Amino-2-hydroxybenzoic acid	153.06	280 d.			
2103	C₇H₇NO₄	6-Nitro-2-methoxyphenol	169.06	62			
2104	C₇H₇NO₄	5-Nitro-2-methoxyphenol	169.06	104			
2105	C₇H₇NO₄	3-Nitro-2-methoxyphenol	169.06	103			
2106	C₇H₇NO₄S	o-Sulfoaminobenzoic acid	201.13	167			
2107	C₇H₇NO₄S	m-Sulfoaminobenzoic acid	201.13	238			
2108	C₇H₇NO₄S	p-Sulfoaminobenzoic acid. .	201.13	280 d.			
2109	C₇H₇NO₄S	p-Nitrotoluene-o-sulfonic acid .	217.13	130			
2110	C₇H₇NS	Thiobenzamide C₆H₅CSNH₂ ..	137.13	116			
2111	C₇H₈	Tropylidene	92.062		118	0.888	686
2112	C₇H₈	Toluene	92.062	−95 1	110.5	0 866	579
2114	C₇H₈BrN	4-Bromo-o-toluidine .	185.99	32	257 d.		
2115	C₇H₈BrN	5-Bromo-o-toluidine .	185.99	59 5	240		
2116	C₇H₈BrN	5-Bromo-m-toluidine .	185.99	36	260	1.144¹⁹	
2117	C₇H₈BrN	6-Bromo-m-toluidine . ..	185.99	78.8	240		
2118	C₇H₈BrN	2-Bromo-p-toluidine	185.99	26	257		
2119	C₇H₈BrN	3-Bromo-p-toluidine	185.99	26	240	1.498	
2120	C₇H₈ClN	4-Chloro-o-toluidine.	141.53	22	238.5		
2120 1	C₇H₈ClN	5-Chloro-o-toluidine	141.53	30	239 2		
2121	C₇H₈ClN	6-Chloro-o-toluidine.	141.53		245		
2122	C₇H₈ClN	2-Chloro-m-toluidine	141.53		229		
2123	C₇H₈ClN	4-Chloro-m-toluidine	141.53	30	230		
2124	C₇H₈ClN	5-Chloro-m-toluidine	141.53		243		
2125	C₇H₈ClN	6-Chloro-m-toluidine . .	141.53	83	241		
2126	C₇H₈ClN	2-Chloro-p-toluidine	141.53	26	245		
2127	C₇H₈ClN	3-Chloro-p-toluidine	141.53		219	1.151	
2128	C₇H₈N₂	Benzalhydrazine C₆H₅CH₂NHNH₂ .	120.08	16	140¹⁴		
2129	C₇H₈N₂	Benzamidine C₆H₅C(:NH)NH₂. ...	120.08	80			
2130	C₇H₈N₂O	o-Aminobenzamide..	136.08	108			
2131	C₇H₈N₂O	m-Aminobenzamide..............	136.08	79			
2132	C₇H₈N₂O	p-Aminobenzamide NH₂C₆H₄CONH₂.	136.08	183			
2133	C₇H₈N₂O	Benzoylhydrazine C₆H₅CONHNH₂..	136.08	112			
2134	C₇H₈N₂O	Nitrosomethylaniline............. ..	136.08	15	225 d.	1 121²⁷·⁷	998
2135	C₇H₈N₂O	Phenylurea C₆H₅NHCONH₂........	136.08	147			1330
2136	C₇H₈N₂O₂	o-Nitromethylaniline	152.08	34			
2137	C₇H₈N₂O₂	m-Nitromethylaniline.................	152.08	66			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2138	C₇H₈N₂O₂	p-Nitromethylaniline	152 08	152		1 201¹⁵¹ ⁹	
2139	C₇H₈N₂O₂	3-Nitro-o-toluidine..	152 08	96		1.190¹⁹⁹	
2140	C₇H₈N₂O₂	4-Nitro-o-toluidine	152 08	105		1.365¹⁵	
2141	C₇H₈N₂O₂	5-Nitro-o-toluidine	152 08	127 5		1.366¹⁵	
2142	C₇H₈N₂O₂	6-Nitro-o-toluidine	152 08	91 5		1.378¹⁵	
2143	C₇H₈N₂O₂	2-Nitro-3-aminotoluene	152 08	53			
2144	C₇H₈N₂O₂	4-Nitro-3-aminotoluene	152 08	109			
2145	C₇H₈N₂O₂	5-Nitro-3-aminotoluene	152 08	98 1			
2146	C₇H₈N₂O₂	6-Nitro-3-aminotoluene...	152 08	138			
2147	C₇H₈N₂O₂	2-Nitro-4-aminotoluene..	152 08	77 5			
2148	C₇H₈N₂O₂	3-Nitro-p-toluidine	152 08	117			
2149	C₇H₈N₂O₃	5-Nitro-3-amino-4-hydroxytoluene	168.08	110		1.312¹⁷	
2150	C₇H₈N₂S	Phenylthiourea C₆H₅NHCSNH₂	152.14	154			
2151	C₇H₈N₄O₂	Theophylline...... . . .	180 09	272			
2152	C₇H₈N₄O₂	Paraxanthine......	180 09	299			
2153	C₇H₈N₄O₂	Theobromine......	180 09	337			
2154	C₇H₈N₄O₃	1, 3-Dimethyluric acid	196.09	410 d.			
2155	C₇H₈N₄O₃	1, 7-Dimethyluric acid	196 09	390 d.			
2156	C₇H₈N₄O₃	1, 9-Dimethyluric acid.	196 09	400 d			
2157	C₇H₈N₄O₃	3, 9-Dimethyluric acid	196 09	340 d.			
2158	C₇H₉N₅O₇	Guanidine picrate	288.11	200			
2159	C₇H₈O	Benzyl alcohol C₆H₅CH₂OH....	108 06	− 15 3	205 8	1.046	713
2160	C₇H₈O	o-Cresol	108 06	30 1	190 8	1 051	727
2161	C₇H₈O	m-Cresol	108 06	10	202 8	1.035	714
2162	C₇H₈O	p-Cresol	108.06	34 8	201 1	1 039¹⁵ ⁵	715
2163	C₇H₈O	Phenyl methyl ether (Anisol) .	108 06	−37 3	155 8	0 994	659
2164	C₇H₈O	4, 6-Dihydrobenzaldehyde......	108 06	< −20	171 5 d.	1 020¹⁴ ⁵	
2165	C₇H₈OS	Thioguaiacol CH₃OC₆H₄SH. .	140.13		219		
2166	C₇H₈O₂	o-Hydroxybenzyl alcohol .	124.06	86		1 161	
2167	C₇H₈O₂	m-Hydroxybenzyl alcohol . .	124 06	67	300 d.		
2168	C₇H₈O₂	p-Hydroxybenzyl alcohol	124.06	110			
2169	C₇H₈O₂	2, 4-Dihydroxytoluene .	124 06	104			
2170	C₇H₈O₂	2, 5-Dihydroxytoluene	124.06	125			
2171	C₇H₈O₂	2, 6-Dihydroxytoluene	124 06	66			
2172	C₇H₈O₂	Homocatechol 3, 4-(HO)₂C₆H₃CH₃	124.06	65	252	1.129¹⁴	1103
2173	C₇H₈O₂	Orcinol 3, 5-(HO)₂C₆H₃CH₃ ..	124.06	108	290	1 290⁴	
2174	C₇H₈O₂	Guaiacol o-HOC₆H₄OCH₃	124.06	28	205 1	1 143¹⁵	1179
2175	C₇H₈O₂	Resorcinol methyl ether	124.06	< −17 5	244 3	>1	
2176	C₇H₈O₂	Hydroquinol methyl ether	124.06	53	243		
2176 1	C₇H₈O₂	Dimethyl-γ-pyrone... .	124 06	132		0 9953¹³⁷	
2178	C₇H₈O₂	Furfurylacetone . .	124.06	40	229		
2179	C₇H₈O₃S	Toluene-o-sulfinic acid........... .	156.13	80			
2180	C₇H₈O₃	2, 5-Dimethylfurfurane-3-carboxylic acid (Uvinic acid)....	140.06	135			
2181	C₇H₈O₃S	Toluene-o-sulfonic acid .	172.13		128 8²⁵		
2183	C₇H₈O₃S	Toluene-p-sulfonic acid	172.13	105	140²⁰		
2184	C₇H₈O₄	Iretol 2, 4, 6-(OH)₃C₆H₂OCH₃	156 06	186			
2185	C₇H₈O₄	Hydrochelidonic anhydride......	156 06	69	210		
2186	C₇H₈O₄S	4-Hydroxytoluene-2-sulfonic acid	188 13	188			
2187	C₇H₈O₄S	2-Hydroxytoluene-6-sulfonic acid	188 13	118			
2188	C₇H₈O₆	Cinchonic acid............... .	188 06	169			
2189	C₇H₈S	Benzyl mercaptan C₆H₅CH₂SH	124.13		195	1 058²⁰	
2190	C₇H₈S	o-Thiocresol o-CH₃C₆H₄SH...	124.13	15	194 3		
2191	C₇H₈S	m-Thiocresol m-CH₃C₆H₄SH .	124.13	< −20	195 4	1 052⁴²	
2192	C₇H₈S	p-Thiocresol p-CH₃C₆H₄SH. ..	124.13	43	195		
2193	C₇H₉AsO₃	Benzylarsonic acid C₆H₅CH₂AsO(OH)₂	216.03	167			
2194	C₇H₉ClN₄O₂	Theobromine hydrochloride .	216.56				1333
2195	C₇H₉N	Benzylamine C₆H₅CH₂NH₂	107.08		184	0.980	720
2196	C₇H₉N	2, 4-Lutidine...	107.08		157	0.949⁰	
2197	C₇H₉N	2, 6-Lutidine....	107.08		143	0.942⁰	
2198	C₇H₉N	3, 4-Lutidine.... . . .	107.08		164 5		
2199	C₇H₉N	2-Ethylpyridine	107.08		148.8	0.950	990
2200	C₇H₉N	3-Ethylpyridine.....	107.08		165 3	0.959	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2201	C_7H_9N	4-Ethylpyridine	107.08		166	0.936	
2202	C_7H_9N	α-Lutidine	107.08		156.5	0.947°	
2203	C_7H_9N	Methylaniline $C_6H_5NHCH_3$	107.08	−57.0	195.70	0.986	757
2204	C_7H_9N	o-Toluidine o-$CH_3C_6H_4NH_2$.	107.08	α −24 4; β −16.3	200 7	0.998	758
2205	C_7H_9N	m-Toluidine m-$CH_3C_6H_4NH_2$	107.08	−31 5	203 3	0.989	989
2206	C_7H_9N	p-Toluidine p-$CH_3C_6H_4NH_2$	107.08	43 7	200 5	1.046	1087
2207	C_7H_9NO	o-Aminobenzyl alcohol	123 08	82	280 s. d.		
2208	C_7H_9NO	p-Aminobenzyl alcohol..	123.08	95			
2209	C_7H_9NO	4-Amino-2-hydroxytoluene	123.08	161			
2210	C_7H_9NO	5-Amino-2-hydroxytoluene	123 08	175			
2211	C_7H_9NO	6-Amino-2-hydroxytoluene	123 08	128			
2212	C_7H_9NO	5-Amino-m-cresol	123 08	79	345		
2213	C_7H_9NO	4-Amino-3-hydroxytoluene	123 08	174			
2214	C_7H_9NO	2-Amino-4-hydroxytoluene	123.08	144 5			
2215	C_7H_9NO	3-Amino-4-hydroxytoluene	123 08	135			
2216	C_7H_9NO	o-Anisidine o-$CH_3OC_6H_4NH_2$	123 08	5.2	224	1.108^{26}	
2217	C_7H_9NO	m-Anisidine m-$CH_3OC_6H_4NH_2$	123.08		251		
2218	C_7H_9NO	p-Anisidine p-$CH_3OC_6H_4NH_2$	123 08	57.7	245	1.071^{55}	
2219	C_7H_9NO	Benzylhydroxylamine $C_6H_5CH_2NHOH$	123 08		123^{50}		
2220	C_7H_9NO	Salicylamine o-$OHC_6H_4CH_2NH_2$	123 08	129			
2221	C_7H_9NO	m-Tolylhydroxylamine ..	123 08	68			
2222	C_7H_9NO	p-Tolylhydroxylamine	123 08	94			
2223	C_7H_9NO	4, 6-Dihydrobenzaldoxime	123 08	44			
2224	$C_7H_9NO_2$	6-Amino-2-methoxyphenol	139 08	127			
2225	$C_7H_9NO_2$	Ammonium benzoate $C_6H_5CO_2NH_4$	139 08	198		$1 262^4$	
2226	$C_7H_9NO_2S$	Toluene-o-sulfoneamide.	171.14	156 3			
2227	$C_7H_9NO_2S$	Toluene-m-sulfoneamide	171 14	108			
2228	$C_7H_9NO_2S$	Toluene-p-sulfoneamide	171 14	137.5			
2229	$C_7H_9NO_3$	Ammonium salicylate	155 08				1333
2234.1	$C_7H_9NO_3S$	Ammonium o-sulfobenzoate	219.14	>250		1 524	1200
2235	$C_7H_8N_4O$	1-Phenylsemicarbazide.	151.09	172			
2236	$C_7H_8N_4O$	4-Phenylsemicarbazide..	151.09	122			
2237	C_7H_{10}	2, 3-Dihydrocycloheptene.	94.077		121		
2238	C_7H_{10}	1, 2-Dihydrotoluene	94.077		108		
2239	C_7H_{10}	1, 3-Dihydrotoluene ...	94 077		110.1	0 835	524
2240	C_7H_{10}	2, 4-Dihydrotoluene ..	94 077		106	0 827	498
2241	C_7H_{10}	1, 3, 5-Heptatriene . .	94 077		114	0.764	
2243	$C_7H_{10}ClN$	o-Toluidine hydrochloride	143 54	214 5	242		
2244	$C_7H_{10}ClN$	m-Toluidine hydrochloride	143.54	228	249 8		
2245	$C_7H_{10}ClN$	p-Toluidine hydrochloride	143 54	239	257 5		
2247	$C_7H_{10}N_2$	Methyl-p-phenylenediamine	122 09	35 5	259 5		
2248	$C_7H_{10}N_2$	Benzylhydrazine $C_6H_5CH_2NHNH_2$.	122 09	26	103^{41}		
2249	$C_7H_{10}N_2$	2, 3-Diaminotoluene	122 09	62	255		
2250	$C_7H_{10}N_2$	2, 4-Diaminotoluene	122 09	99	280		
2251	$C_7H_{10}N_2$	2, 5-Diaminotoluene	122 09	64	274		
2252	$C_7H_{10}N_2$	Toluylene-2, 6-diamine	122 09	105			
2253	$C_7H_{10}N_2$	3, 4-Diaminotoluene	122 09	88 5	265		
2254	$C_7H_{10}N_2$	3, 5-Diaminotoluene	122 09		285		
2255	$C_7H_{10}N_2$	1, 1-Methylphenylhydrazine	122 09		227.5	1.040	766
2256	$C_7H_{10}N_2$	o-Tolylhydrazine o-$CH_3C_6H_4NHNH_2$.	122 09	56			
2257	$C_7H_{10}N_2$	m-Tolylhydrazine....	122 09		224		
2258	$C_7H_{10}N_2$	p-Tolylhydrazine p-$CH_3C_6H_4NHNH_2$	122 09	61			
2259	$C_7H_{10}N_2O_3$	5-Ethyl-5-methylbarbituric acid	170 09	212			
2260	$C_7H_{10}N_2O_3$	Trimethylbarbituric acid....	170 09	165			
2260.1	$C_7H_{10}N_4O_6$	Dimethyl ureindihydroxysuccinate.	234 10	203			1204
2260.2	$C_7H_{10}N_2O_7$	Isohydroxydimethylurea....	230 11	180			1212
2261	$C_7H_{10}O$	1, 2, 3, 4-Tetrahydrobenzaldehyde	110.08		212	1.009°	
2262	$C_7H_{10}O_2$	Δ¹-Tetrahydrobenzoic acid. .	126.08			$1.072^{17.1}$	552
2263	$C_7H_{10}O_2$	Diacetylacetone $CO(CH_2COCH_3)_2$.	142 08	49	121^{10}	1.068^{40}_{40}	1090
2264	$C_7H_{10}O_4$	cis-Pentamethylene-1, 2-dicarboxylic acid	158.08	140			
2265	$C_7H_{10}O_4$	Teraconic acid............	158.08	161 d.			
2266	$C_7H_{10}O_4$	Terebic acid................	158.08	175		0.816	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2267	C₇H₁₀O₄	Dimethyl citraconate	158 08		210 5	1.110	922
2268	C₇H₁₀O₅	3-Ketopimelic acid	174 08	143			
2269	C₇H₁₀O₅	Ethyl mesoxalate (HO)₂C(CO₂C₂H₅)₂.	174 08	< −31	220	1 119²⁰	
2270	C₇H₁₀O₅	Quinic lactone.	174 08	187			
2271	C₇H₁₁BrO₄	Diethyl bromomalonate. .	239 00		235	1 426¹⁸	
2272	C₇H₁₁NO	Nortropinone.	125 09	70			
2273	C₇H₁₁NO₂	Arecaidine.	141 09	224 d.			
2274	C₇H₁₁NO₂	Arecaine.	141 09	214 d.			
2275	C₇H₁₂	n-Amylacetylene C₅H₁₁C⋮CH	96 092	> −70	110 5	0.738¹¹·⁴	160
2276	C₇H₁₂	2, 4-Dimethyl-1, 3-pentadiene	96 092		93 3	0.749¹⁷	815
2277	C₇H₁₂	2, 4-Dimethyl-2, 3-pentadiene	96 092		70		
2278	C₇H₁₂	3-Heptine C₃H₇C⋮CC₂H₅. . .	96 092		106	0.760⁰	
2279	C₇H₁₂	2, 4-Heptadiene	96 092		107	0.731	896
2280	C₇H₁₂	2-Heptine CH₃C⋮CC₄H₉. .	96 092		113 3	0 763⁰	
2281	C₇H₁₂	4-Methylcyclohexene. . .	96 092		102 2	0 800	385
2282	C₇H₁₂	Δ¹-Tetrahydrotoluene.	96 092		111	0.809	431
2283	C₇H₁₂	Δ²-Tetrahydrotoluene	96 092		105	0.805	408
2284	C₇H₁₂	Δ³-Tetrahydrotoluene	96 092		103	0 799	394
2284.1	C₇H₁₂Cl₂O₂	Isobutyl 1, 2-dichloropropionate	199 01.			1 156²¹	
2285	C₇H₁₂N₂O	Sinapoline.	140 11	100			
2286	C₇H₁₂N₄O	Caffeidine.	168 12	94			
2287	C₇H₁₂N₄O₂	Caffoline.	200 12	197			
2288	C₇H₁₂O	Diallyl carbinol (CH₂:CHCH₂)₂CHOH.	112 09		151	0.857	
2289	C₇H₁₂O	Hexahydrobenzaldehyde	112 09		161	0.926	
2289 1	C₇H₁₂O	o-Methylcyclohexanone	112 09		167²⁴⁰	0.930¹⁵·¹	842
2289 2	C₇H₁₂O	m-Methylcyclohexanone	112 09		60¹⁵	0.914²⁵·¹	1027
2289.3	C₇H₁₂O	p-Methylcyclohexanone	112 09		56 4¹⁰·⁵	0 912²⁴·⁴	1021
2290	C₇H₁₂O	Suberone <(CH₂CH₂CH₂)₂> CO	112 09		179 5	0 969⁰	
2291	C₇H₁₂O₂	Pimelic aldehyde OCH(CH₂)₅CHO	128 09		112¹²		
2292	C₇H₁₂O₂	Teracrylic acid	128 09	< −18	218		
2293	C₇H₁₂O₂	Hexahydrobenzoic acid .	128 09	31	233	1 048	1040
2294	C₇H₁₂O₂	1, 2-Isoheptenic acid . . .	128 09	16 5	227	0 942	442
2295	C₇H₁₂O₂	Allyl butyrate C₃H₇CO₂CH₂CH CH₂	128 09		143		
2296	C₇H₁₂O₂	Allyl isobutyrate . . .	128 09		133 5		
2297	C₇H₁₂O₂	Cyclohexyl formate HCO₂C₆H₁₁ -	128 09	<0	162 5	1 010⁰	
2298	C₇H₁₂O₂	Ethyl angelate.	128 09		142	0 918	963
2299	C₇H₁₂O₂	Ethyl tiglate CH₃CH.C(CH₃)CO₂C₂H₅	128 09		152	0 924	904
2300	C₇H₁₂O₃	Hexahydrosalicylic acid	144 09	111			
2301	C₇H₁₂O₃	Ethyl levulinate.	144 09		205 3	1 017¹⁶	263
2302	C₇H₁₂O₃	Ethyl methylacetoacetate	144 09		186 8	1 019	239
2303	C₇H₁₂O₃	Methyl dimethylacetoacetate	144 09		174	0 999²³	
2304	C₇H₁₂O₄	Butylmalonic acid C₄H₉CH(CO₂H)₂	160 09	101 5	150 d.		
2305	C₇H₁₂O₄	Isobutylmalonic acid.	160.09	107			
2306	C₇H₁₂O₄	sec.-Butylmalonic acid	160 09	76			
2307	C₇H₁₂O₄	Diethylmalonic acid (C₂H₅)₂C(CO₂H)₂	160.09	121			
2308	C₇H₁₂O₄	n-Pimelic acid HO₂C(CH₂)₅CO₂H	160 09	103	272¹⁰⁰		
2308.1	C₇H₁₂O₄	Trimethylsuccinic acid	160.09	152		1 242	
2309	C₇H₁₂O₄	Diethyl malonate CH₂(CO₂C₂H₅)₂	160 09	−49 9	198 9	1 054	208
2310	C₇H₁₂O₄	Dimethyl pyrotartrate	160 09		198	1 078	
2311	C₇H₁₂O₄	Methyl ethyl succinate	160 09	< −20	208 2	1.093⁰	
2312	C₇H₁₂O₅	Glycerol diacetate (Diacetin)	176.09		176⁴⁰	1 178¹⁵	
2313	C₇H₁₂O₆	Quinic acid.	192 09	163	d.	1.637	1333
2314	C₇H₁₂O₆	Diethyl mesoxalate .	192.09	57	200		
2315	C₇H₁₃BrN₂O₂	Adalin CH₂BrCONHCON(C₂H₅)₂	237 03	116			
2316	C₇H₁₃BrO₂	Ethyl 1-bromo-n-valerate	209 02		192	1 226¹⁴	
2317	C₇H₁₃BrO₂	Ethyl 1-bromoisovalerate	209.02		186	1 278¹²	
2318	C₇H₁₃ClO₂	Amyl chloroacetate ClCH₂CO₂C₅H₁₁.	164 56		192	1.055	345
2319	C₇H₁₃ClO₂	Isoamyl chloroacetate	164.56		192	1 041²⁵	
2320	C₇H₁₃N	Heptylnitrile C₆H₁₃CN	111 11		183	0.815	240
2321	C₇H₁₃NO	Nortropanol.	127.11	161			
2322	C₇H₁₃NO	Suberoxime (CH₂CH₂CH₂)₂C.NOH.	127.11	23	230	1.023	
2323	C₇H₁₃NO₂	Stachydrine.	143.11	210			
2324	C₇H₁₃NO₄	Quinic amide (OH)₄C₆H₇CONH₂.	191.11	132			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2325	C_7H_{14}	2, 4-Dimethyl-2-pentene.....	98.108		84	0.699²⁵	
2326	C_7H_{14}	3-Ethyl-2-pentene $(C_2H_5)_2C:CHCH_3$	98.108		98	0.725¹⁵	192
2327	C_7H_{14}	Heptamethylene (Cycloheptane) ..	98.108	−12	118.1	0.811	405
2328	C_7H_{14}	Hexahydrotoluene .. .	98 108	−147 5	103	0 764	910
2329	C_7H_{14}	2-Heptene $CH_3CH:CHC_4H_9$.	98 108		98.5		
2330	C_7H_{14}	Methylcyclohexane. . .	98 108	−126 4	100.8	0.764	272
2331	C_7H_{14}	3-Methyl-2(3)-hexene ..	98 108		97.4	0 718	186
2332	C_7H_{14}	1-Heptene $C_5H_{11}CH.CH_2$.	98 108		99		
2333	C_7H_{14}	2, 2, 3-Trimethyl-1-butene	98 108		80		
2334	C_7H_{14}	2, 3-Dimethyl-2-pentene .	98 108		95.1	0.719	
2335	$C_7H_{14}O$	Cycloheptanol....	114 11		185.2	0.958	
2336	$C_7H_{14}O$	2-Heptene-4-ol ..	114 11		63¹¹	0.842¹⁴ ⁴	838
2337	$C_7H_{14}O$	Hexahydrobenzyl alcohol	114 11		181 2	0 916	816
2338	$C_7H_{14}O$	1-Methylcyclohexane-1-ol	114 11	26	168.3	0 919²⁵	1029
2339	$C_7H_{14}O$	o-Hexahydrocresol	114 11		169	0.923	478
2340	$C_7H_{14}O$	m-Hexahydrocresol	114 11	−47	176	0 914	466
2341	$C_7H_{14}O$	dl-m-Hexahydrocresol	114 11		175	0.923	467
2342	$C_7H_{14}O$	p-Hexahydrocresol	114 11		174	0 924¹⁴	833
2343	$C_7H_{14}O$	Heptaldehyde $C_6H_{13}CHO$	114.11	−45 0	155	0.850	202
2344	$C_7H_{14}O$	Dipropyl ketone $(C_3H_7)_2CO$	114 11	−32 6	143.5	0.821¹⁵	173
2345	$C_7H_{14}O$	Diisopropyl ketone $[(CH_3)_2CH]_2CO$	114 11		123.7	0.806	
2346	$C_7H_{14}O$	Ethyl n-butyl ketone $C_2H_5COC_4H_9$	114 11		148.5		
2347	$C_7H_{14}O$	Ethyl isobutyl ketone .	114 11		136	0 815	
2348	$C_7H_{14}O$	Methyl n-amyl ketone $CH_3COC_5H_{11}$	114 11		150	0.822¹⁶	
2349	$C_7H_{14}O$	Methyl isoamyl ketone .	114 11		144	0.821¹⁷	
2350	$C_7H_{14}O_2$	Isoamylacetic acid .	130 11		216 5	0.926¹⁵	
2351	$C_7H_{14}O_2$	Heptylic acid $C_6H_{13}CO_2H$	130 11	−10	223 5	0.922	269
2353	$C_7H_{14}O_2$	n-Amyl acetate $CH_3CO_2C_5H_{11}$	130 11		147 6	0 879²⁰₂₀	130
2354	$C_7H_{14}O_2$	Isoamyl acetate	130.11		142 5	0.875	122
2354 1	$C_7H_{14}O_2$	d-β-Amyl acetate	130 11		131	0 868	100
2355	$C_7H_{14}O_2$	tert.-Amyl acetate.	130 11		124 8	0.874¹⁹	
2356	$C_7H_{14}O_2$	Ethyl n-valerate $C_4H_9CO_2C_2H_5$ ..	130 11		145 5	0.877	1109
2357	$C_7H_{14}O_2$	Ethyl isovalerate .	130 11	−99 3	135	0 866	126
2358	$C_7H_{14}O_2$	n-Hexyl formate $HCO_2C_6H_{13}$	130 11		153 6	0.898⁰	
2359	$C_7H_{14}O_2$	Isobutyl propionate .	130.11	−71 4	138	0 869	108
2359 1	$C_7H_{14}O_2$	d-sec.-Butyl propionate.	130.11		132	0 8657	
2360	$C_7H_{14}O_2$	Methyl n-caproate $C_5H_{11}CO_2CH_3$	130 11		149.5	0 904⁰₀	
2361	$C_7H_{14}O_2$	Propyl n-butyrate $C_3H_7CO_2C_3H_7$	130.11	−95 2	143	0.879¹⁵	123
2362	$C_7H_{14}O_2$	Propyl isobutyrate $(CH_3)_2CHCO_2C_3H_7$	130 11		135 4	0.884⁰₂	97
2363	$C_7H_{14}O_2$	Isopropyl butyrate $C_3H_7CO_2CH(CH_3)_2$	130.11		128	0 865¹³	
2364	$C_7H_{14}O_2$	Isopropyl isobutyrate	130 11		120 8	0.869²₀	
2365	$C_7H_{14}O_3$	Di-n-propyl carbonate $CO(OC_3H_7)_2$	146 11		168 2	0 968²²	
2366	$C_7H_{14}O_3$	Ethyl butyl carbonate .	146 11		169		
2367	$C_7H_{14}O_4$	Glycerol 1-butyrate	162 11		271		
2367.1	$C_7H_{14}O_5$	l-Methyl rhamnoside	178.11	109			1227
2368	$C_7H_{14}O_6$	α-Methyl galactoside	194 11	112			
2369	$C_7H_{14}O_6$	β-Methyl galactoside	194 11	176			
2370	$C_7H_{14}O_6$	α-Methyl glucose	194.11	161			
2371	$C_7H_{14}O_6$	β-Methyl glucose	194.11	135			
2372	$C_7H_{14}O_6$	α-Methyl glucoside	194 11	168	200⁰ ¹		1230
2373	$C_7H_{14}O_6$	β-Methyl glucoside	194 11	104			1171
2373 1	$C_7H_{14}O_6$	α-Methyl mannoside	194 11	194			1217
2374	$C_7H_{14}O_6$	d-Inosite methyl ether (β-Pinite). . .	194.11	187		1 52	
2375	$C_7H_{14}O_6$	l-Inosite methyl ether (Quebrachite)	194 11	191	210ᵛᵃᶜ	1 54	
2376	$C_7H_{14}O_7$	d, β-Galaheptose . . .	210.11	199			
2377	$C_7H_{14}O_7$	d, α-Glucoheptose .	210.11	215 d.			
2378	$C_7H_{14}O_8$	d-Mannoheptonic acid .	226.11	175 d.			
2379	$C_7H_{14}S$	m-Hexahydrothiocresol .	130.17		174		
2380	$C_7H_{15}Br$	n-Heptyl bromide $C_7H_{15}Br$.	179.03		178 8	1 133¹⁶	
2381	$C_7H_{15}Cl$	n-Heptyl chloride $C_7H_{15}Cl$ . ..	134.57		159 5	0 881¹⁶	
2382	$C_7H_{15}F$	n-Heptyl fluoride $C_7H_{15}F$	118.12	−73	119.2	0.804	61
2383	$C_7H_{15}I$	n-Heptyl iodide $C_7H_{15}I$..	226.05		203 8	1.401⁰	469
2384	$C_7H_{15}N$	Ethylpiperidine	113.12		128	0.857²⁵₄	1000

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2385	C₇H₁₅NO	n-Heptylamide C₆H₁₃CONH₂	129 12	96			
2386	C₇H₁₅NO	Heptaldoxime C₆H₁₃CH:NOH..	129 12	55 5	195	0.834¹⁴	1124
2386 1	C₇H₁₅NO₂	Isobutylurethane C₄H₉NHCO₂C₂H₅	145 12	< −65	96¹⁷	0.943	311
2387	C₇H₁₆	2, 4-Dimethylpentane CH₃[CH(CH₃)₂]₂.	100 12		83 9	0.681	45
2388	C₇H₁₆	3, 3-Dimethylpentane	100 12		87	0 711⁰	
2389	C₇H₁₆	n-Heptane CH₃(CH₂)₅CH₃	100 12	−90 0	98 4	0 684	55
2390	C₇H₁₆	2-Methylhexane (CH₃)₂CHC₄H₉	100 12		90 4	0 707⁰₀	
2391	C₇H₁₆	d, 3-Methylhexane C₂H₇CH(CH₃)C₂H₅	100 12		92	0 687	
2392	C₇H₁₆	3-Ethylpentane (C₂H₅)₃CH	100 12		93 8	0 670	89
2393	C₇H₁₆	2, 2, 3-Trimethylbutane	100 12	− 25	80 8	0 695¹³	77
2394	C₇H₁₆	2, 2-Dimethylpentane (CH₃)₃CC₄H₇	100 12		78 6	0.674	
2396	C₇H₁₆O	Dimethylbutyl carbinol	116 12		142 2	0 816	224
2397	C₇H₁₆O	Dimethylisobutyl carbinol ..	116 12		130	0.816	228
2398	C₇H₁₆O	Dimethyl-tert.-butyl carbinol.	116 12	17	132		
2399	C₇H₁₆O	Dipropyl carbinol (C₃H₇)₂CHOH	116 12		155 1	0 820	256
2400	C₇H₁₆O	Diisopropyl carbinol	116 12		140	0 820	265
2400 1	C₇H₁₆O	d-Ethylbutyl carbinol	116 12		66¹⁵	0.823	251
2401	C₇H₁₆O	Ethylisobutyl carbinol.	116 12		148 2		
2402	C₇H₁₆O	Ethyl-sec.-butyl carbinol	116 12		150	0 852⁰	
2403	C₇H₁₆O	n-Heptyl alcohol C₇H₁₅OH	116 12	− 34 6	175 8	0 817¹¹	287
2404	C₇H₁₆O	2-Hydroxy-3-ethylpentane	116 12		152	0.853⁰	
2405	C₇H₁₆O	1-Hydroxy-2-methylhexane	116.12		162 5	0 831¹³	266
2406	C₇H₁₆O	Isoheptyl alcohol.. . . .	116.12		167 2	0 831⁰	291
2407	C₇H₁₆O	Methyl-n-amyl carbinol . .	116 12		158	0 819	259
2407 1	C₇H₁₆O	d-Methylamyl carbinol . .	116 12		73 5²⁰	0 819	253
2408	C₇H₁₆O	Methylisoamyl carbinol	116 12		150	0 819¹⁷ ⁸	
2409	C₇H₁₆O	Methylethylpropyl carbinol	116 12		141	0 823	270
2410	C₇H₁₆O	Methylethylisopropyl carbinol	116 12		140	0 833	
2411	C₇H₁₆O	Propylisopropyl carbinol ..	116 12		141	0.821¹⁷	215
2412	C₇H₁₆O	Triethyl carbinol (C₂H₅)₃COH	116 12		142	0.840	334
2413	C₇H₁₆O	Ethyl isoamyl ether..........	116 12		112	0 764¹⁴	
2414	C₇H₁₆O	Propyl butyl ether C₄H₉OC₃H₇ .	116.12		117 1	0.777⁰	
2415	C₇H₁₆O₂	Ethyl orthoformate HC(OC₂H₅)₃.	148 12	−76.1	145 9	0 897	
2416	C₇H₁₆O₄S₂	Sulfonal (CH₃)₂C(SO₂C₂H₅)₂. . . .	228 25	128	300 d.		
2417	C₇H₁₆O₇	d-Mannoheptitol....... .	212.12	188			
2418	C₇H₁₆O₇	Volemitol.........	212.12	155			
2419	C₇H₁₇N	n-Heptylamine C₇H₁₅NH₂	115.14	− 23 0	155 1	0.777	278
2420	C₈Cl₄O₃	Tetrachloro-o-phthalic anhydride	285.83	257			
2421	C₈H₂Cl₂O₃	3, 6-Dichloro-o-phthalic anhydride	216.93	191	339		
2422	C₈H₂Cl₄O₄	Tetrachloro-o-phthalic acid .	303 85	250			
2422 1	C₈H₄BrNO₂	m-Bromoisatine	225 96	255			
2422.2	C₈H₄ClNO	Isatine chloride.	165 50	180 d.			
2423	C₈H₄Cl₂O₂	o-Phthalyl dichloride o-C₆H₄(COCl)₂...	202 95	0	276 7	1 408	755
2424	C₈H₄Cl₂O₂	Isophthalyl dichloride m-C₆H₄(COCl)₂.	202 95	41	276		
2425	C₈H₄Cl₂O₂	Terephthalyl dichloride p-C₆H₄(COCl)₂.	202 95	78	259		
2426	C₈H₄Cl₂O₄	3, 6-Dichloro-o-phthalic acid . .	234 95	185			
2427	C₈H₄Cl₄O	Trichloromethyl p-chlorophenylketone	257.86	28	181⁴¹		
2428	C₈H₄N₂	Isophthalic nitrile m-C₆H₄(CN)₂	128.05	161			
2429	C₈H₄N₂	Terephthalic nitrile p-C₆H₄(CN)₂	128.05	222			
2430	C₈H₄N₂O₄	Nitroisatine............	192.05	230			
2431	C₈H₄O₃	o-Phthalic anhydride..... .	148.03	130 8	284 5	1.527⁴	
2432	C₈H₅Cl₃O	Dichloromethyl p-chlorophenyl ketone	223.41	51	178⁴⁵		
2433	C₈H₅Cl₄NO	2, 3, 4, 6-Tetrachloroacetanilide	272.88	181			
2434	C₈H₅NO	Benzoyl cyanide C₆H₅COCN	131.05	34	208		
2435	C₈H₅NO₂	o-Cyanobenzoic acid .	147.05	190			
2436	C₈H₅NO₂	m-Cyanobenzoic acid . .	147.05	217			
2437	C₈H₅NO₂	p-Cyanobenzoic acid	147.05	214			
2438	C₈H₅NO₂	Isatine............... .	147.05	201			
2439	C₈H₅NO₂	o-Phthalimide o-C₆H₄(CO)₂NH	147.05	238			
2440	C₈H₅NO₆	3-Nitro-o-phthalic acid	211.05	220			
2441	C₈H₅NO₆	4-Nitro-o-phthalic acid	211.05	164			
2442	C₈H₅NO₆	2-Nitroisophthalic acid	211 05	300			
2443	C₈H₅NO₆	4-Nitroisophthalic acid	211.05	245			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2444	C₈H₅NO₆	5-Nitroisophthalic acid	211.05	255			
2445	C₈H₅NO₆	2-Nitroterephthalic acid	211.05	270			
2446	C₈H₅NO₆	Pyridine-2, 3, 4-tricarboxylic acid	211.05	250 d.			
2447	C₈H₅NO₆	Pyridine 2, 3, 5-tricarboxylic acid.. ..	211.05	323			
2448	C₈H₅NO₆	Pyridine-2, 3, 6-tricarboxylic acid. ...	211.05	100			
2449	C₈H₅NO₆	Pyridine-2, 4, 5-tricarboxylic acid.....	211.05	235			
2450	C₈H₅NO₆	Pyridine-2, 4, 6-tricarboxylic acid.	211.05	227			
2451	C₈H₅NO₆	Pyridine-3, 4, 5-tricarboxylic acid	211.05	261			
2452	C₈H₅N₃O₇	Picryl acetate.	271.06	76	120 d.		
2453	C₈H₆	Phenylacetylene C₆H₅C⋮CH	102.05		143	0.930	820
2454	C₈H₆BrN	Bromobenzyl cyanide C₆H₅CHBrCN.	195.97	> −17	231 7	1.519	1185
2455	C₈H₆Br₂	Styrene-1, 2-dibromide .	261.88	73 5	134¹⁵		
2456	C₈H₆Br₂O	p-Bromophenacyl bromide	277.88	109 7			
2457	C₈H₆Cl₂O₂	Piperonal chloride	204.96		240 s. d.		
2458	C₈H₆Cl₃NO	2, 3, 4-Trichloroacetanilide	238.43	122			
2459	C₈H₆Cl₃NO	2, 4, 5-Trichloroacetanilide	238.43	190			
2460	C₈H₆Cl₃NO	2, 4, 6-Trichloroacetanilide	238.43	204			
2461	C₈H₆I₂O₃	Methyl 3, 5-diiodosalicylate	403.91	110 5			
2462	C₈H₆N₂	Phthalazine	130.06	91	317		
2463	C₈H₆N₂	Quinazoline	130.06	48	243		
2464	C₈H₆N₂	Quinoxaline .	130.06	30 5	226	1 133⁴⁸	1075
2465	C₈H₆N₂O₂	Isatoxime (Nitrosooxindol) .	162.06	202			
2466	C₈H₆N₂O₂	p-Nitrobenzyl cyanide	162.06	117			
2467	C₈H₆N₄O₄	Alloxantin	286.08	170 d.			
2468	C₈H₆O	Coumarone	118.05	> −18	175	1.091	997
2469	C₈H₆O₂	Phenylglyoxal C₆H₅CO.CHO . . .	134.05	73	142¹²⁵		
2470	C₈H₆O₂	o-Phthalic aldehyde o-C₆H₄(CHO)₂	134.05	56			
2471	C₈H₆O₂	Isophthalic aldehyde m-C₆H₄(CHO)₂	134.05	89 5			
2472	C₈H₆O₂	Terephthalic aldehyde p-C₆H₄(CHO)₂	134.05	116	248		
2473	C₈H₆O₂	Phthalide	134.05	73; 65	290		
2474	C₈H₆O₃	Piperonal (Heliotropin) .	150.05	37	263		
2475	C₈H₆O₃	o-Aldehydobenzoic acid	150.05	100 5		1 404	
2476	C₈H₆O₃	m-Aldehydobenzoic acid	150.05	175			
2477	C₈H₆O₃	p-Aldehydobenzoic acid . .	150.05	250			
2478	C₈H₆O₃	Phenylglyoxylic acid .	150.05	66	148⁴		
2479	C₈H₆O₄	o-Phthalic acid o-C₆H₄(CO₂H)₂	166.05	191 d.		1 593	
2480	C₈H₆O₄	Isophthalic acid m-C₆H₄(CO₂H)₂ ..	166.05	330			
2482	C₈H₆O₄	Piperonylic acid CH₂:O₂:C₆H₃.CO₂H	166.05	228			
2483	C₈H₆O₅	2-Hydroxy-o-phthalic acid .	182.05	244			
2485	C₈H₆O₅	4-Hydroxy-o-phthalic acid .	182.05	181 d.			
2486	C₈H₆O₅	2-Hydroxyisophthalic acid	182.05	239			
2487	C₈H₆O₅	4-Hydroxyisophthalic acid	182.05	306			
2488	C₈H₆O₅	5-Hydroxyisophthalic acid	182.05	288			
2489	C₈H₆O₅	Noropianic acid	182.05	171			
2490	C₈H₆S	Thionaphthene .	134.11	32	221	1.165	1049
2491	C₈H₇Br	α-Bromostyrene C₆H₅CBr:CH₂	182.97	−43 5	160⁷⁵	1.4057	770
2492	C₈H₇Br	ω-Bromostyrene (isomer 1)	182.97	7	221	1.4224	786
2493	C₈H₇Br	ω-Bromostyrene (isomer 2) .	182.97	−7 5	108²⁶	1.427	992
2493 1	C₈H₇BrN₂O₃	α-Bromonitroacetanilide	258.99	131		1 765	
2494	C₈H₇BrO	ω-Bromoacetophenone	198.97	50	119	1.647	
2495	C₈H₇Cl	α-Chlorostyrene C₆H₅C.Cl CH₂ . .	138.51		199		
2496	C₈H₇Cl	ω-Chlorostyrene C₆H₅CH.CHCl ..	138.51		198 8	1.112²³	
2497	C₈H₇ClO	ω-Chloroacetophenone...	154.51	59	247	1.324¹⁵	
2498	C₈H₇ClO	p-Chloroacetophenone. . . .	154.51	20	232	1.188	
2499	C₈H₇ClO	Phenylacetyl chloride C₆H₅CH₂COCl	154.51		102.5¹⁷	1.168	
2500	C₈H₇ClO₂	p-Anisyl chloride p-CH₃OC₆H₄COCl.	170.51	27			
2501	C₈H₇ClO₂	Phenyl chloroacetate ClCH₂CO₂C₆H₅	170.51	45	235		
2502	C₈H₇F₂NO	2, 5-Difluoroacetanilide	171.06	122.5			
2503	C₈H₇N	Benzyl cyanide C₆H₅CH₂CN	117.06	−23.8	233.9	1.015¹⁸	679
2504	C₈H₇N	Indole	117.06	52.5	254		1333
2505	C₈H₇N	o-Tolunitrile o-CH₃C₆H₄CN..	117.06		204	0.995¹⁵	1004
2506	C₈H₇N	m-Tolunitrile m-CH₃C₆H₄CN..	117.06		214	0.984¹⁵	
2507	C₈H₇N	p-Tolunitrile p-CH₃C₆H₄CN.	117.06	29.5	217		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2508	C₈H₇NO	p-Anisonitrile p-CH₃OC₆H₄CN.	133 06	60	256		
2509	C₈H₇NO	dl-Mandelonitrile C₆H₅CH(OH)CN.	133 06	−10	d.	1 124	
2510	C₈H₇NO	Indoxyl......................	133 06	85	110		
2511	C₈H₇NO	Oxindol...................... .	133 06	120			
2512	C₈H₇NO₂	Hydrindic acid (Dioxindol) .	149 06	180	195 d.		
2513	C₈H₇NO₂	o-Nitrostyrene o-NO₂.C₆H₄.CH:CH₂	149 06	13 5			
2514	C₈H₇NO₂	m-Nitrostyrene m-NO₂.C₆H₄.CH:CH₂	149 06	−5			
2515	C₈H₇NO₂	p-Nitrostyrene p-NO₂.C₆H₄.CH:CH₂	149 06	29			
2516	C₈H₇NO₃	Oxanilic acid CO₂H.CONHC₆H₅	165 06	150			
2517	C₈H₇NO₃	o-Phthalamic acid .	165 06	149	155 d.		
2518	C₈H₇NO₄	Methyl o-nitrobenzoate	181 06	−8	269	1 284¹⁵	
2519	C₈H₇NO₄	Methyl m-nitrobenzoate	181 06	70	279		
2520	C₈H₇NO₄	Methyl p-nitrobenzoate	181 06	96			
2521	C₈H₇NO₄	Uvitonic acid .. .	181 06	274			
2522	C₈H₇NS	Benzyl isothiocyanate	149 13		243		
2522 1	C₈H₇NS	Benzyl thiocyanate	149 13	41	235		
2523	C₈H₇NS	o-Tolyl isothiocyanate	149 13		239	1 104¹⁵	
2524	C₈H₇NS	m-Tolyl isothiocyanate	149 13		245		
2525	C₈H₇NS	p-Tolyl isothiocyanate	149 13	26	237	1 087¹⁵	
2526	C₈H₇N₃O₅	2, 3-Dinitroacetanilide	225 08	186			
2527	C₈H₇N₃O₅	2, 4-Dinitroacetanilide	225 08	120			
2528	C₈H₇N₃O₅	2, 6-Dinitroacetanilide	225 08	197			
2529	C₈H₇N₃O₅	3, 4-Dinitroacetanilide	225 08	144			
2530	C₈H₇N₃O₅	3, 6-Dinitroacetanilide	225 08	121			
2531	C₈H₇N₃O₆	3, 4, 5-Trinitro-o-xylene	241 08	115			
2532	C₈H₇N₃O₆	3, 4, 6-Trinitro-o-xylene	241 08	72			
2533	C₈H₇N₃O₆	2, 4, 5-Trinitro-m-xylene	241 08	90			
2534	C₈H₇N₃O₆	2, 4, 6-Trinitro-m-xylene	241 08	181 5			
2535	C₈H₇N₃O₆	4, 5, 6-Trinitro-m-xylene	241 08	125			
2536	C₈H₇N₃O₆	2, 3, 6-Trinitro-p-xylene	241 08	140²⁰			
2537	C₈H₇N₃O₇	Ethyl picrate	257 08	78 5			
2538	C₈H₈	Styrene (Phenylethylene)	104 06		146	0 903	907
2539	C₈H₈BrNO	o-Bromoacetanilide	213 99	99			
2540	C₈H₈BrNO	p-Bromoacetanilide..	213 99	165			
2540 1	C₈H₈Br₂	o-Xylenedibromide o-C₆H₄(CH₂Br)₂	263 89	94.5	d.	1.988	
2540 2	C₈H₈Br₂	m-Xylenedibromide m-C₆H₄(CH₂Br)₂	263 89	77	140	1 959	
2541	C₈H₈Br₂	p-Xylenedibromide p-C₆H₄(CH₂Br)₂	263 89	144	245	2 102⁰	
2542	C₈H₈ClNO	o-Chloroacetanilide..............	169 53	88			
2543	C₈H₈ClNO	m-Chloroacetanilide . . .	169 53	72 5			
2544	C₈H₈ClNO	p-Chloroacetanilide .	169 53	172 5			
2544.1	C₈H₈Cl₂	o-Xylenedichloride o-C₆H₄(CH₂Cl)₂	174 98	55	241	1.393	
2544 2	C₈H₈Cl₂	m-Xylenedichloride m-C₆H₄(CH₂Cl)₂	174 98	34 2	255	1 302	
2545	C₈H₈Cl₂	p-Xylenedichloride p-C₆H₄(CH₂Cl)₂	174 98	100 5	120²⁰	1 417⁰	
2546	C₈H₈INO	p-Iodoacetanilide p-CH₃CONHC₆H₄I	261 00	184			
2547	C₈H₈N₂	Apoharmine................ ...	132 08	183			
2548	C₈H₈N₂	1-Methylindazole............	132 08		107¹⁵	1 032¹⁹ ²	1129
2549	C₈H₈N₂OS	Benzoylthiourea C₆H₅CONHCSNH₂	180 14	169			
2550	C₈H₈N₂O₂	Benzoylurea C₆H₅CONHCONH₂	164.08	200			
2551	C₈H₈N₂O₂	o-Phthalic diamide o-C₆H₄(CONH₂)₂	164 08	220			
2552	C₈H₈N₂O₂	Isophthalic diamide m-C₆H₄(CONH₂)₂	164 08	265			
2553	C₈H₈N₂O₂	N-Nitrosoacetanilide	164 08	41			
2554	C₈H₈N₂O₂	Ricinine	164 08	201			
2555	C₈H₈N₂O₃	o-Nitroacetanilide...	180 08	93			
2556	C₈H₈N₂O₃	m-Nitroacetanilide..	180 08	150.5			
2557	C₈H₈N₂O₃	p-Nitroacetanilide.	180 08	214			
2558	C₈H₈N₂O₄	3, 4-Dinitro-o-xylene.	196 08	82			
2559	C₈H₈N₂O₄	3, 6-Dinitro-o-xylene	196 08	56			
2560	C₈H₈N₂O₄	4, 5-Dinitro-o-xylene	196 08	115			
2561	C₈H₈N₂O₄	4, 6-Dinitro-o-xylene	196 08	75			
2562	C₈H₈N₂O₄	2, 5-Dinitro-m-xylene	196.08	101			
2563	C₈H₈N₂O₄	4, 5-Dinitro-m-xylene	196 08	132			
2564	C₈H₈N₂O₄	2, 3-Dinitro-p-xylene	196 08	93			
2565	C₈H₈N₂O₄	2, 5-Dinitro-p-xylene.................	196.08	147			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2566	$C_8H_8N_2O_4$	2, 6-Dinitro-p-xylene	196.08	124			
2566 1	$C_8H_8N_2O_4$	4, 5-Dinitro-1, 2-dimethoxybenzene	228.08	130.5		1.326[131]	
2566 2	$C_8H_8N_4O$	4-Methoxyphenyltetrazole	128 09	228			1306
2567	C_8H_8O	Phenylacetaldehyde $C_6H_5CH_2CHO$	120 06		194	1 027	
2568	C_8H_8O	o-Toluic aldehyde $o-CH_3C_6H_4CHO$	120 06		195 5	1 039	960
2569	C_8H_8O	m-Toluic aldehyde $m-CH_3C_6H_4CHO$	120 06		195.5	1.019	971
2570	C_8H_8O	p-Toluic aldehyde $p-CH_3C_6H_4CHO$	120.06		204	1 020	814; 906
2571	C_8H_8O	Acetophenone $CH_3COC_6H_5$	120 06	19 7	202.3	1 026	705
2572	C_8H_8O	Coumarane	120 06		189 5	1 074	
2573	$C_8H_8O_2$	Phenacyl alcohol $C_6H_5COCH_2OH$	136 06	86		1 013	
2574	$C_8H_8O_2$	5-Hydroxytoluene-2-aldehyde	136.06	108 9			
2575	$C_8H_8O_2$	4-Hydroxytoluene-3-aldehyde	136 06	55 1	21 8		
2576	$C_8H_8O_2$	6-Hydroxytoluene-3-aldehyde	136 06	117 4			
2577	$C_8H_8O_2$	3-Hydroxytoluene-4-aldehyde	136 06	54	223		
2578	$C_8H_8O_2$	o-Methoxybenzaldehyde	136 06	35	242	1 133	745
2579	$C_8H_8O_2$	m-Methoxybenzaldehyde	136 06		230	1 118	836
2580	$C_8H_8O_2$	p-Methoxybenzaldehyde	136 06	2.5	247	1 123	821
2581	$C_8H_8O_2$	o-Hydroxyacetophenone	136 06		213		
2582	$C_8H_8O_2$	m-Hydroxyacetophenone	136 06	95			
2583	$C_8H_8O_2$	p-Hydroxyacetophenone	136 06	109			
2584	$C_8H_8O_2$	Phenylacetic acid $C_6H_5CH_2CO_2H$	136 06	76 7	265 5	1.078[53]	
2585	$C_8H_8O_2$	o-Toluic acid $o-CH_3C_6H_4CO_2H$	136.06	102 4	259 2	1 062[114] *	1157
2586	$C_8H_8O_2$	m-Toluic acid $m-CH_3C_6H_4CO_2H$	136.06	110 5	263	1.054[111] *	640
2587	$C_8H_8O_2$	p-Toluic acid $p-CH_3C_6H_4CO_2H$	136 06	176.8	275		
2588	$C_8H_8O_2$	Benzyl formate $HCO_2CH_2C_6H_5$	136 06		203 4	1.081	
2589	$C_8H_8O_2$	Methyl benzoate $C_6H_5CO_2CH_3$	136.06	−12.5	199 6	1.094	656
2590	$C_8H_8O_2$	Phenyl acetate $CH_3CO_2C_6H_5$	136 06		195 5	1.078	610
2591	$C_8H_8O_2$	o-Xyloquinone 1, 2-$(CH_3)_2C_6H_2O_2$-3, 6	136 06	55			
2592	$C_8H_8O_2$	m-Xyloquinone 1, 3-$(CH_3)_2C_6H_2O_2$-2, 5	136.06	73			
2593	$C_8H_8O_2$	p-Xyloquinone 1, 4-$(CH_3)_2.C_6H_2O_2$-2, 5	136.06	125			
2594	$C_8H_8O_3$	Piperonyl alcohol	152 06	51			
2595	$C_8H_8O_3$	Isovanillin 4, 3-$CH_3OC_6H_3(OH)CHO$	152.06	116		1.196	
2596	$C_8H_8O_3$	Vanillin 3, 4-$CH_3OC_6H_3(OH)CHO$	152.06	81	285		
2597	$C_8H_8O_3$	o-Hydroxymethylbenzoic acid	152.06	120			
2598	$C_8H_8O_3$	m-Hydroxymethylbenzoic acid	152.06	111	190[11]		
2599	$C_8H_8O_3$	p-Hydroxymethylbenzoic acid	152 06	181			
2600	$C_8H_8O_3$	o-Hydroxyphenylacetic acid	152.06	137			
2601	$C_8H_8O_3$	m-Hydroxyphenylacetic acid	152.06	129			
2602	$C_8H_8O_3$	p-Hydroxyphenylacetic acid	152.06	148			
2603	$C_8H_8O_3$	3-Hydroxytoluene-2-carboxylic acid	152.06	167			
2604	$C_8H_8O_3$	4-Hydroxytoluene-2-carboxylic acid	152.06	172.4			
2605	$C_8H_8O_3$	5-Hydroxytoluene-2-carboxylic acid	152 06	178			
2606	$C_8H_8O_3$	6-Hydroxytoluene-2-carboxylic acid	152.06	183			
2607	$C_8H_8O_3$	4-Hydroxytoluene-3-carboxylic acid	152.06	152 5			
2608	$C_8H_8O_3$	5-Hydroxytoluene-3-carboxylic acid	152.06	208			
2609	$C_8H_8O_3$	6-Hydroxytoluene-3-carboxylic acid	152.06	172			
2610	$C_8H_8O_3$	2-Hydroxytoluene-4-carboxylic acid	152.06	207			
2611	$C_8H_8O_3$	3-Hydroxytoluene-4-carboxylic acid	152.06	177 8			
2612	$C_8H_8O_3$	d(l)-Mandelic acid $C_6H_5CH(OH)CO_2H$	152.06	133			
2613	$C_8H_8O_3$	dl-Mandelic acid $C_6H_5CH(OH)CO_2H$	152.06	118		1.361[4]	
2614	$C_8H_8O_3$	o-Methoxybenzoic acid	152 06	98	200		
2615	$C_8H_8O_3$	m-Methoxybenzoic acid	152.06	100			
2616	$C_8H_8O_3$	p-Methyoxybenzoic acid	152.06	184.2	280	1 385[4]	1333
2617	$C_8H_8O_3$	Phenoxyacetic acid $C_6H_5OCH_2CO_2H$	152 06	99	285 s. d.		
2618	$C_8H_8O_3$	Methyl salicylate $HOC_6H_4CO_2CH_3$	152.06	−8.6	223 3	1.184	708
2619	$C_8H_8O_3$	Resorcinol acetate	152 06		283		
2620	$C_8H_8O_4$	Phloroacetophenone	168.06	285			
2621	$C_8H_8O_4$	Berberonic acid 2, 4, 5-$C_5H_2N(CO_2H)_3$	168.06	165			
2622	$C_8H_8O_4$	Dehydracetic acid	168.06	109	270		
2623	$C_8H_8O_4$	Δ[1, 4]-Dihydro-o-phthalic acid	168.06	153			
2624	$C_8H_8O_4$	Δ[2, 4]-Dihydro-o-phthalic acid	168.06	215			
2625	$C_8H_8O_4$	Δ[2, 4]-Dihydro-o-phthalic acid	168.06	215			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2626	C₈H₈O₄	Homogentisinic acid	168 06	147			
2627	C₈H₈O₄	Isovanillic acid	168 06	250			
2628	C₈H₈O₄	Vanillic acid	168.06	207			
2630	C₈H₈O₅	Methyl gallate..	184.06	192 d.			
2631	C₈H₈O₈	Tetramethylene-1, 1, 2, 2-tetracarboxylic acid.....	232 06	203			
2632	C₈H₉Br	o-Xylyl bromide	184 99	21	217 7	1 381¹¹	
2633	C₈H₉Br	4-Bromo-o-xylene	184 99	0 2	214 5	1 369	740
2634	C₈H₉Br	m-Xylyl bromide	184 99		215 8 s. d	1.371¹¹	
2635	C₈H₉Br	2-Bromo-m-xylene	184 99	> −10	206		
2636	C₈H₉Br	4-Bromo-m-xylene	184 99		207		
2637	C₈H₉Br	5-Bromo-m-xylene	184 99	> −20	204	1 362	
2638	C₈H₉Br	p-Xylyl bromide	184 99	38	220 7	1 324	
2639	C₈H₉Br	2-Bromo-p-xylene	184 99	10	205 7	1 356	735
2640	C₈H₉Cl	o-Xylyl chloride..	140 53		199		
2641	C₈H₉Cl	3-Chloro-o-xylene	140 53	> −20	189 5		
2642	C₈H₉Cl	4-Chloro-o-xylene.	140 53	> −20	191 5	1 0692¹⁵	
2643	C₈H₉Cl	m-Xylyl chloride .	140 53		196		
2644	C₈H₉Cl	p-Xylyl chloride...	140 53		202		
2645	C₈H₉N	2-Allylpyridine...	119 08		190	0.959⁰	
2646	C₈H₉NO	o-Aminoacetophenone	135 08		252 s. d.		
2647	C₈H₉NO	m-Aminoacetophenone	135 08	98 5	290		
2648	C₈H₉NO	p-Aminoacetophenone	135 08	106	295		
2649	C₈H₉NO	Acetanilide (Antifebrin)	135 08	114.2	303 8	1.21⁴	
2650	C₈H₉NO	Acetophenoneoxime CH₃C(:NOH)C₆H₅	135 08	58			
2651	C₈H₉NO	Phenylacetamide C₆H₅CH₂CONH₂	135 08	155	284		
2652	C₈H₉NO	o-Toluic amide o-CH₃C₆H₄CONH₂	135 08	138			
2653	C₈H₉NO	m-Toluic amide m-CH₃C₆H₄CONH₂.	135 08	97			
2654	C₈H₉NO	p-Toluic amide p-CH₃C₆H₄CONH₂.	135 08	159			
2655	C₈H₉NO₂	o-Acetoaminophenol	151.08	203			
2656	C₈H₉NO₂	m-Acetoaminophenol	151 08	149			
2657	C₈H₉NO₂	p-Acetoaminophenol	151 08	168			
2658	C₈H₉NO₂	dl-Aminophenylacetic acid. .	151.08	256	265		
2659	C₈H₉NO₂	Homoanthranilic acid	151 08	177 d.			
2660	C₈H₉NO₂	N-Methylanthranilic acid	151 08	179			
2661	C₈H₉NO₂	dl-Phenylaminoacetic acid	151.08	127			
2662	C₈H₉NO₂	Benzyl carbamate C₆H₅CH₂CO₂NH₂ .	151.08	86			
2663	C₈H₉NO₂	Ethyl nicotinate ...	151 08		105⁵		
2664	C₈H₉NO₂	Methyl o-aminobenzoate	151 08	8.2; 24 3	135.5¹⁵	1.168¹⁵	
2665	C₈H₉NO₂	Methyl p-aminobenzoate	151.08	112			
2666	C₈H₉NO₂	3-Nitro-o-xylene	151.08		250 8	1.147¹⁵	
2667	C₈H₉NO₂	4-Nitro-o-xylene.	151.08	30	258	1.139³⁰	
2668	C₈H₉NO₂	2-Nitro-m-xylene	151.08		225 5	1.112¹⁵	
2669	C₈H₉NO₂	4-Nitro-m-xylene	151 08	2	246	1.126¹⁷·⁵	
2670	C₈H₉NO₂	5-Nitro-m-xylene.	151 08	71	273 7		
2671	C₈H₉NO₂	2-Nitro-p-xylene.	151.08		239 9	1.132¹⁵	
2672	C₈H₉NO₂	α-Anisaldoxime CH₃OC₆H₄CH:NOH	151.08	64			
2673	C₈H₉NO₂	β-Anisaldoxime CH₃OC₆H₄CH·NOH	151.08	133			
2674	C₈H₉NO₂	o-Methoxybenzamide	151.08	129			
2675	C₈H₉NO₂	p-Methoxybenzamide	151.08	162.3			
2676	C₈H₉NO₃	3-Nitro-4-methoxytoluene.	167.08	8.5	274 d.		
2677	C₈H₉NO₃	o-Nitrophenetol o-C₂H₅OC₆H₄NO₂	167.08		268	1.190¹⁵	718
2678	C₈H₉NO₃	p-Nitrophenetol p-C₂H₅OC₆H₄NO₂	167.08	60	283		
2679	C₈H₉NO₃	Methyl 3-hydroxy-4-aminobenzoate.	167.08	120			
2680	C₈H₉NO₃	Methyl 3-amino-4-hydroxybenzoate.	167.08	143			
2681	C₈H₉NO₄	Biliverdic acid......	183.08	114			
2682	C₈H₉NS	Thioacetanilide CH₃CSNHC₆H₅ .	151.14	75	d.		
2682.1	C₈H₉N₃O₄	2, 4-Dinitrodimethylaniline	221.09	87		1.476	577
2683	C₈H₁₀	Ethylbenzene C₆H₅CH₂CH₃	106.08	−92.8	136.5⁷⁶·⁷	0.868	577
2684	C₈H₁₀	o-Xylene o-C₆H₄(CH₃)₂	106.08	−27.1	144	0.879	626
2685	C₈H₁₀	m-Xylene m-C₆H₄(CH₃)₂	106.08	−53 6	139.0	0.865	584
2686	C₈H₁₀	p-Xylene p-C₆H₄(CH₃)₂ .	106.08	13.2	137.7	0.861	573
2687	C₈H₁₀ClN	o-Chlorodimethylaniline.	155.54		208.5	1.107	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2688	C₈H₁₀ClN	p-Chlorodimethylaniline.. .	155 54	35.5	231		
2689	C₈H₁₀N₂O	N-Acetyl-o-phenylenediamine	150.09	144.8			
2690	C₈H₁₀N₂O	N-Acetyl-m-phenylenediamine	150.09	279			
2691	C₈H₁₀N₂O	N-Acetyl-p-phenylenediamine	150.09	160.5			
2692	C₈H₁₀N₂O	Benzylurea C₆H₅CH₂NHCONH₂	150.09	147.5			
2693	C₈H₁₀N₂O	Hydracetine CH₃COHN.NHC₆H₅	150.09	128			
2694	C₈H₁₀N₂O	1-Methyl-1-phenylurea	150 09	82			
2695	C₈H₁₀N₂O	p-Nitrosodimethylaniline. . .	150 09	85			
2696	C₈H₁₀N₂O₂	o-Nitrodimethylaniline	166 09		154²⁴	1.179	
2697	C₈H₁₀N₂O₂	m-Nitrodimethylaniline	166 09	66	285	1 313¹⁷	
2698	C₈H₁₀N₂O₂	p-Nitrodimethylaniline..	166 09	163			
2699	C₈H₁₀N₂O₃	3-Amino-4-methoxy-6-nitrotoluene	182 09	131 5			
2700	C₈H₁₀N₂S	Benzylthiourea C₆H₅CH₂NHCSNH₂.	166 16	162			
2701	C₈H₁₀N₄O₂	Caffeine (Theine)	194 11	237		1.23	
2702	C₈H₁₀N₄O₃	1, 3, 9-Trimethyluric acid	210 11	320 d.			
2703	C₈H₁₀N₄O₃	1, 7, 9-Trimethyluric acid	210.11	340			
2704	C₈H₁₀N₄O₃	2, 7, 9-Trimethyluric acid	210 11	380			
2705	C₈H₁₀O	2, 3-Dimethylphenol	122 08	75	218		
2706	C₈H₁₀O	2, 4-Dimethylphenol . .	122 08	26	211 5	1.036	
2707	C₈H₁₀O	2, 6-Dimethylphenol	122 08	49	212		
2708	C₈H₁₀O	3, 4-Dimethylphenol	122 08	65	225 1		
2709	C₈H₁₀O	3, 5-Dimethylphenol	122 08	68	219 5		
2710	C₈H₁₀O	o-Ethylphenol	122 08	> -18	207 5	1.037⁰	
2711	C₈H₁₀O	m-Ethylphenol	122 08	-4	214	1.025⁰	
2712	C₈H₁₀O	p-Ethylphenol	122 08	46	219		
2713	C₈H₁₀O	Methylphenyl carbinol	122 08		205	1.003²⁵	
2713.1	C₈H₁₀O	d-Methylphenyl carbinol	122 08		100¹⁸	1.014	668
2714	C₈H₁₀O	2-Phenylethyl alcohol C₆H₅CH₂CH₂OH	122 08		221	1 024¹⁵	677
2715	C₈H₁₀O	o-Tolyl carbinol o-CH₃C₆H₄CH₂OH	122 08	34	223.3	1.023⁴⁰	
2716	C₈H₁₀O	m-Tolyl carbinol m-CH₃C₆H₄CH₂OH.	122.08	> -20	217	1.036⁰	
2717	C₈H₁₀O	p-Tolyl carbinol p-CH₃C₆H₄CH₂OH.	122.08	59.5	217		
2718	C₈H₁₀O	Benzyl methyl ether C₆H₅CH₂OCH₃..	122 08		174	0 987²⁰	
2719	C₈H₁₀O	o-Cresyl methyl ether o-CH₃C₆H₄OCH₃	122.08		171 3	0.981	619
2720	C₈H₁₀O	m-Cresyl methyl ether...	122 08		177 2	0 978¹³	627
2721	C₈H₁₀O	p-Cresyl methyl ether	122 08		176 5	0.970	646
2722	C₈H₁₀O	Phenetol C₆H₅OC₂H₅	122.08	-30 2	172	0.965	633
2723	C₈H₁₀O₂	Anis alcohol p-CH₃OC₆H₄CH₂OH	138.08	45	258.8	1.109²⁶	
2724	C₈H₁₀O₂	Caffeol	138 08		197		
2725	C₈H₁₀O₂	Creosol 3, 4-CH₃O(OH)C₆H₃CH₃	138.08	5 5	221.8	1 092	709
2726	C₈H₁₀O₂	3, 5-Dimethyl-o-dihydroxybenzene	138 08	74			
2727	C₈H₁₀O₂	4, 5-Dimethyl-o-dihydroxybenzene	138.08	82			
2728	C₈H₁₀O₂	2, 4-Dimethylresorcinol.....	138.08	150			
2729	C₈H₁₀O₂	2, 5-Dimethylresorcinol	138.08	163	280		
2730	C₈H₁₀O₂	4, 5-Dimethylresorcinol	138.08	137			
2731	C₈H₁₀O₂	4, 6-Dimethylresorcinol	138 08	125	279		
2732	C₈H₁₀O₂	2, 3-Dimethylhydroquinone	138.08	221 s. d.			
2733	C₈H₁₀O₂	2, 5-Dimethylhydroquinone . .	138 08	213			
2734	C₈H₁₀O₂	2, 6-Dimethylhydroquinone	138 08	151			
2735	C₈H₁₀O₂	p-Homosaligenin...	138 08	105			
2736	C₈H₁₀O₂	Styrolene alcohol HOCH₂CH₂OC₆H₅	138 08	68	274 2		
2737	C₈H₁₀O₂	o-Dimethoxybenzene o-C₆H₄(OCH₃)₂.	138 08	22 5	206	1 086¹⁸	
2738	C₈H₁₀O₂	o-Ethoxyphenol o-HOC₆H₄OC₂H₅	138 08	28	241		
2739	C₈H₁₀O₂	Hydroquinone dimethyl ether .	138.08	56	212.6	1 053¹¹	
2740	C₈H₁₀O₂	Hydroquinone monoethyl ether ...	138 08	66	247		
2741	C₈H₁₀O₂	Resorcinol dimethyl ether	138 08	-55 3	215	1 080⁰	
2742	C₈H₁₀O₂	Resorcinol monoethyl ether .	138 08		247		
2743	C₈H₁₀O₂S	Ethylphenylsulfone C₂H₅SO₂C₆H₅ .	170.14	42	>300	1.010²²	
2744	C₈H₁₀O₃	3-Methoxy-4-hydroxybenzyl alcohol	154.08	115	d.		
2745	C₈H₁₀O₃	Crotonic anhydride	154.08		247.8	1.040	520
2746	C₈H₁₀O₄	Δ¹-Tetrahydro-o-phthalic acid	170.08	120			
2747	C₈H₁₀O₄	Δ¹-Tetrahydro-o-phthalic acid.	170.08	215			
2748	C₈H₁₀O₄	Diallyl oxalate C₂O₄(C₃H₅)₂.	170 08		217	1.055	
2749	C₈H₁₀O₄	Dimethyl muconate (CH:CH.CO₂CH₃)₂	170.08	75 u.; 156 st.			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2750	C₈H₁₀O₄	Succinic peroxide	234 08	127 d.			
2751	C₈H₁₁BrN₄O₂	Caffeine hydrobromide..	275 03				1333
2752	C₈H₁₁ClN₂O	p-Nitrosodimethylaniline hydrochloride	186 56	177			
2753	C₈H₁₁ClN₄O₂	Caffeine hydrochloride.	230 58				1333
2753.1	C₈H₁₁ClO₄	Ethyl chloromaleate....	206 54		125 5¹⁰	1 191¹⁰	
2754	C₈H₁₁Cl₃O₆	α-Chloralose.	309 46	230			
2755	C₈H₁₁I₃N₄O₂	Caffeine triiodide...	575 91	171			
2756	C₈H₁₁N	Dimethylaniline C₆H₅N(CH₃)₂	121 09	1 67	193 50	0 956	771
2757	C₈H₁₁N	2, 3-Dimethylaniline	121 09	> −15	223 8	0 992	756
2758	C₈H₁₁N	2, 4-Dimethylaniline	121 09		216	0 974	744
2759	C₈H₁₁N	2, 5-Dimethylaniline	121 09	15 5	217	0 980¹⁵	965
2760	C₈H₁₁N	2, 6-Dimethylaniline	121 09		216 9	0 979	748
2761	C₈H₁₁N	3, 4-Dimethylaniline	121 09	49	226	1.076	
2762	C₈H₁₁N	3, 5-Dimethylaniline . .	121 09		221	0.972	742
2763	C₈H₁₁N	N-Ethylaniline C₆H₅NH.C₂H₅	121 09	−63 5	204 72	0 963	739
2764	C₈H₁₁N	o-Ethylaniline o-C₂H₅C₆H₄NH₂	121 09		216	0 983¹¹	
2765	C₈H₁₁N	m-Ethylaniline m-C₂H₅C₆H₄NH₂	121 09		215	0 990⁰	
2766	C₈H₁₁N	p-Ethylaniline p-C₂H₅C₆H₄NH₂	121 09	−5	216 5	0 975¹¹	
2767	C₈H₁₁N	Methyl-o-toluidine CH₃C₆H₄NCH₃	121 09		207	0.977	750
2768	C₈H₁₁N	Methyl-m-toluidine..	121 09		206		
2769	C₈H₁₁N	Methyl-p-toluidine p-CH₃C₆H₄NHCH₃	121 09		206		
2770	C₈H₁₁N	α-Phenylethylamine C₆H₅CH(NH₂)CH₃	121 09		187 4	0 940¹⁵	
2771	C₈H₁₁N	ω-Phenylethylamine C₆H₅CH₂CH₂NH₂	121 09		198 2	0 958¹⁵ ⁴	761
2772	C₈H₁₁N	2-Isopropylpyridine	121 09		159	0 934⁰	
2773	C₈H₁₁N	4-Isopropylpyridine ..	121 09		178	0 944⁰	
2774	C₈H₁₁N	2-Methyl-5-ethylpyridine	121 09		174	0 918¹¹	
2775	C₈H₁₁N	Nicotoine	121 09		208	0 955	643
2776	C₈H₁₁N	2-Propylpyridine (Conyrine)	121 09		165		
2777	C₈H₁₁N	2, 3, 4-Trimethylpyridine	121 09		188	0 913	
2778	C₈H₁₁N	2, 4, 5-Trimethylpyridine	121 09		168	0 906	
2779	C₈H₁₁N	2, 4, 6-Trimethylpyridine	121 09		172	0 917¹⁵	
2780	C₈H₁₁NO	Hydroxyethylaniline	137.09		286	1 110⁰	
2781	C₈H₁₁NO	o-Dimethylaminophenol	137 09	45	200		
2782	C₈H₁₁NO	o-Ethylaminophenol o-HOC₆H₄NHC₂H₅	139.09	107 5			
2783	C₈H₁₁NO	m-Ethylaminophenol	137 09	62	176¹¹		
2784	C₈H₁₁NO	3-Amino-2-methoxytoluene	137 09		223		
2785	C₈H₁₁NO	5-Amino-2-methoxytoluene	137.09	53			
2786	C₈H₁₁NO	o-Phenetidine o-NH₂C₆H₄OC₂H₅	137 09	> −21	229 2		
2787	C₈H₁₁NO	m-Phenetidine m-NH₂C₆H₄OC₂H₅	137.09		248		
2788	C₈H₁₁NO	p-Phenetidine p-NH₂C₆H₄OC₂H₅	137.09	2.4	254.2	1 061	
2789	C₈H₁₁NO	Dimethylaniline oxide C₆H₅N(CH₃)₂O	137 09	153			
2790	C₈H₁₁NO	Tyramine p-HOC₆H₄CH₂CH₂NH₂	137 09	161			
2791	C₈H₁₁NO₃S	m-Dimethylanilinesulfonic acid	201 16	266 d.			
2792	C₈H₁₁NO₃S	p-Dimethylanilinesulfonic acid	201 16	257			
2793	C₈H₁₁NO₃S	m-Ethylaniline sulfonic acid	201 16	294 d.			
2794	C₈H₁₁N₃O	Maretin m-CH₃.C₆H₄NH.NHCONH₂	165.11	184			
2795	C₈H₁₂	Dihydro-o-xylene	108.09		135		
2796	C₈H₁₂	Δ¹·⁵-5-Dihydro-m-xylene.	108 09		130	0 823	497
2797	C₈H₁₂	Δ¹·²-3-Dihydro-p-xylene.	108 09		135 6	0.830	529
2798	C₈H₁₂ClN	ω-Phenylethylamine hydrochloride	157 56	217			
2799	C₈H₁₂N₂	Dimethylketine . .	136.11	86	189		
2800	C₈H₁₂N₂	1, 1-Dimethyl-m-phenylenediamine	136.11		258	0 995¹⁵	
2801	C₈H₁₂N₂	1, 1-Dimethyl-p-phenylenediamine.	136.11	41	262.3	1.036	
2802	C₈H₁₂N₂	2, 6-Dimethylphenylhydrazine	136.11	46			
2803	C₈H₁₂N₂	1-Ethyl-1-phenylhydrazine	136.11		237	1.018¹⁵	
2804	C₈H₁₂N₂	1-Ethyl-2-phenylhydrazine	136 11		240		
2805	C₈H₁₂N₂O₂	Phenylhydrazine acetate	168 11	69			
2806	C₈H₁₂N₂O₃	n-Butylbarbituric acid...... .	184.11	215			
2807	C₈H₁₂N₂O₃	1, 3-Diethylbarbituric acid. ..	184.11	52	167¹⁰		
2808	C₈H₁₂N₂O₃	5, 5-Diethylbarbituric acid .	184 11	191			
2808.1	C₈H₁₂N₂O₄	Tetraacetylhydrazine [(CH₃CO)₂N]₂.	200.11	86			1203
2809	C₈H₁₂O	Amylpropiolic aldehyde.. .	124.09		187	0.89⁰	
2810	C₈H₁₂O₂	Ethyl sorbate CH₃(CH:CH)₂CO₂C₂H₅.	140.09		76.5¹²	0.936	608

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2811	$C_9H_{12}O_4$	Terpenylic acid	172 09	89			
2812	$C_9H_{12}O_4$	Diethyl fumarate ($.CHCO_2C_2H_5)_2$	172 09	0 6	218.5	1.052	377
2813	$C_9H_{12}O_4$	Diethyl maleate ($CHCO_2C_2H_5)_2$	172 09		225	1.067	375
2814	$C_9H_{12}O_4$	Ethyl diacetoacetate	172.09		211 s. d.	1.09	492
2815	$C_9H_{12}O_4$	Dimeric diacetyl	172.09	58		$1.560^{19.8}$	
2816	$C_9H_{12}O_5$	Ethyl oxalacetate	188 09		132^{24}	1.172	
2816 1	$C_9H_{13}BrO_4$	Diethyl bromoisosuccinate	253 02		122^{13}	1 3183^{24}	905
2817	$C_9H_{13}N$	Granatic acid	123 11	270			
2818	$C_9H_{13}N$	Tropidine	123.11		163	0.946	946
2819	$C_9H_{13}NO$	Tropinone	139.11	41	218 5	$0.987^{99.6}$	1141
2820	$C_9H_{13}NO_2$	Arecolidine	155.11	110			
2821	$C_9H_{13}NO_2$	Arecoline	155.11		220		
2822	$C_9H_{13}NO_2$	Scopoline	155 11	110	243	1.016^{104}	
2823	$C_9H_{13}N_2O_2$	Iminodiethylbarbituric acid	183 12	295			
2824	C_8H_{14}	n-Hexylacetylene $C_6H_{13}C:CH$	110 11		125	0.770^0	818
2825	C_8H_{14}	d-Laurolene	110 11		120.5	0 797	397
2826	C_8H_{14}	Methyl-n-amylacetylene	110.11		134		
2827	C_8H_{14}	1, 2, 3, 4-Tetrahydro-m-xylene	110 11		124	0 801	398
2828	$C_8H_{14}BrNO_2$	Arecoline hydrobromide	236 03	168			
2829	$C_8H_{14}ClNO_2$	Arecolidine hydrochloride	191 57	98	250 d.		
2830	$C_8H_{14}O$	1, 1-Dimethylcyclohexene-3-ol	126 11		75^{16}	0 933	926
2831	$C_8H_{14}O$	2, 2-Dimethylcyclohexanone	126 11		172 5	0.913	426
2832	$C_8H_{14}O$	2, 6-Dimethylcyclohexanone	126 11		55 3^{10}	0 914	813
2833	$C_8H_{14}O$	Crotonyl ether ($CH_3CH.CHCH_2)_2O$.	126 11		145	0.890^0	
2834	$C_8H_{14}O$	2-Methyl-2-heptene-6-one	126 11	-67 3	174	0 860	
2835	$C_8H_{14}O$	Homomesityl oxide	126 11		160^{625}	0 863	406
2836	$C_8H_{14}O_2$	Allyl isovalerate $C_4H_9CO_2C_3H_5$	142.11		155		
2837	$C_8H_{14}O_2$	Cyclohexyl acetate $CH_3CO_2C_6H_{11}$	142 11		177		
2838	$C_8H_{14}O_2$	Methyl hexahydrobenzoate	142 11		183	0 995^{15}	
2839	$C_8H_{14}O_3$	Dialdan	158 11	130			
2840	$C_8H_{14}O_3$	n-Butyric anhydride ($C_4H_9CO)_2O$	158 11	-75 0	198 2	0 969	
2841	$C_8H_{14}O_3$	Isobutyric anhydride $[(CH_3)_2CHCO]_2O$	158 11	-53.5	182.5	0 950	
2842	$C_8H_{14}O_3$	1-Ethyl-3-acetylbutyric acid	158 11		158^9		
2843	$C_8H_{14}O_4$	n-Amylmalonic acid $C_5H_{11}CH(CO_2H)_2$	174 11	82	140 d.		
2844	$C_8H_{14}O_4$	2, 2'-Dimethyladipic acid	174.11	76	321		
2845	$C_8H_{14}O_4$	Suberic acid $HO_2C(CH_2)_6CO_2H$.	174.11	140	279^{100}		
2846	$C_8H_{14}O_4$	Diethyl methylmalonate	174.11		201 4	1 018	203
2847	$C_8H_{14}O_4$	Diethyl succinate ($CH_2CO_2C_2H_5)_2$	174 11	-20 8	216 5	1.042	246
2848	$C_8H_{14}O_4$	Di-n-propyl oxalate ($CO_2C_3H_7)_2$	174 11		211	1 018^{22}	
2849	$C_8H_{14}O_4$	Ethyl isopropyl malonate	174.11		217 d.	0.987^{25}	
2849 1	$C_8H_{14}O_5$	Diethyl malate	190 11		253	1.128	355
2850	$C_8H_{14}O_6$	Diethyl d-tartrate $[CH(OH)CO_2C_2H_5]_2$.	206.11	17	280	1 202	421
2851	$C_8H_{15}ClO$	Capryl chloride $C_7H_{15}COCl$	162 57		196	0 975^8	
2852	$C_8H_{15}N$	n-Caprylonitrile $C_7H_{15}CN$	125.12		200	0 $820^{14.3}$	
2853	$C_8H_{15}N$	α-Coniceine	125.12		158	0 893^{16}	
2854	$C_8H_{15}N$	β-Coniceine	125.12	-16	169		
2855	$C_8H_{15}N$	γ-Coniceine	125.12	41	172	0 872	945
2856	$C_8H_{15}N$	δ-Coniceine	125 12	> -50	161 5	0 901^{18}	
2857	$C_8H_{15}N$	Granatinine	125 12	60			
2858	$C_8H_{15}N$	Pseudoconiceine	125 12		172	0 878	
2859	$C_8H_{15}N$	Tropane	125 12		167	0 930	975
2860	$C_8H_{15}NO$	Granatoline	141 12	134			
2861	$C_8H_{15}NO$	Hygrine	141 12		195	0.935	
2862	$C_8H_{15}NO$	Pelletierine	141.12		195 d.	0 988^0	
2863	$C_8H_{15}NO$	Pseudotropine	141 12	108	243		
2864	$C_8H_{15}NO$	Tropine	141 12	63	233	1.016^{100}	1146
2865	C_8H_{16}	Cyclooctane ($CH_2)_8$	112.12	14 4	150 6	0 839	
2866	C_8H_{16}	Diisobutylene ($CH_3)_2C:CHC(CH_3)_3$	112 12		102 6	0.715^{15}	
2867	C_8H_{16}	o-Dimethylcyclohexane	112 12	-57 5	129 4	0.779	317
2868	C_8H_{16}	m-Dimethylcyclohexane	112 12	-85	123 7	0 771	288
2869	C_8H_{16}	p-Dimethylcyclohexane	112 12	-86	120 5	0.769	257
2870	C_8H_{16}	Ethylcyclohexane $C_2H_5.C_6H_{11}$	112 12		128		
2871	C_8H_{16}	2-Methyl-3-ethyl-2-pentene	112.12		117.1		

No.	Formula	Name	Mol. wt.	M. P	B. P.	d	R. I. No.
2872	C₈H₁₆	2-Methyl-2-heptene (CH₃)₂C:CHC₄H₉	112 12		125.2	0 816	
2873	C₈H₁₆	4-Methyl-3-heptene	112 12		120 4	0.724	219
2874	C₈H₁₆	n-Octylene CH₃(CH₂)₅CH:CH₂	112 12		123	0 722¹⁷	
2875	C₈H₁₆BrNO	Pelletierine hydrobromide	222 05	140			
2876	C₈H₁₆ClNO	Pelletierine hydrochloride	177 59	145			
2877	C₈H₁₆N₂O₄	Ethylidene diurethane .	204 14	126			
2878	C₈H₁₆O	1, 2-Dimethylcyclohexanol	128 12		166	0 926¹⁴	834
2879	C₈H₁₆O	d-1, 3-Dimethylcyclohexanol	128 12	72	69¹⁴		
2880	C₈H₁₆O	dl-1, 3-Dimethylcyclohexanol	128 12		169	0 911¹⁴	832
2881	C₈H₁₆O	1, 4-Dimethylcyclohexanol	128 12	50	170		
2882	C₈H₁₆O	2, 2-Dimethylcyclohexanol	128 12	8	72 2¹⁰	0 923	496
2883	C₈H₁₆O	2, 4-Dimethylcyclohexanol	128 12		179	0 912	888
2884	C₈H₁₆O	2, 5-Dimethylcyclohexanol	128 12		178 5	0.907	887
2885	C₈H₁₆O	2, 6-Dimethylcyclohexanol	128 12		174 7		
2886	C₈H₁₆O	3, 3-Dimethylcyclohexanol	128 12	11	99 5¹⁰	0 913¹⁴	468
2887	C₈H₁₆O	3, 4-Dimethylcyclohexanol	128 12		189 2	0 907	889
2888	C₈H₁₆O	cis-3, 5-Dimethylcyclohexanol	128 12		185	0 911	447
2889	C₈H₁₆O	trans-3, 5-Dimethylcyclohexanol	128 12		187 5	0 902¹⁶	463
2890	C₈H₁₆O	2-Methyl-2-heptene-6-ol	128 12		176	0 854	434
2891	C₈H₁₆O	Isoamyl allyl ether	128 12		120		
2892	C₈H₁₆O	n-Caprylic aldehyde C₇H₁₅CHO	128 12		81³²	0 821	261
2893	C₈H₁₆O	Ethyl n-amyl ketone C₂H₅COC₅H₁₁	128 12		168	0 850⁰	
2894	C₈H₁₆O	Ethyl isoamyl ketone	128 12		163 5		
2895	C₈H₁₆O	Methylbutyrone .	128 12		180	0 827¹⁶	
2896	C₈H₁₆O	Methyl hexyl ketone CH₃COC₆H₁₃	128 12	− 21 6	172 7	0 818	225
2897	C₈H₁₆O	Methyl isohexyl ketone	128 12		204	0 817	
2898	C₈H₁₆O	Propyl isobutyl ketone	128 12		155	0 813	
2899	C₈H₁₆O₂	n-Caprylic acid CH₃(CH₂)₆CO₂H	144 12	16	237 5	0 910	296
2900	C₈H₁₆O₂	Triethylacetic acid (C₂H₅)₃CCO₂H	144 12	39 5	202		
2901	C₈H₁₆O₂	Isoamyl propionate	144 12		160 2	0 870	163
2901.1	C₈H₁₆O₂	d-β-Amyl propionate	144 12		58¹⁶	0 866	133
2902	C₈H₁₆O₂	tert.-Amyl propionate	144 12		113 5	0 855¹⁵	
2903	C₈H₁₆O₂	Butyl n-butyrate C₃H₇CO₂C₄H₉	144 12		166 1	0 872²⁰	148
2904	C₈H₁₆O₂	Isobutyl n-butyrate	144 12		156 9	0 866¹⁶	140
2905	C₈H₁₆O₂	Isobutyl isobutyrate	144 12	−80 7	148 7	0 875⁰	120
2906	C₈H₁₆O₂	tert.-Butylethyl acetate	144 12		157		
2907	C₈H₁₆O₂	Ethyl n-caproate C₅H₁₁CO₂C₂H₅	144 12		166 6	0 875¹⁵	
2908	C₈H₁₆O₂	Heptyl formate HCO₂(CH₂)₆CH₃	144 12		176 7	0 894⁰	
2909	C₈H₁₆O₂	n-Hexyl acetate CH₃CO₂(CH₂)₅CH₃	144 12		169 2	0 890⁰	
2909.1	C₈H₁₆O₂	d-β-Hexyl acetate....	144 12		57²⁰	0.864	139
2910	C₈H₁₆O₂	Methyl n-heptylate C₆H₁₁CO₂CH₃	144 12		172 1	0 881¹⁵	187
2911	C₈H₁₆O₂	n-Propyl n-valerate C₄H₉CO₂C₃H₇	144 12		167 5	0 889⁰	
2912	C₈H₁₆O₂	n-Propyl isovalerate	144 12		155 9	0 863	141
2913	C₈H₁₆O₃	1-Hydroxy-n-caprylic acid	160 12	69 5			
2914	C₈H₁₆O₃	Amyl l-lactate CH₃CH(OH)CO₂C₅H₁₁	160 12		110 5²¹ ⁶	0 964⁴	
2915	C₈H₁₆O₄	Metaldehyde (C₂H₄O)₄	176 12		150		1172
2916	C₈H₁₆O₄	Paraldol (C₄H₈O₂)₂ ..	176 12	82			
2916.1	C₈H₁₆O₄	Bismethoxyacetal	176 12	127			1238
2917	C₈H₁₆O₆	Dambonite (Inosite dimethyl ether) .	208 12	195	210		
2918	C₈H₁₆O₆	2, 3-Dimethyl-α-glucose	208 12	87			
2919	C₈H₁₆O₆	2, 3-Dimethyl-β-glucose	208 12	110			
2920	C₈H₁₆O₆	d, α-Ethylglucoside	208 12	114			1197
2921	C₈H₁₆O₇	Ethyl d-gluconate.	224 12	65			
2922	C₈H₁₇Br	n-Octyl bromide CH₃(CH₂)₆CH₂Br	193 05		204	1.116¹⁴	
2922.1	C₈H₁₇Br	l-2-Bromooctane	193 05		71¹⁴	1.091¹⁷	
2923	C₈H₁₇BrN₄	Hexamethylenetetramine bromoethylate (Bromalin) .	249.08	200			
2924	C₈H₁₇Cl	n-Octyl chloride CH₃(CH₂)₆CH₂Cl	148 59		184.6	0.879¹⁵	
2925	C₈H₁₇Cl	2-Chlorooctane C₆H₁₃CHClCH₃	148 59		173	0.871¹⁵	
2926	C₈H₁₇F	n-Octyl fluoride CH₃(CH₂)₆CH₂F	132 13		142.5	0 812¹⁴ ¹	94
2927	C₈H₁₇I	n-Octyl iodide CH₃(CH₂)₆CH₂I	240.06	−45 9	225.5	1.341¹⁴ ⁵	549
2928	C₈H₁₇N	d-Coniine	127.14	− 2 5	166.5	0.845	978
2929	C₈H₁₇N	2, 4, 6-Trimethylpiperidine	127.14		147	0.831	954

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2930	$C_8H_{17}NO$	Conhydrine (Hydroxyconiine)... ...	143.14	118	226		1388
2931	$C_8H_{17}NO$	α-Pseudoconhydrine	143.14	106	236.5		
2932	$C_8H_{17}NO_2$	1-Hydroxy-n-caprylic amide ...	159 14	150			
2933	C_8H_{18}	2, 5-Dimethylhexane ...	114 14	−91 0	109.2	0.693	87
2934	C_8H_{18}	2, 3-Dimethylhexane .	114 14		114 0	0.725⅕	178
2935	C_8H_{18}	2, 4-Dimethylhexane .	114 14		109.9	0.708⅕	138
2936	C_8H_{18}	3, 4-Dimethylhexane	114 14		116 5	0.721	156
2937	C_8H_{18}	Isooctane $(CH_3)_2CH(CH_2)_4CH_3$	114 11		116 0	0.704⅕	103
2938	C_8H_{18}	2-Methyl-3-ethylpentane	114 14		114	0.708¹⁴	134
2939	C_8H_{18}	3-Methylheptane $C_2H_5CH(CH_3)C_4H_9$.	114 14		122 2	0.707	
2940	C_8H_{18}	4-Methylheptane $(C_3H_7)_2CHCH_3$	114 14		118 0	0 722	114
2941	C_8H_{18}	n-Octane $CH_3(CH_2)_6CH_3$.	114 14	−56 5	124.6	0 707⅕	112
2942	C_8H_{18}	2-Ethylhexane $CH_3(C_2H_5)CHC_4H_9$	114 14		118 8	0 717⅕	135
2942.1	C_8H_{18}	3-Ethylhexane $(C_2H_5)_2CHC_3H_7$	114 14		115	0 715	
2943	C_8H_{18}	2, 2, 3, 3-Tetramethylbutane	114 14	104	106 8		
2944	C_8H_{18}	2, 2, 3-Trimethylpentane	114 14		110 8	0 722⅕	233
2945	$C_8H_{18}BrN$	d-Coniine hydrobromide	208 06	211			
2946	$C_8H_{18}ClN$	d-Coniine hydrochloride	163 61	217			
2947	$C_8H_{18}ClNO$	Pseudoconhydrine hydrochloride	179 61	213			
2948	$C_8H_{18}IN$	Coniine hydroiodide	255 08	146			
2949	$C_8H_{18}N_2O$	Nitrosodiisobutylamine.	158 16	−5	221	0 893⅕	
2950	$C_8H_{18}N_2O_3$	Coniine nitrate........	190 16	83			
2951	$C_8H_{18}O$	Dibutyl alcohol....... .	130 14		181 2	0 848⁰	
2952	$C_8H_{18}O$	Diethylpropyl carbinol	130 14		160 5	0 838	339
2953	$C_8H_{18}O$	Dimethyl-n-amyl carbinol	130 14		162	0.879	322
2954	$C_8H_{18}O$	Dimethylisoamyl carbinol	130 14		154	0 823	254
2955	$C_8H_{18}O$	Ethylisoamyl carbinol	130.14	−61	166	0 808	247
2956	$C_8H_{18}O$	1-Hydroxy-2, 5-dimethylhexane	130 14		179.5	0 828	
2957	$C_8H_{18}O$	2-Hydroxy-2, 4-dimethylhexane	130.14		151		
2958	$C_8H_{18}O$	4-Hydroxy-3-ethylhexane ...	130 14		164	0.835⁰	
2959	$C_8H_{18}O$	2-Hydroxy-4-methylheptane	130 14		168		
2960	$C_8H_{18}O$	d-6-Hydroxy-3-methylheptane	130 14		169	0 817	
2961	$C_8H_{18}O$	4-Hydroxy-2, 2, 4-trimethylpentane	130 14	−20	147.5	0 842⁰	
2962	$C_8H_{18}O$	Methyl dipropyl carbinol	130 14		161 5	0.823	297
2963	$C_8H_{18}O$	Methylethylbutylcarbinol	130 14		160 6	0 827	298
2964	$C_8H_{18}O$	Methylethylisobutyl carbinol .	130 14		152 4	0 830¹⁴	308
2965	$C_8H_{18}O$	Methylisohexyl carbinol	130.14		172	0 813	274
2966	$C_8H_{18}O$	n-Octyl alcohol $CH_3(CH_2)_7OH$...	130 14	−16 3	194	0 827	318
2967	$C_8H_{18}O$	d-sec.-Octyl alcohol $C_6H_{13}CH(OH)CH_3$	130 14		86²⁰	0 822	279
2968	$C_8H_{18}O$	dl-sec.-Octyl alcohol $C_6H_{13}CH(OH)CH_3$	130 14	−38 6	178.5	0 819	357
2969	$C_8H_{18}O$	Propylbutyl carbinol	130 14		71¹⁰	0 838⁴₀	
2970	$C_8H_{18}O$	Propylisobutyl carbinol	130 14		164	0 821	248
2971	$C_8H_{18}O$	Isopropylbutyl carbinol .	130 14		154	0 825	249
2972	$C_8H_{18}O$	Isopropylisobutyl carbinol	130 14		163	0 820¹⁴	
2973	$C_8H_{18}O$	n-Butyl ether $C_4H_9OC_4H_9$	130 14		140 9	0 769²⁰₂₀	
2974	$C_8H_{18}O$	Isobutyl ether $[(CH_3)_2CHCH_2]_2O$	130 14		122.5	0 762	
2975	$C_8H_{18}O$	sec.-Butyl ether $(C_2H_5CHCH_3)_2O$.	130 14		121	0 756²¹	
2976	$C_8H_{18}O$	Ethyl hexyl ether $C_2H_5OC_6H_{13}$.	130 14		137		
2977	$C_8H_{18}O$	Methyl n-heptyl ether $CH_3OC_7H_{15}$	130.14		149 8	0 795⁰₀	
2978	$C_8H_{18}O_2S$	n-Butylsulfone $(C_4H_9)_2SO_2$	178 20	43 5			
2979	$C_8H_{18}O_3$	Ethyl orthoacetate $CH_3CH(OC_2H_5)_3$	162 14		142	0.94²²	
2980	$C_8H_{18}O_4S_2$	Trional $C_2H_5(CH_3)C(SO_2C_2H_5)_2$	242 27	76			
2981	$C_8H_{18}S$	Di-n-butyl sulfide $(C_4H_9)_2S$	146 20	−79 7	182	0 852⁰	
2982	$C_8H_{18}S$	Diisobutyl sulfide $[(CH_3)_2CHCH_2]_2S$.	146.20		171	0 836¹⁰	
2983	$C_8H_{18}S$	Di-sec.-butyl sulfide $[C_2H_5CHCH_3]_2S$. .	146 20		165	0 832²²	
2984	$C_8H_{19}N$	Di-n-butylamine $(C_4H_9)_2NH$..........	129.15		161		
2985	$C_8H_{19}N$	Diisobutylamine $[(CH_3)_2CHCH_2]_2NH$.	129.15	−70 0	138 8	0 745	180
2986	$C_8H_{19}N$	n-Octylamine $C_8H_{17}NH_2$..............	129.15		180	0 777²⁷	319
2987	$C_8H_{19}N$	sec.-Octylamine $C_6H_{13}CH(CH_3)NH_2$....	129 15		164	0 771	292
2988	$C_8H_{12}As_2$	Ethylcacodyl $(C_2H_5)_2As_2(C_2H_5)_2$.. .	266 07		190		
2989	$C_8H_{21}NO$	Tetraethylammonium hydroxide. .	147 17	190 d.			
2990	$C_8H_6O_4$	Phthalonic anhydride	176 03	186			
2991	$C_9H_5Cl_2N$	2, 3-Dichloroquinoline	197.96	105			
2992	$C_9H_5Cl_2N$	2, 4-Dichloroquinoline................	197.96	67			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
2993	C₉H₄Cl₂N	5, 6-Dichloroquinoline	197.96	85			
2994	C₉H₄Cl₂N	5, 7-Dichloroquinoline	197 96	117			
2995	C₉H₄Cl₂N	5, 8-Dichloroquinoline	197 96	93			
2996	C₉H₄Cl₂N	6, 8-Dichloroquinoline	197 96	104			
2997	C₉H₄Cl₂N	7, 8-Dichloroquinoline	197 96	85 5			
2998	C₉H₄Br₂O₂	cis-1, 2-Dibromocinnamic acid	216 96	100	124° ᵇ		
2999	C₉H₄Br₂O₂	trans-2, 2-Dibromocinnamic acid	216 96	136	138° ᶜ		
3000	C₉H₆ClN	2-Chloroquinoline	163 51	38	275		
3001	C₉H₆ClN	3-Chloroquinoline	163 51		255 5		
3002	C₉H₆ClN	4-Chloroquinoline	163 51	34	260 4	1.251	
3003	C₉H₆ClN	5-Chloroquinoline	163 51	32	268		
3004	C₉H₆ClN	6-Chloroquinoline	163 51	41	262		
3005	C₉H₆ClN	7-Chloroquinoline	163 51	45	256		
3006	C₉H₆ClN	8-Chloroquinoline	163 51	> −20	288		
3007	C₉H₆Cl₂O₂	cis-1, 2-Dichlorocinnamic acid	216 96	121			
3008	C₉H₆Cl₂O₂	trans-1, 2-Dichlorocinnamic acid	216 96	101			
3009	C₉H₄lNO₄S	Loretin	351 05	d.			
3010	C₉H₆N₂O₂	5-Nitroquinoline	174 06	72			
3011	C₉H₆N₂O₂	6-Nitroquinoline	174 06	150			
3012	C₉H₆N₂O₂	7-Nitroquinoline	174 06	133			
3013	C₉H₆N₂O₂	8-Nitroquinoline	174 06	89			
3014	C₉H₆O₂	Phenylpropiolic acid C₆H₅C:CCO₂H.	146 04	137			
3015	C₉H₆O₂	Chromone	146 04	58			
3016	C₉H₆O₂	Coumarine	146 04	67	301 7	0 935	
3017	C₉H₆O₃	Umbelliferon	162 04	227			
3018	C₉H₆O₄	Daphnetin	178 05	256			
3019	C₉H₆O₄	Esculetin	178 05	270 d.			
3020	C₉H₆O₆	Hemimellitic acid 1, 2, 3-C₆H₃(CO₂H)₃	210 04	190			
3021	C₉H₆O₆	Trimellitic acid 1, 2, 4-C₆H₃(CO₂H)₃	210 05	216			
3022	C₉H₆O₆	Trimesic acid 1, 3, 5-C₆H₃(CO₂H)₃	210 05	350			
3023	C₉H₆O₇	1, 3, 5-Tricarboxyphenol	226 05	180 d.			
3024	C₉H₇BrO₂	cis-Allo-1-bromocinnamic acid	226 97	120	111° ᵇ		
3025	C₉H₇BrO₂	cis-Allo-2-bromocinnamic acid	226 97	160	111° ᵇ		
3026	C₉H₇BrO₂	trans-1-Bromocinnamic acid	226 97	131	121° ᵇ		
3027	C₉H₇BrO₂	trans-2-Bromocinnamic acid	226 97	135	122° ᵇ		
3028	C₉H₇ClO	Cinnamyl chloride C₆H₅CH:CHCOCl.	166.51	36	257 5		
3029	C₉H₇ClO₂	cis-Allo-1-chlorocinnamic acid	182.51	111	99° ᵇ		
3030	C₉H₇ClO₂	cis-Allo-2-chlorocinnamic acid	182 51	132	97° ᵇ		
3031	C₉H₇ClO₂	trans-1-Chlorocinnamic acid	182.51	137	109° ᵇ		
3032	C₉H₇ClO₂	trans-2-Chlorocinnamic acid	182 51	142	113° ᵇ		
3033	C₉H₇ClO₂	o-Chlorocinnamic acid	182 51	211			
3034	C₉H₇Cl₃O₂	Benzyl trichloroacetate	253 43		178 5⁵⁰	1.389⁴	692
3035	C₉H₇N	Cinnamic nitrile C₆H₅CH:CHCN	129.06	11	255	1 037⁰	
3036	C₉H₇N	Isoquinoline	129 06	23	243	1 099	1026
3037	C₉H₇N	Quinoline	129.06	−19.5	237 7	1.093	941
3038	C₉H₇NO	p-Cyanoacetophenone CN.C₆H₄COCH₃	145 06	61			
3039	C₉H₇NO	2-Hydroxyquinoline	145 06	200			
3040	C₉H₇NO	4-Hydroxyquinoline	145 06	201	300		
3041	C₉H₇NO	5-Hydroxyquinoline	145 06	224			
3042	C₉H₇NO	6-Hydroxyquinoline	145 06	193	360		
3043	C₉H₇NO	7-Hydroxyquinoline	145 06	238 d.			
3044	C₉H₇NO	8-Hydroxyquinoline	145 06	76	.266 9		
3045	C₉H₇NO₂	3-Aminocoumarine	161 06	130			
3046	C₉H₇NO₂	Indole-2-carboxylic acid	161.06	203 d.			
3047	C₉H₇NO₂	Indole-3-carboxylic acid	161 06	218 d.			
3048	C₉H₇NO₃	Indoxylic acid	177 06		123		
3049	C₉H₇NO₄	Kynuric acid	177 06	189			
3050	C₉H₇NO₄	o-Nitrocinnamic acid	193 06	240			
3051	C₉H₇NO₄	m-Nitrocinnamic acid	193.06	197			
3052	C₉H₇NO₄	p-Nitrocinnamic acid	193 06	286			
3053	C₉H₇NO₄S	Diaphthol	225.13	295			
3054	C₉H₈	Indene	116 06	−2	182 4	1 006	806
3055	C₉H₈	Phenylallylene C₆H₅C:CCH₃	116.06		185		
3056	C₉H₈Cl₂	Cinnamal chloride C₆H₅CH:CH₂CHCl₂.	186.98	58.5	143³⁰		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3057	$C_9H_8Cl_2O_2$	Benzyl dichloroacetate	218 98		179⁶⁰	1.313	684
3058	$C_9H_8I_2O_2$	Ethyl 3, 5-diiodosalicylate	417 93	132			
3059	$C_9H_8N_2$	2-Aminoquinoline	144 08	129			
3060	$C_9H_8N_2$	3-Aminoquinoline	144 08	94			1319
3061	$C_9H_8N_2$	4-Aminoquinoline	144 08	154			
3062	$C_9H_8N_2$	5-Aminoquinoline	144 08	110			
3063	$C_9H_8N_2$	6-Aminoquinoline	144 08	114			
3064	$C_9H_8N_2$	7-Aminoquinoline	144 08	189			
3065	$C_9H_8N_2$	8-Aminoquinoline	144 08	70			
3066	$C_9H_8N_2$	3-Phenylpyrazolone	144 08	240			
3067	$C_9H_8N_2O$	Cyanoacetanilide CNCH₂CONHC₆H₅	160 08	200			
3068	$C_9H_8N_2O$	Pyrrone (Dipyrryl ketone)	160 08	160			
3069	C_9H_8O	Cinnamic aldehyde C₆H₅CH.CHCHO	132 06	−7.5	251.0	1.049	791
3070	C_9H_8O	α-Hydrindone	132 06	41	244	1.101⁴⁵	
3071	C_9H_8O	β-Hydrindone	132 06	61	225 d.	1.071⁶⁷	1100
3072	$C_9H_8O_2$	o-Coumaric aldehyde	148 06	133			
3073	$C_9H_8O_2$	p-Coumaric aldehyde	148 06	134			
3074	$C_9H_8O_2$	Allocinnamic acid	148 06	68	125¹⁹		
3075	$C_9H_8O_2$	Cinnamic acid C₆H₅CH.CHCO₂H	148 06	133	300	1.284⁴	
3076	$C_9H_8O_2$	Isocinnamic acid	148 06	57	256 d.		
3077	$C_9H_8O_2$	Atropic acid	148 06	107	267 d.		
3078	$C_9H_8O_2$	Melilotic anhydride	148 06	25	272		
3079	$C_9H_8O_2$	Chromanone	148 06	38 5	160⁵⁰		
3080	$C_9H_8O_3$	Acetopiperone	164 06	83			
3081	$C_9H_8O_3$	o-Acetylsalicylic aldehyde	164 06	37	253		
3082	$C_9H_8O_3$	Benzoylacetic acid C₆H₅COCH₂CO₂H	164 06	104			
3083	$C_9H_8O_3$	o-Coumaric acid	164 06	208			
3084	$C_9H_8O_3$	m-Coumaric acid	164 06	191			
3085	$C_9H_8O_3$	p-Coumaric acid	164 06	206			
3086	$C_9H_8O_3$	Phenylpyruvic acid C₆H₅CH₂COCO₂H	164 06	157			
3087	$C_9H_8O_4$	o-Acetylsalicylic acid (Aspirin)	180 06	133 5			1290
3088	$C_9H_8O_4$	Caffeic acid	180 06	195			
3089	$C_9H_8O_4$	Phenylmalonic acid C₆H₅CH(CO₂H)₂	180 06	153			
3090	$C_9H_8O_4$	Uvitic acid 3, 5(CO₂H)₂C₆H₃CH₃	180 06	290			
3091	$C_9H_8O_4$	Methyl phthalate o-CO₂HC₆H₄CO₂CH₃	180 06	82 5			
3092	$C_9H_8O_4$	Benzoyl acetyl peroxide	180 06	36 6	130¹⁹		
3093	$C_9H_8O_5$	Esculetine acid	196 06	168			
3094	$C_9H_8O_5$	Myristicine acid	196 06	210	300		
3095	C_9H_9BrO	Indene oxybromide	212 99	130 5			
3096	$C_9H_9ClO_2$	Benzyl chloroacetate	184 53		147.5⁹	1.222	675
3097	C_9H_9N	Dihydroquinoline	131 08	226			
3098	C_9H_9N	1-Methylindole	131 08		242 4	1 071⁰	
3099	C_9H_9N	2-Methylindole	131 08	60	272 3		
3100	C_9H_9N	3-Methylindole (Scatole)	131.08	95	266 2		
3101	C_9H_9N	5-Methylindole	131 08	58 5			
3102	C_9H_9NO	Cinnamamide C₆H₅CH·CHCONH₂	147.08	141.5			
3103	C_9H_9NO	Hydrocarbostyril	147 08	163			1309
3104	$C_9H_9NO_2$	o-Aminocinnamic acid	163 08	159 d.			
3105	$C_9H_9NO_2$	m-Aminocinnamic acid	163 08	181			
3106	$C_9H_9NO_2$	p-Aminocinnamic acid	163 08	176 d.			
3107	$C_9H_9NO_2$	Benzoylacetaldehydeoxime	163.08	87			
3108	$C_9H_9NO_3$	o-Acetylaminobenzoic acid	179 08	185			
3109	$C_9H_9NO_3$	m-Acetylaminobenzoic acid	179 08	250			
3110	$C_9H_9NO_3$	p-Acetylaminobenzoic acid	179 08	252			
3111	$C_9H_9NO_3$	Hippuric acid C₆H₅CONHCH₂CO₂H	179 08	187 5	d.	1 371	1256
3112	$C_9H_9NO_3$	Methyl oxanilate C₆H₅NHCOCO₂CH₃	179 08	114			
3113	$C_9H_9NO_3$	Acetylsalicylamide	179.08	144			
3114	$C_9H_9NO_4$	Salicyluric acid	195 08	160			
3115	$C_9H_9NO_4$	Ethyl m-nitrobenzoate	195.08	47	298		
3116	$C_9H_9NO_4$	Ethyl p-nitrobenzoate	195.08	57			
3117	$C_9H_9N_3$	5, 8-Diaminoquinoline	159.09	156			
3118	$C_9H_9N_3$	6, 8-Diaminoquinoline	159.09	163			
3119	C_9H_{10}	Benzylethylene C₆H₅CH₂CH:CH₂	118 08		155	0.909	654

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3120	C₉H₁₀	Isoallylbenzene C₆H₅CH.CH·CH₃	118 08		175	0 924[14]	
3121	C₉H₁₀	Hydrindene	118 08		176 5	0 965	970
3122	C₉H₁₀N₂	1-Ethylindazole....	146 09		120[16]	1 064	878
3123	C₉H₁₀O₃	2-Acetamino-4-nitrotoluene	194 09	96			
3124	C₉H₁₀O	Anol p-(CH₃CH:CH)C₆H₄OH	134 08	93	250 d.		
3125	C₉H₁₀O	Chavicol p-(CH₂:CHCH₂)C₆H₄OH	134 08	> -25	237	1 033[14]	935
3126	C₉H₁₀O	Cinnamyl alcohol C₆H₅CH.CHCH₂OH	134 08	33	258 5	1 044	1039
3127	C₉H₁₀O	Allyl phenyl ether C₃H₅OC₆H₅	134 08		192		
3128	C₉H₁₀O	Methyl styryl ether	134 08		213	1 001	877
3129	C₉H₁₀O	2, 4-Dimethylbenzaldehyde	134 08	-8	216		
3130	C₉H₁₀O	Hydrocinnamaldehyde .	134 08	17	280		
3131	C₉H₁₀O	o-Xylene-4-aldehyde . .	134 08		225		
3132	C₉H₁₀O	Ethyl phenyl ketone C₂H₅COC₆H₅	134 08	21	218	1 010	689
3133	C₉H₁₀O	Methyl benzyl ketone CH₃COCH₂C₆H₅	134 08	-15 4	216 7	1 028	
3134	C₉H₁₀O	p-Methylacetophenone (Melilot)	134 08		222	1 013[13]	703
3135	C₉H₁₀O	Chromane	134 08		95[12]	1 064	
3135 1	C₉H₁₀OS	Ethyl thiobenzoate.	166 14		253[761]	1 004[15]	
3136	C₉H₁₀O₂	o-Coumaral alcohol	150 08	119			
3137	C₉H₁₀O₂	Hesperetol.........	150 08	57			
3138	C₉H₁₀O₂	2, 3-Dimethylbenzoic acid	150 08	144			
3139	C₉H₁₀O₂	2, 4-Dimethylbenzoic acid	150 08	126	268		
3140	C₉H₁₀O₂	2, 5-Dimethylbenzoic acid	150 08	132	268	1 069	
3141	C₉H₁₀O₂	2, 6-Dimethylbenzoic acid	150 08	116			
3142	C₉H₁₀O₂	3, 4-Dimethylbenzoic acid	150 08	165			
3143	C₉H₁₀O₂	o-Ethylbenzoic acid .	150 08	68			
3144	C₉H₁₀O₂	m-Ethylbenzoic acid.	150 08	47		1 042[100]	1148
3145	C₉H₁₀O₂	p-Ethylbenzoic acid	150 08	113			
3146	C₉H₁₀O₂	Hydratropic acid C₂H₄(C₆H₅)CO₂H	150 08		265		
3147	C₉H₁₀O₂	Hydrocinnamic acid	150 08	48 6	279 8	1 071[8 7]	
3148	C₉H₁₀O₂	Mesitylinic acid 3, 5-(CH₃)₂C₆H₃CO₂H	150 08	166			
3149	C₉H₁₀O₂	Benzyl acetate CH₃CO₂CH₂C₆H₅. .	150 08	-51 5	213 5	1 058	673
3150	C₉H₁₀O₂	o-Cresyl acetate o-CH₃CO₂C₆H₄CH₃.	150 08		208		
3151	C₉H₁₀O₂	m-Cresyl acetate m-CH₃CO₂C₆H₄CH₃	150 08		212		
3152	C₉H₁₀O₂	p-Cresyl acetate p-CH₃CO₂C₆H₄CH₃	150 08		212 5	1.050	599
3154	C₉H₁₀O₂	Ethyl benzoate C₆H₅CO₂C₂H₅......	150 08	-34 6	213 2	1 047	628
3155	C₉H₁₀O₂	Methyl phenylacetate.......... .	150 08		220	1 044[16]	
3156	C₉H₁₀O₂	Methyl p-toluate p-CH₃C₆H₄CO₂CH₃	150 08	33	217		
3157	C₉H₁₀O₂	Phenyl propionate C₂H₅CO₂C₆H₅...	150 08	20	211	1 054[15]	
3158	C₉H₁₀O₃	Acetovanillone...............	166 08	115	300		
3159	C₉H₁₀O₃	Paeonol 4, 2-CH₃O(OH)C₆H₃COCH₃	166 08	50			
3160	C₉H₁₀O₃	o-Ethoxybenzoic acid . .	166 08	22			
3161	C₉H₁₀O₃	m-Ethoxybenzoic acid	166 08	137			
3162	C₉H₁₀O₃	p-Ethoxybenzoic acid.	166 08	195			
3163	C₉H₁₀O₃	dl-Atrolactic acid	166 08	91			
3164	C₉H₁₀O₃	m-Hydrocoumaric acid	166 08	111			
3165	C₉H₁₀O₃	Melilotic acid. .	166 08	83			
3166	C₉H₁₀O₃	d(l)-2-Phenyllactic acid	166 08	125			
3167	C₉H₁₀O₃	Phloretic acid HOC₆H₄CH(CH₃)CO₂H	166 08	129			
3168	C₉H₁₀O₃	d(l)-Tropic acid........	166 08	128			
3169	C₉H₁₀O₃	dl-Tropic acid..... .. .	166 08	123			
3169.1	C₉H₁₀O₃	Anisyl acetate p-CH₃OC₆H₄O₂CCH₃.	166 08		139[12]	1.101	
3170	C₉H₁₀O₃	Ethyl salicylate OHC₆H₄CO₂C₂H₅.	166 08	1 3	231 5	1 131	670
3171	C₉H₁₀O₃	Guaiacyl acetate (Eucol).	166 08		240	1.138	
3172	C₉H₁₀O₃	Methyl anisate p-CH₃OC₆H₄CO₂CH₃.	166 08	48	256		
3173	C₉H₁₀O₃	Methyl o-cresotinate..	166 08	30	235		
3174	C₉H₁₀O₃	Methyl p-cresotinate	166 08		242		
3175	C₉H₁₀O₃	Methyl dl-mandelate	166 08	58	144[20]		
3176	C₉H₁₀O₄	Hydrocaffeic acid....	182 08	139			
3177	C₉H₁₀O₄	d(l)-Phenylglyceric acid . .	182 08	164			
3178	C₉H₁₀O₄	dl-Phenylglyceric acid . .	182 08	141		1.451	
3179	C₉H₁₀O₄	d(l)-p-Methoxymandelic acid . ..	182 08	105		1.354	
3181	C₉H₁₀O₄	Veratric acid 3, 4-(CH₃O)₂C₆H₃CO₂H .	182 08	181			
3182	C₉H₁₀O₄	Methoxymethyl salicylate.............	182.08		162[42]	1.200[16]	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3183	C₉H₁₀O₄	Methyl vanillate	182.08	63	287		
3184	C₉H₁₀O₄	Glycol salicylate (Spirosal)	182.08		170¹²		
3185	C₉H₁₀O₅	Syringic acid	198.08	245			
3186	C₉H₁₀O₅	Ethyl gallate	198.08	160			
3187	C₉H₁₀O₅	2, 3, 4, 5-Dimethoxydihydroxybenzoic acid	214.08	148			
3187.1	C₉H₁₀S₂	Ethyl dithiobenzoate	182 21		180²⁸	1.1439²⁸	
3188	C₉H₁₁N	Allylaniline C₆H₅NHCH₂CH:CH₂	133 09		209	0.982²⁵	
3189	C₉H₁₁N	Benzylideneethylamine	133 09		195.4		
3190	C₉H₁₁N	Styrylamine C₆H₅CH CHCH₂NH₂	133 09		237		
3191	C₉H₁₁N	1, 2, 3, 4-Tetrahydroisoquinoline	133 09		233	1.064	1012
3192	C₉H₁₁N	1, 2, 3, 4-Tetrahydroquinoline	133 09	20	251	1.055	1013
3193	C₉H₁₁NO	p-Dimethylaminobenzaldehyde	149 09	75			
3194	C₉H₁₁NO	o-Acetotoluide o-CH₃CONHC₆H₄CH₃	149 09	110	296		1255
3195	C₉H₁₁NO	m-Acetotoluide m-CH₃CONHC₆H₄CH₃	149 09	65 5	303		
3196	C₉H₁₁NO	p-Acetotoluide p-CH₃CONHC₆H₄CH₃	149 09	153	307		1276
3197	C₉H₁₁NO	N-Benzylacetamide CH₃CONHC₇H₇	149 09	61	300		
3198	C₉H₁₁NO	N-Ethylbenzamide C₆H₅CONHC₂H₅	149 09	71	290		
3199	C₉H₁₁NO	N-Methylacetanilide (Exalgin)	149 09	102	254.7		1250
3200	C₉H₁₁NO	Propionanilide C₂H₅CONHC₆H₅	149 09	104			
3201	C₉H₁₁NOS	N-Phenylthiourethane	181.16	69			
3202	C₉H₁₁NO₂	4-Acetylamino-2-hydroxytoluene	165 09	225			
3203	C₉H₁₁NO₂	3-Acetylamino-4-hydroxytoluene	165 09	160			
3204	C₉H₁₁NO₂	p-Acetylmethylaminophenol	165 09	240			
3205	C₉H₁₁NO₂	1-Anilinopropionic acid	165 09	162			
3206	C₉H₁₁NO₂	o-Dimethylanthranilic acid	165 09	175			
3207	C₉H₁₁NO₂	m-Ethylaminobenzoic acid	165.09	101			
3208	C₉H₁₁NO₂	l-Phenylalanine	165 09	283 d.			1260
3209	C₉H₁₁NO₂	dl-Phenylalanine	165 09	265 d.			
3210	C₉H₁₁NO₂	o-Tolylaminoacetic acid	165 09	150			
3211	C₉H₁₁NO₂	p-Tolylaminoacetic acid	165 09	118			
3212	C₉H₁₁NO₂	2, 4, 6-Trimethylpyridine-3-carboxylic acid	165.09		155		
3213	C₉H₁₁NO₂	Ethyl p-aminobenzoate	165 09	91			
3214	C₉H₁₁NO₂	Ethyl anthranilate	165.09		260		
3216	C₉H₁₁NO₂	o-Acetaniside o-CH₃OC₆H₄NHCOCH₃	165 09	84	305		
3217	C₉H₁₁NO₂	p-Acetanisidc CH₃CONHC₆H₄OCH₃	165 09	127			
3218	C₉H₁₁NO₂	p-Formylphenetidine	165 09	60			
3219	C₉H₁₁NO₂	Nitrocumene (CH₃)₂CHC₆H₄NO₂	165.09	−35	224 d.		
3220	C₉H₁₁NO₂	Nitromesitylene	165 09	44	255		
3221	C₉H₁₁NO₂	N-Phenylurethane C₂H₅CO₂NHC₆H₅	165 09	52	238		
3222	C₉H₁₁NO₃	l-Tyrosine	181 09	295 d.			
3223	C₉H₁₂	Cumene (CH₃)₂CHC₆H₅	120.09		153 4	1.456	1259
3224	C₉H₁₂	o-Ethyltoluene o-C₂H₅C₆H₄CH₃	120 09	> −17	162	0.864	561
3225	C₉H₁₂	m-Ethyltoluene m-C₂H₅C₆H₄CH₃	120 09		162.5	0.882	615
3226	C₉H₁₂	p-Ethyltoluene p-C₂H₅C₆H₄CH₃	120 09	< −20	162	0.867	585
3227	C₉H₁₂	Hemimellitene 1, 2, 3-(CH₃)₃C₆H₃	120.09		176 5	0 862	568
3228	C₉H₁₂	Mesitylene 1, 3, 5-(CH₃)₃C₆H₃	120.09	−52.7	164 6	0.895	650
3229	C₉H₁₂	n-Propylbenzene CH₃(CH₂)₂C₆H₅	120.09	−101.6	157.5	0.863	580
3230	C₉H₁₂	Pseudocumene 1, 2, 4-(CH₃)₃C₆H₃	120.09	−61.0	169.8	0.862	556
3231	C₉H₁₂N₂O	1-Ethyl-2-phenylurea	164 11	99		0.87	622
3232	C₉H₁₂N₂O₂	p-Phenetylurea C₂H₅OC₆H₄NHCONH₂	180.11	173			
3233	C₉H₁₂N₂O₂	Pilosinine	180.11	79	300²⁵		
3234	C₉H₁₂N₄O₃	1, 3, 7, 9-Tetramethyluric acid	224 12	228	d.		1268
3235	C₉H₁₂O	Benzylmethyl carbinol	136 09		212	0.994	
3235.1	C₉H₁₂O	d-Benzylmethyl carbinol	136 09		125²⁵	0 991	660
3236	C₉H₁₂O	Ethylphenyl carbinol	136 09		219	0.996	
3237	C₉H₁₂O	Hydrocinnamyl alcohol	136 09	< −18	237.4	1.008	706
3238	C₉H₁₂O	Mesitol 2, 4, 6-(CH₃)₃C₆H₂OH	136 09	69	220		
3239	C₉H₁₂O	o-n-Propylphenol o-C₃H₇C₆H₄OH	136 09		226.6		
3240	C₉H₁₂O	m-n-Propylphenol m-C₃H₇C₆H₄OH	136.09	26	228	1.015⁰	
3241	C₉H₁₂O	p-n-Propylphenol p-C₃H₇C₆H₄OH	136.09	61	232.6	1.009⁰	
3242	C₉H₁₂O	Pseudocumenol 2, 4, 5-(CH₃)₃C₆H₂OH	136.09	72	235		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3243	$C_9H_{12}O$	Ethyl benzyl ether $C_7H_7OC_2H_7$	136 09		226	0.998¹⁷·⁴	
3244	$C_9H_{12}O$	Ethyl m-cresyl ether	136 09		192	0.949	648
3245	$C_9H_{12}O$	Ethyl p-cresyl ether $p\text{-}CH_3C_6H_4OC_2H_5$	136 09		189 9	0.874⁰	928
3246	$C_9H_{12}O$	Propyl phenyl ether $C_3H_7OC_6H_5$	136 09		190 5	0 968	
3247	$C_9H_{12}O$	Isopropyl phenyl ether	136 09		177 2	0.946¹⁵	
3248	$C_9H_{12}O_2$	Mesorcinol	152 09	150	275 5		
3249	$C_9H_{12}O_2$	Guaiacyl ethyl ether	152 09		213		•
3250	$C_9H_{12}O_3$	Phloroglucinol trimethyl ether	168 09	52	255 5		
3251	$C_9H_{12}O_3$	Pyrogallol trimethyl ether	168 09	47	241	1 099¹⁵	
3252	$C_9H_{12}O_3$	Metacrolein $(C_3H_4O)_3$	168 09	46			
3253	$C_9H_{12}O_3$	Caryophyllenic acid	168 09			1.140	
3254	$C_9H_{12}O_3S$	Mesitylenesulfonic acid	200 16	77			
3255	$C_9H_{12}O_3S$	Toluene p-ethylsulfonate	200 16	33	173¹⁵	1 174²¹	
3256	$C_9H_{12}O_5$	Anhydrocamphoronic acid	200 09	133			
3257	$C_9H_{13}N$	Cumidine $p\text{-}(CH_3)_2CHC_6H_4NH_2$	135 11	63	225	0.957	1333
3258	$C_9H_{13}N$	Dimethyl-o-toluidine	135 11	−61 0	184 6	0.929	682
3259	$C_9H_{13}N$	Dimethyl-m-toluidine	135 11		212 5	0 941	733
3260	$C_9H_{13}N$	Dimethyl-p-toluidine	135 11		211 5	0 937	726
3261	$C_9H_{13}N$	Ethyl-o-toluidine	135 11		214	0.953¹⁵·⁴	
3262	$C_9H_{13}N$	Ethyl-m-toluidine	135 11		222		
3263	$C_9H_{13}N$	Ethyl-p-toluidine	135 11		217	0 939	
3264	$C_9H_{13}N$	Mesidine 1, 3, 5-$(CH_3)_3C_6H_2NH_2$	135 11		233	0.963	
3265	$C_9H_{13}N$	ω-Mesitylamine	135.11		218 2	0 950	699
3266	$C_9H_{13}N$	Parvoline	135 11		234		
3267	$C_9H_{13}N$	n-Propylaniline $C_6H_5NHC_3H_7$	135 11		222	0 949¹⁴	
3268	$C_9H_{13}N$	Isopropylaniline $C_6H_5NHCH(CH_3)_2$	135 11		213		
3269	$C_9H_{13}N$	Pseudocumidine	135 11	66	235		
3270	$C_9H_{13}NO_2$	Anhydroecgonine	167 11	235 d.			
3271	$C_9H_{13}NO_3$	Adrenaline	183 11	207 d.			
3272	C_9H_{14}	Apocyclene	122 11	43	138 9	0.871⁴⁰	1056
3273	C_9H_{14}	Santene	122 11		142	0.869¹⁵	486
3274	$C_9H_{14}ClNO_2$	Anhydroecgonine hydrochloride	203 57	241			
3275	$C_9H_{14}N_2O_3$	Ethylpropylbarbituric acid	198 12	146			
3276	$C_9H_{14}O$	Nopinone	138 11	0	209		
3277	$C_9H_{14}O$	Phorone	138 11	28	198 5	0 885	598
3278	$C_9H_{14}O_2$	Lauronolic acid	154 11		129¹¹·⁴		
3279	$C_9H_{14}O_2$	Methyl amylpropiolate	154 11		111¹⁶	0 991¹⁵	
3280	$C_9H_{14}O_3$	Castelamarin	170 11	269			
3281	$C_9H_{14}O_4$	cis-Hexahydrohomophthalic acid	186 11	146			
3282 •	$C_9H_{14}O_4$	trans-Hexahydrohomophthalic acid	186.11	157			
3282.1	$C_9H_{14}O_4$	dl-Pinic acid	186.11	102 5	216¹⁰	1 093¹⁰⁹·⁴	1154
3282 2	$C_9H_{14}O_4$	d-Pinic acid	186 11	136	216¹⁰		
3283	$C_9H_{14}O_4$	Diethyl citraconate	186 11		230.3	1.062	847
3284	$C_9H_{14}O_4$	Diethyl glutaconate	186 11		238	1 050	
3285	$C_9H_{14}O_4$	Diethyl itaconate	186 11		227.9	1 045	369
3286	$C_9H_{14}O_4$	Diethyl mesaconate	186 11		229	1.047	594
3287	$C_9H_{14}O_5$	4-Ketoazelaic acid	202 11	102; 109			
3288	$C_9H_{14}O_6$	l-Camphoronic acid	218.11	165			
3289	$C_9H_{14}O_6$	Glycerol triacetate	218 11		259	1.161	326
3290	$C_9H_{14}O_7$	Trimethyl citrate	234 11	79	287 d.		
3291	$C_9H_{15}NO$	Pseudopelletierine	153 12	49	246	1 001⁹⁹·⁴	1138
3292	$C_9H_{15}NO_3$	d-Ecgonine	185 12	257			
3293	$C_9H_{15}NO_3$	l-Ecgonine	185.12	198 d.		1.370⁰¹²	
3294	$C_9H_{15}NO_3$	dl-Ecgonine	185.12	212			
3294 1	$C_9H_{15}N_3O_2S$	Ergothioneine	229 21	290			
3295	C_9H_{16}	Campholene	124 12	> −20	133	0.803	399
3296	C_9H_{16}	Nopinane	124 12		149 5	0.861¹⁷	479
3297	C_9H_{16}	Pulegene	124.12		139	0.791²¹	979
3298	$C_9H_{16}ClNO_3$	l-Ecgonine hydrochloride	221.59	246			
3299	$C_9H_{16}N_2O_2S_2$	Cheiroline	328.33	48	200 d.		
3300	$C_9H_{16}O$	Camphorol	140.12		81¹⁶		
3301	$C_9H_{16}O$	α-Nopinol	140 12	102	205		
3302	$C_9H_{16}O$	dl-Santenol	140.12	98	196	0.987	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3303	$C_9H_{16}O_2$	Amyl l-α-crotonate	156 12			0.896	360
3304	$C_9H_{16}O_2$	Ethyl hexahydrobenzoate	156 12		196.5	0.967_4^{15}	886
3305	$C_9H_{16}O_2$	Methyl cyclohexylacetate	156 12		202	0.990_0^{14}	
3306	$C_9H_{16}O_3$	Ethyl isopropylacetoacetate	172 12		205 d.	$0\ 960_{16}^{25}$	
3307	$C_9H_{16}O_4$	Azelaic acid $HO_2C(CH_2)_7CO_2H$	188 12	106.5	360	1.029	1155
3308	$C_9H_{16}O_4$	n-Butyl ethyl malonate	188 12		130^{12}	0.976_{16}^{25}	284
3309	$C_9H_{16}O_4$	Isobutyl ethyl malonate	188 12		120^8	0.968_{16}^{25}	286
3310	$C_9H_{16}O_4$	sec.-Butyl ethyl malonate	188 12		160^{60}	$0\ 986_{16}^{25}$	310
3311	$C_9H_{16}O_4$	Diethyl dimethylmalonate	188 12		196	0 995	196
3312	$C_9H_{16}O_4$	Diethyl glutarate $CH_2(CH_2COC_2H_5)_2$.	188 12		237	1.025	
3313	$C_9H_{16}O_4$	Dipropyl malonate $CH_2(CO_2C_3H_7)_2$	188 12		228 3	1.027_0^9	
3314	$C_9H_{16}O_4$	Propyl isopropyl malonate	188 12		143^{42}	$0\ 980_{16}^{25}$	293
3314 1	$C_9H_{17}BrO_2$	l-Amyl bromobutyrate	221 05		105^{11}	1.196^{25}	
3315	$C_9H_{17}NO$	Homotropine	155 14	85			
3316	$C_9H_{17}NO$	Methylpelletierine	155 14		215		
3317	$C_9H_{17}NO$	Triacetoneamine	155 14	39 6			
3318	C_9H_{18}	Cyclononane	126 14		172	0.773_4^{16}	
3319	C_9H_{18}	Ethylcycloheptane $C_2H_5C_7H_{13}$.	126 14	< -30	199	0 952	
3320	C_9H_{18}	Hexahydrocumene $(CH_3)_2CHC_6H_{11}$	126 14		150	0.787	
3321	C_9H_{18}	2-Methyl-1-octene $C_6H_{13}C(CH_3):CH_2$..	126 14		143		
3322	C_9H_{18}	Nonylene $C_6H_{13}CH.CHCH_3$ ..	126 14		149 9	$0\ 754_{16}^{15}$	
3323	C_9H_{18}	Propylcyclohexane $C_3H_7C_6H_{11}$	126 14		149 5	0 767	
3324	$C_9H_{18}O$	dl-Pulenol	142 14		187 5	0 908	902
3325	$C_9H_{18}O$	Pelargonic aldehyde $CH_3(CH_2)_7CHO$.	142 14		$93\ 5^{23}$	$0\ 828^{15}$	280
3326	$C_9H_{18}O$	Diisobutyl ketone $[(CH_3)_2CHCH_2]_2CO$	142 14		182	0 833	
3327	$C_9H_{18}O$	Isopropyl isoamyl ketone	142 14		172		
3328	$C_9H_{18}O_2$	Pelargonic acid $CH_3(CH_2)_7CO_2H$.	158 14	12	254	0 907	340
3329	$C_9H_{18}O_2$	Amyl n-butyrate $C_4H_7CO_2C_5H_{11}$	158.14		184 8	0.883_0^9	184
3330	$C_9H_{18}O_2$	Isoamyl n-butyrate	158 14		178 6	$0\ 882_4^9$	
3330 1	$C_9H_{18}O_2$	d-β-Amyl n-butyrate	158 14		71^{16}	0 869	161
3331	$C_9H_{18}O_2$	Amyl isobutyrate $(CH_3)_2CHCO_2C_5H_{11}$	158 14		155	0 859	167
3332	$C_9H_{18}O_2$	Butyl n-valerate $C_4H_9CO_2C_4H_9$	158 14		185 8	$0\ 885^0$	
3333	$C_9H_{18}O_2$	Isobutyl n-valerate	158 14		167	0 854	
3333 1	$C_9H_{18}O_2$	d-sec.-Butyl valerate	158 14		67^{18}	0 860	164
3334	$C_9H_{18}O_2$	Isobutyl isovalerate	158 14		168 5	0 854	162
3335	$C_9H_{18}O_2$	Ethyl n-heptylate $C_6H_{13}CO_2C_2H_5$	158 14		187 1	0.872_{16}^{15}	195
3336	$C_9H_{18}O_2$	n-Heptyl acetate $CH_3CO_2C_7H_{15}$	158 14		191 5	$0\ 874_{16}^{15}$	221
3337	$C_9H_{18}O_2$	Methyl caprylate $C_7H_{15}CO_2CH_3$	158.14	-41	192 9	0.887	
3338	$C_9H_{18}O_2$	d-β-Octylformate	158 14		82^{20}	$0\ 872^{12\ 5}$	209
3339	$C_9H_{18}O_2$	Propyl caproate $C_5H_{11}CO_2C_3H_7$	158.14		185 5	0.884_0^9	
3340	$C_9H_{18}O_3$	Parapropionaldehyde $(C_3H_6O)_3$	174 14		170		
3341	$C_9H_{18}O_3$	Di-n-butyl carbonate $(C_4H_9O)_2CO$	174 14		207.7	0 924	
3342	$C_9H_{18}O_3$	Diisobutyl carbonate	174 14		190.3	0.919^{15}	
3343	$C_9H_{18}O_4$	1, 2-Dihydroxypelargonic acid	190 14	123			
3344	$C_9H_{18}O_7$	Galactite	238 14		142		1214
3345	$C_9H_{19}N$	l-1-Methylconine	141.15		175 5	0.832^{24}	
3346	$C_9H_{19}NO$	N-Diethyl-n-valeramide	157.15		210		
3347	C_9H_{20}	2, 4-Dimethylheptane	128 15		133.3	0.716	143
3348	C_9H_{20}	d-2, 5-Dimethylheptane	128 15		137	$0\ 715^{16}$	
3349	C_9H_{20}	dl-2, 5-Dimethylheptane	128 15		135.9	0.719_{15}^{15}	144
3350	C_9H_{20}	2, 6-Dimethylheptane	128 15		132 0	0.712_{15}^{15}	
3351	C_9H_{20}	4-Ethylheptane $(C_3H_7)_2CHC_2H_5$.	128 15		139	0.741	170
3352	C_9H_{20}	d-3-Methyloctane	128 15		143.4	0.721^{17}	
3353	C_9H_{20}	4-Methyloctane $C_3H_7(CH_3)CHC_4H_9$.	128 15		141 6	0.732_{15}^{15}	147
3354	C_9H_{20}	n-Nonane $CH_3(CH_2)_7CH_3$.	128 15	-51	150 6	0 718	151
3355	$C_9H_{20}O$	Butyl-sec.-butyl carbinol .	144 15		180	0 834	335
3356	$C_9H_{20}O$	Dibutyl carbinol $(C_4H_9)_2CHOH$.	144.15		194	0 823	320
3357	$C_9H_{20}O$	Diisobutyl carbinol	144.15		174 3	$0\ 816_{12}^{12}$	271
3358	$C_9H_{20}O$	Di-sec.-butyl carbinol .	144.15		171	0.836	338
3359	$C_9H_{20}O$	Diethylisobutyl carbinol	144 15		172		
3360	$C_9H_{20}O$	4, 6-Dimethylheptane-2-ol	144 15		195	$0\ 879^0$	
3361	$C_9H_{20}O$	Methylethylisoamyl carbinol	144 15		175	0 829	329
3362	$C_9H_{20}O$	Methylethyl-tert.-amyl carbinol	144.15		166	0.832	348

No.	Formula	Name	Mol. wt.	M P	B. P.	d	R. I. No.
3363	C₉H₂₀O	Methylpropylisobutyl carbinol	144 15		171 3	0 826	330
3364	C₉H₂₀O	n-Nonyl alcohol CH₃(CH₂)₈OH	144 15	−5	215	0 828	344
3365	C₉H₂₀O	Isobutyl-d-amyl ether	144 15		148 2	0 773	125
3366	C₉H₂₀O	Ethyl n-heptyl ether C₂H₅OC₇H₁₅	144 15		166 6	0 790¹⁶	
3367	C₉H₂₀O	Methyl n-octyl ether CH₃OC₈H₁₇	144 15		173	0 802⁵₀	
3368	C₉H₂₀O₂	Propylidene dipropyl ether	136 15		166 2	0 849⁰	
3369	C₉H₂₀O₄	Ethyl orthocarbonate C(OC₂H₅)₄	192 15		159	0 917	90
3370	C₉H₂₀O₄S₂	Tetronal (C₂H₅)₂C(SO₂C₂H₅)₂	256 28	85			
3371	C₉H₂₁N	n-Nonylamine C₉H₁₉NH₂	143 17		195		
3372	C₉H₂₁N	Tri-n-propylamine (C₃H₇)₃N	143 17	−93 5	156	0 757	230
3373	C₁₀H₂Cl₆	Hexachloronaphthalene	334 76	143			
3374	C₁₀H₄Cl₄	α-Tetrachloronaphthalene	265 86	130			
3375	C₁₀H₄Cl₄	β-Tetrachloronaphthalene	265 86	194			
3376	C₁₀H₄Cl₄	γ-Tetrachloronaphthalene	265 86	176			
3377	C₁₀H₄Cl₄	δ-Tetrachloronaphthalene	265 86	141			
3378	C₁₀H₄Cl₄	ε-Tetrachloronaphthalene	265 86	180			
3379	C₁₀H₄Cl₄	ζ-Tetrachloronaphthalene	265 86	160 5			
3380	C₁₀H₄Cl₄	vic.-Tetrachloronaphthalene	265 86	140			
3381	C₁₀H₄N₄O₈	α-Tetranitronaphthalene	308 06	259	exp.		
3382	C₁₀H₄N₄O₈	1, 2, 5, 8-Tetranitronaphthalene	308 06	270 d.			
3383	C₁₀H₄N₄O₈	1, 2, 6, 8-Tetranitronaphthalene	308 06	<300			
3384	C₁₀H₄N₄O₈	1, 3, 5, 8-Tetranitronaphthalene	308 06	195			
3385	C₁₀H₄N₄O₈	1, 3, 6, 8-Tetranitronaphthalene	308 06	203	exp.		
3386	C₁₀H₄N₄O₉	2, 4, 5, 7-Tetranitro-α-naphthol	324 06	180			
3387	C₁₀H₅Cl₃	1, 2, 3-Trichloronaphthalene	231 41	81			
3388	C₁₀H₅Cl₃	1, 2, 4-Trichloronaphthalene	231 41	92			
3389	C₁₀H₅Cl₃	1, 2, 5-Trichloronaphthalene	231 41	78			
3390	C₁₀H₅Cl₃	1, 2, 6-Trichloronaphthalene	231 41	97			
3391	C₁₀H₅Cl₃	1, 2, 7-Trichloronaphthalene	231 41	88			
3392	C₁₀H₅Cl₃	1, 2, 8-Trichloronaphthalene	231 41	83 5			
3393	C₁₀H₅Cl₃	1, 3, 5-Trichloronaphthalene	231 41	103			
3394	C₁₀H₅Cl₃	1, 3, 6-Trichloronaphthalene	231 41	80 5			
3395	C₁₀H₅Cl₃	1, 3, 7-Trichloronaphthalene	231 41	113			
3396	C₁₀H₅Cl₃	1, 3, 8-Trichloronaphthalene	231 41	89 5			
3397	C₁₀H₅Cl₃	1, 4, 5-Trichloronaphthalene	231 41	131			
3398	C₁₀H₅Cl₃	1, 4, 6-Trichloronaphthalene	231 41	66			
3399	C₁₀H₅Cl₃	1, 6, 7-Trichloronaphthalene	231 41	109 5			
3400	C₁₀H₅Cl₃	2, 3, 6-Trichloronaphthalene	231 41	91			
3401	C₁₀H₅Cl₃	2, 3, 7-Trichloronaphthalene	231 41	90			
3402	C₁₀H₅NO₁₀	Pyridinepentacarboxylic acid	299 05	220 d.			
3403	C₁₀H₅N₃O₆	1, 2, 5-Trinitronaphthalene	263 06	113			
3404	C₁₀H₅N₃O₆	1, 3, 5-Trinitronaphthalene	263 06	123			
3405	C₁₀H₅N₃O₆	1, 3, 8-Trinitronaphthalene	263 06	218			
3406	C₁₀H₅N₃O₆	1, 4, 5-Trinitronaphthalene	263 06	247			
3407	C₁₀H₅N₃O₇	2, 4, 5-Trinitro-α-naphthol	279 06	189.5			
3408	C₁₀H₅N₃O₇	2, 4, 7-Trinitro-α-naphthol	279 06	145			
3409	C₁₀H₅N₃O₇	2, 4, 8-Trinitro-α-naphthol	279 06	175			
3410	C₁₀H₆ClNO₂	4-Chloro-1-nitronaphthalene	207 51	84			
3411	C₁₀H₆ClNO₂	7-Chloro-1-nitronaphthalene	207 51	116			
3412	C₁₀H₆Cl₂	1, 2-Dichloronaphthalene	196 96	37	282	1 315⁴⁰ ⁵	1076
3413	C₁₀H₆Cl₂	1, 3-Dichloronaphthalene	196 96	61	289		
3414	C₁₀H₆Cl₂	1, 4-Dichloronaphthalene	196 96	68	287 6	1 300⁷⁶₄	1104
3415	C₁₀H₆Cl₂	1, 5-Dichloronaphthalene	196 96	107			
3416	C₁₀H₆Cl₂	1, 6-Dichloronaphthalene	196 96	48	286	1 261¹⁰⁰₄	1149
3417	C₁₀H₆Cl₂	1, 7-Dichloronaphthalene	196.96	62	d.	1 292¹⁰⁰₄	1150
3418	C₁₀H₆Cl₂	1, 8-Dichloronaphthalene	196 96	88			
3419	C₁₀H₆Cl₂	2, 3-Dichloronaphthalene	196 96	120			
3420	C₁₀H₆Cl₂	2, 6-Dichloronaphthalene	196 96	135	285		
3421	C₁₀H₆Cl₂	2, 7-Dichloronaphthalene	196 96	114			
3422	C₁₀H₆Cl₂O	2, 3-Dichloro-α-naphthol	212 96	101			
3423	C₁₀H₆Cl₂O	2, 4-Dichloro-α-naphthol	212 96	108			
3424	C₁₀H₆Cl₂O	5, 7-Dichloro-α-naphthol	212 96	132			
3425	C₁₀H₆Cl₂O	5, 8-Dichloro-α-naphthol	212 96	115			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3426	$C_{10}H_6Cl_2O$	6, 7-Dichloro-α-naphthol	212.96	151			
3427	$C_{10}H_6Cl_2O$	7, 8-Dichloro-α-naphthol	212.96	95			
3428	$C_{10}H_6Cl_2O$	1, 3-Dichloro-β-naphthol	212.96	81			
3429	$C_{10}H_6Cl_2O$	1, 4-Dichloro-β-naphthol	212.96	124			
3429 1	$C_{10}H_6Cl_2O$	3, 6-(6, 8)-Dichloro-β-naphthol	212 96	125			
3430	$C_{10}H_6Cl_2O_4S_2$	Naphthalene-1, 5-disulfonechloride	325.09	183			
3431	$C_{10}H_6Cl_2O_4S_2$	Naphthalene-1, 6-disulfonechloride	325.09	129			
3432	$C_{10}H_6Cl_2O_4S_2$	Naphthalene-2, 6-disulfonechloride	325.09	226			
3433	$C_{10}H_6Cl_2O_4S_2$	Naphthalene-2, 7-disulfonechloride	325.09	162			
3434	$C_{10}H_6N_2O_3$	Pyrocoll	186 06	269			
3435	$C_{10}H_6N_2O_4$	1, 2-Dinitronaphthalene	218 06	103			
3436	$C_{10}H_6N_2O_4$	1, 3-Dinitronaphthalene	218 06	145			
3437	$C_{10}H_6N_2O_4$	1, 4-Dinitronaphthalene	218 06	129			
3438	$C_{10}H_6N_2O_4$	1, 5-Dinitronaphthalene	218 06	216			
3439	$C_{10}H_6N_2O_4$	1, 6-Dinitronaphthalene	218 06	162			
3440	$C_{10}H_6N_2O_4$	1, 7-Dinitronaphthalene	218 06	156			
3441	$C_{10}H_6N_2O_4$	1, 8-Dinitronaphthalene	218.06	170			
3442	$C_{10}H_6N_2O_5$	2, 4-Dinitro-α-naphthol	234 06	138			
3443	$C_{10}H_6N_2O_5$	4, 5-Dinitro-α-naphthol	234.06	230 d.			
3444	$C_{10}H_6N_2O_5$	4, 8-Dinitro-α-naphthol	234 06	235 d.			
3445	$C_{10}H_6N_2O_5$	1, 6-Dinitro-β-naphthol	234.06	195			
3446	$C_{10}H_6N_2O_5$	1, 8-Dinitro-β-naphthol	234 06	198			
3447	$C_{10}H_6O_2$	1, 2-Naphthoquinone	158.05	120 d.			
3448	$C_{10}H_6O_2$	1, 4-Naphthoquinone	158 05	125			
3449	$C_{10}H_6O_2$	2, 6-Naphthoquinone	158 05	135			
3450	$C_{10}H_6O_8$	Mellophanic acid	254 05	238			
3451	$C_{10}H_6O_8$	Prehnitic acid	254.05	237 d.			
3452	$C_{10}H_6O_8$	Pyromellitic acid	254 05	264			
3453	$C_{10}H_7Br$	α-Bromonaphthalene	206.97	5	281.1	1 476	799
3454	$C_{10}H_7Br$	β-Bromonaphthalene	206 97	59	282	1.605^0	
3455	$C_{10}H_7Cl$	α-Chloronaphthalene	162.51		258	1.191	795
3456	$C_{10}H_7Cl$	β-Chloronaphthalene	162 51	56	264.3	$1.138_4^{70.7}$	1102
3457	$C_{10}H_7ClO$	2-Chloro-α-naphthol	178 51	70			
3458	$C_{10}H_7ClO$	4-Chloro-α-naphthol	178.51	117			
3459	$C_{10}H_7ClO$	5-Chloro-α-naphthol	178 51	131.5			
3460	$C_{10}H_7ClO$	6-Chloro-α-naphthol	178 51	94			
3461	$C_{10}H_7ClO$	7-Chloro-α-naphthol	178.51	123			
3462	$C_{10}H_7ClO$	1-Chloro-β-naphthol	178.51	71			
3463	$C_{10}H_7ClO$	5-Chloro-β-naphthol	178 51	128			
3464	$C_{10}H_7ClO$	6-Chloro-β-naphthol	178.51	115			
3465	$C_{10}H_7ClO$	7-Chloro-β-naphthol	178.51	126.5			
3466	$C_{10}H_7ClO$	8-Chloro-β-naphthol	178.51	101	308		
3467	$C_{10}H_7ClO_2S$	Naphthalene-1-sulfonechloride	226.58	68	195^{12}		
3468	$C_{10}H_7ClO_2S$	Naphthalene-2-sulfonechloride	226 58	76	201^{13}		
3469	$C_{10}H_7F$	α-Fluoronaphthalene	146.05		216 5	1 135°	
3470	$C_{10}H_7F$	β-Fluoronaphthalene	146.05	59	212 5		
3471	$C_{10}H_7IO$	1-Iodo-β-naphthol	269.99	94.5			
3472	$C_{10}H_7NO$	Cinnamyl cyanide $C_6H_5CH:CH_2COCN.$	157.06	115			
3473	$C_{10}H_7NO_2$	α-Nitronaphthalene	173.06	58.8	304	1.331^4	
3474	$C_{10}H_7NO_2$	β-Nitronaphthalene	173.06	79	165^{14}		
3475	$C_{10}H_7NO_2$	2-Nitroso-α-naphthol	173.06	152			
3476	$C_{10}H_7NO_2$	4-Nitroso-α-naphthol	173 06	194			
3477	$C_{10}H_7NO_2$	1-Nitroso-β-naphthol	173.06	109.5			
3478	$C_{10}H_7NO_2$	Cinchoninic acid	173 06	254			
3479	$C_{10}H_7NO_2$	Quinaldinic acid	173.06	156			
3480	$C_{10}H_7NO_2$	Quinoline-3-carboxylic acid	173 06	275			
3481	$C_{10}H_7NO_2$	Quinoline-6-carboxylic acid	173 06	292			
3482	$C_{10}H_7NO_2$	Quinoline-7-carboxylic acid	173 06	249			
3483	$C_{10}H_7NO_2$	Quinoline-8-carboxylic acid	173 06	187.5			
3484	$C_{10}H_7NO_3$	α-Kynurenic acid	189 06	283			
3485	$C_{10}H_7NO_3$	2-Nitro-α-naphthol	189 06	128			
3486	$C_{10}H_7NO_3$	3-Nitro-α-naphthol	189 06	168			
3487	$C_{10}H_7NO_3$	4-Nitro-α-naphthol	189.06	164			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3488	C₁₀H₇NO₃	5-Nitro-α-naphthol..... . .	189 06	171			
3489	C₁₀H₇NO₃	1-Nitro-β-naphthol..	189 06	103			
3490	C₁₀H₇NO₃	5-Nitro-β-naphthol..	189 06	147			
3491	C₁₀H₇NO₃	6-Nitro-β-naphthol. ..	189 06	158			
3492	C₁₀H₇NO₃	8-Nitro-β-naphthol.	189 06	145			
3493	C₁₀H₇NO₄	Indoledicarboxylic acid..	205 06	>250 d			
3494	C₁₀H₈	Naphthalene C₁₀H₈... .	128 06	80 1	217 9	1 145	1143
3495	C₁₀H₈Cl₄	Naphthalenetetrachloride	269 89	182			
3496	C₁₀H₁₀IN	Quinoline methiodide C₉H₇N.CH₃I	271 02	133			
3497	C₁₀H₈N₂	2, 3'-Dipyridyl..	156 08		289		
3498	C₁₀H₈N₂	3, 3'-Dipyridyl	156 08	68	296 5	1 164	
3499	C₁₀H₈N₂	4, 4'-Dipyridyl	156 08	112	304 8		
3500	C₁₀H₈N₂˙	Nicotelline . . .	156 08	148	<300		
3501	C₁₀H₈N₂O₂	3-Nitro-α-naphthylamine	188 08	137			
3502	C₁₀H₈N₂O₂	6-Nitro-α-naphthylamine	188 08	143			
3503	C₁₀H₈N₂O₂	7-Nitro-α-naphthylamine .	188 08	122			
3504	C₁₀H₈N₂O₂	1-Nitro-β-naphthylamine	188 08	127			
3505	C₁₀H₈N₂O₂	5-Nitro-β-naphthylamine	188 08	143			
3506	C₁₀H₈N₂O₂	8-Nitro-β-naphthylamine	188 08	105			
3507	C₁₀H₈O	α-Naphthol C₁₀H₇OH .	144 06	96	280	1 099⁹⁶·³	1126
3508	C₁₀H₈O	β-Naphthol C₁₀H₇OH	144 06	122	286	1 217⁴	1333
3509	C₁₀H₈O₂	1, 2-Dihydroxynaphthalene	160 06	60			
3510	C₁₀H₈O₂	1, 3-Dihydroxynaphthalene	160 06	125			
3511	C₁₀H₈O₂	1, 4-Dihydroxynaphthalene	160 06	176			
3512	C₁₀H₈O₂	1, 5-Dihydroxynaphthalene	160 06	258			
3513	C₁₀H₈O₂	1, 6-Dihydroxynaphthalene	160 06	138			
3514	C₁₀H₈O₂	1, 7-Dihydroxynaphthalene	160 06	178			
3515	C₁₀H₈O₂	1, 8-Dihydroxynaphthalene	160 06	140			
3516	C₁₀H₈O₂	2, 3-Dihydroxynaphthalene	160 06	159			
3517	C₁₀H₈O₂	2, 6-Dihydroxynaphthalene	160 06	216			
3518	C₁₀H₈O₂	2, 7-Dihydroxynaphthalene	160 06	190			
3519	C₁₀H₈O₂S	Naphthalene-1-sulfinic acid	192 13	85			
3520	C₁₀H₈O₂S	Naphthalene-2-sulfinic acid .	192 13	105			
3521	C₁₀H₈O₃	1, 4, 5-Trihydroxynaphthalene	176 06	170			
3522	C₁₀H₈O₃	1, 3, 6-Trihydroxynaphthalene	176 06	97			
3523	C₁₀H₈O₃	2-Benzoylacrylic acid....	176 06	99			
3524	C₁₀H₈O₃S	Naphthalene-1-sulfonic acid ..	208 13	90			
3525	C₁₀H₈O₃S	Naphthalene-2-sulfonic acid	208 13	102			
3526	C₁₀H₈O₄	Anemonin .	192 06	189 s. d.	300 d.		
3527	C₁₀H₈O₄	o-Carboxycinnamic acid	192 06	175			
3528	C₁₀H₈O₄	Furoin	192 06	135			
3529	C₁₀H₈O₄	β-Methylesculetin .	192 06	204			
3530	C₁₀H₈O₄	Scopoletin . ..	192 06	204			
3531	C₁₀H₈O₄	1, 4, 5, 6-Tetrahydroxynaphthalene	192 06	154			
3532	C₁₀H₈O₄S	α-Naphthol-2-sulfonic acid	224 13	<250			
3533	C₁₀H₈O₄S	α-Naphthol-4-sulfonic acid	224 13	170 d.			
3534	C₁₀H₈O₄S	α-Naphthol-5-sulfonic acid	224 13	120			
3535	C₁₀H₈O₄S	α-Naphthol-8-sulfonic acid	224 13	107			
3536	C₁₀H₈O₄S	β-Naphthol-6-sulfonic acid	224 13	125			
3537	C₁₀H₈O₄S	β-Naphthol-7-sulfonic acid	224 13	89			
3538	C₁₀H₈O₅	Fraxetin......... ...	208 06	227			
3539	C₁₀H₈O₆S₂	Naphthalene-1, 5-disulfonic acid	288.19	d.			1303
3540	C₁₀H₈O₆S₂	Naphthalene-1, 6-disulfonic acid	288.19	125 d.			1271
3541	C₁₀H₈O₇	Cotarnic acid...... ...	240 06	178			
3542	C₁₀H₈S	α-Thionaphthol C₁₀H₇SH .	160.13		285 d.	1.146²⁵	
3543	C₁₀H₈S	β-Thionaphthol C₁₀H₇SH	160.13	81	288 s. d.	1.550	
3544	C₁₀H₉Cl₃O₂	Chloralacetophenone......	267.44	77			
3545	C₁₀H₉N	3-Methylquinoline... .	143.08	14	250	1.074	
3546	C₁₀H₉N	4-Methylquinoline (Lepidine)	143.08		262	1.086	
3547	C₁₀H₉N	6-Methylquinoline... .	143.08		255	1.066	1003
3548	C₁₀H₉N	7-Methylquinoline.... .	143.08		252.5	1.072	788
3549	C₁₀H₉N	8-Methylquinoline......... .	143.08		143¹⁴	1.073	789
3550	C₁₀H₉N	α-Naphthylamine C₁₀H₇NH₂....... .	143.08	50	301	1.131	1080

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3551	$C_{10}H_9N$	β-Naphthylamine $C_{10}H_7NH_2$	143 08	110 2	306.1	1.061$^{22}_{4}$	
3552	$C_{10}H_9NO$	3-Amino-β-naphthol	159 08	234			
3553	$C_{10}H_9NO$	7-Amino-β-naphthol	159 08	163			
3554	$C_{10}H_9NO$	2-Hydroxyquinaldine	159 08	205			
3555	$C_{10}H_9NO$	4-Hydroxyquinaldine	159 08	231			
3556	$C_{10}H_9NO$	6-Hydroxyquinaldine	159 08	213			
3557	$C_{10}H_9NO$	7-Hydroxyquinaldine	159 08	234			
3558	$C_{10}H_9NO$	8-Hydroxyquinaldine	159 08	74	267		
3559	$C_{10}H_9NO$	Echinopsine	159 08	152			
3560	$C_{10}H_9NO_2$	α-Scatolecarboxylic acid	175 08	165			
3572	$C_{10}H_9N_4O_4$	Amilalloxan	235 09	248 d.			
3573	$C_{10}H_{10}$	1, 2-Dihydronaphthalene	130 08	−9	84.5^{16}	0.997	
3574	$C_{10}H_{10}$	1, 4-Dihydronaphthalene	130 08	15 5	212	0.998	844
3575	$C_{10}H_{10}$	1-Ethyl-2-phenylacetylene	130 08		203	0 923	
3576	$C_{10}H_{10}$	Phenylcrotonylene $C_6H_5CH.CHC_2H_5$	130 08		190		
3578	$C_{10}H_{10}Cl_3NO_3$	Chloral-p-acetaminophenol	298 46	160 d.			
3579	$C_{10}H_{10}NO_4$	Oxycannabin	208 09	182			
3580	$C_{10}H_{10}N_2$	Naphthylene-1, 2-diamine	158 09	96			
3581	$C_{10}H_{10}N_2$	Naphthylene-1, 4-diamine	158 09	120			
3582	$C_{10}H_{10}N_2$	Naphthylene-1, 5-diamine	158 09	189 5			
3583	$C_{10}H_{10}N_2$	1, 6-Naphthylenediamine	158 09	77 5		1.147$^{99}_{4}$	1137
3584	$C_{10}H_{10}N_2$	1, 8-Naphthylenediamine	158 09	66 5		1.127$^{99}_{4}$	1135
3585	$C_{10}H_{10}N_2O$	N-Phenyl-3-methylpyrazolone	174 09	127	191^{17}		1287
3586	$C_{10}H_{10}N_2O_4S$	N-Sulfophenyl-3-methylpyrazolone	254 16	320 d.			
3587	$C_{10}H_{10}O$	Benzylideneacetone	146 08	42	262	1 008	1068
3588	$C_{10}H_{10}O$	1, 2-Dihydro-β-naphthol	146 08	35	164^{28}		
3589	$C_{10}H_{10}O_2$	cis-Isosafrol	162 08	> −18	243	1.117$^{15}_{4}$	868
3590	$C_{10}H_{10}O_2$	trans-Isosafrol	162 08		248	1 123$^{15}_{4}$	869
3591	$C_{10}H_{10}O_2$	Safrol $CH_2 O_2 C_6H_3C_3H_5$	162 08	11	234 5	1 096	812
3592	$C_{10}H_{10}O_2$	Benzoylpropionaldehyde	162 08		244 4	0 998^{15}	
3593	$C_{10}H_{10}O_2$	Benzoylacetone $C_6H_5COCH_2COCH_3$	162 08	61	262	1 090^{60}	1106
3594	$C_{10}H_{10}O_2$	1-Benzylacrylic acid $CH_2.C(C_7H_7)CO_2H$	162 08	69			
3595	$C_{10}H_{10}O_2$	1-Benzylidenepropionic acid	162 08	74	288		
3596	$C_{10}H_{10}O_2$	2-Benzylidenepropionic acid	162 08	86	302		
3597	$C_{10}H_{10}O_2$	3-Phenylcrotonic acid	162 08	65			
3598	$C_{10}H_{10}O_2$	Allyl benzoate $C_6H_5CO_2C_3H_5$	162 08		230	1 058$^{15}_{15}$	
3599	$C_{10}H_{10}O_2$	Benzyl acrylate $C_2H_3CO_2CH_2C_6H_5$	162 08		110^9	1 069$^8_4$	
3600	$C_{10}H_{10}O_2$	Methyl cinnamate	162 08	36	259 6	1 042^{36}	973
3601	$C_{10}H_{10}O_2$	Phenylvinyl acetate	162 08		121^{10}	1 065	999
3602	$C_{10}H_{10}O_3$	o-Coniferylaldehyde	178 08	131			
3603	$C_{10}H_{10}O_3$	p-Coniferylaldehyde	178 08	82 5			
3604	$C_{10}H_{10}O_3$	m-Methoxycinnamic acid	178 08	115			
3605	$C_{10}H_{10}O_3$	p-Methoxycinnamic acid	178 08	169			
3606	$C_{10}H_{10}O_4$	Methyl benzoylacetate	178 08		265 d.	1 158	712
3607	$C_{10}H_{10}O_4$	1-Benzoyllactic acid	194 08	112			
3608	$C_{10}H_{10}O_4$	Benzylmalonic acid	194 08	117			
3609	$C_{10}H_{10}O_4$	Ferulic acid	194.08	169			
3610	$C_{10}H_{10}O_4$	Hesperetinic acid	194 08	228			
3611	$C_{10}H_{10}O_4$	o-Phenylenediacetic acid	194.08	150			
3612	$C_{10}H_{10}O_4$	m-Phenylenediacetic acid	194 08	170			
3613	$C_{10}H_{10}O_4$	p-Phenylenediacetic acid	194 08	241			
3614	$C_{10}H_{10}O_4$	Dimethyl isophthalate	194 08	68			
3615	$C_{10}H_{10}O_4$	Dimethyl o-phthalate	194 08		282	1 189$^{25}_{25}$	
3616	$C_{10}H_{10}O_4$	Dimethyl terephthalate	194 08	140	>300		
3617	$C_{10}H_{10}O_4$	Ethyl hydrogen o-phthalate	194 08	48			
3618	$C_{10}H_{10}O_4$	Hydroquinone diacetate	194 08	124			
3619	$C_{10}H_{10}O_4$	Methyl acetylsalicylate	194 08	54			
3620	$C_{10}H_{10}O_4$	Resorcinol diacetate	194.08		278 s. d.		
3621	$C_{10}H_{10}O_4$	Meconin	194 08	101	155		
3622	$C_{10}H_{10}O_4$	Salacetol $o-HOC_6H_4CO_2CH_2COCH_3$	194 08	71			
3623	$C_{10}H_{10}O_5$	Larixinic acid	210 08	153			
3624	$C_{10}H_{10}O_5$	Opianic acid	210 08	150			1333
3625	$C_{10}H_{10}O_6$	Apiolic acid	226.08	175			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3626	$C_{10}H_{10}O_6$	Hemipinic acid .	226 08	186			
3627	$C_{10}H_{11}NO_2$	Acetoacetanilide	177 09	85			
3628	$C_{10}H_{11}NO_2$	Diacetanilide $(CH_3CO)_2N.C_6H_5$	177 09	37	142[11]		
3629	$C_{10}H_{11}NO_2$	p-Diacetylaminophenol	193 09	118			
3630	$C_{10}H_{11}NO_3$	Ethyl oxanilate	193 09	67	300		
3631	$C_{10}H_{11}NO_3$	Methyl hippurate	193 09	80 5			
3632	$C_{10}H_{11}NO_3$	dl-Benzoylalanine	193 09	166			
3635	$C_{10}H_{11}NO_4$	Benzacetin	209 09	205			
3636	$C_{10}H_{11}N_3O_4$	4-Nitro-1, 3-diacetylphenylenediamine	237 11	246			
3637	$C_{10}H_{12}$	1, 2, 3, 4-Tetrahydronaphthalene	132 09		207 2	0 971	931
3638	$C_{10}H_{12}$	5, 6, 7, 8-Tetrahydronaphthalene	132 09	- 30	207	0 975	930
3639	$C_{10}H_{12}$	β-Phenyl-β-butylene	132 09		189	0 901[21]	966
3640	$C_{10}H_{12}Br_2O$	2, 4-Dibromothymol	307 92	4	175[15]	1.659[17 4]	
3641	$C_{10}H_{12}Br_2O_2$	Isoeugenol-1, 2-dibromide	323 92	102			
3642	$C_{10}H_{12}N_2$	Isonicoteine	160 11		293	1 098	760
3643	$C_{10}H_{12}N_2$	Nicoteine	160 11		267	1 078[12]	
3643 1	$C_{10}H_{12}N_2O$	1-Allyl-2-phenylurea	176 11	115 5			
3644	$C_{10}H_{12}N_2O_2$	Diacetyl-o-phenylenediamine	192 11	186			
3645	$C_{10}H_{12}N_2O_2$	Diacetyl-m-phenylenediamine	192 11	191			
3646	$C_{10}H_{12}N_2O_2$	Diacetyl-p-phenylenediamine	192 11	160			
3647	$C_{10}H_{12}N_2O_4$	5, 5-Diallylbarbituric acid	208 11	171			
3648	$C_{10}H_{12}O$	p-Anethol p-$CH_3OC_6H_4CH$ $CHCH_3$	148 09	22 5	235 3	0 986	1044
3649	$C_{10}H_{12}O$	1, 2, 3, 4-Tetrahydro-α-naphthol	148 09		140[17]	1 090	917
3650	$C_{10}H_{12}O$	5, 6, 7, 8-Tetrahydro-α-naphthol	148 09	68	265 3		
3651	$C_{10}H_{12}O$	1, 2, 3, 4-Tetrahydro-β-naphthol	148 09		265 5	1 071	
3652	$C_{10}H_{12}O$	5, 6, 7, 8-Tetrahydro-β-naphthol	148 09	57 5	276		
3653	$C_{10}H_{12}O$	Benzyl allyl ether $C_6H_5CH_2OC_3H_5$	148 09		204		
3654	$C_{10}H_{12}O$	Ethyl styryl ether C_6H_5CH $CHOC_2H_5$	148 09		226	0.982	893
3655	$C_{10}H_{12}O$	Methyl chavicyl ether	148 09		216	0.965	676
3656	$C_{10}H_{12}O$	Cumic aldehyde $(CH_3)_2CHC_6H_4CHO$.	148 09		235	0 978	698
3657	$C_{10}H_{12}O$	Mesitylenic aldehyde	148 09		237		
3658	$C_{10}H_{12}O$	3, 4, 5-Trimethylbenzaldehyde	148 09	52			
3659	$C_{10}H_{12}O$	Benzyl acetone $C_6H_5(CH_2)_2COCH_3$.	148 09		236	0 989[24]	
3660	$C_{10}H_{12}O$	Ethyl benzyl ketone	148 09		230 2	1.002[0]	
3661	$C_{10}H_{12}O$	Phenyl isopropyl ketone	148 09		217	0 984	879
3662	$C_{10}H_{12}O$	Phenyl n-propyl ketone	148 09	11	232 3	0.988	
3663	$C_{10}H_{12}O$	p-Tolylacetone	148 09	51	233		
3664	$C_{10}H_{12}O$	p-Tolyl ethyl ketone	148 09		239[763]	0 993	690
3665	$C_{10}H_{12}O_2$	3, 5, 6-Trimethyl-2-hydroxybenzaldehyde	164 09	106			
3666	$C_{10}H_{12}O_2$	Eugenol . . .	164 09		253	1.071[13]	841
3667	$C_{10}H_{12}O_2$	Isoeugenol	164 09	- 10	267 5	1.080	936
3668	$C_{10}H_{12}O_2$	Cumic acid $(CH_3)_2CHC_6H_4CO_2H$	164 09	116 5		1 163[4]	
3669	$C_{10}H_{12}O_2$	o-Isopropylbenzoic acid	164 09	51			
3670	$C_{10}H_{12}O_2$	3-Phenylbutyric acid $C_6H_5(CH_2)_2CO_2H$	164 09	47 5	290		
3671	$C_{10}H_{12}O_2$	o-Propylbenzoic acid $o-C_3H_7C_6H_4CO_2H$.	164 09	58	273		
3672	$C_{10}H_{12}O_2$	p-Propylbenzoic acid	164 09	141			
3673	$C_{10}H_{12}O_2$	3, 4, 5-Trimethylbenzoic acid	164 06	215			
3674	$C_{10}H_{12}O_2$	2, 4, 5-Trimethylbenzoic acid	164 09	149 5			
3675	$C_{10}H_{12}O_2$	2, 4, 6-Trimethylbenzoic acid	164 09	152			
3676	$C_{10}H_{12}O_2$	Benzyl propionate	164 09		220	1 036[17 5]	
3677	$C_{10}H_{12}O_2$	Ethyl phenylacetate $C_6H_5CH_2CO_2C_2H_5$	164 09		226	1 031	589
3678	$C_{10}H_{12}O_2$	Ethyl o-toluate $CH_3C_6H_4CO_2C_2H_5$.	164 09		221 3	1.033	629
3679	$C_{10}H_{12}O_2$	Ethyl m-toluate $CH_3C_6H_4CO_2C_2H_5$	164 09		226.4	1.028	624
3680	$C_{10}H_{12}O_2$	Ethyl p-toluate $CH_3C_6H_4CO_2C_2H_5$	164 09		228	1.026	636
3681	$C_{10}H_{12}O_2$	Isopropyl benzoate . . .	164 09		218 5	1.017[15]	
3681.1	$C_{10}H_{12}O_2$	d-Methylbenzylcarbinyl formate	164 09		110[19]	1.027[22]	595
3682	$C_{10}H_{12}O_2$	Methyl hydrocinnamate	164 09		239	1.018[19]	
3683	$C_{10}H_{12}O_2$	Phenyl n-butyrate $C_3H_7CO_2C_6H_5$	164 09		228	1.027[15]	
3684	$C_{10}H_{12}O_2$	n-Propyl benzoate $C_6H_5CO_2C_3H_7$	164 09	- 51 6	231 2	1 027	
3685	$C_{10}H_{12}O_2$	Thymoquinone	164.09	45 5	232		
3686	$C_{10}H_{12}O_3$	Coniferyl alcohol	180.09	74			
3687	$C_{10}H_{12}O_3$	Benzyl lactate	180.09		130[6]		1025
3688	$C_{10}H_{12}O_3$	Ethyl anisate p-$CH_3OC_6H_4CO_2C_2H_5$.	180.09	7 8	263	1 106	680 .

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3689	$C_{10}H_{12}O_3$	Ethyl mandelate . . .	180 09	34	255		
3690	$C_{10}H_{12}O_3$	Propyl salicylate o-HOC$_6$H$_4$CO$_2$C$_3$H$_7$	180 09		240	1.099^{15}	
3691	$C_{10}H_{12}O_4$	Cantharic acid	196 09	278			
3692	$C_{10}H_{12}O_4$	Ethyl vanillate	196 09	44	293		
3693	$C_{10}H_{12}O_4$	Cantharidin	196 09	212			
3694	$C_{10}H_{12}O_4$	Guaiacyl methyl glycollate .	196 09		156^{15}	1.180	
3695	$C_{10}H_{12}O_4$	Sparassol	196 09	68			
3696	$C_{10}H_{12}O_5$	Asaronic acid	212 09	144	300		
3697	$C_{10}H_{12}O_5$	Glycerol monosalicylate	212 09	76		1.366	
3698	$C_{10}H_{12}O_6$	β-Anemonic acid	228 09	189			
3699	$C_{10}H_{13}ClO$	4-Chlorothymol	184 56	64			
3700	$C_{10}H_{13}ClO$	6-Chlorothymol	184 56	64			
3701	$C_{10}H_{13}N$	Kairoline (1-Methyl-1, 2, 3, 4-tetrahydro-quinoline)	147 11		245.5	1 021	1005
3702	$C_{10}H_{13}N$	5, 6, 7, 8-Tetrahydro-α-naphthylamine	147 11		276.8	1 054$^{18.1}$	1006
3703	$C_{10}H_{13}N$	5, 6, 7, 8-Tetrahydro-β-naphthylamine	147 11	38	278 5	1 029$^{22.2}$	986
3704	$C_{10}H_{13}NO$	o-Acetylmethyltoluidine	163 11	56			
3705	$C_{10}H_{13}NO$	p-Acetylmethyltoluidine	163 11	80			
3706	$C_{10}H_{13}NO$	N-Butyranilide C$_6$H$_5$NHOCC$_3$H$_7$. . .	163 11	92	189^{15}		
3707	$C_{10}H_{13}NO$	3, 5-Dimethylacetanilide .	163 11	174			
3708	$C_{10}H_{13}NO$	ω-Dimethylaminoacetophenone	163 11	59			
3709	$C_{10}H_{13}NO$	N-Ethylacetanilide . .	163 11	54 5	259	0.994$^{40}_{4}$	
3710	$C_{10}H_{13}NO$	Thalline	163 11	43	283 8		
3711	$C_{10}H_{13}NO_2$	1-Anilinobutyric acid .	179 11	141			
3712	$C_{10}H_{13}NO_2$	Propyl p-aminobenzoate	179 11	76			
3713	$C_{10}H_{13}NO_2$	o-Acetphenetidine	179 11	79	<250		
3714	$C_{10}H_{13}NO_2$	m-Acetphenetidine.	179 11	96			
3715	$C_{10}H_{13}NO_2$	2-Nitrocymene	179.11		152^{15}	1 085^{15}	
3716	$C_{10}H_{13}NO_2$	Phenacetin C$_2$H$_5$OC$_6$H$_4$NHCOCH$_3$	179 11	135	d.		1246
3717	$C_{10}H_{13}NO_3$	Damascenine	195 11	27	168		
3718	$C_{10}H_{13}NO_3$	2-Nitrothymol	195 11	119			
3719	$C_{10}H_{13}NO_3$	4-Nitrothymol	195 11	142			
3720	$C_{10}H_{13}NO_3$	Ratanhine	195 11	252			
3721	$C_{10}H_{13}NO_4$	Surinamine (N-Methyltyrosine)	195 11	280 d.			
3722	$C_{10}H_{13}N_3O_4$	2, 4-Dinitro-N-diethylaniline. .	239 12	80			
3723	$C_{10}H_{13}N_3O_5$	Vernine	283 14	240			
3724	$C_{10}H_{14}$	n-Butylbenzene CH$_3$(CH$_2$)$_3$C$_6$H$_5$.	134 11		180	0 862	554
3725	$C_{10}H_{14}$	$sec.$-Butylbenzene C$_2$H$_5$(CH$_3$)CHC$_6$H$_5$. .	134 11		175	0 860	550
3726	$C_{10}H_{14}$	$tert.$-Butylbenzene (CH$_3$)$_3$C.C$_6$H$_5$.	134 11		168.7	0 867	582
3727	$C_{10}H_{14}$	o-Cymene o-CH$_3$(CH$_2$)$_2$C$_6$H$_4$CH$_3$. . . .	134.11		157	0 858^{15}	601
3728	$C_{10}H_{14}$	m-Cymene m-CH$_3$(CH$_3$)$_2$C$_6$H$_4$CH$_3$	134 11	> −25	175	0 860	559
3728 1	$C_{10}H_{14}$	p-Cymene p-CH$_3$(CH$_3$)$_2$C$_6$H$_4$CH$_3$	134 11	−73 5	176	0 857	1022
3729	$C_{10}H_{14}$	o-Diethylbenzene o-(C$_2$H$_5$)$_2$C$_6$H$_4$ . .	134 11	< −20	184 5	0 866	
3730	$C_{10}H_{14}$	m-Diethylbenzene m-(C$_2$H$_5$)$_2$C$_6$H$_4$.	134 11	< −20	182	0 860	
3731	$C_{10}H_{14}$	p-Diethylbenzene p-(C$_2$H$_5$)$_2$C$_6$H$_4$	134 11	−35	183	0 865	569.1
3732	$C_{10}H_{14}$	1, 2, 4, 5-Tetramethylbenzene	134 11	80	195	0 838$^{81.3}_{4}$	1273
3733	$C_{10}H_{14}$	4-Ethyl-m-xylene C$_2$H$_5$C$_6$H$_3$(CH$_3$)$_2$.	134 11	< −20	183	0.878	
3734	$C_{10}H_{14}$	5-Ethyl-m-xylene C$_2$H$_5$C$_6$H$_3$(CH$_3$)$_2$	134 11	< −20	185	0 861	
3735	$C_{10}H_{14}$	Hexahydronaphthalene	134 11		205 5	0 934	
3736	$C_{10}H_{14}$	Isobutylbenzene (CH$_3$)$_2$CHCH$_2$C$_6$H$_5$.	134.11		171.4	0 858^{15}	562
3739	$C_{10}H_{14}$	1, 2, 3, 5-Tetramethylbenzene	134.11		197	0 896$^{0}_{4}$	
3740	$C_{10}H_{14}$	1, 2, 3, 4-Tetramethylbenzene . .	134 11	−4	204	0.901	662
3741	$C_{10}H_{14}$	Verbenene	134.11		159	0 886^{15}	593
3742	$C_{10}H_{14}Br_2O$	d-α, α'-Dibromocamphor.	309 94	61			1209
3743	$C_{10}H_{14}ClN$	Thermin (Tetrahydro-β-naphthylamine hydrochloride).	183 57	237			
3744	$C_{10}H_{14}Cl_2O$	α-Dichlorocamphor .	221.02	96	200 d.	4 2	
3745	$C_{10}H_{14}Cl_2O$	β-Dichlorocamphor .	221 02	77			
3746	$C_{10}H_{14}N_2$	Isonicotine .	162 12	78	260 d.		
3747	$C_{10}H_{14}N_2$	Nicotine	162.12		274.3	1.009	695
3748	$C_{10}H_{14}N_2$	Nicotimine	162.12		250		
3749	$C_{10}H_{14}N_2O_2$	6-Nitroso-3-(diethylamino) phenol . .	194.12	84			
3750	$C_{10}H_{14}N_2O$	p-Nitroso-N-diethylaniline.	178.12	84			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3751	C₁₀H₁₄N₂O₂	Phenocoll p-C₂H₅OC₆H₄NHCOCH₂NH₂	194 12	100 5			
3752	C₁₀H₁₄O	Carvacrol	150 11	0 5	237.9	0 976	678
3753	C₁₀H₁₄O	d-Carvol	150 11		225	0 960	940
3754	C₁₀H₁₄O	Cuminal alcohol	150 11		246 6	0 978¹⁵	
3754 1	C₁₀H₁₄O	Methyl d-methylbenzyl carbinol	150 11		85¹¹	0 927²⁷	
3754 2	C₁₀H₁₄O	Methyl l-phenylethyl carbinol	150 11		132¹⁴	0 9767	658
3755	C₁₀H₁₄O	3-Methyl-2-hydroxyisopropylbenzene	150 11		226	0 987¹⁵·⁷	669
3756	C₁₀H₁₄O	Thymol (CH₃)₂CHC₆H₃(OH)CH₃	150 11	51 5	231 8	0 969	1170
3757	C₁₀H₁₄O	5-Methyl-2-hydroxyisopropylbenzene	150 11	36	229	0 982¹⁷·⁸	674
3758	C₁₀H₁₄O	Benzyl propyl ether C₆H₅CH₂OC₃H₇	150 11		196		
3759	C₁₀H₁₄O	n-Butyl phenyl ether C₆H₅OC₄H₉	150 11		210 3	0 950⁰	
3760	C₁₀H₁₄O	Isobutyl phenyl ether.	150 11		198	0 939¹⁴	
3761	C₁₀H₁₄O	Myrtenal (Myrtenic aldehyde)	150 11		90¹⁰	0 988	616
3762	C₁₀H₁₄O	Eucarvol .	150 11		106²⁰	0 952	845
3763	C₁₀H₁₄O	Pinocarvol	150 11		224	0 984	620
3764	C₁₀H₁₄O	d(l)-Piperitone	150 11		235	0 934ᵛᵃᶜ	542
3765	C₁₀H₁₄O	Umbellulone	150 11		220	0 958	551
3766	C₁₀H₁₄O₂	o-Diethoxybenzene o-(C₂H₅O)₂C₆H₄	166 11	45			
3767	C₁₀H₁₄O₂	Coerulignol .	166 11		246	1 049¹⁵	
3768	C₁₀H₁₄O₂	Hydroquinone diethyl ether	166 11	72			
3769	C₁₀H₁₄O₂	Resorcinol diethyl ether	166 11	12 4	235 2		
3770	C₁₀H₁₄O₂	d-Camphorquinone..	166 11	198			
3771	C₁₀H₁₄O₂	Thymohydroquinone	166 11	143	290		
3772	C₁₀H₁₄O₂	Crocetin .	166 11	101			
3773	C₁₀H₁₄O₃	dl-Camphoric anhydride	182 11	221	270		
3774	C₁₀H₁₄O₄	1, 2, 3, 5-Tetramethoxybenzene	198 11	47	271		
3775	C₁₀H₁₄O₄	Guaiamar..	198 11	75			
3776	C₁₀H₁₄O₄	Diethyl muconate	198 11	13; 62	64	0 983²²·¹	
3777	C₁₀H₁₄O₅	Pinoylformic acid	214 11	80			
3777.1	C₁₀H₁₄O₆	Diallyl tartrate	230 11		191²⁰	1 187²⁴·⁴	
3778	C₁₀H₁₅BrO	α-Bromocamphor	231 03	78	274	1 449	1252
3779	C₁₀H₁₅BrO	β-Bromocamphor	231 03	61	130¹⁰		
3780	C₁₀H₁₅Cl	Myrtenyl chloride	170 57		90¹²	1 015	586
3782	C₁₀H₁₅ClO	α-Chlorocamphor	186 57	125	220 ₈ d		
3783	C₁₀H₁₅ClO	β-Chlorocamphor	186 57	92 5	247		
3784	C₁₀H₁₅ClO	γ-Chlorocamphor	186 57	100	237 ₈ d.		
3785	C₁₀H₁₅N	n-Butylaniline C₆H₅NHC₄H₉	149 12		240 9		
3786	C₁₀H₁₅N	2-Dimethylamino-m-xylene	149 12		196 2	0 915	649
3787	C₁₀H₁₅N	4-Dimethylamino-m-xylene	149 12		232 2	0 939	730
3788	C₁₀H₁₅N	4-Dimethylamino-o-xylene	149 12		205	0 916	663
3789	C₁₀H₁₅N	Diethylaniline C₆H₅N(C₂H₅)₂	149 12	−34 4	216 27	0 934	717
3790	C₁₀H₁₅N	Isobutylaniline C₆H₅NHCH₂CH(CH₃)₂	149 12		242	0.940	
3791	C₁₀H₁₅N	Prehnidine 1, 2, 3, 4-C₆H₂(CH₃)₄	149 12	70	260		
3792	C₁₀H₁₅NO	m-Diethylaminophenol	165 12	78	278		
3793	C₁₀H₁₅NO	Ephedrine.	165 12	40	255		
3794	C₁₀H₁₅NO	Hordenine	165 12	118	174¹¹		
3795	C₁₀H₁₅NO	Pseudoephedrine	165 12	117			
3796	C₁₀H₁₅NO₃S	Diethylaniline-m-sulfonic acid	229 19	270 d.			
3797	C₁₀H₁₅N₃O₆	Pilocarpidine nitrate	257 14	137			1333
3800	C₁₀H₁₆	l-Bornylene	136 12	111	147		
3801	C₁₀H₁₆	dl-Camphene	136 12	50	160	0 822	1116
3802	C₁₀H₁₆	d(l)-Camphene	136 12	42 7	159		1074
3803	C₁₀H₁₆	Camphilene .	136 12		156	0 87¹¹	
3804	C₁₀H₁₆	d(l)-Δ⁴-Carene	136 12		167⁷⁰⁷	0 855¹⁰	1037
3805	C₁₀H₁₆	Cyclofenchene	136 12		144	0 861	445
3806	C₁₀H₁₆	Dipentene..	136 12		176	0 865¹⁴	515
3807	C₁₀H₁₆	d(l)-Fenchene	136 12		150	0 869	955
3808	C₁₀H₁₆	Fenchylene.	136 12		142	0 840	435
3809	C₁₀H₁₆	Geraniene .	136 12		164	0.843	
3810	C₁₀H₁₆	d(l)-Limonene	136 12	−96 9	177	0 842	510
3811	C₁₀H₁₆	Myrcene	136 12		167	0 802	503
3812	C₁₀H₁₆	Ocimene...	136.12		74²¹	0.799	835
3813	C₁₀H₁₆	cis-β-Octalin............ . . .	136.12		73¹⁶	0.915	984

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3814	C₁₀H₁₆	trans-β-Octalin	136 12		190	0.909¹¹	
3815	C₁₀H₁₆	d(l)-α-Phellandrene	136.12		175	0.843	983
3816	C₁₀H₁₆	β-Phellandrene	136 12		171	0.852	527
3817	C₁₀H₁₆	dl-α-Pinene	136 12	−55	154	0.878	
3818	C₁₀H₁₆	l-β-Pinene	136 12		164	0.873¹⁵	824
3819	C₁₀H₁₆	Sabinene.	136 12		165	0.842	914
3820	C₁₀H₁₆	d(l)-Sylvestrene	136 12		177	0.863	919
3821	C₁₀H₁₆	α-Terpinene	136 12		175	0.834	915
3822	C₁₀H₁₆	β-Terpinene	136 12		174	0.840	982
3823	C₁₀H₁₆	Δ¹·⁸-Terpinene	136.12		182	0.855	541
3824	C₁₀H₁₆	Terpinolene.	136 12		185	0 855	537
3825	C₁₀H₁₆	Terpinylene	136 12		175		
3826	C₁₀H₁₆	α-Thujene	136 12		151	0.830	440
3827	C₁₀H₁₆	β-Thujene	136.12		147.7	0.821	420
3828	C₁₀H₁₆ClNO	Ephedrine hydrochloride	201.59	210			
3829	C₁₀H₁₆ClNO	α-Limonene nitrosylchloride	201 60	104			
3830	C₁₀H₁₆ClNO	Pseudoephedrine hydrochloride	201.59	175			
3831	C₁₀H₁₆Cl₂	α-Camphordichloride	207 04	148			
3832	C₁₀H₁₆Cl₂	β-Camphordichloride ...	207 04	178			
3833	C₁₀H₁₆N₂	p-Aminodiethylaniline	164.14		262		
3834	C₁₀H₁₆N₂	o-Tetramethylphenylenediamine	164 14		218		
3835	C₁₀H₁₆N₂	m-Tetramethylphenylenediamine	164 14	−2	262	0 988¹⁵ ⁸	
3836	C₁₀H₁₆N₂	p-Tetramethylphenylenediamine	164 14	51	260		
3837	C₁₀H₁₆N₂O₂	α-Camphordioxime	196 14	182 d.			
3838	C₁₀H₁₆N₂O₂	γ-Camphordioxime	196 14	132			
3839	C₁₀H₁₆N₂O₃	5, 5-n-Butylethylbarbituric acid.. .	212.14	128			
3840	C₁₀H₁₆N₂O₃	5, 5-sec.-Butylethylbarbituric acid .	212 14	157			
3841	C₁₀H₁₆N₂O₃	5, 5-Dipropylbarbituric acid	212.14	145			
3842	C₁₀H₁₆N₂O₃	5, 5-Isobutylethylbarbituric acid ...	212 14	176			
3843	C₁₀H₁₆N₂O₃	5, 5-n-Propylisopropylbarbituric acid	212.14	162			
3844	C₁₀H₁₆O	Alantol	152.12		200		
3845	C₁₀H₁₆O	dl-Camphor	152 12	174			
3846	C₁₀H₁₆O	d-Camphor	152 12	179	209 1	0 990²⁵	
3847	C₁₀H₁₆O	Carvenone	152 12		233	0 926	897
3848	C₁₀H₁₆O	Caryophyllin	152 12	295			
3849	C₁₀H₁₆O	α-Citral .	152.12		229	0 893¹⁵	920
3850	C₁₀H₁₆O	β-Citral	152.12		104¹²	0 888	956
3851	C₁₀H₁₆O	Cyclocitral	152 12		114²⁹	0 957¹⁴	825
3852	C₁₀H₁₆O	d-Fenchone	152 12	6	195	0 944	839
3853	C₁₀H₁₆O	Hartin	152.12	230		1 120	
3854	C₁₀H₁₆O	Isopulegon	152 12		90¹²	0.921¹⁷ ⁵	499
3855	C₁₀H₁₆O	Myristicol	152 12		218		
3856	C₁₀H₁₆O	Myrtenol	152 12		224	0 976	581
3857	C₁₀H₁₆O	Phellandral	152 12		230	0 945	553
3858	C₁₀H₁₆O	Pinol	152 12		184	0 942	507
3859	C₁₀H₁₆O	Pulegon	152.12		224	0 937	861
3860	C₁₀H₁₆O	Sabinol. ...	152.12		209	0 943	546
3861	C₁₀H₁₆O	α-Thujone	152.12		200	0 913	827
3862	[C₁₀H₁₆O]ₓ	Urson	[152 12]ₓ	264			
3863	C₁₀H₁₆O₂	Acetylmethylheptenone	168.12	−6	234	0 945¹⁵	860
3864	C₁₀H₁₆O₂	Ascaridol	168.12		84⁵	1 008¹⁵	518
3865	C₁₀H₁₆O₂	Geranic acid .	168.12		119²⁰	0 952	544
3866	C₁₀H₁₆O₂	Hydroxycamphor.......	168.12	205			
3867	C₁₀H₁₆O₃	d(l)-Pinonic acid......	184 12	99	180¹²		
3867 1	C₁₀H₁₆O₃	dl-Pinonic acid........ .	184.12	105		1.216	
3868	C₁₀H₁₆O₄	dl-Camphoric acid	200.12	202			
3869	C₁₀H₁₆O₄	d-Camphoric acid . .	200.12	187			
3870	C₁₀H₁₆O₄	Cyclohexyl acid succinate	200.12	44			
3871	C₁₀H₁₆O₄	dl-Isocamphoric acid.	200.12	191			
3872	C₁₀H₁₆O₄	d-Methyl pinate	200.12		130⁸	1.055	
3873	C₁₀H₁₆O₅	l-Cineolic acid........	216.12	196			1325
3874	C₁₀H₁₆O₅	Diethyl acetylsuccinate	216.12		256 d.	1.081	884
3875	C₁₀H₁₇Br	d-Pinene hydrobromide...............	217.05	80			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3876	$C_{10}H_{17}Cl$	Camphene hydrochloride	172 59	156 5			
3877	$C_{10}H_{17}Cl$	cis-β-Chlorodecalin	172 59		112[15]		
3878	$C_{10}H_{17}Cl$	Fenchyl chloride	172 59		85[14]	0 983	
3879	$C_{10}H_{17}Cl$	Geranyl chloride	172 59		103[14]	0 918[15]	517
3880	$C_{10}H_{17}Cl$	Isobornyl chloride	172 59	161 5			
3881	$C_{10}H_{17}Cl$	d-Pinene hydrochloride	172 59	128	207 1		
3882	$C_{10}H_{17}N$	Camphenamine	151 14		205 5	0 940	564
3883	$C_{10}H_{17}N$	Pinylamine	151 14		207	0 940	613
3884	$C_{10}H_{17}NO$	Camphoroxime	167 14	119 5	249		
3885	$C_{10}H_{17}NO$	d-Fenchoneoxime	167 14	165	240		
3886	$C_{10}H_{17}NO_2$	l-Ecgonine methyl ester	199 14			1 147	547
3886 1	$C_{10}H_{17}NO_2$	dl-α-Pinone oxime	199 14	150		1 210	
3887	$C_{10}H_{17}NO_6$	Phaseolunatin	217 14	144			
3888	$C_{10}H_{18}$	Camphane	138 14	152	160		
3889	$C_{10}H_{18}$	Carane	138 14		50[9]	0.838^{20}_{20}	459
3890	$C_{10}H_{18}$	cis-Decahydronaphthalene	138 14	− 125	193 3	0 898	539
3891	$C_{10}H_{18}$	trans-Decahydronaphthalene	138 14		185 3	0 872	504
3892	$C_{10}H_{18}$	d-Menthene	138 14		168	1 4481	423
3893	$C_{10}H_{18}$	d-Pinane	138 14	− 45	169 4	0 839	448
3894	$C_{10}H_{18}$	Pinocamphane	138 14		164 9	0 856	477
3895	$C_{10}H_{18}$	Thujane	138 14		157	0 814	363
3896	$C_{10}H_{18}Cl_2N_2$	o-Tetramethylphenylenediamine hydrochloride	237 07	180			
3897	$C_{10}H_{18}O$	Apopinol	154 14		199	0 804[14]	
3899	$C_{10}H_{18}O$	Aurantiol	154 14		95[15]	0 869[20]	
3900	$C_{10}H_{18}O$	dl-Borneol	154 14	210 5			
3901	$C_{10}H_{18}O$	d(l)-Borneol	154 14	208 6	213 5	1 .011	
3902	$C_{10}H_{18}O$	Cineol	154 14	− 1	176 4	0 901[15]	474
3903	$C_{10}H_{18}O$	d-Citronellal	154 14		208	0 856	
3904	$C_{10}H_{18}O$	dl-Fenchyl alcohol	154 14	33	204 6	0 953	
3905	$C_{10}H_{18}O$	dl, (d)-Fenchyl alcohol	154 14	42	201	0 935[40]	
3906	$C_{10}H_{18}O$	dl, (l)-Fenchyl alcohol	154 14	47	201	0 933[40]	
3907	$C_{10}H_{18}O$	d, (l)-Fenchyl alcohol	154.14	49	209		
3908	$C_{10}H_{18}O$	Geraniol	154 14	< − 15	229	0 881	531
3909	$C_{10}H_{18}O$	dl-Isoborneol	154 14	212			
3910	$C_{10}H_{18}O$	d(l)-Isoborneol	154 14	216			
3911	$C_{10}H_{18}O$	dl-Isofenchyl alcohol	154 14		204		
3912	$C_{10}H_{18}O$	l-Isofenchyl alcohol	154 14	62	202	0 961[15]	859
3913	$C_{10}H_{18}O$	Isopulegol	154 14		102[12]	0 915	513
3913 1	$C_{10}H_{18}O$	l-Isopulegol	154 14		94[11]	0.9110	509
3914	$C_{10}H_{18}O$	Lavendol	154 14		199	0 873[15]	
3915	$C_{10}H_{18}O$	d-Linalool	154 14		198 3	0 875	480
3916	$C_{10}H_{18}O$	l-Linalool	154 14		195	0 866[15]	981
3917	$C_{10}H_{18}O$	dl-Menthone	154 14		210	0 897	441
3918	$C_{10}H_{18}O$	l-Menthone	154 14		207	0 896	
3919	$C_{10}H_{18}O$	Myrcenol	154 14		101[10]	0 901[14] [6]	840
3920	$C_{10}H_{18}O$	Nerol	154 14		225 2	0 881	
3921	$C_{10}H_{18}O$	Pinen hydrate (Homopinol)	154 14	59	205		
3922	$C_{10}H_{18}O$	dl, α-Terpineol	154 14	35	219 8	0 936	538
3923	$C_{10}H_{18}O$	d(l), α-Terpineol	154 14	40	217 7	0 919	890
3924	$C_{10}H_{18}O$	β-Terpineol	154 14	33	210 3	$0 819^{20}_{20}$	521
3925	$C_{10}H_{18}O$	γ-Terpineol	154 14	70			
3926	$C_{10}H_{18}O$	dl-Terpinen-4-ol	154 14		214	0 929	533
3927	$C_{10}H_{18}O$	d-Terpinen-4-ol (Origanol)	154 14		212	0 926	526
3928	$C_{10}H_{18}O$	Thujyl alcohol	154 14		212	0 921	923
3929	$C_{10}H_{18}O_2$	Acetylmethyl hexyl ketone	170 14	− 6	237 d.	$0 907^{21}_{21}$	
3930	$C_{10}H_{18}O_2$	d(l)-Campholic acid	170 14	107	260		
3931	$C_{10}H_{18}O_2$	d-Citronellic acid	170 14		257	0 931	
3932	$C_{10}H_{18}O_2$	9, 10-Decylenic acid	170 14	< 0	142[4]		
3933	$C_{10}H_{18}O_2$	Fencholic acid	170 14	18	255	0.970[10] [9]	462
3934	$C_{10}H_{18}O_3$	Pinol glycol	186 14	129			
3935	$C_{10}H_{18}O_3$	n-Valeric anhydride $(C_4H_9CO)_2O$	186 14		215	0.929	
3936	$C_{10}H_{18}O_3$	Isovaleric anhydride	186.14		215	0.933	229

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3937	$C_{10}H_{18}O_3$	Ethyl diethylacetoacetate	186 14		158.2	1.282	327
3938	$C_{10}H_{18}O_4$	Sebacic acid $HO_2C(CH_2)_8CO_2H$	202 14	127	294.5[100]		1161
3939	$C_{10}H_{18}O_4$	Isoamyl ethyl malonate	202 14		150[20]	0.954[22/15]	306
3940	$C_{10}H_{18}O_4$	n-Butyl isopropylmalonate	202 14		136[14]	0.974[25/15]	331
3941	$C_{10}H_{18}O_4$	Di-n-butyl oxalate $(CO_2C_4H_9)_2$	202 14		243.4	1.0108	
3942	$C_{10}H_{18}O_4$	Diisobutyl oxalate	202 14		229	1 002[14]	
3943	$C_{10}H_{18}O_4$	Dipropyl succinate	202 14		250 8	1 006[15]	
3944	$C_{10}H_{18}O_5$	Dipropyl malate	218 14	10 5	151[10]	1.075	366
3945	$C_{10}H_{18}O_6$	Dipropyl d-tartrate $[HOCHCO_2C_3H_7]_2$	234 11		303	1.139	
3945 1	$C_{10}H_{18}O_6$	Di-sec.-propyl tartrate	234 11		158[16]	1 116[13.7]	
3946	$C_{10}H_{18}O_9$	Arabin	282 14	260			
3947	$C_{10}H_{19}Cl$	sec.-Menthyl chloride	174 60		215	0 941	485
3948	$C_{10}H_{19}Cl$	tert -Menthyl chloride	174 60		94[14 5]	0 948	488
3949	$C_{10}H_{19}N$	Bornylamine	153 15	163	200		
3950	$C_{10}H_{19}N$	Camphylamine	153 15		198		
3951	$C_{10}H_{19}N$	l-Fenchylamine	153 15		195	0 910[22]	
3952	$C_{10}H_{19}N$	Geranylamine	153 15		105[19]	0.829[25]	511
3953	$C_{10}H_{19}NO$	Lupinine	169 15	68	257		
3954	$C_{10}H_{19}NO_3$	Sebamic acid	201 15	170			
3955	$C_{10}H_{20}$	α-Decylene $CH_2.CH(CH_2)_7CH_3$	140 15		172	0.763[0]	912
3956	$C_{10}H_{20}$	γ-Decylene $C_3H_7CH CHC_6H_{11}$	140 15		161		
3957	$C_{10}H_{20}$	2, 3-Dimethyl-2-octene	140 15		162[650]	0.748	
3958	$C_{10}H_{20}$	2, 6-Dimethyl-1(2)-octene	140 15		169	0.789[0]	993
3959	$C_{10}H_{20}$	o-Menthane	140 15		171	0.814	965
3960	$C_{10}H_{20}$	m-Menthane	140 15		168 2	0 790	387
3961	$C_{10}H_{20}$	p-Menthane	140 15		170	0 793	358
3962	$C_{10}H_{20}$	2-Methyl-5-ethyl-5-heptene	140 15		158 4	0 761[0]	302
3963	$C_{10}H_{20}$	3, 3, 5-Trimethyl-4-heptene	140 15		157.5	0.788[0/0]	
3964	$C_{10}H_{20}ClNO$	Lupinine hydrochloride	205 62	213			1244
3965	$C_{10}H_{20}N_2O_6$	Lycetol (Dimethylpiperazine tartrate)	264 17	250			
3966	$C_{10}H_{20}O$	α-Carvacromenthol	156 15		219		
3967	$C_{10}H_{20}O$	β-Carvacromenthol	156 15		222	0.918[0]	
3968	$C_{10}H_{20}O$	d-Citronellol	156 15		221.7	0 857[15]	410
3969	$C_{10}H_{20}O$	l-Citronellol	156 15		114[15]	0 861	464
3970	$C_{10}H_{20}O$	d-Isomenthol	156 15	83			
3971	$C_{10}H_{20}O$	o-Menthan-2-ol	156 15		95[25]		
3972	$C_{10}H_{20}O$	p-Menthan-8-ol	156 15	36	207 4		
3973	$C_{10}H_{20}O$	l-α-Menthol	156 15	42 5	212	0 890[15/15]	1168
3974	$C_{10}H_{20}O$	l-β-Menthol	156 15	35 5	212	0 890[15/15]	
3974 1	$C_{10}H_{20}O$	l-Neomenthol	156 15	< −15	105[21]	0.8995	473
3975	$C_{10}H_{20}O$	n-Capric aldehyde $CH_3(CH_2)_8CHO$.	156 15		209 2	0.828[15]	307
3976	$C_{10}H_{20}O$	Isocapric aldehyde	156 15		169 6	0.828[0]	
3977	$C_{10}H_{20}O$	Isopropyl n-hexyl ketone	156 15		210	0.841[17]	
3978	$C_{10}H_{20}O$	Methyl n-octyl ketone $CH_3COC_8H_{17}$	156 15	3.5	211	0 825	
3978 1	$C_{10}H_{20}O$	Propyl hexyl ketone $C_3H_7COC_6H_{13}$	156 15	−9	207	0 824	
3979	$C_{10}H_{20}O_2$	cis-Terpine	172 15	104.7	258		
3980	$C_{10}H_{20}O_2$	trans-Terpine	172.15	158	265		
3981	$C_{10}H_{20}O_2$	n-Capric acid $CH_3(CH_2)_8CO_2H$	172.15	31	268 4	0 895[30]	1038
3981 1	$C_{10}H_{20}O_2$	Di-n-butylacetic acid	172.15		140[16]	0 898[18 4]	
3982	$C_{10}H_{20}O_2$	n-Amyl valerate $C_4H_9CO_2C_5H_{11}$	172 15		203 7	0 881[0]	213
3983	$C_{10}H_{20}O_2$	n-Butyl caproate $C_5H_{11}CO_2C_4H_9$	172.15		204 3	0 882[9/9]	
3984	$C_{10}H_{20}O_2$	Ethyl n-caprylate $C_7H_{15}CO_2C_2H_5$	172.15	−44 8	205 8	0 878[17]	
3985	$C_{10}H_{20}O_2$	n-Heptyl propionate $C_2H_5CO_2C_7H_{15}$.	172.15		208	0 885[0]	
3986	$C_{10}H_{20}O_2$	Isoamyl isovalerate	172.15		194	0 870[0]	198
3987	$C_{10}H_{20}O_2$	Methyl pelargonate $C_8H_{17}CO_2CH_3$	172 15		214	0 877[17 5]	
3988	$C_{10}H_{20}O_2$	d-γ-Nonyl formate	172.15		95[22]	0 869	258
3989	$C_{10}H_{20}O_2$	n-Octyl acetate $CH_3CO_2C_8H_{17}$	172.15	−38.5	210	0.885[0/4]	250
3991	$C_{10}H_{20}O_3$	1-Hydroxycapric acid	188.15	70 5			
3992	$C_{10}H_{21}N$	l-Menthylamine	155.17		208 2	0.860	475
3993	$C_{10}H_{22}$	n-Decane $CH_3(CH_2)_8CH_3$	142.17	−32.0	174	0.747	220
3994	$C_{10}H_{22}$	2, 6-Dimethyloctane	142.17		159	0 734	185
3995	$C_{10}H_{22}$	2, 7-Dimethyloctane	142.17	−52.8	160	0 722	171
3996	$C_{10}H_{22}$	dl, 3, 6-Dimethyloctane	142 17		162		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
3997	C₁₀H₂₂	d, 3, 6-Dimethyloctane	142 17		160 8	0.735¹³	
3998	C₁₀H₂₂	2-Methylnonane (CH₃)₂CH(CH₂)₆CH₃	142 17		160	0 728¹⁸·¹	174
3999	C₁₀H₂₂	3-Methylnonane C₂H₅(CH₃)CHC₆H₁₃..	142 17		166 9	0 735	197
4000	C₁₀H₂₂	5-Methylnonane (C₄H₉)₂CHCH₃.	142 17		166 2	0 732	189
4001	C₁₀H₂₂	Tripropylmethane (C₃H₇)₃CH	142 17		161 7	0 740¹⁴·²	210
4002	C₁₀H₂₂O	n-Decyl alcohol CH₃(CH₂)₉OH	158 17	7	231	0 829	
4003	C₁₀H₂₂O	3, 7-Dimethyl-n-octyl alcohol	158 17		118¹⁵	0 849⁰₁	
4004	C₁₀H₂₂O	Methylethylisohexyl carbinol	158 17		89¹⁴	0 834¹⁸₄	851
4005	C₁₀H₂₂O	Propyl-n-hexyl carbinol	158 17		211	0 826	
4006	C₁₀H₂₂O	n-Amyl ether (C₅H₁₁)₂O	158 17		190	0 774	
4007	C₁₀H₂₂O	Isoamyl ether [(CH₃)₂CHCH₂CH₂]₂O.	158 17		172 2	0 783¹¹·⁸	172
4008	C₁₀H₂₂O₂	cis-Terpine hydrate .	190 15	117 1			1210
4009	C₁₀H₂₂O₆S₂	d-Glucosediethylmercaptal	286 30	128			
4010	C₁₀H₂₂S	Diisoamyl sulfide . .	174 23		216	0 843	443
4011	C₁₀H₂₃N	n-Decylamine CH₃(CH₂)₉NH₂.	157 19	17	218		
4012	C₁₀H₂₃N	Diisoamylamine.	157 19		190	0 767	281
4013	C₁₀H₁₅Sb	Pentaethyl stibine (C₂H₅)₅Sb	266 96		100		
4014	C₁₀H₂₀O	α(β)-Lactucerol.	166 23	181			
4015	C₁₀H₂₀O₆	Agaric acid . .	230 23	142 d.			
4016	C₁₁H₆O₁₀	Benzenepentacarboxylic acid	298 05	233 d.			
4017	C₁₁H₇ClO	α-Naphthoyl chloride C₁₀H₇COCl	190 51		297 5		
4018	C₁₁H₇ClO	β-Naphthoyl chloride C₁₀H₇COCl	190 51	43	306		
4019	C₁₁H₇N	α-Naphthylcyanide . . .	153 06	33 5	296 5	1 117⁵₁	
4020	C₁₁H₇N	β-Naphthylcyanide	153 06	66 5	305	1 094⁶⁰₆₀	
4021	C₁₁H₇NO₄	Quinoline-2, 3-dicarboxylic acid	217 06	130 d.			
4022	C₁₁H₇NO₄	Quinoline-2, 4-dicarboxylic acid	217 06	246			
4023	C₁₁H₈O	α-Naphthaldehyde.	156 06		291 6	1 148	962
4024	C₁₁H₈O	β-Naphthaldehyde. . .	156 06	60 5		1 078⁹⁹·⁴	1133
4025	C₁₁H₈N₂O₄	Benzoylbarbituric acid	232 08	275			
4026	C₁₁H₈O₂	2-Hydroxy-α-naphthaldehyde	172 06	81	192⁹⁷		
4027	C₁₁H₈O₂	4-Hydroxy-α-naphthaldehyde	172 06	178			
4028	C₁₁H₈O₃	8-Hydroxy-α-naphthoic acid	188 06	169			
4029	C₁₁H₈O₂	α-Naphthoic acid	172 06	160	300		
4030	C₁₁H₈O₂	β-Naphthoic acid .	172 06	185	>300	1 077¹⁰⁰₄	
4031	C₁₁H₈O₃	3-Hydroxy-β-naphthoic acid	188 06	219			
4032	C₁₁H₉N	2-Phenylpyridine .	155 08		270	>1	
4033	C₁₁H₉N	3-Phenylpyridine	155 08		270 4	>1	
4034	C₁₁H₉N	4-Phenylpyridine .	155 08	78	275		
4035	C₁₁H₉NO₂	Aniluvitonic acid .	187 08	241			
4036	C₁₁H₉NO₃	Quininic acid.	203 08	280			
4037	C₁₁H₉NO₆	Hydrastininic acid	251 08	164			
4038	C₁₁H₁₀	α-Methylnaphthalene	142 08	−22	243	1 025	790
4039	C₁₁H₁₀	β-Methylnaphthalene	142 08	35 1	245	1 029	1062
4040	C₁₁H₁₀I₃NO₃	Thyroxin. .	584 88	250			
4041	C₁₁H₁₀O	Methyl α-naphthyl ether	158 08	< −10	258	1 096¹¹·⁹	831
4042	C₁₁H₁₀O	Methyl β-naphthyl ether	158 08	72	274		
4043	C₁₁H₁₀O₂	Ethyl phenylpropiolate	174 08		270 d.		
4043.1	C₁₁H₁₁BrN₂O	4-Bromoantipyrine	267 02	117			1181
4044	C₁₁H₁₁N	2, 4-Dimethylquinoline	157 09		264		
4045	C₁₁H₁₁N	2, 6-Dimethylquinoline	157 09	58	261		
4046	C₁₁H₁₁N	2, 7-Dimethylquinoline	157 09	61	265		
4047	C₁₁H₁₁N	3, 4-Dimethylquinoline	157 09	65	291		
4048	C₁₁H₁₁N	4, 6-Dimethylquinoline	157 09		256		
4049	C₁₁H₁₁N	4, 7-Dimethylquinoline	157 09	55	259		
4050	C₁₁H₁₁N	Methyl-α-naphthylamine	157 09		293		
4051	C₁₁H₁₁NO	Physostigmol . .	173 09	108			
4052	C₁₁H₁₁NO₂	Indole-2-propionic acid	189 09	136			
4053	C₁₁H₁₁NO₄	Ethyl o-nitrocinnamate	221 09	44			
4054	C₁₁H₁₁NO₄	Ethyl p-nitrocinnamate	221 09	141			
4055	C₁₁H₁₃BrNO₂S	p-Bromophenylmercapturic acid . .	318 08	153			
4056	C₁₁H₁₃IN	Quinaldine methiodide	285 03	190			
4057	C₁₁H₁₃IN	Quinoline ethiodide	285 03	157	d.		
4058	C₁₁H₁₃N₂O	Antipyrine.	188.11	109; 113	319¹⁷⁴		1307

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4059	$C_{11}H_{12}N_2O_2$	4, 4-Phenylethylhydantoin	204.11	199			
4060	$C_{11}H_{12}N_2O_2$	l-Tryptophane	204 11	289			
4060 1	$C_{11}H_{12}O$	Benzylidene methyl ethyl ketone	160 09	37 5		0.987[10]	1061
4061	$C_{11}H_{12}O_2$	Ethyl atropate.	176 09		124.4[16]	1.051	
4062	$C_{11}H_{12}O_2$	trans-Ethyl cinnamate	176 09	6.5	271	1.049	746
4063	$C_{11}H_{12}O_3$	3-Benzoylbutyric acid	192 09	126			
4064	$C_{11}H_{12}O_3$	Ethyl benzoylacetate	192 09		270 d.	1 122	704
4065	$C_{11}H_{12}O_3$	α-Ethyl phenylpyruvate	192 09	52	154 5[15]		
4066	$C_{11}H_{12}O_3$	β-Ethyl phenylpyruvate	192 09		152[15]		
4067	$C_{11}H_{12}O_3$	γ-Ethyl phenylpyruvate	192 09	79			
4068	$C_{11}H_{12}O_3$	Eugenol formate	192 09		150[20]		
4069	$C_{11}H_{12}O_3$	Isoeugenol formate	192 09		160[20]		
4071	$C_{11}H_{12}O_4$	Benzylsuccinic acid	208 09	161			
4072	$C_{11}H_{12}O_4$	α-Hydropiperic acid	208 09	76			
4073	$C_{11}H_{12}O_5$	Sinapic acid	224 09	191			
4074	$C_{11}H_{13}BrN_2O$	Antipyrine hydrobromide	269 03	150			
4075	$C_{11}H_{13}ClN_2O$	Antipyrine hydrochloride	224 57	160			
4076	$C_{11}H_{13}N$	Lilolidine	159 11		156[15]		
4077	$C_{11}H_{13}NO_2$	Hydrastinine	207 11	116			
4077 1	$C_{11}H_{13}NO_3$	Ethyl hippurate	207 11	60 5	180	1 043[23]	
4078	$C_{11}H_{13}NO_4$	Benzacetin	223 11	190			
4079	$C_{11}H_{13}NO_4$	Neurodin	223 11	87			
4080	$C_{11}H_{13}N_3O$	4-Aminoisoantipyrine	203 12	109			
4081	$C_{11}H_{13}N_3O$	Benzylcreatinine	203 12	225			
4082	$C_{11}H_{13}N_3O_6$	2, 4, 6-Trinitro-tert.-butyltoluene	283 12	97			
4083	$C_{11}H_{14}ClNO_3$	Hydrastinine hydrochloride	243 57	210			
4084	$C_{11}H_{14}N_2$	Calycanthine	174 12	243			
4085	$C_{11}H_{14}N_2$	Isocalycanthine	174 12	235			
4086	$C_{11}H_{14}N_2O$	Cytisine	190.12	153			1333
4087	$C_{11}H_{14}N_2O_2$	Antithermine (Acetopropionylphenylhydrazone)	206 12	108			
4088	$C_{11}H_{14}O$	Butyl phenyl ketone $C_6H_5COC_4H_9$	162 11		239 5		
4089	$C_{11}H_{14}O$	Isobutyl phenyl ketone	162 11		225	0 967	
4090	$C_{11}H_{14}O$	Isopropyl benzyl ketone	162 11		237	0 985$^0_4$	
4090 1	$C_{11}H_{14}O$	p-Methylbutyrophenone	162 11		252[739]	1 026	683
4091	$C_{11}H_{14}O$	Propyl benzyl ketone	162 11		244	0 984$^0_4$	
4091 1	$C_{11}H_{14}O$	2, 4, 6-Trimethylacetophenone	162.11		240 5[738]	0 975	661
4092	$C_{11}H_{14}O_2$	Eugenol methyl ether	178 11		249	1 055[15]	
4093	$C_{11}H_{14}O_2$	Isoeugenol methyl ether	178.11		264	1.055	
4094	$C_{11}H_{14}O_2$	p-Isopropylphenylacetic acid	178 11	52			
4095	$C_{11}H_{14}O_2$	n-Butyl benzoate $C_6H_5CO_2C_4H_9$	178 11	−22 4	250.3	1 000[20]	
4096	$C_{11}H_{14}O_2$	Benzyl butyrate $C_3H_7CO_2CH_2C_6H_5$	178 11		240	1 016[17 4]	
4097	$C_{11}H_{14}O_2$	Benzyl isobutyrate	178.11		228	1.016[18]	557
4097 1	$C_{11}H_{14}O_2$	d-β-Butyl benzoate	178.11		120[20]	1 000	563
4098	$C_{11}H_{14}O_2$	Ethyl hydrocinnamate	178 11		249	1 015	571
4099	$C_{11}H_{14}O_2$	Isobutyl benzoate	178.11		237	1.002[15]	
4100	$C_{11}H_{14}O_2$	Phenyl isovalerate	178.11		226		
4101	$C_{11}H_{14}O_3$	n-Butyl salicylate	194 11		155[15]		
4102	$C_{11}H_{14}O_3$	Propyl anisate $p-CH_3OC_6H_4CO_2C_3H_7$	194 11		176[45]	1 09	653
4103	$C_{11}H_{14}O_3$	Zingerone	194 11	41	188[14]		
4104	$C_{11}H_{15}NO$	p-Diethylaminobenzaldehyde	177.12	41	174[7]		
4105	$C_{11}H_{15}NO$	Isovaleroanilide	177.12	115			
4106	$C_{11}H_{15}NO$	n-Valeroanilide	177 12	49	267		
4107	$C_{11}H_{15}NO_2$	p-Diethylaminobenzoic acid	193 12	193			
4108	$C_{11}H_{15}NO_2$	Isobutyl p-aminobenzoate	193 12	65			
4109	$C_{11}H_{15}NO_2$	Methylacetophenetidine	193 12	40	300		
4110	$C_{11}H_{15}NO_2$	Triphenin.	193 12	120			
4111	$C_{11}H_{15}NO_3$	Anhalamine	209 12	188			
4112	$C_{11}H_{15}NO_3$	Lactophenine	209.12	118			
4113	$C_{11}H_{15}NO_3$	Methoxyacetophenetidin	209.12	98			
4114	$C_{11}H_{15}NO_7S$	Hydrastinine bisulfate	305.19	216			
4115	$C_{11}H_{16}$	n-Amylbenzene $CH_3(CH_2)_4C_6H_5$	148 12		202.1	0 860	514
4116	$C_{11}H_{16}$	tert.-Amylbenzene	148.12		189.3	0.874[15]	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4117	$C_{11}H_{16}$	3, 5-Diethyltoluene	148 12		200	0 870	
4118	$C_{11}H_{16}$	Isoamylbenzene $(CH_3)_2CH(CH_2)_2C_6H_5$	148 12		194	0 885	
4119	$C_{11}H_{16}$	Pentamethylbenzene $(CH_3)_5C_6H$	148 12	53	230	0 847$^{107.3}$	1152
4120	$C_{11}H_{16}$	4-Propyl-o-xylene $C_3H_7C_6H_3(CH_3)_2$	148 12	< −20	209		
4121	$C_{11}H_{16}$	4-Propyl-m-xylene $C_3H_7C_6H_3(CH_3)_2$	148 12	< −20	208 5		
4122	$C_{11}H_{16}$	2-Propyl-p-xylene $C_3H_7C_6H_3(CH_3)_2$	148 12	< −20	207		
4123	$C_{11}H_{16}Br_2N_2O_3$	N-2, 3-Dibromopropyl-5, 5-diethylbarbituric acid	383 97	125			
4124	$C_{11}H_{16}ClNO_2$	Anhalamine hydrochloride	245 59	258			
4125	$C_{11}H_{16}N_2O_2$	Pilocarpine	208 14	34			
4126	$C_{11}H_{16}N_2O_2$	Isopilocarpine	208 14		261^{10}		
4127	$C_{11}H_{16}O$	p-Isoamylphenol	164 12	93	255		
4128	$C_{11}H_{16}O$	Pentamethylphenol	164 12	125	267		
4129	$C_{11}H_{16}O$	Benzyl n-butyl ether $C_6H_5CH_2OC_4H_9$	164 12		216		
4130	$C_{11}H_{16}O$	Benzyl isobutyl ether	164 12		213	0 928$^{16.3}$	
4131	$C_{11}H_{16}O$	Phenyl isoamyl ether	164 12		225	0 920	545
4132	$C_{11}H_{16}O$	Thymyl methyl ether	164 12		216 2	0 954	
4133	$C_{11}H_{17}BrN_2O_2$	Isopilocarpine hydrobromide	289 06	147			
4134	$C_{11}H_{17}BrN_2O_2$	Pilocarpine hydrobromide	289 06	185			1333
4135	$C_{11}H_{17}ClN_2O_2$	Isopilocarpine hydrochloride	244 61	127			
4136	$C_{11}H_{17}ClN_2O_2$	Pilocarpine hydrochloride	244 61	196 7			1333
4137	$C_{11}H_{17}N$	o-Diethyltoluidine	163 14		206		
4138	$C_{11}H_{17}N$	m-Diethyltoluidine	163 14		228		
4139	$C_{11}H_{17}N$	p-Diethyltoluidine	163 14		229	0 924$^{16.3}$	
4140	$C_{11}H_{17}N$	Isoamylaniline	163 14		254 5	0 928^{16}	
4141	$C_{11}H_{17}NO_3$	Mescaline	211 14	151			
4142	$C_{11}H_{17}N_3O_6$	Isopilocarpine nitrate	271 16	159			
4143	$C_{11}H_{17}N_3O_6$	Pilocarpine nitrate	271 16	173			1333
4144	$C_{11}H_{18}O_2$	Citronellyl formate	181 13		98^{11}	0 884	453
4145	$C_{11}H_{18}N_2O_3$	5, 5-n-Butylisopropylbarbituric acid	226 16	210			
4146	$C_{11}H_{18}N_2O_3$	5, 5-Isoamylethylbarbituric acid	226 16	156			
4147	$C_{11}H_{18}O_2$	d-Bornyl formate	182 14		230	1.009	858
4148	$C_{11}H_{18}O_2$	Geranyl formate	182 14		98^{11}	0.909	491
4149	$C_{11}H_{18}O_2$	Isobornyl formate	182 14		100^{14}	1.017^{16}	
4150	$C_{11}H_{18}O_2$	Methyl geranate	182 14		117^{14}	0 922	961
4151	$C_{11}H_{18}O_2$	d, α-Terpinyl formate	182 14		136^{40}	0 999^0	
4152	$C_{11}H_{18}O_4$	Ethyl camphorate	214 14	87			
4153	$C_{11}H_{18}O_5$	Diethyl ethylacetylmalonate	230 14		137.5^{20}	1 053	316
4154	$C_{11}H_{19}N_3O$	d-Camphor semicarbazone	209 17	238			
4155	$C_{11}H_{20}O$	Geranyl methyl ether	168 15		212		
4156	$C_{11}H_{20}O$	Methyl d-bornyl ether	168 15		195 3	0 916	1011
4157	$C_{11}H_{20}O_2$	l-Menthyl formate	184 15	9	217	0 936	
4158	$C_{11}H_{20}O_2$	Undecylenic acid	184 15	24 5	295	0 907	
4159	$C_{11}H_{20}O_3$	Isoamyl ethylacetoacetate	200 15		236 d.	0 951^{15}	
4160	$C_{11}H_{20}O_4$	Di-n-butyl malonate $CH_2(CO_2C_4H_9)_2$	216 15		251 5	1 005^0	
4161	$C_{11}H_{20}O_4$	Diethyl diethylmalonate	216 15		223	0.990	282
4162	$C_{11}H_{20}O_4$	Isoamyl isopropyl malonate	216 15		140^{14}	0.958^{15}	314
4163	$C_{11}H_{20}O_5$	Glycerol 1, 2-dibutyrate	232 15		282		
4164	$C_{11}H_{21}NO_2$	Menthyl carbamate	199 17	165	>200 d.		
4165	$C_{11}H_{22}$	α-Undecylene $CH_2:CH(CH_2)_8CH_3$	154 17		188	0 763	
4166	$C_{11}H_{22}$	β-Undecylene $CH_3CH:CH(CH_2)_7CH_3$	154 17		193	0.774^{15}	341
4167	$C_{11}H_{22}N_2O_4$	Clavine	260 19	263			
4168	$C_{11}H_{22}O$	Methyl l-menthyl ether	170 17			0 861	
4169	$C_{11}H_{22}O$	Undecylic aldehyde	170 17	−4	117^{18}	0 825^{21}	342
4170	$C_{11}H_{22}O$	Diamyl ketone $(C_5H_{11})_2CO$	170.17	14 6	226 3	0.826^{20}	
4171	$C_{11}H_{22}O$	Diisoamyl ketone	170 17		226		
4172	$C_{11}H_{22}O$	Methyl n-nonyl ketone	170 17	12 1	228	0.826	312
4173	$C_{11}H_{22}O_2$	Umbellulic acid	186.17	23	280		
4174	$C_{11}H_{22}O_2$	Undecylic acid $CH_3(CH_2)_{11}CO_2H$	186.17	29 3	228^{100}		1066
4175	$C_{11}H_{22}O_2$	Ethyl pelargonate $C_8H_{17}CO_2C_2H_5$	186.17	−44 5	219	0 866$^{17.5}$	
4176	$C_{11}H_{22}O_2$	Methyl caprate $C_9H_{19}CO_2CH_3$	186 17	−18	224		
4177	$C_{11}H_{22}O_3$	Diisoamyl carbonate	202 17		228.7	0.912^{15}	
4178	$C_{11}H_{24}$	n-Undecane $CH_3(CH_2)_9CH_3$	156.18	−26.5	197	0.741	234

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4178 1	$C_{11}H_{24}$	e-Ethylnonane	156 18		71[16]	0.751[16]	
4179	$C_{11}H_{24}O$	n-Undecyl alcohol $CH_3(CH_2)_9CH_2OH$	172 19	19	146[30]	0.833	374
4179 1	$C_{11}H_{24}O$	n-Undecan-6-ol	172 19	16	235[754]	0.833	
4180	$C_{11}H_{25}N$	n-Undecylamine $CH_3(CH_2)_9CH_2NH_2$	171 20	16 5	234		
4181	$C_{12}H_5N_7O_{12}$	Dipicrylamine [2, 4, 6-$(NO_2)_3C_6H_2]_2NH$	439 10	250 d.			
4182	$C_{12}H_6O_{12}$	Mellitic acid $C_6(CO_2H)_6$	342 05	286			
4183	$C_{12}H_7N_3O_7$	Phenyl picrate	305 08	153			
4184	$C_{12}H_8$	Acenaphthylene	152 06	93	275		1192
4185	$C_{12}H_8AsN$	Phenarsazine	241 03	310			
4185 1	$C_{12}H_8Br_2$	p, p'-Di-(bromophenyl)	311 89	164		1.897	
4186	$C_{12}H_8Cl_2$	1, 2-Dichloracenaphthene	222 98	115			
4187	$C_{12}H_8N_2$	Phenanthroline	180 08	78.5	>360		
4188	$C_{12}H_8N_2$	Phenazine	180 08	171	>360		
4189	$C_{12}H_8N_2$	Phenazone	180 08	156	>360		
4190	$C_{12}H_8N_2$	Pseudophenanthroline	180 08	173			
4191	$C_{12}H_8N_2O_4$	Dinitroacenaphthene	244 08	206 d.			
4192	$C_{12}H_8N_2O_4$	o, o'-Dinitrodiphenyl	244 08	124			
4193	$C_{12}H_8N_2O_4$	m, m'-Dinitrodiphenyl	244 08	198			
4194	$C_{12}H_8N_2O_4$	p, p'-Dinitrodiphenyl	244 08	233			
4195	$C_{12}H_8O$	Diphenylene oxide	168 06	87	288		
4196	$C_{12}H_8O_2$	2-Phenylbenzoquinone	184 06	107			
4197	$C_{12}H_8O_4$	1, 8-Naphthalic acid	216 06	270			
4198	$C_{12}H_8O_4$	Bergaptene	216 06	188			
4199	$C_{12}H_8O_4$	Paracotoin	216 06	152			
4200	$C_{12}H_8O_4$	Xanthotoxin	216 06	146			
4201	$C_{12}H_8S_2$	Thianthrene	216 19	160	366		
4202	$C_{12}H_8AsClN$	Phenarsazine chloride	277 50	193			
4203	$C_{12}H_9Br$	3-Bromoacenaphthene	232 99	51 2	336 4	1 437[15]	
4204	$C_{12}H_9Cl$	3-Chloroacenaphthene	188 53	69 8	319		
4205	$C_{12}H_9Cl$	o-Chlorodiphenyl $o-ClC_6H_4C_6H_5$	188 53	34	268		
4206	$C_{12}H_9Cl$	m-Chlorodiphenyl $m-ClC_6H_4C_6H_5$	188 53	89			
4207	$C_{12}H_9Cl$	p-Chlorodiphenyl $p-ClC_6H_4C_6H_5$	188 52	75 5	282		
4208	$C_{12}H_9ClN_2$	m-Chloroazobenzene	216 54	67 5			
4209	$C_{12}H_9ClN_2$	p-Chloroazobenzene $p-ClC_6H_4NNC_6H_5$	216 54	89			
4210	$C_{12}H_9I$	3-Iodoacenaphthene	280 00	65	180 d.	1 674[62]	
4211	$C_{12}H_9N$	Carbazole	167 08	244 8	354 8		1333
4212	$C_{12}H_9NO_2$	o-Nitrodiphenyl $o-NO_2C_6H_4C_6H_5$	199 08	37	320		
4213	$C_{12}H_9NO_2$	m-Nitrodiphenyl $m-NO_2C_6H_4C_6H_5$	199 08	61			
4214	$C_{12}H_9NO_2$	p-Nitrodiphenyl $p-NO_2C_6H_4C_6H_5$	199 08	113	340		
4215	$C_{12}H_9NS$	Thiodiphenylamine	199 14	180	371 d.		
4216	$C_{12}H_9N_3O_2$	p-Nitroazobenzene	227 09	129 9			
4217	$C_{12}H_9N_3O_5$	2, 4-Dinitro-4'-hydroxydiphenylamine	275 09	190			
4218	$C_{12}H_{10}$	Acenaphthene	154 08	95	277 5	1 024[99 2]	1127, 1193
4219	$C_{12}H_{10}$	Diphenyl $C_6H_5C_6H_5$	154 08	69 0	254 9	1 041	1105
4220	$C_{12}H_{10}AsCl$	Diphenyl arsine chloride	264 50	42 8	327 d.	1 583[40]	
4221	$C_{12}H_{10}As_2$	Arsenobenzene $C_6H_5AsAsC_6H_5$	304 00	196			
4221 1	$C_{12}H_{10}ClI$	Diphenyliodonium chloride	316 47	d. 230		1 67	
4222	$C_{12}H_{10}Cl_2N_2$	Dichlorobenzidine [2, 4-Cl(NH_2)C_6H_3]_2	253 01	163			
4223	$C_{12}H_{10}Cl_2N_2$	p, p-Dichlorbenzidine	253 01	60			
4224	$C_{12}H_{10}N_2$	Aribine	182 09	237			
4225	$C_{12}H_{10}N_2$	Azobenzene $C_6H_5NNC_6H_5$	182 09	67	297 4	1 203	
4226	$C_{12}H_{10}N_2O$	Azoxybenzene	198 09	36		1 246	1031
4227	$C_{12}H_{10}N_2O$	p-Hydroxyazobenzene	198 09	152			
4228	$C_{12}H_{10}N_2O$	N-Nitrosodiphenylamine $(C_6H_5)_2NNO$	198 09	66 5			
4229	$C_{12}H_{10}N_2O$	p-Nitrosophenylaniline	198 09	143			
4230	$C_{12}H_{10}N_2O_2$	o, o'-Azophenol	214 09	172			
4231	$C_{12}H_{10}N_2O_2$	m, m'-Azophenol $HOC_6H_4NNC_6H_4OH$	214 09	205			
4232	$C_{12}H_{10}N_2O_2$	p, p'-Azophenol	214 09	215			
4233	$C_{12}H_{10}N_2O_2$	o-Nitrodiphenylamine	214 09	75			
4234	$C_{12}H_{10}N_2O_2$	p-Nitrodiphenylamine	214 09	133			
4235	$C_{12}H_{10}N_2O_2S$	Benzidinesulfone	246 16	>350			
4236	$C_{12}H_{10}N_2O_3$	o, o'-Azoxyphenol	288 17	102			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4237	$C_{12}H_{10}N_2O_2$	p, p'-Azoxyphenol	288 17	156; 107			
4238	$C_{12}H_{10}O$	o-Phenylphenol $C_6H_5C_6H_4OH$	170 08	56	275		
4239	$C_{12}H_{10}O$	m-Phenylphenol $C_6H_5C_6H_4OH$	170 08	78	>300		
4240	$C_{12}H_{10}O$	p-Phenylphenol $C_6H_5C_6H_4OH$	170 08	165	308		
4241	$C_{12}H_{10}O$	Phenyl ether $C_6H_5OC_6H_5$	170 08	26 9	259	1 072	1019
4242	$C_{12}H_{10}OS$	Diphenyl sulfoxide $(C_6H_5)_2SO$	202 14	70 5	340		
4243	$C_{12}H_{10}O_2$	o, o'-Diphenol $OHC_6H_4.C_6H_4OH$	186 08	109	326		
4244	$C_{12}H_{10}O_2$	o, p'-Diphenol $OHC_6H_4.C_6H_4OH$	186 08	161	342		
4245	$C_{12}H_{10}O_2$	m, m'-Diphenol $OHC_6H_4.C_6H_4OH$	186 08	123 5			
4246	$C_{12}H_{10}O_2$	p, p'-Diphenol $OHC_6H_4.C_6H_4OH$	186 08	272			
4247	$C_{12}H_{10}O_2$	α-Naphthyl acetate $CH_3CO.C_{10}H_7$	186 08	44 8			
4248	$C_{12}H_{10}O_2$	β-Naphthyl acetate $CH_3CO.C_{10}H_7$	186 08	68 5			
4249	$C_{12}H_{10}O_2S$	Diphenyl sulfone $(C_6H_5)_2SO_2$	218 14	129	377 8		
4250	$C_{12}H_{10}O_3S$	Phenyl benzenesulfonate	234 14	35			
4251	$C_{12}H_{10}O_4$	2, 2'-Diresorcinol	218 08	268			
4252	$C_{12}H_{10}O_4$	4, 4'-Diresorcinol	218 08	222			
4253	$C_{12}H_{10}O_4$	5, 5'-Diresorcinol	218 08	310			
4254	$C_{12}H_{10}O_4$	Piperic acid	218 08	217	220 d.		
4255	$C_{12}H_{10}O_4$	Quinhydrone	218 08	171			
4256	$C_{12}H_{10}O_4S$	4, 4'-Dihydroxydiphenylsulfone	250 14	239			
4257	$C_{12}H_{10}O_6$	Paracotoic acid	234 08	108			
4258	$C_{12}H_{10}O_5S_2$	Benzenesulfonic anhydride	298 21	90	210¹⁰ d		
4259	$C_{12}H_{10}P_2$	Phosphobenzene $C_6H_5P.PC_6H_5$	216 13	149			
4260	$C_{12}H_{10}S$	Diphenyl sulfide $(C_6H_5)_2S$	186 11		293	1.119	948
4261	$C_{12}H_{10}S_2$	Diphenyl disulfide $(C_6H_5)_2S_2$	218 21	61	310		
4262	$C_{12}H_{10}Se$	Diphenyl selenide $(C_6H_5)_2Se$	233 28		302	1 356¹⁵	
4263	$C_{12}H_{10}Te$	Diphenyl telluride $(C_6H_5)_2Te$	281 58		320	1 556¹⁵	800
4264	$C_{12}H_{11}As$	Diphenylarsine $(C_6H_5)_2AsH$	230 05		155⁴⁷		
4265	$C_{12}H_{11}AsO_2$	Diphenylarsonic acid $(C_6H_5)_2AsOOH$	262 05	178			
4266	$C_{12}H_{11}N$	o-Aminodiphenyl $C_6H_5C_6H_4NH_2$	169 09	45 5	299		
4267	$C_{12}H_{11}N$	2-Benzylpyridine	169 09		276		
4268	$C_{12}H_{11}N$	3-Benzylpyridine	169 09	34	286		
4269	$C_{12}H_{11}N$	4-Benzylpyridine	169 09		287		
4270	$C_{12}H_{11}N$	Diphenylamine $(C_6H_5)_2NH$	169 09	53	302	1 159	1333
4271	$C_{12}H_{11}NO$	m-Phenylaminophenol	185 09	82	340		
4272	$C_{12}H_{11}NO_2S$	Benzenesulfanilide	233 16	110			1183
4273	$C_{12}H_{11}N_3$	m-Aminoazobenzene	197 11	59			
4274	$C_{12}H_{11}N_3$	p-Aminoazobenzene $C_6H_5N_2C_6H_4NH_2$	197 11	126	>360		
4275	$C_{12}H_{11}N_3$	Diazoaminobenzene $C_6H_5N_2NHC_6H_5$	197 11	96	exp.		
4276	$C_{12}H_{11}N_3O_2$	o-Nitrobenzidine	229 11	143			
4277	$C_{12}H_{11}N_3O_2$	m-Nitrobenzidine	229 11	190			
4278	$C_{12}H_{11}P$	Diphenylphosphine $(C_6H_5)_2PH$	186 11		280	1 07¹⁴	
4279	$C_{12}H_{12}$	1, 4-Dimethylnaphthalene	156 09	< −18	264 3	1.016	900
4280	$C_{12}H_{12}$	2, 3-Dimethylnaphthalene	156 09		266		
4281	$C_{12}H_{12}$	2, 6-Dimethylnaphthalene	156 09	111			
4282	$C_{12}H_{12}$	α-Ethylnaphthalene	156 09	< −14	258 d.	1 064	
4283	$C_{12}H_{12}$	β-Ethylnaphthalene	156 09	−19	251	1 008⁰	
4284	$C_{12}H_{12}ClN$	Diphenylamine hydrochloride	205 56				1333
4285	$C_{12}H_{12}N_2$	p-Aminodiphenylamine	184 11	75	354		
4286	$C_{12}H_{12}N_2$	Benzidine $(p-NH_2C_6H_4)_2$	184 11	128 7	401 7		
4287	$C_{12}H_{12}N_2$	β-Benzidine	184 11	45	363		
4288	$C_{12}H_{12}N_2$	1, 1-Diphenylhydrazine $(C_6H_5)_2NNH_2$	184 11	36	220⁵⁰		
4289	$C_{12}H_{12}N_2$	Hydrazobenzene $C_6H_5NHNHC_6H_5$	184 11	131	d.		
4290	$C_{12}H_{12}N_2O$	Harmalol	200 11	212 d.			
4291	$C_{12}H_{12}N_2O_3$	Luminal (5,5-Phenylethylbarbituric acid)	232 11	173			
4292	$C_{12}H_{12}N_2O_6S_2$	Benzene-o, o'-disulfonic acid	344 24	> 175 d.			
4293	$C_{12}H_{12}N_4$	Chrysoidine	212 12	117 5			1333
4294	$C_{12}H_{12}N_4$	p, p'-Diaminoazobenzene	212 12	241			
4295	$C_{12}H_{12}N_4O_4$	Urocanic acid	276 12	213 d.			
4296	$C_{12}H_{12}O$	Ethyl α-naphthyl ether	172.09	5 5	276.4	1 061	770
4297	$C_{12}H_{12}O$	Ethyl β-naphthyl ether	172 09	37 5	282	1 064	1071
4297.1	$C_{12}H_{12}O$	l-Methyl-α-naphthyl carbinol	172 09	47	110¹¹	1 115	
4298	$C_{12}H_{12}O_2$	Benzylideneacetylacetone	188.09		188¹⁵		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4299	$C_{12}H_{12}O_2$	Allyl cinnamate	188.09		286 d.	1.052_{15}^{22}	
4300	$C_{12}H_{12}O_3$	Benzoylacetylacetone	204.09	35	167^{22}	1.152_{15}^{14}	
4301	$C_{12}H_{12}O_4$	Brasilic acid	252 09	129			
4302	$C_{12}H_{12}O_6$	Phloroglucinol triacetate	252 09	106			
4303	$C_{12}H_{12}O_6$	Pyrogallol triacetate	252.09	165			
4304	$C_{12}H_{13}N$	Dimethyl-α-naphthylamine	171.11		276	1.045_{15}^{15}	810
4305	$C_{12}H_{13}N$	Dimethyl-β-naphthylamine	171 11	46	305	$1 028_{15}^{52.3}$	1081
4306	$C_{12}H_{13}N$	Ethyl α-naphthylamine	171 11		176^{15}	1 060	871
4307	$C_{12}H_{13}N$	Ethyl β-naphthylamine	171 11		183^{15}	1 057	969
4308	$C_{12}H_{13}N$	2, 6, 8-Trimethylquinoline	171 11	46	261 4		
4309	$C_{12}H_{13}NO_3$	Pyrantin	219 11	155			
4310	$C_{12}H_{13}N_3$	p, p'-Diaminodiphenylamine	199 12	158			
4311	$C_{12}H_{14}As_2Cl_2N_2O_2$	Arsphenamine	438 96	160 d.			
4312	$C_{12}H_{14}IN$	Quinaldine ethiodide	299.05	234			
4313	$C_{12}H_{14}N_2O$	p-Tolylantipyrine	202.12	137			
4314	$C_{12}H_{14}N_2O_4S_2$	Benzidine-o, o'-disulfoneamide	342 27	278			
4315	$C_{12}H_{14}N_4O_6$	Desoxyamalic acid	310 14	260 s. d.			
4316	$C_{12}H_{14}N_4O_4$	Amalic acid (Tetramethylalloxantine)	342 14	221 d.			
4317	$C_{12}H_{14}O_2$	n-Propyl cinnamate	190 11		285 1	1.044^0	
4318	$C_{12}H_{14}O_3$	Eugenol acetate	206 11	31	282 4	1 084	665
4318 1	$C_{12}H_{14}O_3$	Ethyl p-methoxycinnamate	206.11	52			1232
4319	$C_{12}H_{14}O_3$	Isoeugenol acetate	206 11	80	283		
4322	$C_{12}H_{14}O_4$	Apiol	222 11	29 5	294	1 015	1310
4323	$C_{12}H_{14}O_4$	Isoapiol	222 11	56	304	$1 197^{12}$	817
4324	$C_{12}H_{14}O_4$	Diethyl o-phthalate $o\text{-}C_6H_4(CO_2C_2H_5)_2$	222.11		296 1	1.122	607
4325	$C_{12}H_{15}N$	Carbazoline	173 12	99	297		
4326	$C_{12}H_{15}N$	Diallylaniline $C_6H_5N(CH_2CH:CH_2)_2$	173.12		245	0.954	
4327	$C_{12}H_{15}N$	Julolidine	173 12	40	280		
4328	$C_{12}H_{15}NO$	Benzoylpiperidine	189 12	48	184^{17}		
4329	$C_{12}H_{15}NO$	Naphthalanmorpholine	189 12	63	312		
4330	$C_{12}H_{15}NO_2$	Dipropionanilide $C_6H_5N(OCC_2H_5)_2$	205 12	44	179.5^{30}		
4330.1	$C_{12}H_{15}NO_3$	Ethyl phenaceturate	221 12	79			1280
4331	$C_{12}H_{15}NO_3$	Anhalonidine	221 12	160			
4332	$C_{12}H_{15}NO_3$	Anhalonine	221 12	85 5			
4333	$C_{12}H_{15}NO_3$	Hydrocotarnine	221 12	55	100 d.		
4334	$C_{12}H_{15}NO_4$	Cotarnine	237 12	133			
4335	$C_{12}H_{16}N_2O$	Methylcytisine (Caulophylline)	204 14	137			
4336	$C_{12}H_{16}N_2O_4S$	Aniline sulfate $(C_6H_5NH_2)_2H_2SO_4$	284 20			1.377^4	
4337	$C_{12}H_{16}O$	Isoamyl phenyl ketone	176 12		242 5		
4338	$C_{12}H_{16}O$	Isobutyl benzyl ketone	176 12		250 5	$0 969_4^0$	
4339	$C_{12}H_{16}O_2$	Eugenol ethyl ether	192 12		254	$1 021^{9.5}$	808
4340	$C_{12}H_{16}O_2$	Isoeugenol ethyl ether	192.12	64			
4341	$C_{12}H_{16}O_2$	Pentamethylbenzoic acid	192 12	210 5			
4342	$C_{12}H_{16}O_2$	Amyl benzoate $C_6H_5CO_2C_5H_{11}$	192.12		d.	0 989	566
4343	$C_{12}H_{16}O_2$	Benzyl isovalerate	192 12		136^{25}		
4344	$C_{12}H_{16}O_2$	Benzyl d-valerate	192 12		250^{730}	$0 982^{22}$	558
4345	$C_{12}H_{16}O_2$	Isoamyl benzoate	192 12		262	0.993	
4345.1	$C_{12}H_{16}O_2$	Isopropyl hydrocinnamate	192 12		126^{11}	0.986^{25}	
4346	$C_{12}H_{16}O_2$	Thymyl acetate	192 12		243	$1 009^0$	
4347	$C_{12}H_{16}O_3$	n-Amyl salicylate $o\text{-}HOC_6H_4CO_2C_5H_{11}$	208 12		265	$1 065^{15}$	
4348	$C_{12}H_{16}O_3$	Butyl anisate $p\text{-}CH_3OC_6H_4CO_2C_4H_9$	208 12		183^{40}	1 054	635
4349	$C_{12}H_{16}O_3$	Isoamyl salicylate	208 12		273	$1 045_{15}^{15}$	
4350	$C_{12}H_{16}O_3$	Isobutyl anisate	208 12		170^{46}	1.052	634
4351	$C_{12}H_{16}O_3$	Guaiacyl valerate $C_4H_9CO_2C_6H_4OMe$	208 12		265		
4352	$C_{12}H_{16}O_3$	Asaron	208 12	67	296	1.165	1333
4353	$C_{12}H_{16}O_3$	Elemicin	208 12		147^{10}	1 063	694
4354	$C_{12}H_{16}O_4$	Aspidinol	224 12	161			
4355	$C_{12}H_{16}O_6$	Diethyl succinylsuccinate	256 12	128			
4356	$C_{12}H_{16}O_6$	d, β-Phenylglucoside	256 12	175			
4357	$C_{12}H_{16}O_7$	Arbutin	272 12	195			1333
4358	$C_{12}H_{17}AsN_2O_4$	Aniline arsenate $(C_6H_5NH_2)_2H_3AsO_4$	328 11	140			
4359	$C_{12}H_{17}NO$	N-n-Butylacetanilide	191.14		276.5		
4360	$C_{12}H_{17}NO$	Caproanilide $CH_3(CH_2)_4CONHC_6H_5$	191 14	95			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4361	C₁₂H₁₇NO	C-Diethylacetanilide	191 14	124			
4362	C₁₂H₁₇NO₂	Ethyl-N-phenacetine	207 14	38	298		
4363	C₁₂H₁₇NO₂	Ethyl-o-tolylurethane	207 14		255		
4364	C₁₂H₁₇N₄O₉	Lysine picrate...	375 17	252 d			
4365	C₁₂H₁₈	Hexamethylbenzene	162 14	166	265		
4365.1	C₁₂H₁₈	1-Methyl-3-tert.-amylbenzene	162 14		208	0.8673	
4366	C₁₂H₁₈	1, 2, 4-Triethylbenzene	162 14		218	0.882	583
4367	C₁₂H₁₈	1, 3, 5-Triethylbenzene	162 14		218	0 863	565
4367 1	C₁₂H₁₈N₂O₄	Rhamnose phenylhydrazone	254 16	159			
4367.2	C₁₂H₁₈N₂O₅	d, α-Glucosephenylhydrazone	270 16	160			
4367 3	C₁₂H₁₈N₂O₅	d, β-Glucosephenylhydrazone	270 16	141			
4367.4	C₁₂H₁₈N₄O	Phenylhydrazine hydrate	234 17	24			
4367 5	C₁₂H₁₈N₄O₃	Hexamethylenetetramineresorcinol.	250 17	200 d			
4367 6	C₁₂H₁₈O	Benzyl isoamyl ether.	178 14		237 5		
4367.7	C₁₂H₁₈O	Thymyl ethyl ether.	178.14		226 9	0 933⁰₈	
4367 8	C₁₂H₁₈O	Mellithyl alcohol (CH₃)₅C₆CH₂OH	178 14	160 5			
4367 9	C₁₂H₁₈O₃	Phloroglucinol triethyl ether	210 14	43	175²⁴		
4368	C₁₂H₁₈O₃	Pyrogallol triethyl ether	210 14	39			
4368 1	C₁₂H₁₈O₄	Cascarillin..... .	226 14	205			
4368 2	C₁₂H₁₈O₆	Trimeric diacetyl	258 14	105	280 1		
4368 3	C₁₂H₁₈O₆	Diethyl 1, 1′-diacetylsuccinate	258.14	88		1.209 (at.) 1.176 (met.)	1106, 1201
4368 4	C₁₂H₁₈O₆	Triethyl aconitate..	258 14		253⁷⁵⁰	1.106	454
4368 41	C₁₂H₁₈O₆	Diethyl diacetyltartrate	290 14	68	170¹⁵	1 109⁷¹	
4368.5	C₁₂H₁₉Br₂O₂	Bromal d-borneolate.	434 89	109		1 868⁰	
4368.6	C₁₂H₁₉ClO₂	d-Bornyl chloroacetate	230 60		147³⁰		
4368 7	C₁₂H₁₉Cl₃O₂	Chloral-d-borneolate	301 52	56			
4368 8	C₁₂H₁₉N	n-Dipropylaniline C₆H₅N(C₃H₇)₂	177 15		241	0 910	
4368 9	C₁₂H₂₀N₂O₃	Isoamylisopropylbarbituric acid	240 17	175			
4369	C₁₂H₂₀N₂O₃	Isoamylpropylbarbituric acid .	270 17	132			
4369.1	C₁₂H₂₀N₄O₇	Hexamethylenetetraminemethylene citrate	332 19	175			
4369.2	C₁₂H₂₀O	Ballanophorin. . .	180 15	56			
4370	C₁₂H₂₀O	Homophorone	180 15		210⁸²⁵	0 886	530
4371	C₁₂H₂₀O₂	Geranylacetic acid	196 15		179¹⁹	0 938	516
4372	C₁₂H₂₀O₂	dl-Bornyl acetate	196 15		114²²	0 985	483
4373	C₁₂H₂₀O₂	d-Bornyl acetate	196 15	29	226	0 991¹⁴	994
4374	C₁₂H₂₀O₂	Geranyl acetate	196 15		242	0 917¹⁴	493
4375	C₁₂H₂₀O₂	Isobornyl acetate	196 15		89⁹	0 981	1010
4375.1	C₁₂H₂₀O₂	Isopulegyl acetate	196 15		103¹⁴	0 935¹⁴	934
4376	C₁₂H₂₀O₂	l-Linalyl acetate	196 15		220	0 895	414
4377	C₁₂H₂₀O₂	Neryl acetate	196 15		134¹⁵	0 916¹⁴	
4378	C₁₂H₂₀O₂	dl, α-Terpinyl acetate	196 15	< −50	220 d.	0.957	
4379	C₁₂H₂₀O₂	d (l), α-Terpinyl acetate	196 15		140⁴⁰	0 983⁰₀	
4380	C₁₂H₂₀O₅	Diethyl 1-ethyl-1′-acetylsuccinate	244 15		263	1 064¹⁶₁₇,₆	
4381	C₁₂H₂₀O₇	Triethyl citrate	276 15		294	1.137	409
4382	C₁₂H₂₀O₁₀	Maltosan..	324 15	150 (?)			
4383	C₁₂H₂₁ClO₂	l-Menthyl chloroacetate	232 62	38	137¹²	1 056	
4384	C₁₂H₂₁N₃	Kyanpropine... ...	207.19	116			
4385	C₁₂H₂₂O	Ethyl d-bornyl ether	182 17		205	0.901	1023
4386	C₁₂H₂₂O	Hexenyl ether	182 17		118		
4387	C₁₂H₂₂O₂	d-Citronellyl acetate	198 17		121¹⁵	0 903¹⁵₄	402
4388	C₁₂H₂₂O₂	l-Menthyl acetate (HOCHCO₂C₃H₉)₂	198.17		227	0 919	418
4389	C₁₂H₂₂O₃	Lanolic acid..	214.17	77			
4390	C₁₂H₂₂O₃	l-Menthyl glycollate	214 17	87			
4391	C₁₂H₂₂O₄	Diisoamyl oxalate	230 17		265	0.968¹¹	
4392	C₁₂H₂₂O₆	Di-n-butyl d-tartrate	262.17	22 5	203¹⁴	1 098¹⁴	
4393	C₁₂H₂₂O₆	Diisobutyl d-tartrate	262.17	69	325		
4393.1	C₁₂H₂₂O₆	Diisobutyl l-tartrate	262.17	74	185²¹	1 029⁷⁹	
4394	C₁₂H₂₂O₁₁	Lactose. . .	342.17	201 6	d.	1 525	1229
4395	C₁₂H₂₂O₁₁(H₂O)	Maltose	360 19			1.540	1333
4396	C₁₂H₂₂O₁₁	Saccharose	342.17	186		1.588¹⁶₄	1242
4397	C₁₂H₂₂O₁₁	Trehalose (2H₂O)	342.17	210			1195

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4398	$C_{12}H_{23}ClO$	Lauryl chloride $CH_3(CH_2)_{10}COCl$	218 64	−17	145[18]		
4399	$C_{12}H_{23}N$	Lauronitrile $CH_3(CH_2)_{10}CN$	181 19	4	198[100]	0.827[15]	
4400	$C_{12}H_{24}$	n-Dodecylene $CH_2{:}CH(CH_2)_9CH_3.$	168 19	−31 5	96[15]	0.762[15]	
4401	$C_{12}H_{24}N_2O_{10}$	d-Glucosealdazine	356 20	100			
4402	$C_{12}H_{24}O$	n-Amyl hexyl ketone $C_5H_{11}COC_6H_{13}$	184 19	9	112[9]		
4403	$C_{12}H_{24}O$	Ethylmenthol	184 19		85[4]	0.904[17]	
4404	$C_{12}H_{24}O$	l-Ethyl menthyl ether	184 19		212 9	0.854	918
4405	$C_{12}H_{24}O$	Lauric aldehyde $CH_3(CH_2)_{10}CHO$	184 19	44 5	185[100]		
4406	$C_{12}H_{24}O_2$	Lauric acid $CH_3(CH_2)_{10}CO_2H$	200 19	48 0	225[100]	0 883	1123
4407	$C_{12}H_{24}O_2$	n-Decyl acetate $CH_3CO_2C_{10}H_{21}$	200 19		191 5		1082
4408	$C_{12}H_{24}O_2$	Ethyl n-caprate $C_9H_{19}CO_2C_2H_5$	200 19		245	0 862	
4409	$C_{12}H_{24}O_4$	n-Parabutyraldehyde	216 19		100[35]		
4410	$C_{12}H_{25}NO$	Lauramide $CH_3(CH_2)_{10}CONH_2$	199 20	102	200[12 5]		
4411	$C_{12}H_{26}$	n-Dodecane $CH_3(CH_2)_{10}CH_3$	170 20	−12	216	0 768	255
4412	$C_{12}H_{26}$	5-Propylnonane $(C_4H_9)_2CHC_3H_7$	170 20		205	0 756	268
4413	$C_{12}H_{26}$	2, 4, 5, 7-Tetramethyloctane	170 20		210		
4414	$C_{12}H_{26}O$	n-Amylhexyl carbinol	186 20	30	119[9]		
4415	$C_{12}H_{26}O$	n-Dodecyl alcohol $CH_3(CH_2)_{10}CH_2OH$	186 20	24	259	0.831	
4416	$C_{12}H_{26}O$	n-Hexyl ether $(C_6H_{13})_2O$	186 20		208 8		
4417	$C_{12}H_{27}N$	Dodecylamine $C_{12}H_{25}NH_2$	185 22	28	135[15]		
4418	$C_{12}H_{27}N$	Tri-n-butylamine $(C_4H_9)_3N$	185 22		214	0 778[20/20]	
4419	$C_{12}H_{27}N$	Triisobutylamine $[(CH_3)_2CHCH_2]_3N..$	185 22	−21 8	191 5	0 766[25/25]	294
4420	$C_{12}H_{28}N_2O_4$	Ethylenediamine isovalerate	264 23	129			
4421	$C_{13}H_7Br_3O_3$	Tribromosalol	450 80	195			
4422	$C_{13}H_8Cl_2O$	p, p'-Dichlorobenzophenone	250 98	145			
4423	$C_{13}H_8N_2O_5$	p, p'-Dinitrobenzophenone	272 08	190			
4424	$C_{13}H_8N_4O_9$	o, o', p, p'-Tetranitrodiphenylurea	392 11	189			
4425	$C_{13}H_8O$	Fluorenone	180 06	84	341 5		
4426	$C_{13}H_8O$	Pyrene ketone	180 06	142			
4427	$C_{13}H_8O_2$	Xanthone	196 06	174	351		
4428	$C_{13}H_8O_3S$	Benzophenonesulfone	244 13	187			
4429	$C_{13}H_8O_4$	Euxanthone	228 06	240			
4430	$C_{13}H_9BrO_2$	p-(p-Bromophenyl) benzoic acid	276 99	194			
4431	$C_{13}H_9ClO$	o-Chlorobenzophenone	216 53	45 5	330		
4432	$C_{13}H_9ClO$	m-Chlorobenzophenone	216 53	83			
4433	$C_{13}H_9ClO$	p-Chlorobenzophenone	216 53	78	>300		
4434	$C_{13}H_9N$	Acridine	179 08	108	346		
4435	$C_{13}H_9N$	α-Naphthoquinoline	179 08	52	351		
4436	$C_{13}H_9N$	β-Naphthoquinoline	179 08	93	351		
4437	$C_{13}H_9N$	Phenanthridine	179 08	104	360		
4438	$C_{13}H_9NO$	9-Acridone	195 08	354			
4439	$C_{13}H_{10}$	Fluorene	166 08	116	295		
4440	$C_{13}H_{10}AsN$	Diphenylcyanoarsine $(C_6H_5)_2AsCN....$	255 05	30			
4441	$C_{13}H_{10}Cl_2$	Benzophenone chloride	236 99		305	1 235[18 5]	
4442	$C_{13}H_{10}Cl_2$	m, m'-Dichlorodiphenylmethane	236 99		318	1 234[21]	
4443	$C_{13}H_{10}Cl_2$	p, p'-Dichlorodiphenylmethane	236 99	55	210[15]		
4444	$C_{13}H_{10}N_2O_3$	Benzeneazosalicylic acid	242 09	218 d.			
4445	$C_{13}H_{10}O$	p-Diphenylaldehyde $p-C_6H_5C_6H_4CHO.$	182 08	60			
4446	$C_{13}H_{10}O$	Fluorenol	182 08	156			
4447	$C_{13}H_{10}O$	α-Benzophenone $(C_6H_5)_2CO$	182 08	48 5	305 4	1 083[53 5]	
4448	$C_{13}H_{10}O$	β-Benzophenone	182 08	26 5	306	1 108[23]	1014
4449	$C_{13}H_{10}O$	γ-Benzophenone	182 08	45 48			
4450	$C_{13}H_{10}O$	δ-Benzophenone	182 08	−51			
4451	$C_{13}H_{10}O$	Xanthene	182 08	100.5	315		
4452	$C_{13}H_{10}O_2$	o-Hydroxybenzophenone	198 08	41	250[630]		
4453	$C_{13}H_{10}O_2$	m-Hydroxybenzophenone	198 08	116			
4454	$C_{13}H_{10}O_2$	p-Hydroxybenzophenone	198 08	134			
4455	$C_{13}H_{10}O_2$	o-Phenylbenzoic acid	198 08	111	344		
4456	$C_{13}H_{10}O_2$	m-Phenylbenzoic acid	198 08	161			
4457	$C_{13}H_{10}O_2$	p-Phenylbenzoic acid	198 08	219			
4458	$C_{13}H_{10}O_4$	Phenyl benzoate $C_6H_5CO_2C_6H_5$	198 08	70	314	1.235[21]	
4459	$C_{13}H_{10}O_3$	2, 5-Dihydroxybenzophenone	214 08	122			
4460	$C_{13}H_{10}O_3$	2, 2'-Dihydroxybenzophenone	214.08	59	340		

No.	Formula	Name	Mol wt.	M. P.	B. P.	d	R. I. No.
4461	C₁₃H₁₀O₃	2, 3'-Dihydroxybenzophenone	214 08	126			
4462	C₁₃H₁₀O₃	2, 4'-Dihydroxybenzophenone	214 08	144			
4463	C₁₃H₁₀O₃	3, 4'-Dihydroxybenzophenone	214 08	197			
4464	C₁₃H₁₀O₃	4, 4'-Dihydroxybenzophenone	214 08	210			
4465	C₁₃H₁₀O₃	o-Phenoxybenzoic acid .	214 08	114 5	355 d.		
4466	C₁₃H₁₀O₃	Diphenyl carbonate (C₆H₅O)₂CO	214 08	81	302		
4467	C₁₃H₁₀O₃	Salol o-HOC₆H₄CO₂C₆H₅	214 08	43	173¹¹	1.250	
4468	C₁₃H₁₀O₄	2, 6, 2'-Trihydroxybenzophenone	230 08	133			
4469	C₁₃H₁₀O₄	Pimpinellin ...	246 08	119			
4470	C₁₃H₁₀O₆	Maclurin.	262 08	220 d			
4471	C₁₃H₁₀O₆	Sordidin	294 08	210			
4472	C₁₃H₁₀S	Thiobenzophenone (C₆H₅)₂CS	198 14	146 5			
4473	C₁₃H₁₁N	Benzylideneaniline C₆H₅N.CHC₆H₅	181 09	54	300		
4474	C₁₃H₁₁N	5, 10-Dihydroacridine	181 09	169			
4475	C₁₃H₁₁NO	o-Aminobenzophenone	197 09	108			
4476	C₁₃H₁₁NO	m-Aminobenzophenone	197 09	86			
4477	C₁₃H₁₁NO	p-Aminobenzophenone	197 09	124			
4478	C₁₃H₁₁NO	Benzanilide C₆H₅NHCOC₆H₅	197 09	161		1.321⁴	
4479	C₁₃H₁₁NO	Benzophenoneoxime (C₆H₅)₂C·NOH	197 09	142			
4480	C₁₃H₁₁NO	N-Phenylformanilide (C₆H₅)₂NOCH.	197 09	74	220	1 230	
4481	C₁₃H₁₁NO₂	o-Benzoylaminophenol	213 09	167 d.			
4482	C₁₃H₁₁NO₂	m-Benzoylaminophenol .	213 09	174			
4483	C₁₃H₁₁NO₂	p-Benzoylaminophenol	213 09	227			
4484	C₁₃H₁₁NO₂	p-Nitrodiphenylmethane	213 09	31			
4485	C₁₃H₁₁NO₂	Salicylanilide o-OHC₆H₄CONHC₆H₅.	213 09	135			
4486	C₁₃H₁₁NO₃	p-Aminosalol	229 09	152			
4487	C₁₃H₁₁NO₄	Gallanilide	245 09	205			
4488	C₁₃H₁₁N₃	2, 8-Diaminoacridine	209 11	284			
4489	C₁₃H₁₁O₅	Gelsemic acid	247 09	206			
4490	C₁₃H₁₂	Diphenylmethane (C₆H₅)₂CH₂	168 09	27	262	1.006	1030
4491	C₁₃H₁₂	o-Phenyltoluene CH₃C₆H₄C₆H₅	168 09		260		
4492	C₁₃H₁₂	m-Phenyltoluene CH₃C₆H₄C₆H₅	168 09		277	1.031⁰	
4493	C₁₃H₁₂	p-Phenyltoluene CH₃C₆H₄C₆H₅	168 09	−3	267	1 015¹⁷	
4494	C₁₃H₁₂N₂	Benzaldehyde phenylhydrazone	196 11	156			
4495	C₁₃H₁₂N₂O	1-Benzoyl-1-phenylhydrazine	212 11	70			
4496	C₁₃H₁₂N₂O	1-Benzoyl-2-phenylhydrazine	212 11	168			
4497	C₁₃H₁₂N₂O	o, o'-Diaminobenzophenone	212 11	135			
4498	C₁₃H₁₂N₂O	m, m'-Diaminobenzophenone	212 11	174			
4499	C₁₃H₁₂N₂O	p, p'-Diaminobenzophenone	212 11	237			
4500	C₁₃H₁₂N₂O	1, 2-Diphenylurea CO(NHC₆H₅)₂	212 11	235	260		1329
4501	C₁₃H₁₂N₂O	1, 1-Diphenylurea (C₆H₅)₂NCONH₂.	212 11	189			
4502	C₁₃H₁₂N₂O	Harmine	212 11	257 d.			
4503	C₁₃H₁₂N₂O₃	o-Nitrobenzylaniline	228 11	44; 57			
4504	C₁₃H₁₂N₂S	1, 2-Diphenylthiourea	228.17	154	d.	1.321⁴	
4505	C₁₃H₁₂O	o-Benzylphenol C₆H₅CH₂C₆H₄OH	184 09	21	312		
4506	C₁₃H₁₂O	p-Benzylphenol C₆H₅CH₂C₆H₄OH..	184 09	84	322		
4507	C₁₃H₁₂O	Diphenyl carbinol (C₆H₅)₂CHOH	184 09	68	298.5		
4508	C₁₃H₁₂O	Benzyl phenyl ether C₆H₅OCH₂C₆H₅	184 09	39	287		
4509	C₁₃H₁₂O₃S	Phenyl-p-toluenesulfonate	248.16	96			
4512	C₁₃H₁₃N	Benzylaniline C₆H₅NHCH₂C₆H₅	183 11	37	300	1 038¹⁸	
4513	C₁₃H₁₃N	N-Methyldiphenylamine (C₆H₅)₂NCH₃.	183 11	−7 6	293 4	1.047²²	
4514	C₁₃H₁₃NO	m-(o-Tolylamino) phenol ..	199 11		375		
4515	C₁₃H₁₃NO	p-(m-Tolylamino) phenol	199.11	91	350		
4517	C₁₃H₁₃NO₂S	Toluene-p-sulfoneanilide	247 17	103			
4518	C₁₃H₁₃N₃	Diphenylguanidine......	211 12	148			
4519	C₁₃H₁₄N₂	o, p'-Diaminodiphenylmethane	198 12	88			
4520	C₁₃H₁₄N₂	m, m'-Diaminodiphenylmethane	198 12	48			
4521	C₁₃H₁₄N₂	m, p'-Diaminodiphenylmethane	198.12	90			
4522	C₁₃H₁₄N₂	p, p'-Diaminodiphenylmethane	198 12	89			
4523	C₁₃H₁₄N₂	1-Phenyl-2-benzylhydrazine .	198 12	26			
4524	C₁₃H₁₄N₂O	Harmaline.............................	214.12	238			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4525	$C_{13}H_{14}N_2O_2$	Analgen (5-Acetylamino-8-ethoxyquino-line)	230 12	155			
4526	$C_{13}H_{14}N_2S$	1, 2-Di(p-aminophenyl) thiourea . .	258 21	195			
4526 1	$C_{13}H_{14}O_2$	Isobutyl phenylpropiolate	202 11		176^{12}	1.158^{25}	
4527	$C_{13}H_{14}O_4$	Drimine	234 11	256			
4528	$C_{13}H_{15}Cl_3N_2O_3$	Chloralantipyrine	353 51	68			
4529	$C_{13}H_{15}N$	2, 5, 6, 8-Tetramethylquinoline	185 12	20	300		
4530	$C_{13}H_{16}IN$	2, 4-Dimethylquinoline ethiodide	313 06	225			
4530 1	$C_{13}H_{16}N_2O$	4-Ethyl antipyrine . .	216 14	68			1237
4530 2	$C_{13}H_{16}N_2O$	1-Phenyl-2-propyl-3-methylpyrazolone..	216 14	93			1262
4530 3	$C_{13}H_{16}O$	Benzalpinacoline	188 12	39.5		0.939^{60}	1048
4531	$C_{13}H_{16}O_3$	Ethyl benzylacetoacetate	220 12		290 d.	1 036$^{15}_{16}$	
4532	$C_{13}H_{16}O_3$	Isoeugenol propionate	220 12		292		
4533	$C_{13}H_{16}O_4$	Ethyl phenylmalonate	236 12		285 d.	1 095$^{15}_{25}$	
4534	$C_{13}H_{16}O_7$	l-Hebein	284.12	175			
4535	$C_{13}H_{16}O_7$	Salinigrin	284 12	195			
4536	$C_{13}H_{17}NO_4$	Thermodin	251.14	88			1333
4537	$C_{13}H_{17}N_3O$	Pyramidon	231 16	108			
4538	$C_{13}H_{18}BrNO_2$	Phenoval	300 06	150			
4539	$C_{13}H_{18}N_2O$	Eseroline	218.16	127			
4541	$C_{13}H_{18}N_4O_6S$	Hexamethylenetetramine salicylsulfonic acid (Hexal)	358.24	190 d.			
4542	$C_{13}H_{18}O$	Phenyl hexyl ketone $C_6H_5COC_6H_{13}$.	190.14	17	271 5		
4543	$C_{13}H_{18}O_2$	Eugenol propyl ether	206 14		270.5	1 002	
4544	$C_{13}H_{18}O_2$	Phenyl heptylate $C_6H_{13}CO_2C_6H_5$	206 14		282 3	0 982$^{15}_{15}$	
4545	$C_{13}H_{18}O_3$	Isoamyl anisate . .	222 14		188^{30}	1 040	638
4546	$C_{13}H_{18}O_7$	Methylarbutin	286 14	175			
4547	$C_{13}H_{18}O_7$	Salicin .	286 14	201.5	240	1.434^{24}	
4548	$C_{13}H_{18}O_8$	Calmatambetin .	302.14	148			
4549	$C_{13}H_{19}NO$	Heptanilide $CH_3(CH_2)_5CONHC_6H_5$	205.15	71			
4550	$C_{13}H_{19}NO_2$	Benzalaminoacetal	221 15		220^{160}		
4551	$C_{13}H_{19}NO_3$	Dioscorine	221 15	43 5			
4552	$C_{13}H_{19}NO_3$	Pellotine	237.15	111			1333
4553	$C_{13}H_{19}NO_9$	Gynocardine	333 15	162			
4554	$C_{13}H_{19}O_8$	Aucubine	303.15	181			
4555	$C_{13}H_{20}ClNO_3$	Dioscorine hydrochloride .	257.62	204			
4556	$C_{13}H_{20}ClNO_3$	Gujasanol (Diethylaminoacetic acid guaiacol hydrochloride)	273 62	184			
4557	$C_{13}H_{20}N_2O_2$	Novocaine..	236.17	60			
4558	$C_{13}H_{20}N_2O_2(2H_2O)$	Novocaine....	272.19	51			
4559	$C_{13}H_{20}O$	α-Ionone	192.15		147 5^{28}	0 930	988
4560	$C_{13}H_{20}O$	β-Ionone..... .	192.15		140^{18}	0.944	667, 951
4561	$C_{13}H_{20}O$	Irone	192.15		144^{16}	0.939	605
4562	$C_{13}H_{20}O$	Lactucol . .	192.15	160			
4563	$C_{13}H_{20}O$	Pseudoionone	192.15		170^{28}	0 897	1001
4564	$C_{13}H_{20}O_2$	Galbanic acid	208 15	156			
4565	$C_{13}H_{21}ClN_2O_2$	Novocaine hydrochloride	272 64	156			
4566	$C_{13}H_{21}ClN_2O_2$	Procaine	272 64	155			
4567	$C_{13}H_{21}N$	N-Ethyl-isoamylaniline	191.17		262		
4568	$C_{13}H_{21}NO_4$	Meteloidine	255 17	141			
4569	$C_{13}H_{22}BrNO_4$	Meteloidine hydrobromide	336 09	250			
4570	$C_{13}H_{22}N_2O_3$	Ethylheptylbarbituric acid	254.19	119			
4571	$C_{13}H_{22}O$	Zeorin...... . .	194.17	251			
4572	$C_{13}H_{22}O_2$	d-Bornyl propionate	210 27		110^{11}	0.979^{15}	857
4573	$C_{13}H_{22}O_3$	l-Menthyl pyruvate	226.17		140^{22}	0 985	
4574	$C_{13}H_{22}O_7$	Taxicatin.... .	290.17	171			
4575	$C_{13}H_{24}NO_3$	Cuscohygrine ..	226.19		170^{23}		
4576	$C_{13}H_{24}O$	Allyl l-menthyl ether	196.19		104^{13}	0.876	
4577	$C_{13}H_{24}O$	Geranylacetone	196.19		139^{19}		
4578	$C_{13}H_{24}O_2$	l-Menthyl propionate	212.19		118^{15}	0 918	
4579	$C_{13}H_{24}O_3$	l-Menthyl dl-lactate	228.19	32	142^{15}	0.984	
4580	$C_{13}H_{24}O_4$	Brassylic acid......................	244.19	114			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4580.1	$C_{13}H_{24}O_4$	Di-*l*-amyl malonate	244 19		154[13]	0.962[25]	
4581	$C_{13}H_{26}$	Tridecylene	182 20		232 7	0.845[0]	
4582	$C_{13}H_{26}O_2$	Tridecylic acid $CH_3(CH_2)_{11}CO_2H$	214 20	51	236[100]		
4583	$C_{13}H_{26}O_2$	Isoamyl caprylate.....	214 20		136[10]		
4584	$C_{13}H_{26}O_2$	Methyl laurate $C_{11}H_{23}CO_2CH_3$	214 20	5	148[18]		
4585	$C_{13}H_{28}$	Dipropylhexylmethane $(C_3H_7)_2CHC_6H_{13}$	184 22		221 2	0 765[14 4]	299
4586	$C_{13}H_{28}$	Tributylmethane $(C_4H_9)_3CH$	184 22			0.760	300
4587	$C_{13}H_{28}$	*n*-Tridecane $CH_3(CH_2)_{11}CH_3$	184 22	- 6 2	234	0 757	908
4588	$C_{13}H_{28}O$	Di-*n*-hexylcarbinol $(C_6H_{13})_2CHOH$	200 22	42			
4589	$C_{13}H_{28}O$	*n*-Tridecyl alcohol $CH_3(CH_2)_{11}CH_2OH$	200 22	30 5	156[15]	0 822[11]	
4590	$C_{13}H_{29}N$	Tridecylamine $CH_3(CH_2)_{11}CH_2NH_2$	199 23	27	265		
4591	$C_{14}H_2Cl_8$	Octachloroanthracene	453 68	>350			
4592	$C_{14}H_3Cl_7$	Heptachloroanthracene	419 23	>350			
4593	$C_{14}H_4Cl_4O_2$	1, 2, 3, 4-Tetrachloroanthraquinone.	345 86	191			
4594	$C_{14}H_4Cl_4O_2$	β-Tetrachloroanthraquinone	345 86	330			
4595	$C_{14}H_4Cl_6$	Hexachloroanthracene	384 78	330			
4596	$C_{14}H_6Cl_2O_2$	α-1, 2-Dichloroanthraquinone	276 96	161			
4597	$C_{14}H_6Cl_2O_2$	β-1, 2-Dichloroanthraquinone	276 96	207			
4598	$C_{14}H_6Cl_2O_2$	1, 4-Dichloroanthraquinone	276 96	187 5			
4599	$C_{14}H_6Cl_2O_2$	1, 5-Dichloroanthraquinone	276 96	232			
4600	$C_{14}H_6Cl_2O_2$	1, 6-Dichloroanthraquinone	276 96	204			
4601	$C_{14}H_6Cl_2O_2$	1, 8-Dichloroanthraquinone	276 96	199			
4602	$C_{14}H_6Cl_2O_2$	2, 3-Dichloroanthraquinone	276 96	267			
4603	$C_{14}H_6Cl_2O_2$	2, 6-Dichloroanthraquinone	276 96	282			
4604	$C_{14}H_6Cl_2O_2$	2, 7-Dichloroanthraquinone	276 96	211			
4605	$C_{14}H_6Cl_4$	1, 2, 3, 4-Tetrachloroanthracene	315 88	149			
4606	$C_{14}H_6Cl_4$	α-Tetrachloroanthracene...	315 88	220			
4607	$C_{14}H_6Cl_4$	β-Tetrachloroanthracene.....	315 88	152			
4608	$C_{14}H_6N_2O_6$	1, 3-Dinitroanthraquinone	298 06	240			
4609	$C_{14}H_6O_8$	Ellagic acid	302 05			1.667[18]	
4610	$C_{14}H_7ClO_2$	1-Chloroanthraquinone . .	242 51	162			
4611	$C_{14}H_7ClO_2$	2-Chloroanthraquinone .	242 51	208			
4612	$C_{14}H_7ClO_2$	3-Chloroanthraquinone	242 51	204			
4613	$C_{14}H_7NO_4$	1-Nitroanthraquinone .	253 06	230			
4614	$C_{14}H_7NO_4$	2-Nitroanthraquinone	253 06	181			
4615	$C_{14}H_7NO_6$	4-Nitro-α-alizarin	285 06	289			
4616	$C_{14}H_7NO_6$	3-Nitro-β-alizarin ...	285.06	244			
4617	$C_{14}H_8Br_2$	9, 10-Dibromoanthracene	335 89	221			
4618	$C_{14}H_8Cl_2$	1, 2-Dichloroanthracene.	246 98	255			
4619	$C_{14}H_8Cl_2$	9, 10-Dichloroanthracene	246 98	209			
4620	$C_{14}H_8O_2$	Anthraquinone $C_6H_4:(CO)_2.C_6H_4$	208 06	285	379.8	1.438	
4621	$C_{14}H_8O_2$	Isoanthraquinone .	208 06	212			
4622	$C_{14}H_8O_2$	Phenanthraquinone	208 06	207	360	1.405	
4623	$C_{14}H_8O_2$	3, 4-Phenanthraquinone . .	208 06	133			
4624	$C_{14}H_8O_3$	2-Hydroxyanthraquinone	224 06	302			
4625	$C_{14}H_8O_3$	Diphenic anhydride	224 06	219			
4626	$C_{14}H_8O_4$	Alizarin	240 06	290	430		
4627	$C_{14}H_8O_4$	Anthraflavic acid ..	240 06	330			
4628	$C_{14}H_8O_4$	Anthrarufin... ...	240.06	280			
4629	$C_{14}H_8O_4$	1, 6-Dihydroxyanthraquinone	240 06	272			
4630	$C_{14}H_8O_4$	1, 7-Dihydroxyanthraquinone	240 06	292			
4631	$C_{14}H_8O_4$	Chrysazin.. ...	240.06	191			
4632	$C_{14}H_8O_4$	Hystazarin (2, 3-Dihydroxyanthraquinone)...... ...	240.06	>280			
4633	$C_{14}H_8O_4$	Quinizarin. . .	240 06	195			
4634	$C_{14}H_8O_4$	Xanthopurpurin	240 06	263			
4635	$C_{14}H_8O_5$	Anthragallol	256 06	310	s. 290		
4636	$C_{14}H_8O_5$	Anthrapurpurin .	256 06	330	402		
4637	$C_{14}H_8O_5$	Flavopurpurin	256 06	>360 '	459		
4638	$C_{14}H_8O_5$	Purpurin...	256 06	256			
4639	$C_{14}H_8O_5$	1, 4, 6-Trihydroxyanthraquinone .	256 06	>300		1.171[20 4]	1140
4640	$C_{14}H_9Cl$	1-Chloroanthracene . .	212 53	82			
4641	$C_{14}H_9Cl$	9-Chloroanthracene......	212 53	103			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4642	C₁₄H₉NO₂	1-Aminoanthraquinone	223 08	256			
4643	C₁₄H₉NO₂	2-Aminoanthraquinone	223 08	302			
4644	C₁₄H₉NO₂	9-Nitroanthracene	223 .08	146			
4645	C₁₄H₉NO₂	2-Nitrophenanthrene	223 08	99			
4646	C₁₄H₉NO₂	3-Nitrophenanthrene	223 08	170			
4647	C₁₄H₉NO₂	4-Nitrophenanthrene	223 08	80			
4648	C₁₄H₉NO₂	9-Nitrophenanthrene	223 08	116			
4649	C₁₄H₁₀	Anthracene C₆H₄(CH)₂.C₆H₄	178 08	218	342	1.25₄²⁷	
4650	C₁₄H₁₀	Diphenylacetylene C₆H₅C.C.C₆H₅	178 08	60	300		
4651	C₁₄H₁₀	Isoanthracene	178 08	134 5			
4652	C₁₄H₁₀	Phenanthrene	178 08	99 6	340 2	1 025	1158
4653	C₁₄H₁₀Cl₂	Dichlorostilbene	248 99	170			
4654	C₁₄H₁₀Cl₂	α-Tolane dichloride	248 99	143	183¹⁵		
4655	C₁₄H₁₀Cl₂	β-Tolane dichloride	248 99	63	178¹⁸		
4656	C₁₄H₁₀Cl₄	Tolane tetrachloride	319 91	163			
4656 1	C₁₄H₁₀N₂O₂	Phthalylphenylhydrazine	238 09	179		1.356	
4657	C₁₄H₁₀N₂O₂	α-Diaminoanthraquinone	238 09	236			
4658	C₁₄H₁₀N₂O₂	β-Diaminoanthraquinone	238 09	>300			
4659	C₁₄H₁₀N₂O₃	p, p'-Azoxybenzaldehyde	254 09	194			
4660	C₁₄H₁₀N₂O₄	o, o'-Azobenzoic acid	270 09	237			
4661	C₁₄H₁₀N₂O₄	m, m'-Azobenzoic acid	270 09	340			
4662	C₁₄H₁₀N₂O₄	α-p, p'-Dinitrostilbene	270 09	285			
4663	C₁₄H₁₀N₂O₄	β-p, p'-Dinitrostilbene	270 09	216			
4664	C₁₄H₁₀N₂O₅	o, o'-Azoxybenzoic acid	286 09	240			
4665	C₁₄H₁₀N₂O₅	m, m'-Azoxybenzoic acid	286 09	320			
4666	C₁₄H₁₀N₂O₅	p, p'-Azoxybenzoic acid	286 09	240 d.			
4667	C₁₄H₁₀O	Anthranol	194 08	170 d.			
4668	C₁₄H₁₀O	1-Anthrol (1-Hydroxyanthracene)	194 08	153			
4669	C₁₄H₁₀O	2-Anthrol	194 08	200 d.			
4670	C₁₄H₁₀O	Diphenylketene (C₆H₅)₂C:CO	194 08		146¹²	1.104	
4671	C₁₄H₁₀O	Phenanthrone	194 08	152			
4672	C₁₄H₁₀O₂	Benzil C₆H₅COCOC₆H₅	210 08	95 2	348	1.521₄¹² ³	1186
4673	C₁₄H₁₀O₂	Chrysazol	210 08	220 d.			
4674	C₁₄H₁₀O₂	Flavene	210 08	270			
4675	C₁₄H₁₀O₂	3, 4-Dihydroxyphenanthrene	210 08	143			
4676	C₁₄H₁₀O₃	Benzoic anhydride (C₆H₅CO)₂O	226 08	43	360	1.199₄¹⁵	
4677	C₁₄H₁₀O₃	o-Benzoylbenzoic acid	226 08	127			
4678	C₁₄H₁₀O₃	m-Benzoylbenzoic acid	226 08	162			
4679	C₁₄H₁₀O₃	p-Benzoylbenzoic acid	226 08	194			
4680	C₁₄H₁₀O₃	Desoxyalizarin	226 08	208			
4681	C₁₄H₁₀O₃	Disalicylic aldehyde	226 08	128			
4682	C₁₄H₁₀O₄	Benzoylsalicylic acid	242 08	207			
4683	C₁₄H₁₀O₄	1, 8-Diphenic acid	242 08	252			
4684	C₁₄H₁₀O₄	1, 9-Diphenic acid	242 08	216			
4685	C₁₄H₁₀O₄	1, 10-Diphenic acid	242 08	228			
4686	C₁₄H₁₀O₄	2, 9-Diphenic acid	242 08	340			
4687	C₁₄H₁₀O₄	Diphenyl oxalate (CO₂C₆H₅)₂	242 08	136 d.	325 s. d.		
4688	C₁₄H₁₀O₄	Benzoyl peroxide (C₆H₅CO₂)₂	242 08	104	d.		1235
4689	C₁₄H₁₀O₄S₂	Dithiosalicylic acid	306 21	290			
4690	C₁₄H₁₀O₅	Gentianin	258 08	267	400		
4691	C₁₄H₁₀O₅	Gentienin	258 08	225			
4692	C₁₄H₁₀O₅	Salicylosalicylic acid	258 08	148			
4693	C₁₄H₁₀O₆	Aponic acid	274 08	252 d.			
4694	C₁₄H₁₀O₉	Tannin	322 08	200 d.			
4695	C₁₄H₁₁N	α-Anthramine C₆H₄:(CH)₂:C₆H₃NH₂	193 09	130			
4696	C₁₄H₁₁N	β-Anthramine C₆H₄.(CH)₂.C₆H₃NH₂	193 09	238			
4697	C₁₄H₁₁N	o-Benzylbenzonitrile	193 09	19	314		
4698	C₁₄H₁₁N	1-Methylacridine	193 09	88			
4699	C₁₄H₁₁N	3-Methylacridine	193 09	134			
4700	C₁₄H₁₁N	5-Methylacridine	193 09	114	360⁷⁴⁰		
4701	C₁₄H₁₁N	α-Naphthoquinaldine	193 09		>300		
4702	C₁₄H₁₁N	β-Naphthoquinaldine	193 09	82	>300		
4703	C₁₄H₁₁N	γ-Naphthoquinaldine	193 .09	92			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4704	$C_{14}H_{11}NO_2$	α-Benziloxime $C_6H_5COC(:NOH)C_6H_5$	225 09	138			
4705	$C_{14}H_{11}NO_3$	Dibenzohydroxamic acid	241 09	161			
4706	$C_{14}H_{11}NO_4$	Disalicylamide	257 09	200 d.			
4707	$C_{14}H_{12}$	1, 1-Diphenylethylene $(C_6H_5)_2C.CH_2$	180 09	9	277	1 038¹⁴	837
4708	$C_{14}H_{12}$	Stilbene $C_6H_5CH:CHC_6H_5$	180 09	124	307	0 970¹⁵	
4709	$C_{14}H_{12}N_2$	Benzalazine $C_6H_5CH:N.NCH C_6H_5$	208 11	93			
4710	$C_{14}H_{12}N_2$	Orexine .	208 .11	95		1 290⁴	
4711	$C_{14}H_{12}N_2$	Tolazone . .	208 11	187	>360		
4712	$C_{14}H_{12}N_2O_2$	α-Benzildioxime $(C_6H_5C:NOH)_2$	240 11		237 d.		
4713	$C_{14}H_{12}N_2O_2$	β-Benzildioxime . .	240 11	105			
4714	$C_{14}H_{12}N_2O_2$	γ-Benzildioxime	240 11	165			
4715	$C_{14}H_{12}N_2O_2$	Oxanilide $(CONHC_6H_5)_2$	240 11	250	320		
4716	$C_{14}H_{12}N_2O_4$	Di-o-aminophenyl oxalate	272 11	167 5 d.			
4717	$C_{14}H_{12}N_2O_4$	Di-m-aminophenyl oxalate	272 11	180 d.			
4718	$C_{14}H_{12}N_2O_4$	Di-p-aminophenyl oxalate	272 11	220 d.			
4719	$C_{14}H_{12}N_2O_4$	Hydrazo-o-benzoic acid	272 11	205			
4722	$C_{14}H_{12}N_2S$	Dehydrothio-p-toluidine	240 17	191	434		
4723	$C_{14}H_{12}O$	Diphenylacetaldehyde	196 09		193²⁷	1 100	775
4724	$C_{14}H_{12}O$	Phenyl benzyl ketone	196 09	60	322		
4725	$C_{14}H_{12}O$	Phenyl o-tolyl ketone	196 09	> −18	316		
4726	$C_{14}H_{12}O$	Phenyl m-tolyl ketone	196 09		316 5	1 088¹⁷ ⁵	
4727	$C_{14}H_{12}O$	Phenyl p-tolyl ketone	196 09	60	326 5		1188
4728	$C_{14}H_{12}O_2$	Benzoin $C_6H_5COCH(OH)C_6H_5$	212 09	133	344		
4729	$C_{14}H_{12}O_2$	o-Benzylbenzoic acid	212 09	114			
4730	$C_{14}H_{12}O_2$	m-Benzylbenzoic acid	212 09	108			
4731	$C_{14}H_{12}O_2$	p-Benzylbenzoic acid	212 09	155			
4732	$C_{14}H_{12}O_2$	Diphenylacetic acid $(C_6H_5)_2CHCO_2H$	212 09	148			
4733	$C_{14}H_{12}O_2$	Benzyl benzoate $C_6H_5CO_2CH_2C_6H_5$	212 09	18 5	321	1 114¹⁵ ⁵	
4734	$C_{14}H_{12}O_2$	p-Cresyl benzoate p-$CH_3C_6H_4O_2CC_6H_5$	212 09	71 5	316		
4735	$C_{14}H_{12}O_3$	Benzyl salicylate	228 09		214²² ⁵		
4736	$C_{14}H_{12}O_2$	m-Cresyl benzoate $C_6H_5CO_2C_6H_4CH_3$	212 09	55			
4737	$C_{14}H_{12}O_3$	Trihydroxydihydroanthracene	228 09	256			
4738	$C_{14}H_{12}O_3$	Benzilic acid $(C_6H_5)_2C(OH)CO_2H$	228 09	150			
4739	$C_{14}H_{12}O_3$	Amyrolin	228 09	124		1.351¹⁸	1312
4740	$C_{14}H_{12}O_3$	Benzosol $C_6H_5CO_2C_6H_4(OCH_3)$-o	228 09	61			
4741	$C_{14}H_{12}O_3$	o-Cresyl salicylate	228 09	35			
4742	$C_{14}H_{12}O_3$	m-Cresyl salicylate	228 09	74			
4743	$C_{14}H_{12}O_3$	p-Cresyl salicylate	227 09	39			
4744	$C_{14}H_{12}O_4$	Cotoin . . .	224 09	129			
4745	$C_{14}H_{12}O_4$	Isocotoin . .	244 09	162			
4746	$C_{14}H_{12}O_4$	Guaiacyl salicylate	244 09	65			
4747	$C_{14}H_{12}O_5$	Gardenin .	276 .09	164			
4748	$C_{14}H_{13}NO$	N-Benzoyl-o-toluidine	211 .11	143			1296
4749	$C_{14}H_{13}NO$	N-Benzoyl-m-toluidine	211 .11	125			1209
4750	$C_{14}H_{13}NO$	N-Benzoyl-p-toluidine	211 11	158	232		1291
4751	$C_{14}H_{13}NO$	o-Benzylbenzamide	211 11	163			
4752	$C_{14}H_{13}NO$	N-Diphenylacetamide	211 .11	103			1281
4753	$C_{14}H_{13}NO$	Phenylacetanilide .	211 .11	117			
4754	$C_{14}H_{13}NO_2$	Benzoylanisidine	227 11	154			
4755	$C_{14}H_{13}N_3O$	m-Acetylaminoazobenzene	239 12	131			
4756	$C_{14}H_{14}$	Dibenzyl $(C_6H_5CH_2)_2$	182 11	52 .5	284	0 942⁴⁰ ⁵	1118
4757	$C_{14}H_{14}$	1, 1-Diphenylethane $(C_6H_5)_2CHCH_3$	182.11		272	1 006²¹	763
4758	$C_{14}H_{14}$	o, o'-Ditolyl $(CH_3C_6H_4)_2$ . .	182.11	17 8	272	0 955¹⁰	
4759	$C_{14}H_{14}$	o, m'-Ditolyl $(CH_3C_6H_4)_2$ . .	182 11		287 5		
4760	$C_{14}H_{14}$	o, p'-Ditolyl $(CH_3C_6H_4)_2$ . .	182.11		281		
4761	$C_{14}H_{14}$	m, m'-Ditolyl $(CH_3C_6H_4)_2$	182.11	7	288	0.999	
4762	$C_{14}H_{14}$	p, p'-Ditolyl $(CH_3C_6H_4)_2$	182.11	121	295		
4763	$C_{14}H_{14}N_2$	o, o'-Azotoluene $(o$-$CH_3C_6H_4N)_2$	210 12	55			
4764	$C_{14}H_{14}N_2$	o', p'-Azotoluene . . .	210 12	71			
4765	$C_{14}H_{14}N_2$	m, m'-Azotoluene $(m$-$CH_3C_6H_4)_2N_2$	210.12	55			
4766	$C_{14}H_{14}N_2$	p, p'-Azotoluene $(p$-$CH_3C_6H_4)_2N_2$	210 12	144			
4767	$C_{14}H_{14}N_2$	o, o'-Diaminostilbene	210.12	170			
4768	$C_{14}H_{14}N_2$	p, p'-Diaminostilbene	210.12	231			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4769	$C_{14}H_{14}N_2O$	Agathin o-OHC$_6$H$_4$CH:N.N(CH$_3$)C$_6$H$_5$.	226.12	74			
4770	$C_{14}H_{14}N_2O$	o, o'-Azoxytoluene	226.12	59			
4771	$C_{14}H_{14}N_2O$	m, m'-Azoxytoluene	226 12	37			
4772	$C_{14}H_{14}N_2O$	p, p'-Azoxytoluene	226 12	70			
4773	$C_{14}H_{14}N_2O_2$	o, o'-Azoanisol (o-CH$_3$OC$_6$H$_4$)$_2$N$_2$.	242 12	164 0			
4774	$C_{14}H_{14}N_2O_2$	p, p'-Azoxyanisol (p-CH$_3$OC$_6$H$_4$)$_2$N$_2$.	258 12	117.4			
4775	$C_{14}H_{14}N_4$	"Cyanaline"	238 14	220			
4776	$C_{14}H_{14}N_4O_4$	Theobromine salicylate	318 14				1333
4777	$C_{14}H_{14}O$	Benzyl ether (C$_6$H$_5$CH$_2$)$_2$O	198 11		298	1 036^{16}	
4778	$C_{14}H_{14}O$	o-Cresyl ether (CH$_3$C$_6$H$_4$)$_2$O	198.11		278	1.047^{24} ^{3}	
4779	$C_{14}H_{14}O$	m-Cresyl ether (CH$_3$C$_6$H$_4$)$_2$O	198.11		288		
4780	$C_{14}H_{14}O$	p-Cresyl ether (p-CH$_3$C$_6$H$_4$)$_2$O	198 11	50			
4781	$C_{14}H_{14}O_2$	dl-Hydrobenzoin [C$_6$H$_5$CH(OH)]$_2$	214 11	139	>300		
4782	$C_{14}H_{14}O_2$	Guaiacyl benzyl ether	214 11	62			
4783	$C_{14}H_{14}O_2$	Isohydrobenzoin	214 11	121			
4784	$C_{14}H_{14}O_2S$	Dibenzylsulfone (C$_6$H$_5$CH$_2$)$_2$SO$_2$	246 17	150	290 s. d.		
4785	$C_{14}H_{14}O_2S$	p-Ditolylsulfone (CH$_3$C$_6$H$_4$)$_2$SO$_2$	246 17	158	405^{714}		
4786	$C_{14}H_{14}S_2$	Dibenzyl disulfide (C$_6$H$_5$CH$_2$)$_2$S$_2$	246 24	72			
4787	$C_{14}H_{14}S$	Dibenzylsulfide (C$_6$H$_5$CH$_2$)$_2$S	214.17	49		1 071$^{50}_{50}$	
4788	$C_{14}H_{14}Se$	Dibenzyl selenide (C$_6$H$_5$CH$_2$)$_2$Se	261 31	45 5			
4789	$C_{14}H_{15}N$	Dibenzylamine (C$_6$H$_5$CH$_2$)$_2$NH	197.12	−26 0	300	1 026$^{21}_{4}$ ^{6}	976
4790	$C_{14}H_{15}N$	o-Ditolylamine (o-CH$_3$C$_6$H$_4$)$_2$NH.	197.12		313 4		
4791	$C_{14}H_{15}N$	m-Ditolylamine (m-CH$_3$C$_6$H$_4$)$_2$NH	197 12		320		
4792	$C_{14}H_{15}N$	p-Ditolylamine (p-CH$_3$C$_6$H$_4$)$_2$NH	197 12	79	330 5		
4793	$C_{14}H_{15}N$	Ethyldiphenylamine (C$_6$H$_5$)$_2$NC$_2$H$_5$	197 12		297		
4794	$C_{14}H_{15}N$	N-Methylbenzylaniline	197 12	9 2	306		
4795	$C_{14}H_{15}NO_2S$	p-Toluenesulfonemethylanilide	261 19	95			
4796	$C_{14}H_{15}N_3$	4-Amino-2, 4'-dimethylazobenzene	225 14	127			
4797	$C_{14}H_{15}N_3$	4'-Amino-2, 3'-dimethylazobenzene	225 14	100			
4798	$C_{14}H_{15}N_3$	4-Amino-2, 3'-dimethylazobenzene	225 14	80			
4799	$C_{14}H_{15}N_3$	4-Amino-3, 4'-dimethylazobenzene	225 14	127			
4800	$C_{14}H_{15}N_3$	o, o'-Diazoaminotoluene	225 14	51			
4801	$C_{14}H_{15}N_3$	p, p'-Diazoaminotoluene	225 14	116			
4802	$C_{14}H_{16}$	Hexahydroanthracene	184 12	63	290		
4803	$C_{14}H_{16}N_2$	o-Hydrazotoluene (o-CH$_3$C$_6$H$_4$NH)$_2$	212 14	165			
4805	$C_{14}H_{16}N_2$	p-Hydrazotoluene (CH$_3$C$_6$H$_4$NH)$_2$	212 14	126	d.	0.957	
4806	$C_{14}H_{16}N_2$	o-Tolidine [4, 3-H$_2$N(CH$_3$)C$_6$H$_3$]$_2$	212 14	129			
4807	$C_{14}H_{16}N_2$	m-Tolidine [4, 2-H$_2$N(CH$_3$)C$_6$H$_3$]$_2$	212 14	107			
4808	$C_{14}H_{16}N_2O$	3-Ethoxybenzidine	228 14	139			
4809	$C_{14}H_{16}N_2O_2$	3, 3'-Dimethoxybenzidine	244 14	172			
4810	$C_{14}H_{16}N_4$	2, 2'-Diamino-4, 4'-azotoluene	240 16	203			
4811	$C_{14}H_{16}N_4$	3, 3'-Diamino-2, 2'-azotoluene	240.16	a, 145; b, 133; c, 159			
4812	$C_{14}H_{16}N_4O_9$	Oscine picrate	384.16	238			
4813	$C_{14}H_{17}N$	Diethyl-α-naphthylamine	199.14		160 6^{18}	1 005	937
4814	$C_{14}H_{17}N$	Diethyl-β-naphthylamine	199.14		192^{19}	1 026	977
4815	$C_{14}H_{17}NO$	Etheserolene.	215 14	48			
4816	$C_{14}H_{17}NO_6$	Indican	295.14	57			
4817	$C_{14}H_{17}NO_6$	l-Mandelonitrile glucoside	295 14	147			
4818	$C_{14}H_{17}NO_6$	Prulaurasin	295 14	122			
4819	$C_{14}H_{17}NO_6$	Sambunigrin	295 14	152			
4820	$C_{14}H_{18}O_5$	Apocynamarin	234 14	175 d.			
4821	$C_{14}H_{18}O_7$	Picein	298 14	194			
4822	$C_{14}H_{20}N_2O_6S$	Methylamino-p-phenol sulfate	344 24	260 d.			
4823	$C_{14}H_{20}O_2$	Isanic acid	220 15	41			
4823 1	$C_{14}H_{20}O_2$	l-Amyl hydrocinnamate	220.15		172^{28}	0 9721	
4824	$C_{14}H_{20}O_3$	Helleboretin	236 15	>200			
4825	$C_{14}H_{21}ClN_2O_4$	Nirvanin	316 64	185			
4826	$C_{14}H_{21}NO_2$	Thymacetine	235 17	136			
4827	$C_{14}H_{22}$	1, 2, 3, 4-Tetraethylbenzene	190.17		254	0 887	637
4828	$C_{14}H_{22}$	1, 2, 4, 5-Tetraethylbenzene	190 17	13	250	0 888	609
4829	$C_{14}H_{22}ClNO_2$	Stovain	271.64	175			
4830	$C_{14}H_{22}O_2$	Longifolic acid	222 17	153	234^{54}		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4831	$C_{14}H_{22}O_4$	Dicyclohexyl oxalate	254 17	45	191¹⁴		
4831.1	$C_{14}H_{22}ClO_4$	Di-l-amyl chlorofumarate	290 65		185¹⁴	1 052²⁴	
4832	$C_{14}H_{23}N$	N-Dibutylaniline C₆H₅N(C₄H₉)₂	205 19		262 8		
4832 1	$C_{14}H_{23}N$	Diisobutylaniline	205 19		146⁷¹	0 909¹⁶	
4833	$C_{14}H_{24}O_2$	Kersyl alcohol.	224 19	85	156¹¹		
4834	$C_{14}H_{24}O_2$	d-Bornyl n-butyrate.	224 19		121¹¹	0 966¹⁵	856
4835	$C_{14}H_{24}O_2$	Geranyl butyrate .	224 19		153¹⁸	0 901	
4836	$C_{14}H_{24}O_2$	l-Menthyl crotonate	224 19		140 5¹⁴	0 833	
4837	$C_{14}H_{24}O_3$	l-Menthyl acetoacetate	240 19	45	145¹¹	0 986¹⁵₁₅	
4837 1	$C_{14}H_{24}O_4$	Di-l-amyl maleate	256 19		165²⁵	0 9708¹⁴	
4838	$C_{14}H_{24}O_4$	l-Menthyl acid succinate	256 19	62	300 d.		
4839	$C_{14}H_{25}NO_2$	Carpaine..	239 20	121			1333
4840	$C_{14}H_{26}ClNO_2$	Carpaine hydrochloride	275 67	225			
4841	$C_{14}H_{26}O_2$	l-Menthyl n-butyrate	226 20		129¹⁵	0 911	
4842	$C_{14}H_{26}O_2$	l-Menthyl isobutyrate	226 20		117¹²	0 906	
4843	$C_{14}H_{26}O_3$	n-Heptylic anhydride (C₆H₁₃CO)₂O	242 20	17	258	0 932	332
4844	$C_{14}H_{26}O_3$	Menthyl ethyl glycollate	242 20		155²⁰		
4845	$C_{14}H_{26}O_4$	Diamyl succinate	258 20		293	0 952¹⁵₁₅	
4845 1	$C_{14}H_{26}O_4$	Di-l-amyl succinate	258 20		129¹	0 957¹⁵	
4846	$C_{14}H_{26}O_4$	Diethyl sebacate..	258 20	1	308	0 965¹⁶	
4846.1	$C_{14}H_{26}O_6$	Diisoamyl tartrate .	290 20		195¹⁶	1 063¹⁶	
4847	$C_{14}H_{27}ClO$	Myristyl chloride CH₃(CH₂)₁₂COCl.	246 67	−1	168¹⁵		
4848	$C_{14}H_{27}N$	Myristic nitrile CH₃(CH₂)₁₂CN	209 22	19	226¹⁰⁰	0 828	
4849	$C_{14}H_{28}$	n-Tetradecylene	196 22	−12	246	0 775	
4850	$C_{14}H_{28}O$	Myristic aldehyde CH₃(CH₂)₁₂CHO	212 22	52 5	166²⁴		
4851	$C_{14}H_{28}O_2$	Myristic acid CH₃(CH₂)₁₂CO₂H	228 22	58	250 5¹⁰⁰	0 858⁶⁰₆₀	1088
4852	$C_{14}H_{28}O_2$	Ethyl laurate C₁₁H₂₃CO₂C₂H₅	228 22	−10 7	269	0.868¹⁸₄	337
4853	$C_{14}H_{28}O_3$	Hydroxymyristic acid........	244 22	51			
4854	$C_{14}H_{28}O_4$	Ipurolic acid	260 22	101			
4855	$C_{14}H_{29}NO$	Myristic amide CH₃(CH₂)₁₂CONH₂	227 23	103			
4856	$C_{14}H_{30}$	n-Tetradecane CH₃(CH₂)₁₂CH₃	198 23	5.5	252 5	0.765	412
4857	$C_{14}H_{30}O$	n-Heptyl ether (C₇H₁₅)₂O...	214 23		260	0 815⁰	
4858	$C_{14}H_{30}O$	n-Tetradecyl alcohol C₁₃H₂₇CH₂OH	214 23	38	167¹⁵	0 824⁴¹₄	
4859	$C_{14}H_{31}N$	Tetradecyl amine C₁₃H₂₇CH₂NH₂	213 25	37	162¹⁵		
4860	$C_{15}H_8O_4$	Anthraquinone-α-carboxylic acid	252 06	294			
4861	$C_{15}H_8O_4$	Anthraquinone-β-carboxylic acid	252 06	288			
4862	$C_{15}H_8O_4$	Anthraquinone-γ-carboxylic acid	252 06	285			
4863	$C_{15}H_8O_6$	Alizarin-β-carboxylic acid	284 06	305			
4864	$C_{15}H_8O_7$	Pseudopurpurin.	300 06	220			
4865	$C_{15}H_9N$	Thebenidine.....	203 08	148			
4866	$C_{15}H_{10}$	Fluoranthene. .	190 08	110	251⁶⁰		
4867	$C_{15}H_{10}$	Succisterene	190 08	160	300		
4868	$C_{15}H_{10}O_2$	Flavone .	222 08	97			
4869	$C_{15}H_{10}O_2$	Anthracene-1-carboxylic acid	222 08	260			
4870	$C_{15}H_{10}O_2$	Anthracene-2-carboxylic acid	222 08	280			
4871	$C_{15}H_{10}O_2$	Anthracene-9-carboxylic acid	222 08	206			
4872	$C_{15}H_{10}O_2$	1-Methylanthraquinone	222 08	171			
4873	$C_{15}H_{10}O_2$	2-Methylanthraquinone	222 08	175			
4874	$C_{15}H_{10}O_4$	Chrysine.....	254 08	275			
4875	$C_{15}H_{10}O_4$	Chrysophanic acid	254 08	193			
4876	$C_{15}H_{10}O_4$	α-Methylalizarin	254 08	229			
4877	$C_{15}H_{10}O_4$	β-Methylalizarin	254 08	179			
4878	$C_{15}H_{10}O_4$	Rumicin	254 08	182			
4879	$C_{15}H_{10}O_5$	Aloe-emodin. .	270 08	218			
4880	$C_{15}H_{10}O_5$	Emodin..	270 08	250			
4881	$C_{15}H_{10}O_5$	Galangin. .	270 08	217			
4882	$C_{15}H_{10}O_5$	Morindon .	270 08	275			
4883	$C_{15}H_{10}O_6$	Fisetin.	286 08	360			
4884	$C_{15}H_{10}O_6$	Kaempferol ..	286 08	274			
4885	$C_{15}H_{10}O_6$	Luteolin.	286 08	320			
4886	$C_{15}H_{10}O_6$	Rhein	286 08	314			
4887	$C_{15}H_{10}O_6$	Scutellarein..	286 08	300 d.			
4888	$C_{15}H_{10}O_7$	Morin.................	302 08	285			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
4889	$C_{15}H_{10}O_7$	Quercetin......	302.08	310			
4890	$C_{15}H_{10}O_8$	Gossypetin	318 08	230			
4891	$C_{15}H_{10}O_8$	Quercetagetin.	318 08	318			
4892	$C_{15}H_{11}N$	2-Phenylquinoline	205 09	86	363		
4893	$C_{15}H_{11}N$	4-Phenylquinoline	205 09	62			
4894	$C_{15}H_{11}N$	6-Phenylquinoline	205 09	111	260^{77}	1.195	
4895	$C_{15}H_{11}N$	8-Phenylquinoline	205 09		283^{187}		
4896	$C_{15}H_{11}NO$	Benzoylphenylacetonitrile	221 09	99			
4897	$C_{15}H_{12}$	α-Methylanthracene	192 09	86	200	$1 047^{99\ 4}$	1134
4898	$C_{15}H_{12}$	2-Methylanthracene	192 09	207			
4899	$C_{15}H_{12}$	9-Methylanthracene	192 09	80		$1 066^{99\ 4}$	1136
4900	$C_{15}H_{12}N_2O_3$	Furfuramide	268 11	121	250 d.		
4901	$C_{15}H_{12}N_2O_3$	Furfurine	268 11	116			
4902	$C_{15}H_{12}O$	Benzylideneacetophenone	208 09	62	348	1.071_4^{62}	
4903	$C_{15}H_{12}O_2$	Benzoylacetophenone	224 09	81	>200		
4904	$C_{15}H_{12}O_3$	p-Toluyl-o-benzoic acid ..	240 09	139			
4905	$C_{15}H_{12}O_4$	Chrysophanol	240 09	204			
4906	$C_{15}H_{12}O_4$	Acetylsalol $o\text{-}CH_3CO_2C_6H_4CO_2C_6H_5$	256 09	97	198		
4907	$C_{15}H_{12}O_4$	Benzosalin	256 09	85	385		
4908	$C_{15}H_{12}O_4$	Diphenyl malonate $CH_2(CO_2C_6H_5)_2$	256 09	50	210^{15} d.		
4909	$C_{15}H_{12}O_6$	Eriodictyol	288.09	267			
4910	$C_{15}H_{12}O_5$	Methylenedisalicylic acid	288 09	238 d.			
4911	$C_{15}H_{13}NO_4$	Salophen	271.11	188			
4912	$C_{15}H_{14}O$	Benzylacetophenone..	210.11	73	360		
4913	$C_{15}H_{14}O$	Benzyl p-tolyl ketone	210 11	109	360		
4914	$C_{15}H_{14}O$	Dibenzyl ketone $(C_6H_5CH_2)_2CO$	210 11	33 9	330 5		
4915	$C_{15}H_{14}O$	p, p'-Dimethylbenzophenone	210 11	92	335.1		
4916	$C_{15}H_{14}O_2$	Benzyl o-toluate	226.11		315	$1 12^{17}$	
4917	$C_{15}H_{14}O_3$	Benzyl phenylacetate	226 11		319	1 101	
4918	$C_{15}H_{14}O_3$	Benzyl mandelate	242.11	93			
4919	$C_{15}H_{14}O_3$	Methyl benzilate	242 11	73			
4920	$C_{15}H_{14}O_3$	Lapachol	242 11	140			
4921	$C_{15}H_{14}O_4$	Hydrocotoin	258.11	95 5			
4922	$C_{15}H_{14}O_4$	Peucedanin	258 11	109			
4923	$C_{15}H_{14}O_4$	N-Xanthoxyllin	258.11	132 5			
4924	$C_{15}H_{14}O_5$	Guaiacyl carbonate $(o\text{-}CH_3OC_6H_4O)_2CO.$	274 11	86			
4925	$C_{15}H_{14}O_5$	Kavaiin (Methysticin)	274 11	137			
4926	$C_{15}H_{14}O_5$	Phloretin	274.11	255 d.			1333
4927	$C_{15}H_{15}NO$	p-Dimethylaminobenzophenone	225 12	90			
4928	$C_{15}H_{15}NO_3$	Malakin	257 12	92			
4929	$C_{15}H_{15}NO_3$	Narceine acid	337 12	184			
4930	$C_{15}H_{16}$	Dibenzylmethane $(C_6H_5CH_2)_2CH_2$	196 12	< −20	299	1 007	762
4931	$C_{15}H_{16}N_2O$	sym.-Di-o-tolylurea ..	240 14	256			
4932	$C_{15}H_{16}N_2O$	sym.-Di-m-tolylurea	240 14	203			
4933	$C_{15}H_{16}N_2O$	sym.-Di-p-tolylurea	240.14	263			
4934	$C_{15}H_{16}N_2S$	1, 2-Di-o-tolylthiourea	256 20	156	218		
4935	$C_{15}H_{16}N_2S$	sym -Di-m-tolylthiourea	256 20	111 5			
4936	$C_{15}H_{16}O_3$	Santinic acid	228 12	132 5			
4936.1	$C_{15}H_{16}O_4$	Picrotoxinin	292 12	206			1265
4937	$C_{15}H_{16}O_9$	Daphnin	340 12	200			
4938	$C_{15}H_{16}O_9$	Esculin	340 12	205			
4939	$C_{15}H_{17}N$	Ethylbenzylaniline	211 14		298	$1 034^{18\ 5}$	
4940	$C_{15}H_{17}N_3$	Di-o-tolylguanidine	239 16	179			
4941	$C_{15}H_{18}$	Azulene	198 14		168 4¹¹	0 988	
4942	$C_{15}H_{18}N_2$	p, p'-Diamino-o, o'-ditolylmethane	226 16	149			
4943	$C_{15}H_{18}O_3$	Santonin .	246 14	170		1 187	1282
4944	$C_{15}H_{18}O_4$	Artemisin	262 14	202			1333
4944.1	$C_{15}H_{18}O_4$	Coriamyrtin	262.14	225			
4945	$C_{15}H_{18}O_7$	Hyenanchin.	310 14	234 d.			
4946	$C_{15}H_{18}O_7$	Pierotin	310.14	250			
4947	$C_{15}H_{19}NO_2$	Tropacocaine	245 15	49	d.	1.043_4^{100}	1147
4948	$C_{15}H_{19}NO_9$	Lithuric acid	357 15	204 5			
4949	$C_{15}H_{20}ClNO_2$	Tropacocaine hydrochloride...........	281 62	271			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
4950	$C_{15}H_{20}O_2$	Alantolactone ...	232 15	76	192^{10}		
4951	$C_{15}H_{20}O_3$	Perezone	248 15	105			
4952	$C_{15}H_{20}O_3$	Pipitzol	248 15	141			
4953	$C_{15}H_{20}O_4$	Absinthiin . . .	264 15	68			
4954	$C_{15}H_{20}O_4$	Isosantonic acid...	264 15	155	160^4		
4955	$C_{15}H_{20}O_4$	dl-Santonic acid...	264 15	120 d.			
4956	$C_{15}H_{20}O_4$	d(l)-Satonic acid .	264 15	179	260^5	1 251	1333
4957	$C_{15}H_{20}O_8$	Androsin	328 15	220			
4958	$C_{15}H_{21}NO_2$	β-Eucaine	247 17	91			
4959	$C_{15}H_{21}NO_4$	Ajacine	279 17	143			
4960	$C_{15}H_{21}N_3O_2$	Physostigmine	275 19	105			1263
4961	$C_{15}H_{21}N_3O_3$	Geneserine	291 19	129			
4962	$C_{15}H_{22}BrN_3O_2$	Physostigmine hydrobromide	356 11				1333
4963	$C_{15}H_{22}ClNO_2$	β-Eucaine hydrochloride	283 64	268			
4964	$C_{15}H_{22}ClNO_4$	Ajacine hydrochloride	315 64	93			
4965	$C_{15}H_{22}ClN_3O_2$	Physostigmine hydrochloride	311 65				1333
4966	$C_{15}H_{22}O_2$	Santalic acid	234 17		195^9		
4967	$C_{15}H_{22}O_2$	Eugenol isoamyl ether	234 17		302 2 d.	0 976	846
4968	$C_{15}H_{22}O_2$	Thymyl isovalerate	234 17		249	$0 959^{15}_{15}$	
4969	$C_{15}H_{22}O_3$	Alantic (Alantolic) acid	250 17	94			
4970	$C_{15}H_{23}Cl$	Santalyl chloride	238 64		155^{10}	1 040	
4971	$C_{15}H_{24}$	Atractylene	204 19		$111^{14 \cdot}$	0 927	625
4972	$C_{15}H_{24}$	l-Cadinene .	204 19		275	0 918	631
4973	$C_{15}H_{24}$	Cannibene	204 19		259	$0 897^{15}$	
4974	$C_{15}H_{24}$	α-Caryophyllene	204 19		260	0 906	596
4975	$C_{15}H_{24}$	Cedrene. ...	204 19		264	0 929	590
4976	$C_{15}H_{24}$	Clovene . .	204 19		263	0 930	603
4977	$C_{15}H_{24}$	Guajene	204 19		124^9	0 908	602
4978	$C_{15}H_{24}$	Patschoulene	204 19		256	0 930	591
4979	$C_{15}H_{24}$	α-Santalene	204 19		252	$0 913^{15}$	862
4980	$C_{15}H_{24}$	β-Santalene	204 19		126^7	0 894	569
4981	$C_{15}H_{24}$	γ-Santalene .	204 19		120^{10}	0.936	617
4982	$C_{15}H_{24}$	α-Selinene	204.19		135^{16}	0 014	
4983	$C_{15}H_{24}$	Zingiberene..	204.19		270	0.872^{15}	574
4984	$C_{15}H_{24}N_2O$	d(l)-Lupanine	218 20	44			
4985	$C_{15}H_{24}N_2O$	Oxysparteine . .	248 20	84	$209^{12 \, b}$		
4986	$C_{15}H_{24}O$	Betulol	220 19		158^{13}	$0 978^{16}$	865
4987	$C_{15}H_{24}O$	α-Santalol....	220 19		300	0.979^{15}	957
4988	$C_{15}H_{24}O$	β-Santalol	220 19		309	$0 973^{15}$	958
4989	$C_{15}H_{25}BrO_2$	Bornyl bromoisovalerate	317 11		163^{10}		
4990	$C_{15}H_{25}NO_7$	Seneciofolidine . .	331 20	212			
4991	$C_{15}H_{26}$	Elemone.. .	206 20		119^{10}	0 883	
4992	$C_{15}H_{26}$	Ferulene . .	206 20		126^7	0.870	
4993	$C_{15}H_{26}N_2$	Isosparteine	234 22		$179^{16 \, b}$	$1 028^{17}$	916
4994	$C_{15}H_{26}N_2$	Sparteine	234 22		325 2	1 023	959
4995	$C_{15}H_{26}N_2O$	Retamine	250 22	162			
4996	$C_{15}H_{26}O$	Atractylol	222 20	59	292	1.511	
4997	$C_{15}H_{26}O$	Cedrol.....	222 20	87	294		
4998	$C_{15}H_{26}O$	α-Elemol...	222 20	46	143^{10}	$0 941^{21 \, 3}$	967
4999	$C_{15}H_{26}O$	β-Elemol... . .	222 20		144^{10}	0.942^{15}	611
5000	$C_{15}H_{26}O$	Eudesmol .	222 20	78	156^{10}	0.988	657
5001	$C_{15}H_{26}O$	Farnesol.... .	222 20		$120^{0 \, 2}$	0 895	548
5002	$C_{15}H_{26}O$	Guajol	222 20	93	289 s. d.		1175
5003	$C_{15}H_{26}O$	Nerolidol	222 20		277	0 880	891
5004	$C_{15}H_{26}O$	Zingiberol	222 20		$157^{14 \, b}$		
5005	$C_{15}H_{26}O_2$	Bornyl isovalerate	238 20		260	0 949	985
5006	$C_{15}H_{26}O_2$	Isobornyl isovalerate	238.20		138^{12}	$0 957^{15}$	
5007	$C_{15}H_{26}O_2$	d-Bornyl n-valerate .	238.20		130^{11}	0.956^{15}	855
5008	$C_{15}H_{26}O_2$	l-Menthyl angelate..	238 20		141^{10}		
5009	$C_{15}H_{26}O_3$	l-Menthyl levulinate... .	254.20		169^{12}	0.977	
5010	$C_{15}H_{26}O_6$	Tributyrin..........	302.20	< −75	310	1.027	351
5011	$C_{15}H_{27}ClN_2$	Sparteine hydrochloride	270 68				1333
5012	$C_{15}H_{27}IN_2$	Sparteine hydroiodide	362.16				1333

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5013	$C_{16}H_{28}O_2$	l-Menthyl isovalerate	240 22		127[11]	0.907[15]	427
5014	$C_{16}H_{28}O_2$	Cimicic acid	240 22	44 2			
5015	$C_{16}H_{28}O_2$	l-Menthyl n-valerate	240 22		141[15]	0 907	
5016	$C_{16}H_{30}O_2$	Pentadecylic acid	242 23	54	257[100]		
5017	$C_{16}H_{30}O_2$	Methyl myristate	242 23	19	295 3		
5018	$C_{16}H_{32}$	n-Pentadecane $CH_3(CH_2)_{13}CH_3$	212 25	10	270 5	0 772	
5019	$C_{16}H_{32}O$	n-Pentadecyl alcohol $CH_3(CH_2)_{14}OH$	228 25	46			
5020	$C_{16}H_{33}N$	Pentadecylamine	227 26	36 5	301		
5021	$C_{16}H_{33}N$	Triisoamylamine	227 26		237	0 785[21]	
5022	$C_{16}H_8O_6$	Anthraquinone-1, 3-dicarboxylic acid	296 06	330			
5023	$C_{16}H_8O_6$	Anthraquinone-1, 4-dicarboxylic acid	296 06	300			
5024	$C_{16}H_8O_6$	Anthraquinone-2, 3-dicarboxylic acid	296 06	340			
5025	$C_{16}H_{10}$	Diphenyldiacetylene	202 08	88			
5026	$C_{16}H_{10}$	Pyrene	202 08	150	>360		
5027	$C_{16}H_{10}N_2$	α, β-Naphthophenazine	230 09	142 5	>360		
5028	$C_{16}H_{10}N_2O_2$	Indigotin	262 09	392 d.		1 35	
5028 1	$C_{16}H_{10}O_3$	Diphenylmalene anhydride	250 08	155		1 340	1211
5029	$C_{16}H_{10}O_4$	Anthracene-1, 3-dicarboxylic acid	266 08	330			
5030	$C_{16}H_{10}O_4$	Anthracene-1, 4-dicarboxylic acid	266 08	320			
5031	$C_{16}H_{10}O_4$	Anthracene-2, 3-dicarboxylic acid	266 08	345			
5032	$C_{16}H_{10}O_6$	Trifoltin	298 08	275			
5033	$C_{16}H_{11}N$	Amaron	217 09	240			
5034	$C_{16}H_{11}N$	Aminopyrene	217 09	116			
5035	$C_{16}H_{11}NO_2$	Atophan (2-Phenylquinoline-4-carboxylic acid	249.09	209			
5036	$C_{16}H_{11}N_3O_2$	Indigoxime	277 11	205			
5037	$C_{16}H_{12}$	α-Phenylnaphthalene	204 09		325		
5038	$C_{16}H_{12}$	β-Phenylnaphthalene	204 09	102.5	345		
5039	$C_{16}H_{12}$	Pseudophenanthrene	204 09	115			
5040	$C_{16}H_{12}ClNO_2$	Chloroxyl (Phenylcinchoninic acid hydrochloride)	285 56	223			
5041	$C_{16}H_{12}N_2O_4$	Isatid	296 11	237 5			
5042	$C_{16}H_{12}N_4O$	Azoxytolunitrile	276 12	182			
5043	$C_{16}H_{12}O$	Phenyl α-naphthyl ether	220 09	55	340		
5044	$C_{16}H_{12}O$	Phenyl β-naphthyl ether	220 09	45; 93	335 8		
5045	$C_{16}H_{12}O_3S$	Atronylenesulfonic acid	284 16	258			
5046	$C_{16}H_{12}O_4$	α-Ethylalizarin	268 09	189			
5047	$C_{16}H_{12}O_4$	Pratol	268 09	253			
5048	$C_{16}H_{12}O_5$	Physcion (Physcic acid)	284 09	207			
5049	$C_{16}H_{12}O_6$	Chrysoeriol	300 09	>337			
5050	$C_{16}H_{12}O_5$	Emodine methyl ether	300 09	195			
5051	$C_{16}H_{12}O_5$	Hematein	300 09	250 d.			
5052	$C_{16}H_{12}O_6$	Laccainic acid	332 09		180 d.		
5053	$C_{16}H_{13}N$	Flavoline	219 11	65	375		
5054	$C_{16}H_{13}N$	N-Phenyl-α-naphthylamine	219 11	62	335[258]		
5055	$C_{16}H_{13}N$	N-Phenyl-β-naphthylamine	219 11	108	399 5		
5056	$C_{16}H_{13}NO_7$	Papaveric acid	331 11	233 d.			
5057	$C_{16}H_{13}N_3$	Galegine	233 12	65			
5058	$C_{16}H_{13}N_3$	Hydrazoindole	247 12	140			
5059	$C_{16}H_{14}$	Atronene	206 11		326		
5060	$C_{16}H_{14}$	2, 3-Dimethylanthracene	206 11	246			
5061	$C_{16}H_{14}$	2, 4-Dimethylanthracene	206.11	71			
5062	$C_{16}H_{14}$	2, 6-Dimethylanthracene	206 11	231			
5062 1	$C_{16}H_{14}$	Distyrene $C_6H_5CH:CHCH:CHC_6H_5$.	206 11	124			
5063	$C_{16}H_{14}$	9-Ethylanthracene	206 11	59		1 041[99 2]	1130
5064	$C_{16}H_{14}Cl_2N_2O_2$	3, 3'-Dichlorodiacetylbenzidine	337 04	302			
5065	$C_{16}H_{14}N_2$	α-Flavamine	234 12	97			
5066	$C_{16}H_{14}N_2$	Indolin	234 12		245		
5066 1	$C_{16}H_{14}N_2$	1, 5-Diphenyl-3-methylpyrazole	234 12	63			1199
5067	$C_{16}H_{14}O$	Dypnone	222 11		225[21]		
5067 1	$C_{16}H_{14}O$	Benzylidene-p-tolyl ketone	222 11	77			1289
5068	$C_{16}H_{14}O_2$	Benzyl cinnamate	238 11	34	244[15]		
5069	$C_{16}H_{14}O_3$	Diphenacyl $C_6H_5COCH_2CH_2COC_6H_5$	238.11	145			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5070	C₁₆H₁₄O₃	Guaiacyl cinnamate	254 11	130			
5071	C₁₆H₁₄O₃	Phenylacetic anhydride	254 11	117 5			
5072	C₁₆H₁₄O₃	o-Toluic anhydride (o-CH₃C₆H₄CO)₂O	254 11	39	325		
5073	C₁₆H₁₄O₄	Dibenzyl oxalate (CO₂CH₂C₆H₅)₂	270 11	81	235¹⁴		
5074	C₁₆H₁₄O₄	Diphenyl succinate (CH₂CO₂C₆H₅)₂	270 11	121	330		
5075	C₁₆H₁₄O₅	Brasilin	286 11	250			
5076	C₁₆H₁₄O₅	Sakuranetin	286 11	150			
5077	C₁₆H₁₄O₆	Diphenyl tartrate (CHOHCO₂C₆H₅)₂	302 11	102			
5078	C₁₆H₁₄O₆	Hematoxylin	302 11	140			1333
5079	C₁₆H₁₄O₆	Hesperetin	302 11	226			
5080	C₁₆H₁₄O₆	Homoeriodyctiol	302 11	223			
5081	C₁₆H₁₆N₂O₂	Anisaldazine	254 12	169	180	1 031¹⁸⁵	
5082	C₁₆H₁₆N₂O₂	Diacetylbenzidine (p-CH₃CONHC₆H₄)₂	268 14	331			
5082 1	C₁₆H₁₆N₂O₆	o-Aminophenyl tartrate	332 14	211 d.			
5082 2	C₁₆H₁₆N₂O₆	m-Aminophenyl tartrate	332 14	175 d.			
5082 3	C₁₆H₁₆N₂O₆	p-Aminophenyl tartrate	332 14	220 d.			
5082 4	C₁₆H₁₆N₂O₂	Diacetylhydrazobenzene	268 15	105			1293
5083	C₁₆H₁₆N₂S	Dehydrothioxylidine	268 20		197		
5084	C₁₆H₁₆N₄O₁₀	Damascenine picrate	424 16	159			
5085	C₁₆H₁₆O₂	p-Dimethylbenzoin	240 12	89			
5086	C₁₆H₁₆O₅	Anisilic acid	288 12	164			
5087	C₁₆H₁₆O₃	Ethyl benzilate	256 12	34	201²¹		
5088	C₁₆H₁₇NO₃	Amygdophenine	271 14	141			
5089	C₁₆H₁₇NO₄	Lycorine	287 14	235 d.			
5090	C₁₆H₁₇NO₄	Phenetidine salicylacetate	287 14	182			
5091	C₁₆H₁₈ClNO₄	Lycorine hydrochloride	323 61	208			
5092	C₁₆H₁₈N₂	Azo-o-ethylbenzene	238 16	46 5			
5093	C₁₆H₁₈N₂	Azo-p-ethylbenzene	238 16	63	>340		
5094	C₁₆H₁₈N₂	3, 3'-Azo-o-xylene	238 16	111			
5095	C₁₆H₁₈N₂	4, 4'-Azo-o-xylene	238 16	141			
5096	C₁₆H₁₈N₂	4, 4'-Azo-m-xylene	238 16	129			
5097	C₁₆H₁₈N₂	4, 5'-Azo-m-xylene	238 16	47			
5098	C₁₆H₁₈N₂	5, 5'-Azo-m-xylene	238 16	137			
5099	C₁₆H₁₈N₂	2, 2'-Azo-p-xylene	238 16	119			
5100	C₁₆H₁₈N₂	Diphenylpiperazine	238 16	163 5	212²⁰		
5101	C₁₆H₁₈N₂O	Paricine	254 16	130			
5102	C₁₆H₁₈N₂O₂	o-Azophenetol (C₂H₅OC₆H₄N:)₂	270 16	131	240		
5103	C₁₆H₁₈N₂O₂	p-Azophenetol (C₂H₅OC₆H₄N:)₂	270 16	160 2			
5104	C₁₆H₁₈N₂O₃	3, 3'-Azoxy-4-methoxytoluene	286 16	149			
5105	C₁₆H₁₈N₂O₃	p-Azoxyphenetol	286 16	136 9			
5106	(C₁₆H₁₈N₂O₃)ₓ	Bilirubin	[286 16]ₓ	192 5			
5107	C₁₆H₁₈N₂O₃	Carpiline	286 16	185			
5108	C₁₆H₁₈N₂O₃	Hematoporphyrin	286 16	<100 d.			
5109	C₁₆H₁₈N₂O₃	Pilosine	286 16	187			
5110	C₁₆H₁₈O	Thymyl phenyl ether	226 14		296 8	1 011	
5111	C₁₆H₁₈O₂S	Di-m-xylylsulfone	274 20	121			
5112	C₁₆H₁₈O₇	Barbaloin	322 14	148			
5113	C₁₆H₁₉NO₄	Benzoylecgonine	289 15	195			
5114	C₁₆H₂₀N₂	3-Hydrazo-o-xylene	240 17	141			
5115	C₁₆H₂₀N₂	4-Hydrazo-o-xylene	240 17	107			
5116	C₁₆H₂₀N₂	4-Hydrazo-m-xylene	240 17	122			
5117	C₁₆H₂₀N₂	5-Hydrazo-m-xylene	240 17	125			
5118	C₁₆H₂₀N₂	2-Hydrazo-p-xylene	240 17	145			
5119	C₁₆H₂₀N₂O₂	o-Hydrazophenetol (o-C₂H₅OC₆H₄NH)₂	272 17	89			
5123	C₁₆H₂₀N₄	m-Tetramethyldiaminoazobenzene	268 19	118			
5124	C₁₆H₂₀O₄	Phenyl acid camphorate	276 15	100			
5125	C₁₆H₂₀O₉	Gentiopicrin	356 15	191			
5126	C₁₆H₂₁N₃	p-(Tetramethyldiamino)diphenylamine	255 19	119			
5127	C₁₆H₂₁NO₂	Camphoranilic acid	275 17	204			1333
5128	C₁₆H₂₁NO₃	Homoatropine	275 17	97 5			
5129	C₁₆H₂₁NO₃	Noratropine	275 17	114			
5130	C₁₆H₂₁NO₃	Norhyoscyamine	275 17	140 5			1333
5131	C₁₆H₂₂BrNO₃	Homoatropine hydrobromide	356 09	212 d.			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5132	$C_{16}H_{22}ClNO_3$	Homoatropine hydrochloride	311.64	217			1333
5133	$C_{16}H_{22}N_4$	m-Hydrazodimethylaniline	270 20	100			
5134	$C_{16}H_{22}N_4O_6S$	Caffeine sulfate	486 30				1333
5135	$C_{16}H_{22}O_4$	Di-n-butyl phthalate	278 17		340		
5135 1	$C_{16}H_{22}O_4$	Methyl santoate	278 17		86	1.167	1321
5136	$C_{16}H_{22}O_6$	Bilme acid	310 17	190			
5137	$C_{16}H_{22}O_8$	Coniferin	342 17	185			
5138	$C_{16}H_{22}O_{11}$	d-Glucose pentacetate	390 17	113			
5139	$C_{16}H_{23}NO_4$	Bakankosin	357 19	157			
5140	$C_{16}H_{24}O_2$	Methyl santalate	248 19		164[10]	1 002	
5141	$C_{16}H_{24}$	Pentaethylbenzene	218 20	< -20	277	0 896	655
5142	$C_{16}H_{26}O$	Patchouli alcohol	234 20	56	271 d.	0 994$_4^{70}$	
5142 1	$C_{16}H_{26}O$	Guaiol	234 20	91			1176
5143	$C_{16}H_{26}O_2$	Menthyl l-sorbinate	250 20		173[14]		
5143 1	$C_{16}H_{26}O_8$	Disobutyl d-diacetyl tartrate	346 20		157[3.5]	1 0864[17]	
5144	$C_{16}H_{27}ClN_2O_2$	Alypin hydrochloride	314 68	169			
5145	$C_{16}H_{27}N_3O_5$	Alypin nitrate.	341 23	152			
5146	$C_{16}H_{28}N_2$	Genisteine	248 23	60 5	178[22]		
5147	$C_{16}H_{28}O_2$	Hydrocarpic acid	252 22	60			
5148	$C_{16}H_{28}O_2$	Palmitolic acid	252 22	47	240[15]		
5149	$C_{16}H_{28}O_4$	Palmitoxylic acid	284 22	67			
5150	$C_{16}H_{30}O_2$	Gaidic acid	254 23	39			
5151	$C_{16}H_{30}O_2$	Hypogaeic acid	254 23	33	236[15]		
5152	$C_{16}H_{30}O_2$	l-Menthyl n-capronate.	254 23		153[15]	0 903	
5153	$C_{16}H_{30}O_3$	n-Caprylic anhydride $(C_7H_{15}CO)_2O$.	270 23	-1	285		
5154	$C_{16}H_{30}O_3$	7-Ketopalmitic acid	270 23	74			
5155	$C_{16}H_{31}N$	Palmitonitrile $CH_3(CH_2)_{13}CH_2CN$	237 25	29	251.5[100]	0 822^{11}	
5156	$C_{16}H_{32}$	α-Hexadecylene $CH_2:CH(CH_2)_{13}CH_3$	224 25	4	274	0.789	388
5157	$C_{16}H_{32}N_2O_6S$	Pelletierine sulfate.	380 33	133			
5158	$C_{16}H_{32}O$	Palmitic aldehyde $C_{15}H_{31}CHO$	240 25	58 5	202[29]		
5159	$C_{16}H_{32}O_2$	Palmitic acid $C_{15}H_{31}CO_2H$	256 25	64	215[15]	0 853$_4^{62}$	1113
5160	$C_{16}H_{32}O_2$	Ethyl myristate $C_{13}H_{27}CO_2C_2H_5$	256 25	10 5	295		
5161	$C_{16}H_{32}O_3$	Jalapinolic acid	272 25	68			
5162	$C_{16}H_{32}O_3$	Juniperic acid	272 25	95			
5163	$C_{16}H_{32}O_3$	Lanopalmic acid	272 25	88			
5164	$C_{16}H_{33}I$	n-Cetyl iodide $C_{15}H_{31}CH_2I$	352 19	22	212 5[15]	1 123	535
5165	$C_{16}H_{33}NO$	Palmitic amide $C_{15}H_{31}CONH_2$	255 26	106	236[12]		
5166	$C_{16}H_{34}$	7, 8-Dimethyltetradecane	226 26		267 5	0.792[14]	
5167	$C_{16}H_{34}$	n-Hexadecane	226 26	20	287 5	0 775	
5168	$C_{16}H_{34}O$	Cetyl alcohol $C_{15}H_{31}CH_2OH$	242 26	49.3	344	0 798$_4^{78.9}$	1108
5169	$C_{16}H_{34}O$	n-Octyl ether $(C_8H_{17})_2O$	242 26		291 8	0.820	
5171	$C_{17}H_{10}O$	Benzanthrone	230 08	170			
5172	$C_{17}H_{11}N$	α-Anthraquinoline	229 09	170	446		
5173	$C_{17}H_{12}O$	Phenyl α-naphthyl ketone	232 09	75 5	385		
5174	$C_{17}H_{12}O$	Phenyl β-naphthyl ketone	232 09	82			
5175	$C_{17}H_{12}O_2$	Chrysenic acid	248 09	186 5			
5176	$C_{17}H_{12}O_2$	α-Naphthyl benzoate	248 09	56			
5177	$C_{17}H_{12}O_2$	β-Naphthyl benzoate	248 09	110			
5178	$C_{17}H_{12}O_3$	α-Naphthyl salicylate	264 09	83			
5179	$C_{17}H_{12}O_3$	β-Naphthyl salicylate	264 09	95			
5180	$C_{17}H_{12}O_5$	Alpinin	296 09	174			
5181	$C_{17}H_{12}O_5$	Pratensol	296 09	225			
5182	$C_{17}H_{13}NO_2$	6-Methyl-2-phenylquinoline-4-carboxylic acid	263 11	228			
5183	$C_{17}H_{14}$	α-Benzylnaphthalene	218 11	59	350	1 165⁰	
5184	$C_{17}H_{14}$	β-Benzylnaphthalene	218 11	35 5	350	1 176⁰	
5185	$C_{17}H_{14}O$	Dibenzylideneacetone	234 11	112			
5186	$C_{17}H_{14}O_2$	Atronic acid	250 11	164			
5187	$C_{17}H_{14}O_2$	Isatronic acid..	250 11	157			
5188	$C_{17}H_{14}O_4$	Nepalin	282.11	136			
5189	$C_{17}H_{15}N_5O_9$	Tryptophane picrate	433.16	196 s. d.			
5190	$C_{17}H_{16}$	1, 2, 4-Trimethylanthracene	220.12	243			
5191	$C_{17}H_{16}$	1, 3, 6-Trimethylanthracene	220.12	222			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5192	C₁₇H₁₆	1, 4, 6-Trimethylanthracene	220 12	227			
5193	C₁₇H₁₆O₂	Eugenol benzoate ..	268 12	70	360		
5194	C₁₇H₁₆O₂	Isoeugenol benzoate	268 12	104			
5195	C₁₇H₁₆O₄	Dibenzyl malonate.	284 12		234.5¹⁴ d.		
5196	C₁₇H₁₇NO₂	Apomorphine......	267 14	170 d.			
5197	C₁₇H₁₈ClNO₂	Apomorphine hydrochloride	303 61	210			1333
5198	C₁₇H₁₈N₂O₂	Antipyrine resorcinate	298 16	115			
5199	C₁₇H₁₈O	Dibenzylacetone CO(CH₂CH₂C₆H₅)₂	238 14		224¹⁴		
5200	C₁₇H₁₈O₂	Eugenol benzyl ether	254 14	30	235 d.		
5201	C₁₇H₁₈O₂	Isoeugenol benzyl ether	254 14	59			
5202	C₁₇H₁₉NO₃	Morphine	285 15	d.	193 ᵛᵃᶜ	1 317	1277
5203	C₁₇H₁₉NO₃	α-Isomorphine........	285 15	247			
5204	C₁₇H₁₉NO₃	Piperine	285 15	129 5			
5205	C₁₇H₂₀BrNO₃	Morphine hydrobromide	366 08				1333
5206	C₁₇H₂₀ClNO₃	Morphine hydrochloride	321 62	250 d			1333
5207	C₁₇H₂₀N₂O	Tetramethyldiaminobenzophenone	268 17	174	>360 s. d		
5208	C₁₇H₂₀N₂O₃	Nicotine salicylate	300 17	117 5			1333
5209	C₁₇H₂₀N₂O₄	l-Arabinose diphenylhydrazone	316 17	218			
5211	C₁₇H₂₀N₂S	3, 3-Tetramethyldiaminothiobenzophe-none	284 24	202			
5212	C₁₇H₂₀N₄O₃	l-Arabinosazone	340 19	166	200 d.		
5213	C₁₇H₂₀N₄O₃	d-Xylosephenylosazone	328 19	164	167 d.		
5213.1	C₁₇H₂₀O₃	Di-(p-dianisyl)dimethylmethane	256 15	60 5		1 150	1294
5214	C₁₇H₂₀O₇	Tutin .	336 15	208			
5215	C₁₇H₂₀O₁₀	Patellaric acid	384 15	100			
5216	C₁₇H₂₁NO₂	Apoatropine	271 17	62			
5217	C₁₇H₂₁NO₃	Dihydromorphine	287 17	157			
5218	C₁₇H₂₁NO₄	Atroscine .	303 17	50			
5219	C₁₇H₂₁NO₄	α-Cocaine. . .	303 17	88			
5220	C₁₇H₂₁NO₄	dl-Cocaine . .	303 17	80			
5221	C₁₇H₂₁NO₄	d(l)-Cocaine .	303.17	98			1326
5222	C₁₇H₂₁NO₄	Hyoscine	303.17	55			1333
5223	C₁₇H₂₁NO₄	dl-Pseudococaine	303 17	81 5		1 103⁹⁹ ᵇ	1139
5224	C₁₇H₂₁NO₄	d-Pseudococaine	303 17	41		1.102⁹⁹·⁶	1142
5225	C₁₇H₂₁N₃	Auramine . .	267 19	136			
5226	C₁₇H₂₂BrNO₄	Hyoscine hydrobromide	384 09	194			1333
5227	C₁₇H₂₂ClNO₂	Apoatropine hydrochloride	307 64	239			1333
5228	C₁₇H₂₂ClNO₄	Cocaine hydrochloride	339 64	187			1257
5229	C₁₇H₂₂ClNO₄	Hyoscine hydrochloride ..	339 64				1333
5230	C₁₇H₂₂N₂	p-(Tetramethyldiamino)-diphenyl-methane.	254 19	91			
5231	C₁₇H₂₂N₂O	p-(Tetramethyldiamino)-diphenyl carbi-nol [p-(CH₃)₂NC₆H₄]₂CHOH	270 19	96			
5232	C₁₇H₂₂O₃	Podocarpic acid..	274 17	188			
5233	C₁₇H₂₂O₆	Guaiacyl acid camphorate	306 17	112			
5234	C₁₇H₂₂O₉	Syringin........ ..	370 17	192			
5235	C₁₇H₂₃NO₃	Atropine......	289 19	115.5			1333
5236	C₁₇H₂₃NO₃	d-Hyoscyamine.	289 19	106			
5237	C₁₇H₂₃NO₃	Pseudoatropine .	289 19	120			
5238	C₁₇H₂₄BrNO₃	Atropine hydrobromide	370 11	162			1333
5239	C₁₇H₂₄BrNO₃	Hyoscyamine hydrobromide	370 11	152			1333
5240	C₁₇H₂₄ClNO₃	Atropine hydrochloride	325 65	165			1333
5241	C₁₇H₂₄ClNO₃	Hyoscyamine hydrochloride	325 65				1333
5242	C₁₇H₂₄N₂O₅S	Sinapine thiocyanate	368 22	176			
5243	C₁₇H₂₄N₂O₆	Atropine nitrate	352 20				1333
5244	C₁₇H₂₄O₂	Menthyl benzoate .	260.19	54 5	288	0 808	
5244.1	C₁₇H₂₄O₄	Ethyl santoate	292 19	89		1.148	1322
5245	C₁₇H₂₄O₁₀	Verbenalin........ .	388.19	181.6			
5246	C₁₇H₂₅NO₃	Euphthalmine........	291 20	113			
5247	C₁₇H₂₆O₄	Scillitin................ . .	325 19	154			
5248	C₁₇H₂₆ClNO₃	Euphthalmine hydrochloride .	327.67	183			
5249	C₁₇H₂₆O	Benzylmenthol...............	246.20	111	183¹⁰		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I No.
5250	$C_{17}H_{28}O$	Phellyl alcohol	248 22	100			
5251	$C_{17}H_{29}NO_2$	Ajaconine..	279 23	163			
5252	$C_{17}H_{26}O_5$	Jalapic acid	378 23	120			
5253	$C_{17}H_{32}O_2$	l-Menthyl heptylate	268 25		165^{14}	0 901	
5254	$C_{17}H_{34}$	8-Heptadecene $C_7H_{15}CH.CHC_8H_{17}$	238 26		$160^{9\ 5}$	0.798^{10}	
5255	$C_{17}H_{34}O$	Margaric aldehyde $C_{16}H_{33}CHO$	254 26	36	204^{26}		
5256	$C_{17}H_{34}O_2$	Daturic acid	270 26	60	227^{100}		
5257	$C_{17}H_{34}O_2$	Margaric acid $C_{16}H_{33}CO_2H$	270 26	59 9	227^{100}	0.853^{60}	
5258	$C_{17}H_{34}O_2$	Methyl palmitate $C_{15}H_{31}CO_2CH_3$	270 26	29 5	196^{15}		1119
5259	$C_{17}H_{35}NO_2$	Sphingosine	285 28	244	250 d.		
5260	$C_{17}H_{36}$	n-Heptadecane $CH_3(CH_2)_{15}CH_3$	240 28	22 5	303	0 778	359
5261	$C_{17}H_{36}O$	Heptadecane-9-ol $C_8H_{17}CH(OH)C_8H_{17}$	256 28	61			
5262	$C_{17}H_{37}N$	Heptadecylamine $C_{17}H_{35}NH_2$	255 29	49	340		
5263	$C_{18}H_{12}$	Benzanthrene	228 09	84			
5264	$C_{18}H_{12}$	Chrysene	228 09	251	448		
5265	$C_{18}H_{12}$	Triphenylene ..	228 09	198 5			
5266	$C_{18}H_{12}$	Truxene	228 09	>360			
5267	$C_{18}H_{12}N_2$	2, 3'-Diquinolyl	256 11	176			
5268	$C_{18}H_{12}N_2$	2, 7'-Diquinolyl	256 11	193			
5269	$C_{18}H_{12}N_2$	6, 6'-Diquinolyl	256 11	178			
5270	$C_{18}H_{12}N_2$	8, 8'-Diquinolyl	256 11	205			
5271	$C_{18}H_{12}O_3$	o-(α-Naphthoyl) benzoic acid	276 09	173 5			
5272	$C_{18}H_{12}O_6$	Calycin	308 09	240			
5273	$C_{18}H_{13}N$	Aminochrysene	243 11	203			
5274	$C_{18}H_{14}$	p-Diphenylbenzene $C_6H_4(C_6H_5)_2$	230 11	205	427		
5275	$C_{18}H_{14}O_3$	Cinnamic anhydride $(C_6H_5CH:CHCO)_2O$	278 11	135			
5276	$C_{18}H_{14}O_4$	Epicarin	294 11	195			
5277	$C_{18}H_{14}O_7$	Xanthoeridol ...	342 11	258			
5278	$C_{18}H_{14}O_8$	Diaspirin (Succinyldisalicylic acid)	358 11	178			
5279	$C_{18}H_{15}As$	Triphenylarsine $(C_6H_5)_3As$	306 08	60			
5280	$C_{18}H_{15}Bi$	Triphenyl bismuthine $(C_6H_5)_3Bi$	410 16	78		1 585^{20}	
5281	$C_{18}H_{15}N$	Triphenylamine $(C_6H_5)_3N$. .	245 12	126 5	365	0 774$^{0}_{0}$	
5282	$C_{18}H_{15}O_3P$	Triphenyl phosphite $(C_6H_5O)_3P$....	310 14		220^{11}	1 184^{18}	
5283	$C_{18}H_{15}O_4P$	Triphenyl phosphate $(C_6H_5O)_3PO$...	326 14	49 9	245^{11}		
5284	$C_{18}H_{15}P$	Triphenylphosphine $(C_6H_5)_3P$.	262 14	79	>360	1 194	
5285	$C_{18}H_{15}Sb$	Triphenylstibine $(C_6H_5)_3Sb$	352 89	48	>360	1 500^{11}	
5286	$C_{18}H_{15}NO_2$	Aporheine	278 13	89	290 d.		
5287	$C_{18}H_{16}N_2$	Diphenyl-m-phenylenediamine	260 14	95			
5288	$C_{18}H_{16}N_2$	Triphenylhydrazine $(C_6H_5)_2NNHC_6H_5$..	260 14	142		0 869$^{70}_{4}$	
5289	$C_{18}H_{16}N_2O_3$	Analgen	292 14	210			
5290	$C_{18}H_{16}N_2O_3$	5, 5'-Dibenzylbarbituric acid	308 14	222			
5291	$C_{18}H_{16}N_2O_6S$	Chinosol	388 20	177 5			
5292	$C_{18}H_{16}O_2$	Cinnamyl cinnamate	264 12	44		1 085$^{16\ 5}$	
5293	$C_{18}H_{16}O_4$	α-Isatropic acid	296 12	237			
5294	$C_{18}H_{16}O_4$	β-Isatropic acid	296 12	206			
5295	$C_{18}H_{16}O_4$	α-Truxillic acid	296 12	272			
5296	$C_{18}H_{16}O_4$	Isotruxillic acid	296.12	206			
5297	$C_{18}H_{16}O_4$	γ-Truxillic acid	296 12	228			
5298	$C_{18}H_{16}O_4$	δ-Truxillic acid	296.12	174			
5299	$C_{18}H_{16}O_4$	ϵ-Truxillic acid	296 12	192			
5300	$C_{18}H_{16}O_4$	η-Truxillic acid	296 12	260			
5301	$C_{18}H_{16}O_4$	Dibenzyl fumarate	296 12	59 5	211^{5}		
5302	$C_{18}H_{16}O_4$	Nepodin	296 12	158			
5303	$C_{18}H_{16}O_7$	dl-Usnic acid	344 12	193			
5304	$C_{18}H_{16}O_7$	d(l)-Usnic acid	344.12	203			1295
5305	$C_{18}H_{16}O_{14}$	Igasuric acid (Chlorogenic acid) ..	456 12	207			
5306	$C_{18}H_{18}$	Retene	234 14	98 5	394	1 13^{14}	
5307	$C_{18}H_{18}$	1, 3, 5, 7-Tetramethylanthracene	234 14	280 d.			
5308	$C_{18}H_{18}N_2O_4$	Antipyrine salicylate ..	326 16	92			
5308 1	$C_{18}H_{18}N_3$	Vesuvin	346 20	143 5			
5310	$C_{18}H_{18}O_4$	Dibenzyl succinate	298 14	45	238^{14}		
5312	$C_{18}H_{19}NO_3$	Berbamine	297 15	200			
5313	$C_{18}H_{19}N_3O_2$	Dimazon (Diacetylaminoazotoluene)..	309 17	75			
5314	$C_{18}H_{20}BrNO_3$	Apomorphine methobromide	362.08	180			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5315	$C_{18}H_{20}N_2O_2$	Cinchotenine	312.17	198			
5316	$C_{18}H_{21}NO_3$	Bebeerine	299 17	214			
5317	$C_{18}H_{21}NO_3$	Codeine	299 17	155	179	1 315¹⁴	1283, 1286
5318	$C_{18}H_{21}NO_3$	Isobebeerine	299 17	297			
5319	$C_{18}H_{21}NO_3$	Isocodeine	299 17	144	d		1288
5320	$C_{18}H_{21}NO_3$	Pseudocodeine	299 17	181		1 290¹⁸⁰	1264
5321	$C_{18}H_{22}BrNO_3$	Codeine hydrobromide	380 09				1333
5322	$C_{18}H_{22}BrNO_3$	Morphine methylbromide	380 09	265 d			
5323	$C_{18}H_{22}ClNO_3$	Bebeerine hydrochloride	335 64	260			
5324	$C_{18}H_{22}ClNO_3$	Codeine hydrochloride	335 64	264			1333
5325	$C_{18}H_{22}N_2O_2$	Holocaine	298 19	117			
5325 1	$C_{16}H_{22}N_2O_5$	Pilocarpine salicylate	346 19	120			1333
5326	$C_{18}H_{22}N_4O_4$	Galactosazone	358 20	201	202 d.		
5327	$C_{18}H_{22}N_4O_4$	d-Glucosazone	358 20	208 d			
5328	$C_{18}H_{22}N_4O_4$	l-Glucosazone	358 20	205 d			
5329	$C_{18}H_{22}N_4O_4$	Gulososazone	358 20	168	180 d.		
5330	$C_{18}H_{22}O_{10}$	Murrayin	398 17	170			
5331	$C_{18}H_{23}ClN_3O_2$	Holocaine hydrochloride	334 65	189			
5332	$C_{18}H_{23}NO_6$	Cocaine formate	349 19	42			
5333	$C_{18}H_{24}NO_7P$	Codeine phosphate	397 22	235			1333
5334	$C_{18}H_{26}O_2$	Menthyl phenylacetate	274 20		205 5²⁵	1 002	
5335	$C_{18}H_{26}O_4$	Diamyl phthalate ..	306 20		344		
5336	$C_{18}H_{27}NO_3$	Capsaicin	305 22	65			1226
5337	$C_{18}H_{27}NO_6$	Senecifoline	385 22	194			
5338	$C_{18}H_{28}ClNO_6$	Senecifoline hydrochloride	421 68	260			
5339	$C_{18}H_{28}O_4$	Embellic acid .	308 22	142			
5340	$C_{18}H_{30}$	Hexaethylbenzene $C_6(C_2H_5)_6$	246 23	129	298	0 831⁴¹⁰·⁴	1159
5341	$C_{18}H_{30}O$	Sycoceryl alcohol	262 23	90			
5342	$C_{18}H_{30}O_2$	Linolenic acid	278 23		232¹⁷	0 914	
5343	$C_{18}H_{31}ClN_2O_6$	dl-Ecgonine hydrochloride	406 71	247			
5343.1	$C_{18}H_{32}$	Fichtelite.	248 25	46		1 010	1247
5344	$C_{18}H_{32}O_2$	Chaulmoogric acid	280 25	69	248²⁰		
5345	$C_{18}H_{32}O_2$	α-Eleostearic acid	280 25	49	235¹⁷		
5346	$C_{18}H_{32}O_2$	Linoleic acid	280 25	< −18	230¹⁶	0 903	
5347	$C_{18}H_{32}O_2$	Stearolic acid $C_6H_{17}C:C(CH_2)_7CO_2H$	280 25	48	260		
5348	$C_{18}H_{32}O_2$	Tariric acid.	280 25	50 5			
5349	$C_{18}H_{32}O_4$	Stearoxylic acid	312 25	86			
5350	$C_{18}H_{32}O_{16}$	Raffinose . . .	504 25	119	130 d.	1 465	
5351	$C_{18}H_{32}O_{16}$	Procellose	504 25	210			
5352	$C_{18}H_{33}N_2O_{12}$	Piperazine quinate (Sidonal)	469 27	171			
5353	$C_{18}H_{34}$	Hexadecylacetylene $C_{16}H_{33}C:CH$	250 26	26	180¹⁵	0 798²⁶	
5354	$C_{18}H_{34}$	1-Methyl-2-pentadecylacetylene	250 26	30	184¹⁵	0 802	
5355	$C_{18}H_{34}O$	Chaulmoogryl alcohol	266 26	36			
5356	$C_{18}H_{34}O$	Oleic aldehyde	266 26		169⁴	0.851¹⁵	456
5357	$C_{18}H_{34}O_2$	Elaidic acid	282 26	51 5	288¹⁰⁰	0 851⁷⁹·⁴	
5358	$C_{18}H_{34}O_2$	Gynocardic acid	282 26	67 5			
5359	$C_{18}H_{34}O_2$	Oleic acid $C_8H_{17}CH:CH(CH_2)_7CO_2H$	282 26	14	286¹⁰⁰	0 895⁴₁⁷·⁷	929
5360	$C_{18}H_{34}O_2$	Petroselinic acid	282 26	34		0 868⁴⁰	1057
5361	$C_{18}H_{34}O_2$	Rapic acid	282 26	14		0 897¹⁵	
5362	$C_{18}H_{34}O_2$	l-Menthyl n-caprylate	282.26		175¹⁵	0 898	
5363	$C_{18}H_{34}O_3$	3-Ketostearic acid.	298.26	97			
5364	$C_{18}H_{34}O_3$	6-Ketostearic acid	298 26	75			
5365	$C_{18}H_{34}O_3$	8-Ketostearic acid	298.26	83			
5366	$C_{18}H_{34}O_3$	9-Ketostearic acid	298 26	76			
5367	$C_{18}H_{34}O_3$	10-Ketostearic acid	298 26	65			
5368	$C_{18}H_{34}O_3$	Ricinelaidic acid	298 26	53	250¹⁵		
5369	$C_{18}H_{34}O_3$	Ricinic acid	298 26	81	252¹⁵		
5370	$C_{18}H_{34}O_3$	Ricinoleic acid	298 26	17	250¹⁵	0 945¹⁵	
5371	$C_{18}H_{34}O_5$	Oleic acid ozonide	330 26			1 022	472
5371.1	$C_{18}H_{34}O_6$	Di-n-heptyl tartrate .	346 26	35	235¹⁴	0.999⁶¹	
5372	$C_{18}H_{34}O_{16}$	Clavisepsin	506 26	198			
5373	$C_{18}H_{35}ClO$	Stearyl chloride $C_{17}H_{35}COCl$.	302 73	23	215¹⁵		
5374	$C_{18}H_{35}N$	Stearonitrile $C_{17}H_{35}CN$	265 28	41	214¹²		

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5375	$C_{18}H_{35}NO$	Oleicamide	281 28	76			
5376	$C_{18}H_{35}NO_2$	Oleohydroxamic acid	297 28	61			
5377	$C_{18}H_{36}$	n-Octodecylene	252 28	18	179^{15}	0.791	
5378	$C_{18}H_{36}O$	Stearic aldehyde $C_{17}H_{35}CHO$	268 28	63.5	261^{100}		
5379	$C_{18}H_{36}O_2$	Stearic acid $C_{17}H_{35}CO_2H$	284 28	69.3	383	$0 847^{69 3}$	1117
5380	$C_{18}H_{36}O_2$	Cetyl acetate $CH_3CO_2C_{16}H_{33}$.	284 28	18 5	$200 5^{15}$	0 858	1041
5381	$C_{18}H_{36}O_2$	Ethyl palmitate $C_{15}H_{31}CO_2C_2H_5$	284 28	24 2	$185 5^{10}$		1043
5382	$C_{18}H_{36}O_2$	Methyl margarate	284 28	29			
5383	$C_{18}H_{36}O_3$	1-Hydroxystearic acid	300 28	85			
5384	$C_{18}H_{36}O_3$	dl-2-Hydroxystearic acid	300 28	85			
5385	$C_{18}H_{36}O_3$	9-Hydroxystearic acid	300 28	81.5			
5386	$C_{18}H_{36}O_3$	10-Hydroxystearic acid	300 28	79			
5387	$C_{18}H_{36}O_3$	11-Hydroxystearic acid	300 28	78			
5388	$C_{18}H_{36}O_4$	4, 9-Dihydroxystearic acid	316 28	136.5			
5389	$C_{18}H_{37}I$	n-Octodecyl iodide ...	380 22	34	$170^{0.5}$		
5390	$C_{18}H_{37}NO$	Stearic amide $C_{17}H_{35}CONH_2$	283 29	109	251^{12}		
5391	$C_{18}H_{38}$	n-Octadecane $CH_3(CH_2)_{16}CH_3$	254 29	28	317	0 777	1047
5392	$C_{18}H_{38}O$	n-Octadecyl alcohol	270 29	58.5	$210 5^{15}$	$0 812^{59}_4$	
5394	$C_{19}H_{12}O$	Benzylideneacenaphthenone	256 09	107			
5395	$C_{19}H_{13}N$	9-Phenylacridine	255 11	181	404		
5396	$C_{19}H_{15}N_3O_6$	Tri-p-nitrophenylmethane	379 12	207			
5397	$C_{19}H_{14}O_3$	Aurine	290 11	>220			
5398	$C_{19}H_{14}O_5$	Oroxylin .	338 11	225			
5399	$C_{19}H_{15}$	Triphenylmethyl $(C_6H_5)_3C$	243 12	147			
5400	$C_{19}H_{15}Cl$	Triphenylchloromethane $(C_6H_5)_3CCl$	278 57	112	310		
5401	$C_{19}H_{15}N_3$	Chrysaniline	285 14	270			
5402	$C_{19}H_{16}$	Triphenylmethane $(C_6H_5)_3CH$	244 12	92 5	359.2	$1 014^{99}_4$	1128
5403	$C_{19}H_{16}N_2$	Benzophenone phenylhydrazone....	272 14	137			
5404	$C_{19}H_{16}O$	Triphenyl carbinol $(C_6H_5)_3COH...$	260 12	162 5	>360	1 188	
5405	$C_{19}H_{16}O_3$	Triphenyl orthoformate $HC(OC_6H_5)_3$.	292 12	77	277^{55}		
5406	$C_{19}H_{17}N$	m-Aminotriphenylmethane .	259.14	120			
5407	$C_{19}H_{17}N$	p-Aminotriphenylmethane	259 14	84			
5408	$C_{19}H_{17}N$	Diphenylbenzylamine	259 14	87			
5409	$C_{19}H_{17}N$	Triphenylmethylamine $(C_6H_5)_3C.NH_2$	259 14	105			
5410	$C_{19}H_{17}NO_2$	Novatophan	291 14	76			
5411	$C_{19}H_{17}NO_4$	Cusparidine	307 14	79			
5412	$C_{19}H_{17}NO_4$	Cusparine	307 14	92			
5413	$C_{19}H_{17}NO_5$	Isocusparine ...	307 14	194			
5414	$C_{19}H_{17}N_3$	α-Triphenylguanidine	287 16	145	d.		
5415	$C_{19}H_{17}N_3$	β-Triphenylguanidine	287 16	131			
5416	$C_{19}H_{18}ClN_3$	α-Triphenylguanidine hydrochloride	323 62	241		$0 875^{70}_4$	
5417	$C_{19}H_{18}N_2$	p, p'-Diaminotriphenylmethane	274.16	140			
5418	$C_{19}H_{18}O_3$	Eugenol cinnamate	294 14	90			
5419	$C_{19}H_{18}O_7$	Eriodonol	358 14	199			
5420	$C_{19}H_{18}O_8$	Atranoric acid	374 14	197			
5421	$C_{19}H_{18}O_{11}$	Euxanthic acid	422 14	162	d.		
5422	$C_{19}H_{19}NO_2$	Ditamine	293 15	75			
5423	$C_{19}H_{19}NO_3$	Galipidine	309 15	111			
5424	$C_{19}H_{19}NO_4$	Bulbocapnine	325 15	199			1332
5425	$C_{19}H_{19}NO_5$	Stylopine	341 15	202			
5426	$C_{19}H_{19}N_3$	o-Leucaniline $(NH_2C_6H_4)_3CH$	289 17	165			
5427	$C_{19}H_{19}N_3$	p-Leucaniline $(NH_2C_6H_4)_3CH$	289 17	148			
5428	$C_{19}H_{19}N_3O$	Pararosaniline $(NH_2C_6H_4)_3C(OH)$.	305 17	189			
5428.1	$C_{19}H_{20}N_2O$	Cinchoninone . . .	292.17	127		1 226	1301
5429	$C_{19}H_{20}N_2O_4$	Antipyrine mandelate	340 17	53			
5430	$C_{19}H_{20}N_2O_4$	dl-Ornithuric acid........	340 17	183			
5431	$C_{19}H_{20}O_4$	Diethyl diphenylmalonate	312 15	59			
5432	$C_{19}H_{20}O_5$	Guaiaconic acid	328 15	100			
5433	$C_{19}H_{21}NO_3$	Isothebaine	311.17	204			
5434	$C_{19}H_{21}NO_3$	Oxyacanthine	311 17	210			
5435	$C_{19}H_{21}NO_3$	Thebaine	311 17	193		1.305	
5436	$C_{19}H_{21}NO_5$	Eupyrin . .	343.17	88			
5437	$C_{19}H_{22}N_2$	Desoxycinchonidine	278.19	61			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5438	$C_{19}H_{22}N_2$	Desoxycinchonine	278 19	92			
5439	$C_{19}H_{22}N_2O$	Apocinchonine	294 19	228			
5440	$C_{19}H_{22}N_2O$	Cinchonicine	294 19	59			
5441	$C_{19}H_{22}N_2O$	Cinchonidine	294 19	210			1278
5442	$C_{19}H_{22}N_2O$	α-Cinchonine	294 19	264 3			1304
5443	$C_{19}H_{22}N_2O$	Homocinchonidine	294 19	207 6			
5444	$C_{19}H_{22}N_2O$	β-Isocinchonine	294 19	126			
5445	$C_{19}H_{22}N_2O_2$	Apoconquinine	310 19	137			
5446	$C_{19}H_{22}N_2O_2$	Apoquinine	310 19	210 d.			
5447	$C_{19}H_{22}N_2O_2$	Cupreine	310 19	202			
5448	$C_{19}H_{22}N_2O_4$	Chitenine	342 19	286 d.			
5451	$C_{19}H_{23}ClN_2O$	Cinchonidine hydrochloride	330 65	242 d			
5452	$C_{19}H_{23}ClN_2O$	Cinchonine hydrochloride	330 65	218 d.			1333
5453	$C_{19}H_{23}NO_3$	Codethyline	313 19	93			
5454	$C_{19}H_{23}NO_4$	Cinnamylcocaine	329 19	121			
5455	$C_{19}H_{23}NO_4$	Corytuberine	329 19	240			
5456	$C_{19}H_{23}NO_4$	Porphyroxine	329 19	135			
5457	$C_{19}H_{23}NO_4$	Sinomenine	329 19	161			
5458	$C_{19}H_{23}NO_5$	Morphine acetate	345 19	200 d			
5459	$C_{19}H_{23}N_3O_4$	Cinchonine nitrate	357 20				1333
5460	$C_{19}H_{24}BrNO_3$	Eucodine (Methylcodeine bromide)	394 11	261			
5461	$C_{19}H_{24}ClNO_3$ $(2H_2O)$	Dionine	349 65	123	170 d		
5462	$C_{19}H_{24}N_2O$	Cinchamidine (Hydrocinchonidine)	296 20	230			
5463	$C_{19}H_{24}N_2O$	Cinchonamine	296 20	185			
5464	$C_{19}H_{24}N_2O$	Cinchotine	296 20	286			
5465	$C_{19}H_{24}N_2O$	Pereirine	296 20	124			
5466	$C_{19}H_{24}N_2O_2$	Conquinamine	312 20	123			
5467	$C_{19}H_{24}N_2O_2$	Geissospermine	312 20	189			
5468	$C_{19}H_{24}N_2O_2$	Hydrocupreine	312 20	230			
5469	$C_{19}H_{24}N_2O_2$	Quinamine	312 20	172			
5473	$C_{19}H_{25}N_4O_4$	Ionidine	373 23	156			
5474	$C_{19}H_{26}N_2O$	Aspidosine	298 22	245			
5475	$C_{19}H_{27}NO_4$	α-Eucaine	333 22	103			
5476	$C_{19}H_{28}ClNO_4$	α-Eucaine hydrochloride	369 68	200			
5477	$C_{19}H_{28}O_2$	Abietic acid	288 22	161			1251
5478	$C_{19}H_{28}O_4$	Convallaretin	320 22	>255			
5479	$C_{19}H_{28}O_{13}$	Calmatambin	464 22	144			
5480	$C_{19}H_{30}O_2$	Benzyl laurate $C_{11}H_{23}CO_2CH_2C_6H_5$	290 23	8 5	211^{12}	$0\ 946^{25}_{15}$	540
5481	$C_{19}H_{34}O_2$	Methyl chaulmoograte	294 26	22	227^{70}	$0\ 912^{25}_{25}$	
5482	$C_{19}H_{36}O_3$	Methyl ricinolate	312 28		245^{10}	0 924	465
5483	$C_{19}H_{38}O_2$	Nondecylic acid $CH_3(CH_2)_{17}CO_2H$	298 29	66	299^{100}		
5484	$C_{19}H_{38}O_2$	Ethyl margarate $CH_3(CH_2)_{15}CO_2C_2H_5$	298 29	27			
5485	$C_{19}H_{38}O_2$	Methyl stearate $C_{17}H_{35}CO_2CH_3$	298 29	38	215^{14}		
5486	$C_{19}H_{40}$	n-Nondecane $CH_3(CH_2)_{17}CH_3$	268 31	32	330	$0\ 777^{17}_{4}$	1045
5487	$C_{20}H_{10}I_4O_4$	Nosophen (Tetraiodophenolphthalein)	821 81	225			
5488	$C_{20}H_{12}$	Perylene	252 09	264			
5489	$C_{20}H_{12}O_2$	Fluoran	300 09	175			
5490	$C_{20}H_{12}O_5$	Fluorescein	332 09		290 d.		
5491	$C_{20}H_{14}$	α, α'-Dinaphthyl $C_{10}H_7.C_{10}H_7$	254 11	160.5	360		
5492	$C_{20}H_{14}$	α, β'-Dinaphthyl	254 11	80			
5493	$C_{20}H_{14}$	β, β'-Dinaphthyl $C_{10}H_7.C_{10}H_7$	254 11	187.8	452		
5494	$C_{20}H_{14}$	9-Phenylanthracene	254 11	153	417		
5495	$C_{20}H_{14}N_2$	α, α'-Azonaphthalene	282.12	190			
5496	$C_{20}H_{14}N_2$	β, β'-Azonaphthalene	282 12	204			
5497	$C_{20}H_{14}N_2O$	α, α'-Azoxynaphthalene	298.12	127			
5498	$C_{20}H_{14}N_2O$	β, β'-Azoxynaphthalene	298.12	167			
5499	$C_{20}H_{14}O$	α-Naphthyl ether $(C_{10}H_7)_2O$	270.11	110	>360		
5500	$C_{20}H_{14}O$	β-Naphthyl ether $(C_{10}H_7)_2O$	270 11	105	250^{12}		
5501	$C_{20}H_{14}O$	α, β'-Naphthyl ether	270 11	81	264^{15}		
5502	$C_{20}H_{14}O_2$	α-Dinaphthol	286 11	300			
5503	$C_{20}H_{14}O_2$	β-Dinaphthol	286.11	218			
5504	$C_{20}H_{14}O_4$	Phenolphthalein	318.11	261		1.277^{17}_{4}	

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5505	$C_{20}H_{14}O_5$	Fluorescin	334 11	127			
5506	$C_{20}H_{14}O_9$	Psoromic acid	398 11	264			
5507	$C_{20}H_{14}S$	α, α'-Dinaphthyl sulfide $(C_{10}H_7)_2S$..	286 17	110	290^{15}		
5508	$C_{20}H_{15}N$	β, β'-Dinaphthylamine $(C_{10}H_7)_2NH$.	269.12	172 2	471		
5509	$C_{20}H_{13}NO_4$	Sanguinarine	333.12	213			
5510	$C_{20}H_{15}NO_3$	Berulic acid	397.12	200			
5511	$C_{20}H_{15}N_3$	p-Amino-α-azonaphthalene	297.14	175			
5512	$C_{20}H_{15}N_3$	Amino-β-azonaphthalene	297 14	156			
5513	$C_{20}H_{16}N_2$	α, α'-Hydrazonaphthalene	284 14	α 271; β 274			
5514	$C_{20}H_{16}N_2$	β, β'-Hydrazonaphthalene	284 14	164			
5515	$C_{20}H_{16}N_2O$	Benzilphenylhydrazone	300.14	134			
5516	$C_{20}H_{16}N_4$	Nitron	312 16	189 d.			
5517	$C_{20}H_{16}O_2$	Triphenylacetic acid $(C_6H_5)_3C.CO_2H$	288 12	265			
5518	$C_{20}H_{16}O_3$	Rosolic acid	304 12	270	d.		
5519	$C_{20}H_{17}N_3O_2$	Rubazonic acid	359 17	181			
5520	$C_{20}H_{18}$	Diphenyl-m-tolylmethane	258 14	61 5	356	1 07^{15}	
5521	$C_{20}H_{18}$	1, 1, 2-Triphenylethane	258 14	54	349 4		
5522	$C_{20}H_{18}ClNO_4$	Berberine hydrochloride	371 61			1 397	1333
5523	$C_{20}H_{18}N_2O$	α-Benzoinphenylhydrazone	302 16	155			
5524	$C_{20}H_{18}N_2O$	β-Benzoinphenylhydrazone	302 16	106			
5525	$C_{20}H_{18}N_4S$	Triphenylguanylthiourea	346 24	157			
5526	$C_{20}H_{19}N$	Dibenzylaniline $C_6H_5N(CH_2C_6H_5)_2$.	273 15	70			
5527	$C_{20}H_{19}NO_5$	Chelidonine	353 15	136			
5528	$C_{20}H_{19}NO_5$	Papaveraldine	353 15	210			
5529	$C_{20}H_{19}NO_5$	Protopine	353 15	207			
5530	$C_{20}H_{19}NO_9$	Berberilic acid	417 15	182			
5532	$C_{20}H_{20}N_2O_4$	Antipyrine acetylsalicylate	368 17	65			
5533	$C_{20}H_{20}O_5$	Cubebinol	340 15	92			
5534	$C_{20}H_{20}O_6$	Cubebin	356 15	132			
5535	$C_{20}H_{20}O_7$	Coccelic acid	372 15	178			
5536	$C_{20}H_{20}O_{10}$	Scoparin	420 15	219 d.			
5537	$C_{20}H_{20}O_{13}$	Luteic acid	452.15	274			
5538	$C_{20}H_{21}NO_3$	Galipeine	323.17	115			
5539	$C_{20}H_{21}NO_4$	l-Canadine	339.17	134			
5540	$C_{20}H_{21}NO_4$	Dicentrine	339 17	169			
5541	$C_{20}H_{21}NO_4$	Papaverine	339 17	147	d.	1 337	1331
5542	$C_{20}H_{21}NO_4$	dl-Canadine	339 17	167			
5544	$C_{20}H_{22}ClNO_4$	Papaverine hydrochloride	375 64	221 d.			
5545	$C_{20}H_{22}N_2O$	Quinene	306.19	82			
5546	$C_{20}H_{22}N_2O_2$	Dehydroquinine	322 19	181			
5547	$C_{20}H_{22}N_2O_2$	Jelsemine ...	322.19	178			
5548	$C_{20}H_{22}N_2O_4$	Lysuric acid	354 19	145			
5549	$C_{20}H_{22}O_8$	Populin	390 17	180			
5550	$C_{20}H_{23}ClN_2O_2$	Jelsemine hydrochloride	358 65	300			
5551	$C_{20}H_{23}NO_4$	Acetylcodeine	341.19	133.5			
5552	$C_{20}H_{23}NO_4$	Corypalmine	341.19	236			
5553	$C_{20}H_{23}N_3O_4$	Pyramidon salicylate	369 20	70			
5554	$C_{20}H_{23}O_4$	Naphthyl acid camphorate	327.18	122			
5555	$C_{20}H_{24}Cl_2N_2O_2$	Quinene dichloride	395.12	97			
5556	$C_{20}H_{24}NO_4$	Staphisagroine	342 19	275			
5557	$C_{20}H_{24}N_2O$	Desoxyquinine ..	308 20	52			
5558	$C_{20}H_{24}N_2O_2$	Isoconquinine ..	324.20	142			
5559	$C_{20}H_{24}N_2O_2$	Isoquinine	324.20	185			
5560	$C_{20}H_{24}N_2O_2$	Quinicine ...	324 20	60			
5561	$C_{20}H_{24}N_2O_2$	Quinidine.	324.20	168			1298
5562	$C_{20}H_{24}N_2O_2$	Quinine . ..	324.20	175			1279
5563	$C_{20}H_{24}N_2O_2$	Quinine (isomer A)	324.20	193 5			
5564	$C_{20}H_{24}N_2O_2$	Quinine (isomer B)	324.20	189			
5566	$C_{20}H_{25}BrN_2O_2$	Quinine hydrobromide	405.13	200			
5567	$C_{20}H_{25}ClN_2O_2$	Quinidine hydrochloride	360.67	259 d.			
5568	$C_{20}H_{25}ClN_2O_2$	Quinine hydrochloride	360.67	160	259 d.		
5570	$C_{20}H_{25}NO_3$	Lobelidine	311.20	106			
5571	$C_{20}H_{25}NO_4$	Codamine	343.20	121			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5572	$C_{20}H_{25}NO_4$	Laudanidine	343 20	177			
5573	$C_{20}H_{25}NO_4$	Laudanine	343 20	164 5		1 256	
5575	$C_{20}H_{24}N_2O_6S$	Quinine disulfate	422 28	160 d.			
5577	$C_{20}H_{26}N_2O_2$	Hydroquinidine	326 22	167			
5578	$C_{20}H_{26}N_2O_2$	Hydroquinine	326 22	172 3			
5579	$C_{20}H_{27}NO_4$	Diversine	361 22	93			
5580	$C_{20}H_{27}NO_{11}$	Amygdalin	457 22	200			
5581	$C_{20}H_{27}N_2O_4P$	Quinine hypophosphite	390 25	181			
5583	$C_{20}H_{28}O_4$	Thymyl acid camphorate	332 22	89			
5584	$C_{20}H_{22}O_5$	Eugenol acid camphorate	318 22	116			
5585	$C_{20}H_{28}O_4$	Cholanic acid ...	364 22	285			
5586	$C_{20}H_{28}O_{13}$	Primeverin	476 22	206			
5587	$C_{20}H_{30}NO_6$	Quinine hydrate	378 25	57	d.		
5588	$C_{20}H_{30}O_2$	d-Pimaric acid	302 23	212	282¹⁰		
5589	$C_{20}H_{30}O_4$	Onoceric acid	334 23	120			
5590	$C_{20}H_{30}O_5$	Andrographolide	350 23	218			
5591	$C_{20}H_{32}O_6$	Andrapholic acid	368 25	188			
5592	$C_{20}H_{33}NO$	Myristic anilide	303 26	84			
5593	$C_{20}H_{33}N_3$	Ormosine	315 28	87			
5594	$C_{20}H_{33}N_3$	Ormosinine	315 28	205			
5595	$C_{20}H_{34}O$	Ambrosterol	290 26	147			
5596	$C_{20}H_{34}O$	Cinchol	290 26	139			
5597	$C_{20}H_{34}O$	Cupreol	290 26	140			
5598	$C_{20}H_{34}O$	Quebrachol	290 26	125			
5599	$C_{20}H_{34}O_{10}$	Cyclamin	434 26	236			1333
5600	$C_{20}H_{36}N_6O_{16}$	Vicine	628 34	242 d.			
5601	$C_{20}H_{36}O$	Excretin	292 28	96			
5602	$C_{20}H_{36}O_2$	Eicosinic acid	308 28	69	270¹⁵		
5603	$C_{20}H_{36}O_2$	Ethyl chaulmoograte	308 28		230²⁰	0 906	1036
5604	$C_{20}H_{38}O_2$	Eicosenic acid	310 29	50	267¹⁵		
5605	$C_{20}H_{38}O_3$	Ethyl ricinoleate	326 29		258¹²	0 914	481
5606	$C_{20}H_{40}O$	Phytol	296 31		204¹⁰	0 856	484
5607	$C_{20}H_{40}O_2$	Arachidic acid	312 31	77	328		
5608	$C_{20}H_{40}O_2$	Ethyl stearate $C_{17}H_{35}CO_2C_2H_5$	312 31	33 7	224		
5609	$C_{20}H_{41}I$	n-Eicosyl iodide	408 25	42	192⁰ ˢ		
5610	$C_{20}H_{42}$	n-Eicosane $CH_3(CH_2)_{18}CH_3$	282 32	38	205¹⁵	0 778¹⁶·⁷	1065
5611	$C_{20}H_{42}O$	Eicosyl alcohol $CH_3(CH_2)_{18}CH_2OH$	298 32	71	220⁵		
5612	$C_{21}H_{14}O$	α, β'-Dinaphthyl ketone	282 11	135			
5613	$C_{21}H_{14}O$	β, β'-Dinaphthyl ketone	282 11	a 125 5 b 164 5			
5614	$C_{21}H_{14}O_2$	Picenic acid	298 11	201			
5615	$C_{21}H_{15}Bi_3O_9$	Bismuth salicylate	829 12	135 d.			
5616	$C_{21}H_{16}$	α, α'-Dinaphthylmethane	268 12	109	360		
5617	$C_{21}H_{16}$	α, β'-Dinaphthylmethane $(C_{10}H_7)_2CH_2$	268.12	95			
5618	$C_{21}H_{16}$	β, β'-Dinaphthylmethane $(C_{10}H_7)_2CH_2$	268 12	93			
5619	$C_{21}H_{16}N_2$	Lophine	296 14	275			
5620	$C_{21}H_{16}O_{11}$	Methylenecitrylsalicylic acid	444 12	154			
5621	$C_{21}H_{18}N_2$	Amarin	298 16	129			
5622	$C_{21}H_{18}N_2$	Hydrobenzamide	298.16	101			
5623	$C_{21}H_{18}O_{12}$	Scutellarin	462.14	200 d.			
5624	$C_{21}H_{19}NO_4$	Fumarine	349.15	199			
5625	$C_{21}H_{20}$	Phenylditolylmethane	272 15	56			
5626	$C_{21}H_{20}N_2O_4$	Alstonine (Chlorogenine)	364.17	195			
5627	$C_{21}H_{20}O_6$	Curcumin	368.15	183			1333
5628	$C_{21}H_{20}O_9$	Aloin......	416.15	147 9			
5629	$C_{21}H_{20}O_9$	1, 2-Dihydro-3, 5-dihydroxy-4-(α, 3, 4-trihydroxybenzylbenzofuran)*	416.15	217			
5630	$C_{21}H_{20}O_9$	Frangulin	416.15	226			
5631	$C_{21}H_{20}O_{11}$	Quercitrin	448.15	185			
5632	$C_{21}H_{20}O_{13}$	Incarnatrin	464.15	245			
5633	$C_{21}H_{21}N$	Tribenzylamine $(C_6H_5CH_2)_3N$	287.17	92		0.991¹⁵	
5634	$C_{21}H_{21}NO_6$	d-Corycavamine ...	367 17	149			
5635	$C_{21}H_{21}NO_6$	Hydrastine ...	383.17	132			

* Also commonly known as Catechol, Pyrocatechol, Catechin, Pyrocatechin. See #1414.

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5636	$C_{21}H_{21}NO_6$	Rhoeadine	383.17	232 d.			
5637	$C_{21}H_{21}N_3$	Anhydroformaldehydeaniline	315 19	45.5	185		
5638	$C_{21}H_{21}O_4P$	Tri-p-cresyl phosphate	368 19	77			
5639	$C_{21}H_{21}O_6P$	Triguaiacyl phosphite	400.19	78			
5640	$C_{21}H_{21}O_7P$	Triguaiacyl phosphate	416 19	98			
5641	$C_{21}H_{22}N_2O_2$	Isostrychnine	334 19	214 5			
5642	$C_{21}H_{22}N_2O_2$	Strychnine	334 19	268	270[5]	1 359[18]	
5645	$C_{21}H_{22}Cl_2N_2O_2$	Benzamide hydrochloride	436 12	178			
5646	$C_{21}H_{23}NO_4$	Meconidine	353 19	58			
5647	$C_{21}H_{23}NO_5$	Cryptopine	369 19	218		1 351	
5648	$C_{21}H_{23}NO_5$	Diacetylmorphine	369 19	172			1260
5649	$C_{21}H_{23}NO_5$	α-Homochelidonine	369 19	182			
5650	$C_{21}H_{23}NO_5$	β-Homochelidonine	369 19	159			
5651	$C_{21}H_{23}NO_5$	γ-Homochelidonine	369.19	171			
5652	$C_{21}H_{23}NO_6$	Colchiceine	385 19	172			
5653	$C_{21}H_{23}N_3O_5$	Strychnine nitrate	397 20				1333
5654	$C_{21}H_{24}ClNO_4$	Diacetylmorphine hydrochloride	405 65	230			
5655	$C_{21}H_{24}N_2O$	Paytine	320 20	156			
5656	$C_{21}H_{24}N_2O$	Strychnidine	320 20	250 5	295[14]		
5657	$C_{21}H_{24}N_6O_{10}$	Geneserine picrate	520 23	175			
5658	$C_{21}H_{24}O_9$	Glycyphylline	420 19	180			
5659	$C_{21}H_{24}O_{10}$	Phloridzin	436 19	170 d.		1 430	
5660	$C_{21}H_{24}O_{11}$	Datiscin	452 19	180			
5661	$C_{21}H_{24}O_{12}$	Saponarin	468 19	232			
5663	$C_{21}H_{25}NO_4$	Corybulbine	355 20	239			
5664	$C_{21}H_{25}NO_4$	Corydine	355 20	105			
5665	$C_{21}H_{25}NO_4$	Glaucine	355 20	120			1165
5666	$C_{21}H_{25}NO_4$	Isocorybulbine	355 20	180			
5667	$C_{21}H_{25}N_3O_2$	Porphyrine	351 22	97			
5668	$C_{21}H_{26}N_2O$	Desoxystrychnine	322 22	172			
5669	$C_{21}H_{26}N_2O_3$	Corynanthine	354 22	242			
5670	$C_{21}H_{26}N_2O_3$	Quebrachine	354 22	248			1333
5671	$C_{21}H_{26}N_2O_4$	Quinine formate	370 22	113			
5672	$C_{21}H_{27}ClN_2O_3$	Quebrachine hydrochloride	390 68	290			
5673	$C_{21}H_{27}NO_4$	d(l)-Laudanosine	357 22	89			
5674	$C_{21}H_{27}NO_{10}$	d-Cocaine bitartrate	453 22	112			
5675	$C_{21}H_{28}N_2O$	Tetraethyldiaminobenzophenone	324 23	96			
5676	$C_{21}H_{31}O_4$	Marrubiin	344 22	154.5	297[15]		
5677	$C_{21}H_{30}N_2O_4$	Struxine	374 25	250 d.			
5678	$C_{21}H_{30}O_2$	Cannabinol	314 23		315[100]	1 042[18]	
5679	$C_{21}H_{30}O_4$	Euonymol	346 23	250			
5680	$C_{21}H_{30}O_9$	Antiarin	410 23	215			
5681	$C_{21}H_{34}O$	Pyrethrol	302 27	199	290		
5682	$C_{21}H_{34}O_2$	Benzyl myristate $C_{13}H_{27}CO_2CH_2C_6H_5$	318 26	20 5	231[11]	0 932[25]	536
5683	$C_{21}H_{34}O_3$	Di-d-bornyl carbonate	334.26	216			
5684	$C_{21}H_{34}O_4$	Ipurganol	350 26	225			
5685	$C_{21}H_{34}O_{10}$	Helleborein	446.26	230 d.			
5686	$C_{21}H_{36}O_4$	Trifolianol	352 28	300			
5687	$C_{21}H_{38}O_3$	Di-l-menthyl carbonate	338 29	106			
5688	$C_{21}H_{38}O_6$	Tricaproin	386 29	−25		0 988	392
5689	$C_{21}H_{40}O_2$	Dimentholformal	324 31	57	337		
5690	$C_{21}H_{42}$	9-Heneicosene $C_8H_{17}CH:CHC_{11}H_{23}$	294 32	3	202[11]	0.805[15]	
5691	$C_{21}H_{42}O_2$	Cluytinic acid	326 32	69			
5692	$C_{21}H_{42}O_2$	Heneicosonic acid $CH_3(CH_2)_{19}CO_2H$	326 32	74			
5693	$C_{21}H_{43}NO$	Heneicosamide $CH_3(CH_2)_{19}CONH_2$	325 34	110			
5694	$C_{21}H_{44}$	n-Heneicosane $CH_3(CH_2)_{19}CH_3$	296.34	40 4	215[16]	0.775[40.5]	1067
5695	$C_{22}H_{14}$	Picene	278 11	364	520		
5696	$C_{22}H_{14}N_2O$	Rosindon (Rosindulon)	322.12	262			
5697	$C_{22}H_{15}NO_5$	Colchinine	389.12	146			
5698	$C_{22}H_{15}N_3$	Rosinduline	321.14	199			
5699	$C_{22}H_{18}O_4$	o-Cresolphthalein	346.14	216			
5700	$C_{22}H_{20}O_{13}$	Carminic acid	492 15	136 d.			
5701	$C_{22}H_{22}O_{11}$	Isotrifolin	462.17	250			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5702	C₂₂H₂₂O₁₁	Trifolin	462.17	260			
5703	C₂₂H₂₃NO₇	Gnoscopine	413.19	233			
5704	C₂₂H₂₃NO₇	Narcotine.	413.19	175		1 371	
5705	C₂₂H₂₃N₃O₇	Pyrene picrate	431.12	218			
5706	C₂₂H₂₄O₁₀	Sakuranin .	448.19	212			
5707	C₂₂H₂₅NO₄	Corycavidine .	367.20	213			
5708	C₂₂H₂₅NO₆	l-Colchicine...	399.20	146			1333
5709	C₂₂H₂₆N₂O₂	Apoyohimbine	350.22	252			
5710	C₂₂H₂₆N₂O₃	Acetylquinine	366.22	108			
5711	C₂₂H₂₆N₂O₃	Gelsemine .	366.22	178			
5712	C₂₂H₂₆N₂O₄	Chaimaridine	382.22	128			
5713	C₂₂H₂₆N₂O₄	Chaimarine	382.22	233			
5714	C₂₂H₂₆N₂O₄	Conchamarine	382.22	120			
5715	C₂₂H₂₆N₂O₄	Conchairamidine	382.22	115			
5716	C₂₂H₂₆N₂O₄	Mitraversine	382.22	237			
5718	C₂₂H₂₆O₁₂	Hesperidin	482.20	171	251 d.		
5719	C₂₂H₂₇AsNO₆	Strychnine methylarsinate	460.18	60 d			
5720	C₂₂H₂₇BrN₂O₃	Gelsemine hydrobromide	447.14				1333
5721	C₂₂H₂₇ClN₂O₂	Apoyohimbine hydrochloride	386.68	300			
5722	C₂₂H₂₇ClN₂O₃	Gelseminine hydrochloride	402.68	330			1333
5723	C₂₂H₂₇NO₄	dl-Corydaline . .	369.22	136			
5724	C₂₂H₂₇N₃O₅	Physostigmine salicylate	413.23	178.9			1333
5725	C₂₂H₂₈N₂O₂	Aspidosamine	352.23	100			
5726	C₂₂H₂₈N₂O₂	Aspidospermatine	352.23	162			
5727	C₂₂H₂₈N₂O₄	Ditaine (Echitamine)	384.23	206			1333
5728	C₂₂H₂₈N₂O₄	Quinine acetate ..	384.23	126			
5729	C₂₂H₂₈N₄	Camphorosazone	348.25	55			
5730	C₂₂H₂₈O₃	Santalyl salicylate	340.22		126.6²⁰	1 070¹⁰	
5732	C₂₂H₂₉IO₂	Europhen (Diisobutyl-p-cresol iodide)	452.16	110			
5733	C₂₂H₃₀N₂O₂	Aspidospermine	354.25	208	220²		
5734	C₂₂H₃₁NO₅ (?)	Mitragynine .	389.25	106	240⁶		
5735	C₂₂H₃₂O₃	Anacardic acid	344.25	26			
5736	C₂₂H₃₂O₄	Digitoxigenin . .	360.25	230			
5737	C₂₂H₃₂O₆	Genin	392.25	206			
5738	C₂₂H₃₃NO₅	Atropine isovalerate	391.26	32			
5739	C₂₂H₃₃NO₅	Atropine valerate	391.26	42			1333
5741	C₂₂H₃₄N₄O₈S	Pilocarpine sulfate	514.36	132			1333
5742	C₂₂H₃₅NO₆	Delphinine .	409.28	187.5			
5743	C₂₂H₃₆O₄	Bryonol .. .	364.28	212			
5744	C₂₂H₃₆O₅	Capsularin....	428.28	176			
5745	C₂₂H₃₇NO	Palmitic anilide	331.29	90.5	284¹⁷		
5746	C₂₂H₃₈O	Cholestol... ..	318.29	139	360		
5747	C₂₂H₃₈O	Ilicyl alcohol	318.29	175	350		
5748	C₂₂H₃₈O₄	Citrullol..	366.29	290			
5759	C₂₂H₃₈O₄	Di-l-menthyl oxalate	366.29	68	225¹²		
5760	C₂₂H₃₉ClO	Behenolyl chloride C₂₁H₃₉COCl	354.76	29			
5761	C₂₂H₄₀O₂	Behenolic acid C₂₁H₃₉CO₂H	336.31	57.5			
5762	C₂₂H₄₁NO	Behenolyl amide C₂₁H₃₉CONH₂	335.32	90			
5763	C₂₂H₄₂O₂	Brassidic acid	338.32	61.5	282³⁰	0.859⁵⁷·¹	1085
5764	C₂₂H₄₂O₂	Erucic acid	338.32	33.5	281⁵⁰	0.860⁵⁵·⁴	
5765	C₂₂H₄₂O₃	14-Ketobehenic acid	354.32	84			
5765.1	C₂₂H₄₂O₃	Isobutyl ricinoleate	354.32		262⁹	0.903²²	980
5766	C₂₂H₄₃NO	Erucamide C₂₁H₄₁CONH₂	337.34	83			
5767	C₂₂H₄₄O	Erucyl alcohol. .	324.34	34.6	200⁰·²		
5768	C₂₂H₄₄O₂	Behenic acid . .	340.34	84	306⁶⁰		
5769	C₂₂H₄₄O₂	Methyl heneicosate C₂₀H₄₁CO₂CH₃	340.34	49			
5770	C₂₂H₄₅I	Docosyl iodide CH₃(CH₂)₂₀CH₂I	436.28	49			
5771	C₂₂H₄₅NO	Behenamide C₂₁H₄₃CONH₂ .	339.36	112			
5772	C₂₂H₄₆	n-Docosane CH₃(CH₂)₂₀CH₃	310.35	44.4	224.5¹⁵	0 778⁴⁴·⁴	
5773	C₂₂H₄₆O	Docosyl alcohol CH₃(CH₂)₂₀CH₂OH .	326.35	74			
5774	C₂₃H₂₀O₂	Amaric anhydride	328.15	140.5			
5775	C₂₃H₂₃NO₄	Corycavine................	409.19	216			
5776	C₂₃H₂₄N₂O₆	Buphnatine.......●.............	424.20	240			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R I. No.
5777	$C_{22}H_{24}N_4O_2$	Methylenediantipyrine	388.22	177			
5778	$C_{22}H_{24}N_4O_{11}$	Hyoscine picrate	532.22	188			
5779	$C_{23}H_{24}O_4$	o-Cresol orthoacetate	348.19	89			
5780	$C_{23}H_{24}O_9$	Picropodophyllin	444.19	227			
5781	$C_{23}H_{24}O_9$	Podophyllotoxin	444.19	94			
5782	$C_{23}H_{25}NO_4$	Lanthopine	379.20	200			
5783	$C_{23}H_{26}ClN_2O_2$	Acom	427.68	178			
5784	$C_{23}H_{26}N_2O_4$	Aricine	394.22	188 d.			
5785	$C_{23}H_{26}N_2O_4$	Brucine	394.22	178			
5786	$C_{23}H_{26}N_2O_4$	Concuseonine	394.22	208			
5787	$C_{23}H_{26}N_2O_4$	Cusconine	394.22	110			
5788	$C_{23}H_{26}N_2O_5$	Allobrucine oxide	410.22	189			
5789	$C_{23}H_{27}NO_6$	Homoatropine salicylate	413.22				1333
5790	$C_{23}H_{27}NO_8$	Narceine	445.22	170			
5791	$C_{23}H_{27}N_3O_7$	Brucine nitrate	457.23	230 d.			
5792	$C_{23}H_{28}ClNO_8$	Narceine hydrochloride	481.68	192			1333
5793	$C_{23}H_{28}N_2O_4$	Vellosine	396.23	189 d.			
5794	$C_{23}H_{29}NO_2$	Lobeline	351.23	131			
5795	$C_{23}H_{30}N_2O_4$	Quinine propionate	398.25	111			
5796	$C_{23}H_{30}N_2O_4$	dl-Quinine lactate	414.25	165.5			
5797	$C_{23}H_{30}N_2O_5$	d-Quinine lactate	414.25	175			
5798	$C_{23}H_{30}N_2O_5$	l-Quinine lactate	414.25	171			
5799	$C_{23}H_{31}NO_2$	Atisine	353.25	85			
5801	$C_{23}H_{32}N_2O_4$	Quinine ethyl carbonate (Equinine)	401.27	91			
5802	$C_{23}H_{33}N_3O_5$	Pyramidon acid camphorate	431.28	94			
5803	$C_{23}H_{36}O_2$	Lactucon (Lactucol acetate)	344.28	184			
5804	$C_{23}H_{36}O_4$	Calabarol	376.28	245			
5804.1	$C_{23}H_{38}N_2$	Conessine	342.31	125			1333
5805	$C_{23}H_{38}O_2$	Benzyl palmitate	346.29	36		0.914_{25}^{36}	1079
5806	$C_{23}H_{38}O_4$	Anonol	378.29	298			
5807	$C_{23}H_{38}O_4$	Grindelol (Phytosterol glucoside)	378.29	257			
5808	$C_{23}H_{40}O$	Ambrein	332.31	82			
5809	$C_{23}H_{40}O$	Xanthosterin	332.31	214			
5810	$C_{23}H_{40}O_4$	Di-l-menthyl malonate	380.31	62	170^1	0.944_4^{76}	
5811	$C_{23}H_{40}O_4$	Ipuranol	380.31	290			
5812	$C_{23}H_{44}O_2$	Methyl behenolate $C_{21}H_{39}CO_2CH_3$	350.32	22			
5813	$C_{23}H_{44}O_2$	Methyl erucate $C_{21}H_{41}CO_2CH_3$	352.34		222^5	0 870	457
5814	$C_{23}H_{46}O$	Laurone $(C_{11}H_{23})_2CO$	338.35	69		$0.789_4^{90.9}$	1111
5815	$C_{23}H_{46}O_2$	Methyl behenate $C_{21}H_{43}CO_2CH_3$	354.35	54.5	225		
5816	$C_{23}H_{48}$	n-Tricosane $CH_3(CH_2)_{21}CH_3$ ...	324.37	47.7	320.7	$0.779_4^{47.7}$	1120
5817	$C_{24}H_{16}$	Crackene	306.14	308	500		
5818	$C_{24}H_{18}$	1, 3, 5-Triphenylbenzene	306.14	170		1 206	1317
5819	$C_{24}H_{18}As_2N_2O$	Phenarsazine oxide	500.08	350			
5820	$C_{24}H_{18}N_2$	p, p'-Diphenylazobenzene	334.16	250			
5821	$C_{24}H_{18}N_2O$	p, p'-Diphenylazoxybenzene	350.16	205			
5822	$C_{24}H_{20}N_2$	p, p'-Diphenylhydrazobenzene	336.17	247			
5823	$C_{24}H_{20}O_6$	Glycerol tribenzoate	404.15	76.5			
5824	$C_{24}H_{20}O_9$	Glycerol trisalicylate	452.15	79			
5826	$C_{24}H_{25}N_3O$	Benzoylauramine	371.22	179			
5829	$C_{24}H_{28}O_6$	Diguaiacyl camphorate	412.22	124			
5830	$C_{24}H_{28}O_8$	α-Flavaspidic acid	444.22	92			
5831	$C_{24}H_{28}O_8$	β-Flavaspidic acid	444.22	156			
5832	$C_{24}H_{29}NO_6$	Atropine salicylate	427.23				1333
5834	$C_{24}H_{30}O_5$	Elaterone	398.23	300			
5835	$C_{24}H_{30}O_7$	Anthamantin	430.23	79			
5836	$C_{24}H_{30}O_{11}$	Scopolin	558.23	218			
5837	$C_{24}H_{32}N_2O_4$	Quinine butyrate	412.26	77.5			
5838	$C_{24}H_{32}N_4O_9$	Maltosazone	520.28	206			
5839	$C_{24}H_{38}N_2O$	Holarrhenine	370.31	198			
5840	$C_{24}H_{38}O_4$	Di-d-bornyl succinate	390.29	83.7			
5841	$C_{24}H_{40}N_2$	Conessine	356.32	125			
5842	$C_{24}H_{40}O_4$	Choleic acid	392.31	190			
5843	$C_{24}H_{40}O_4$	Cucurbitol ...	392.31	260			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5844	$C_{24}H_{40}O_5$	Cholic acid .	408 31	195			
5845	$C_{24}H_{41}NO$	Stearic amlide $CH_3(CH_2)_{16}CONHC_4H_5$	359 32	93 6			
5846	$C_{24}H_{42}O_4$	Di-*l*-menthyl succinate	394 32	63	220 d	0 947$_4^{71}$	
5847	$C_{24}H_{42}O_6$	Di-*l*-menthyl *d*-tartrate	426 32	75		1 054	
5848	$C_{24}H_{42}O_6$	Di-*l*-menthyl *l*-tartrate	426 32	42		1 045^{14}	
5849	$C_{24}H_{44}O_5$	Lithofellinic acid	412 34	206			
5850	$C_{24}H_{44}I_2O_2$	Ethyl diiodobrassidate	648 20	37			
5851	$C_{24}H_{46}O_2$	Ethyl behenolate $C_{21}H_{39}CO_2C_2H_5$	364 34	15			
5852	$C_{24}H_{46}O_2$	Ethyl brassidate .	366 35	30 5			1046
5853	$C_{24}H_{46}O_2$	Ethyl erucate $C_{21}H_{41}CO_2C_2H_5$	366 35		230	0 865	449
5854	$C_{24}H_{48}O_2$	Carnaubic acid	368 37	72			
5855	$C_{24}H_{48}O_2$	Lignoceric acid $C_{23}H_{47}CO_2H$	368 37	81			
5856	$C_{24}H_{48}O_2$	Paraffinic acid $C_{23}H_{47}CO_2H$	368 37	46			
5857	$C_{24}H_{48}O_2$	Pisangceryhc acid $C_{23}H_{47}CO_2H$	368 37	72			
5858	$C_{24}H_{48}O_2$	Tetracosic acid $CH_3(CH_2)_{22}CO_2H$	368 37	85 5			
5859	$C_{24}H_{48}O_2$	Ethyl behenate $C_{21}H_{43}CO_2C_2H_5$	368 37	50 5	231		
5860	$C_{24}H_{50}$	Isotetracosane .	338 39	51	243^{15}		
5861	$C_{24}H_{50}$	*n*-Tetracosane $CH_3(CH_2)_{22}CH_3$	338 39	51	324 1	0 779$_4^{51.1}$	
5862	$C_{24}H_{50}O$	Carnaubyl alcohol $C_{24}H_{49}OH$	354 39	69			
5863	$C_{25}H_{20}$	Tetraphenylmethane $C(C_6H_5)_4$	320 15	285	431		
5864	$C_{25}H_{21}N_3$	Tetraphenylguanidine	363 19	131			
5865	$C_{25}H_{28}O_{11}$	Ononin	502 20	210			
5866	$C_{25}H_{28}O_{14}$	Gentiin	552 22	274			
5867	$C_{25}H_{29}NO_8S$	Codeine *o*-guaiacolsulfonate	503 30	165			
5868	$C_{25}H_{32}O_8$	Albaspidin	460 25	147			
5869	$C_{25}H_{32}O_8$	Aspidin	460 25	124			
5871	$C_{25}H_{34}O_{14}$	Loganin	558 26	215			
5872	$C_{25}H_{39}NO_8$	Pseudoaconine	481 31	95			
5873	$C_{25}H_{40}O$	Fungisterin	356 31	144			
5874	$C_{25}H_{40}O$	Homotaraxasterol	356 31	164			1024
5875	$C_{25}H_{40}O_2$	Benzyl oleate .	372 31		237^7	0 933$_{15}^{14}$	1078
5876	$C_{25}H_{42}O_2$	Benzyl stearate $C_{17}H_{35}CO_2CH_2C_6H_5$	374 32	45 8		0 908$_{16}^{40}$	
5877	$C_{25}H_{44}O_4$	Di-*l*-menthyl glutarate	408 34		243^{20}		
5878	$C_{25}H_{50}O_2$	Neocerotic acid	382 39	77 8			
5879	$C_{25}H_{50}O_2$	Hyenic acid .	382 39	78			
5880	$C_{25}H_{50}O_3$	Cerebronic acid	398 39	100			
5881	$C_{25}H_{52}$	Pentacosane $CH_3(CH_2)_{23}CH_3$	352 40	54	284^{40}	0.779	
5882	$C_{26}H_{14}$	Rubicene	326 11	306			
5883	$C_{26}H_{20}$	Tetraphenylethylene	332 15	221	425		
5884	$C_{26}H_{20}O$	α-Benzopinacoline	348 15	205			
5885	$C_{26}H_{20}O$	β-Benzopinacoline	348 15	181			
5886	$C_{26}H_{21}NO_{11}$	Aconine	523 17	132			
5887	$C_{26}H_{22}$	1, 1, 2, 2-Tetraphenylethane	334 17	209	383	1 182	
5888	$C_{26}H_{22}N_4$	Benzilosazone	390 20	225			
5889	$C_{26}H_{22}O_2$	Benzopinacone	366 17	186 d.			
5890	$C_{26}H_{23}N_5$	Tetraphenyldiguanidine	405 22	136			
5891	$C_{26}H_{26}N_2O_2$	Benzoylcinchonine	398 22	106			
5892	$C_{26}H_{27}ClN_2O_2$	Benzoylcinchonine hydrochloride	434 68	207			
5893	$C_{26}H_{28}N_2O_4$	Cinchonidine salicylate	432 23	70			
5895	$C_{26}H_{28}O_{14}$	Ruberythric acid	564 22	260			
5896	$C_{26}H_{28}O_{14}$	Morindin	564 22	245	247		
5897	$C_{26}H_{30}N_2O_6S$	Quinine phenolsulfonate	498 31				1333
5898	$C_{26}H_{30}O_4$	Bixin .	406 23	189			
5899	$C_{26}H_{32}N_2O_2$	Ibogine	404 26	152			
5900	$C_{26}H_{37}NO_2$	Jervine	411 29	241			
5901	$C_{26}H_{38}$	Carotin	350 29	167 8			
5902	$C_{26}H_{40}O$	Ergosterin	368 31	154	185^{20}	1.040	
5903	$C_{26}H_{40}O_7$	Laserpitin	464 31	117 5	240^{10} d.		
5904	$C_{26}H_{41}NO_{10}$	Japaconine	527 32	97			
5905	$C_{26}H_{42}O_3$	Sarsasapogenin	402 32	183			
5906	$C_{26}H_{42}O_3$	Smilacin .	402.32	160 d.			
5907	$C_{26}H_{43}NO_2$	Rubijervine	401 34	236			
5908	$C_{26}H_{43}NO_6$	Glycocholic acid .	465 34	134			

No	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No
5909	$C_{24}H_{44}O$	Caulosterol	372 34	159			
5910	$C_{26}H_{44}O_2$	Onocerin	388 34	232			
5911	$C_{26}H_{44}O_4$	Gitogenin	420 34	272			
5912	$C_{26}H_{44}O_{10}$	Parillin	516 34	176 1			
5913	$C_{26}H_{45}NO_5$	Protoveratridine	499 36	265			
5914	$C_{26}H_{46}O$	Mochyl alcohol $C_{26}H_{45}OH$	374 35	234			
5915	$C_{26}H_{46}O_4$	Di-l-menthyl adipate	122 35	61			
5916	$C_{26}H_{52}O_2$	Cerotic acid	396 40	82 5			0 836$^{79}_4$
5917	$C_{26}H_{52}O_2$	Ethyl lignocerate	396 40	56	310^{20}		
5918	$C_{26}H_{54}$	n-Hexacosane $CH_3(CH_2)_{24}CH_3$	366 42	60	296^{40}	0 779	
5919	$C_{26}H_{54}$	Isohexacosane	366 42	61	207$^{9'}$		
5920	$C_{26}H_{54}O$	Ceryl alcohol $C_{26}H_{53}OH$	382 42	80			
5921	$C_{27}H_{29}Br_2N_2O_6$	Quinine dibromosalicylate	620 06	198			
5922	$C_{27}H_{24}N_6S_3$	Diphenylguanidine trithiocarbonate	532 16	89			
5925	$C_{27}H_{30}N_2O_6$	Quinine salicylate	462 25	187			1333
5926	$C_{27}H_{26}O_{15}$	Apiin	594 23	228			
5927	$C_{27}H_{30}O_{16}$	Sophorin	610 23	166			
5928	$C_{27}H_{32}O_{16}$	Rutin	612 25	183	d.		
5929	$C_{27}H_{38}O_7$	Strophantidin	474 29	195			
5930	$C_{27}H_{39}N_3O_5$	Paucine	513 34	126			
5931	$C_{27}H_{40}O_8$	Cerberin	492 31	192			
5932	$C_{27}H_{42}O$	Ergosterin	382 32	165			
5933	$C_{27}H_{46}O$	Cholesterin	386 35	148	>360	1 067	
5934	$C_{27}H_{46}O$	Phytosterol	386 35	136			
5935	$C_{27}H_{46}O$	Sitosterol	386 35	140			
5936	$C_{27}H_{46}O_5$	Atropurol	402 35	285			
5937	$C_{27}H_{47}N$	Cholesterylamine	385 37	104			
5938	$C_{27}H_{47}NO_9$	Indaconine	529 37	94			
5939	$C_{27}H_{48}O$	Coprosterol	388 37	105			
5940	$C_{27}H_{50}O_6$	Tricaprylin	470 39	8		0 954	425
5941	$C_{27}H_{54}O$	Myristone $(C_{13}H_{27})_2CO$	394 42	76		0 792$^{90}_4$ 9	
5942	$C_{27}H_{56}$	n-Heptacosane $CH_3(CH_2)_{25}CH_3$	380 43	59 5	270^{15}	0 779$^{59}_4$ 5	
5943	$C_{28}H_{14}$	9, 9'-Dianthranyl	354 14	300			
5944	$C_{28}H_{20}N_2$	Amaron (Tetraphenylpyrazine)	384 17	240			
5945	$C_{28}H_{22}N_2O$	Benzoylamarin	402 19	180			
5946	$C_{28}H_{22}O_2$	Anthrapinacone	390 17	182 d.			
5947	$C_{28}H_{24}N_2$	Benzylamarin	388 20	124			
5948	$C_{28}H_{24}N_2O_5$	Strychnine salicylate	472 23				1333
5949	$C_{28}H_{30}O_5$	Columbin ...	398 23	182			
5950	$C_{28}H_{34}O_{11}$	Phillrin	546 26	160			
5951	$C_{28}H_{36}N_2O_4$	Ipecamine	464 29	90			
5952	$C_{28}H_{36}N_2O_4$	Psychotrine	464 29	138			
5953	$C_{28}H_{36}O_6$	Digitogenic acid	484 28	210			
5954	$C_{28}H_{38}N_2O_4$	Cephaeline	466 31	99			
5955	$C_{28}H_{38}N_2O_4$	Hydroipecamine	466 31	92			
5956	$C_{28}H_{38}O_7$	α-Elaterin	486 29	232			
5957	$C_{28}H_{38}O_7$	β-Elaterin	486 29	195			
5958	$C_{28}H_{44}O_2$	Lactucerin	412 34	210			
5959	$C_{28}H_{45}NO$	Behenolic anilide $C_{21}H_{39}CONHC_6H_5$	411 36	72			
5960	$C_{28}H_{46}NO_9$	Isopyroine	540 36	160			
5961	$C_{28}H_{46}O_2$	Cholesteryl formate	414 35				1216
5962	$C_{28}H_{47}NO$	Brassidic anilide $C_{21}H_{41}CONHC_6H_5$	413 37	78			
5963	$C_{28}H_{47}NO$	Erucic anilide $C_{21}H_{41}CONHC_6H_5$	413 37	66			
5964	$C_{28}H_{48}O_{10}$	Gitalin	544 37	253			
5965	$C_{28}H_{49}NO$	Behenic anilide $CH_3(CH_2)_{20}CONHC_6H_5$	415 39	102			
5966	$C_{28}H_{54}O_2$	l-Menthyl stearate	422 42	39			
5967	$C_{28}H_{58}$	Octocosane $CH_3(CH_2)_{26}CH_3$	394 45	65	318^{40}	0.779	
5968	$C_{28}H_{58}O$	Cluytyl alcohol	410 45	82.5			
5969	$C_{29}H_{24}O_4$	Fortoin (Methylenedicotoine)	500 19	213			
5970	$C_{29}H_{26}O_{12}$	Aromadendrin	566 20	216			
5971	$C_{29}H_{32}N_2O_6$	Quinine acetylsalicylate	504 26	157			
5972	$C_{29}H_{35}NO_7$	Paniculatine .	509 28	263			
5973	$C_{29}H_{36}N_2O_4$	Emetamine..	476.29	156			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I. No.
5974	C₂₉H₄₀N₂O₄	Isoemetine	480.32	98			
5975	C₂₉H₄₂Cl₂N₂O₄	Isoemetine hydrochloride	553.26	310 d			
5976	C₂₉H₄₃NO₇	Pseudojervine	517.31	307			
5977	C₂₉H₄₃NO₈	Sabadenine	533.31	160	197 d		1333
5978	C₂₉H₄₈	Spinacene	396.37	20	260ª	0.8592²⁰	570
5979	C₂₉H₄₈O	Taraxasterol	412.37	222			
5980	C₂₉H₄₉O₃	Phytosterol acetate	445.38	122			
5981	C₂₉H₅₀O₄	Cluytianol	478.39	300			
5982	C₂₉H₅₁NO₈	Sabadine	541.40	240			
5983	C₂₉H₅₂O₂₀	Sapotin	720.40	240			
5984	C₂₉H₅₈O₂	Montanic acid	438.45	86.8			
5985	C₂₉H₆₀	Nonacosane CH₃(CH₂)₂₇CH₃	408.46	63.6	318⁶⁰	0.780	
5986	C₃₀H₂₀NO₉	Adlumidine	538.16	234			
5987	C₃₀H₂₈O₁₀	Santalin	548.22	226	195ª		
5989	C₃₀H₃₄O₁₃	Picrotoxin	602.26	200			
5990	C₃₀H₃₈O₄	Hellesboresin	462.29	150 d			
5991	C₃₀H₄₀N₂O₅	Emetine	508.32	74			
5993	C₃₀H₄₂Cl₂N₂O₅	Emetine dihydrochloride	581.26	53			1333
5994	C₃₀H₄₂I₂N₂O₅	Emetine dihydroiodide	764.20	238			
5995	C₃₀H₄₂N₂O₁₆S₂	Sinalbin	731.17	138.5			
5996	C₃₀H₄₄N₆O₆S	Physostigmine sulfate	648.45	140			
5997	C₃₀H₄₄O₉	Cymarin	548.31	138 d			
5998	C₃₀H₄₆O₁₂	Ouabain	598.35	185			
5999	C₃₀H₄₈O₂	Echicerin	440.37	157			
6000	C₃₀H₄₈O₂	Mycosterol	440.37	160			
6001	C₃₀H₄₈O₈	β-Quinovin	536.37	235			
6002	C₃₀H₅₀O	α-Amyrin	426.39	185	>300		
6003	C₃₀H₅₀O	β-Amyrin	426.39	195			
6004	C₃₀H₅₀O	Androsterol	426.39	208			
6005	C₃₀H₅₀O	Stigmasterol	426.39	140			
6006	C₃₀H₅₀O₂	Betulin	442.39	252			
6007	C₃₀H₅₀O₂	Cholesterol propionate	442.39	98.7			
6008	C₃₀H₅₂O₄	Menthyl camphorate	476.40	86			
6009	C₃₀H₅₄N₄O₆S	Sparteine sulfate	566.51				1333
6010	C₃₀H₆₀	Melene	420.46	63	380	0.890	
6011	C₃₀H₆₀O₂	Melissic acid CH₃(CH₂)₂₈CO₂H	452.46	91			
6012	C₃₀H₆₀O₄	Lanoceric acid	484.46	105			
6013	C₃₀H₆₂	Melissane	422.48	71	222⁶⁴		
6014	C₃₀H₆₂	n-Triacontane CH₃(CH₂)₂₈CH₃	422.48	70	235¹⁵	0.780	
6015	C₃₀H₆₂O	Melissyl alcohol	438.48	88		0.777⁹⁵	
6016	C₃₀H₆₂O₂	Cocceryl alcohol	454.48	104			
6017	C₃₁H₁₅NO₄	Apomorphine dibenzoate	465.12	156			
6018	C₃₁H₂₆O₁₀	Tephrosin	558.20	187			
6019	C₃₁H₂₇NO₅	Dibenzoylmorphine	493.22	190.5			
6020	C₃₁H₃₈O₁₀	Kosin	570.29	142			1333
6021	C₃₁H₄₃NO₁₁	Napelline	603.36	165			
6022	C₃₁H₄₈O	Lupeon	431.33	170			
6023	C₃₁H₅₀O	Lupeol	438.39	215			
6024	C₃₁H₅₂O₂	Cholesterol butyrate	456.40	92.8			
6025	C₃₁H₅₂O₂	Euonysterol	456.40	138			
6026	C₃₁H₆₂O	Palmitone (C₁₅H₃₁)₂CO	450.48	83		0.795²⁰·⁹	1125
6027	C₃₁H₆₂O₃	Cocceric acid	482.48	93			
6028	C₃₁H₆₄	n-Hentriacontane CH₃(CH₂)₂₉CH₃	436.49	68.1	302¹⁵	0.781²⁸·¹	
6029	C₃₂H₂₂O₁₀	Heraclin	566.17	185			
6030	C₃₂H₂₆	Pentaphenylethane	410.20	173			
6031	C₃₂H₂₇N₂O	Benzacine	469.23	150			
6032	C₃₂H₄₁NO₉	Pyraconitine	583.32	171			
6032.1	C₃₂H₄₂N₂O₅	Lappaconitine	598.34	223			
6033	C₃₂H₄₄N₂O₁₀S	Homoatropine sulfate	648.42				1333
6034	C₃₂H₄₄O₁₀	Quassiin	588.34	211			
6035	C₃₂H₄₅NO₈	Indobenzaconine	587.36	130			
6036	C₃₂H₄₆BrNO₁₀	Benzaconine hydrobromide	684.28	282			
6037	C₃₂H₄₆ClNO₁₀	Benzaconine hydrochloride	639.82	α 217; β 208			

No.	Formula	Name	Mol. wt.	M. P.	B. P.	d	R. I No
6038	$C_{22}H_{43}N_2O_{16}S$	Sinapine sulfate	716 45	193			
6039	$C_{32}H_{49}NO_9$	Veratrine	591 39	205			
6040	$C_{32}H_{51}NO_{11}$	Protoveratrine	625 40	250			
6041	$C_{32}H_{52}N_2O_2$	Lycopodine	512 42	115			
6042	$C_{32}H_{52}O_2$	Echitin	468 40	170			
6043	$C_{32}H_{54}O_2$	Cholesterol valerate	470 42	89 6			
6044	$C_{32}H_{54}O_2$	Phytosterol valerate	470 42	30			
6045	$C_{32}H_{62}O_3$	Palmitic anhydride $(C_{15}H_{31}CO)_2O$	494 48	64			
6046	$C_{32}H_{62}O_{14}$	Convolvulin (Rhodeoretin)	702 48	158			
6047	$C_{32}H_{64}O_2$	Cetyl palmitate $C_{15}H_{31}CO_2C_{16}H_{34}$	480 49	54		0 832$_4^{50}$	
6048	$C_{32}H_{66}$	n-Dotriacontane $CH_3(CH_2)_{30}CH_3$	450 51	75	310^{15}	0 775$^{79.4}$	1110
6049	$C_{33}H_{40}O_{19}$	Robinin	740 39	195			
6050	$C_{33}H_{43}NO_{11}$	Anhydroaconitine	629 34	186			
6051	$C_{33}H_{46}N_2O_5$	Septentrionaline	614 37	131			
6052	$C_{33}H_{50}O_{10}$	Tormentol	606 39	228			
6053	$C_{33}H_{53}NO_7$	Solangustine	575 42	235 d.			
6054	$C_{33}H_{56}O_2$	Cholesterol capronate	484 44	91 2			
6055	$C_{33}H_{56}O_4$	Phytosteroline	548 43	290			
6056	$C_{33}H_{62}O_6$	Tricaprin	554 48	31 1		0 921$_4^{40}$	1054
6057	$C_{33}H_{66}O_2$	Psyllostearic acid	494 51	95			
6058	$C_{33}H_{68}O$	Psyllostearyl alcohol	480 52	69 5			
6059	$C_{34}H_{32}O_4$	Isoeugenol dibenzoate	536 25	161			
6060	$C_{34}H_{36}N_2O_6$	Pseudomorphine	568 29	327 d.			
6061	$C_{34}H_{36}N_2O_9$	Sekisanine	616 29	200			
6062	$C_{34}H_{40}N_2O_{10}S$	Morphine sulfate	668 39	250 d.			1333
6063	$C_{34}H_{40}N_2O_{12}S_2$	Quinine diguaiacolsulfonate	732 45	130 d.			
6064	$C_{34}H_{44}N_2O_8S$	Apoatropine sulfate	640 42				1333
6065	$C_{34}H_{46}O_4$	d-Camphor salicylate	580 34	60			
6066	$C_{34}H_{47}NO_{10}$	Indaconitine	629 37	203			
6067	$C_{34}H_{47}NO_{11}$	Aconitine	645 37	195			
6068	$C_{34}H_{48}BrNO_{11}$	Aconitine hydrobromide	726 29	163			1333
6069	$C_{34}H_{48}ClNO_{11}$	Aconitine hydrochloride	681 84	149			1333
6070	$C_{34}H_{48}N_2O_{10}S$	Atropine sulfate	676 45	194			1333
6071	$C_{34}H_{48}N_2O_{10}S$	Hyoscyamine sulfate	676 45	206			1333
6072	$C_{34}H_{48}N_2O_{14}$	Aconitine nitrate	708 39				1333
6073	$C_{34}H_{49}NO_{11}$	Japaconitine	647 39	204 2			
6074	$C_{34}H_{50}ClNO_{11}$	Japaconitine hydrochloride	683 85	149			
6075	$C_{34}H_{50}O_2$	Cholesterol benzoate	490 39	145 5			
6076	$C_{34}H_{50}O_3$	Cholesterol salicylate	506 39	180			1180
6077	$C_{34}H_{54}O_{11}$	Digitoxin	638 42	244			
6078	$C_{34}H_{56}O_{16}$	Jalapin	720 43	150			
6079	$C_{34}H_{57}NO_2$	Solanidine	511 45	215			
6080	$C_{34}H_{70}$	n-Tetratriacontane	478 54	76 5	255$^{1 0}$	0 781	
6081	$C_{34}H_{70}O$	Incarnatryl alcohol	494 54	74			
6082	$C_{35}H_{38}O_{17}$	Filixic acid	650 29	184			
6083	$C_{35}H_{39}N_5O_5$	Ergotinine	609 .34	229 d.			1333
6084	$C_{35}H_{41}N_5O_6$	Ergotoxine	627 36	164			
6085	$C_{35}H_{44}N_5O_{10}P$	Ergotoxine phosphate	725 40	187			
6086	$C_{35}H_{56}O_2$	Echiretin	508 43	52			
6087	$C_{35}H_{56}O_{14}$	Digitalin	700 43	217			
6088	$C_{35}H_{58}O_3$	Phytosterolene acetate	607 45	160			
6089	$C_{35}H_{60}NO_4$	Imperaline	558 47	254 d.			
6090	$C_{35}H_{70}O$	Stearone $(C_{17}H_{35})_2CO$	506 54	88		0 793$_4^{88}$	
6091	$C_{35}H_{72}$	n-Pentatriacontane	492 55	74 7	331^{15}	0.782$_4^{74.7}$	
6092	$C_{36}H_5O_6$	Lophopetalin	533 01	230			
6093	$C_{36}H_{34}N_2O_6S$	Aporheine sulfate	654 34	75			
6094	$C_{36}H_{34}N_2O_{13}$	Cynoctonine	702 28	137			
6095	$C_{36}H_{42}O_6$	Helleborin	570 32	>250 d			
6096	$C_{36}H_{42}O_{11}$	Filicic acid	682 32	125			
6097	$C_{36}H_{44}N_2O_{10}S$	Codeine sulfate	696 42	278			1333
6098	$C_{36}H_{48}O_{10}$	α-Picrasmin	640 37	204			
6099	$C_{36}H_{48}O_{10}$	β-Picrasmin	640 37	212			
6100	$C_{36}H_{50}N_4O_4$	Pyramidon camphorate	662 43	90			

No.	Formula	Name	Mol. wt	M. P.	B P.	d	R. I. No.
6101	C₃₆H₅₁NO₁₁	Bikhaconitine	673 40	113			
6102	C₃₆H₅₁NO₁₂	Pseudaconitine	689 40	211			
6104	C₃₆H₆₁O₃₁	Inulin	990 48	178 d		1 35	
6105	C₃₆H₆₆O₃	Oleic anhydride	546 51	22 2			
6106	C₃₆H₇₀O₃	Stearic anhydride [CH₃(CH₂)₁₆CO]₂O.	550 54	72			
6107	C₃₆H₇₄	Hexatriacontane	506 57	76 5	265¹⁰	0 782⁷⁶	
6108	C₃₇H₃₆N₂O₆	Xanthaline	652 29	208			
6109	C₃₇H₅₁NO₁₁	Taxine	685 40	82 d.			
6110	C₃₇H₆₄O₂	Cholesterol caprinate	540 49	82 2			
6111	C₃₈H₄₄N₂O₁₂	Morphine tartrate	720 36				1333
6112	C₃₈H₄₄N₄O₃	Dicinchonine	588 37	40			
6113	C₃₈H₄₆N₂O₄	α-Truxilline	658 37	80			
6114	C₃₈H₄₆N₂O₄	β-Truxilline	658 37	45			
6115	C₃₈H₄₆N₄O₆S	Cinchonidine sulfate	686 45	242			
6116	C₃₈H₄₆N₄O₆S	Cinchonine sulfate	686 45	198 5			
6117	C₃₈H₄₆N₄O₈S	Cupreine sulfate	718 45	257 d.			
6119	C₃₉H₄₁NO₁₂	Adlumine...	715 32	188			
6120	C₃₉H₆₃NO₁₀	Zygadenine	705 49	200			
6120.1	C₃₉H₇₄O₆	Trilaurin	638 57	46 5		0 891⁶⁵	
6122	C₄₀H₄₀N₂O₁₀S₂	Quinine-β-naphtholsulfonate	772 45	186			
6124	C₄₀H₅₀N₄O₈S	Quinine sulfate	746 48	235 2			
6125	C₄₀H₅₆O₁₅	Strophantin	776 43	179			
6126	C₄₀H₇₀O₂	Homocuonysterol	582 54	134			
6127	C₄₁H₅₀N₄O₇	Quinine carbonate	710 42	169			
6129	C₄₂H₄₆N₄O₈S	Strychnine sulfate	766 45	200			
6131	C₄₂H₅₄N₂O₇	Tritopine	698 43	182			
6133	C₄₂H₆₆O₆	Caulosapogenin	666 51	315			
6135	C₄₂H₇₀O₂	Echitein..	606 54	195			
6136	C₄₃H₄₅N₃O₂₄	Quinoline tartrate	987 37	125			
6137	C₄₃H₅₇N₄O₁₀P	Quinine glycerophosphate	820 50	181			
6138	C₄₄H₅₄N₄O₈	Quinine succinate	766 45	192			
6139	C₄₄H₅₄N₄O₉	Quinine malate .	782 45	177 5			
6141	C₄₄H₅₄N₄O₁₀	Quinine tartrate.	798 45	202 5			1333
6142	C₄₄H₆₆NO₁₉	Glycyrrhizic acid	910 50	220			
6143	C₄₄H₇₆O₂₀	Sarsasaponin	924 59	248			
6144	C₄₄H₈₂O₃	Brassidic anhydride	658 63	64		0.835⁷⁰	1145
6145	C₄₄H₈₂O₃	Erucic anhydride	658 63	48			1144
6147	C₄₅H₈₆O₆	Trimyristin	722 66	55		0.885⁴⁰	1089
6148	C₄₆H₅₀N₄O₁₀	Strychnine d-tartrate	818 42	228		1.420	
6150	C₄₆H₅₆N₂O₂₀S	Narceine sulfate	988 51				1333
6151	C₄₇H₅₄O₁₆	Filmaron..	874 42	60			
6153	C₄₈H₉₃NO₉	Phrenosin	827 72	215 s. d.			
6154	C₄₉H₈₀O₂₃	Gitonin. ..	1036 6	272 d.			
6155	C₅₀H₆₆O₂₀	Hyssopin..	1146 5	275			
6156	C₅₀H₇₀O₈	Lupulinic acid	798 54	93			
6157	C₅₁H₉₈O₆	Tripalmitin	806 76	65.1; 46		0 866⁴⁰	1114
6158	C₅₂H₉₁NO₁₈	Solanine .	1017 7	254 d.			
6159	C₅₂H₉₂ClNO₁₈	Solanine hydrochloride	1054 2	212			
6160	C₅₂H₁₀₄O₂	Ceryl cerotate.	760 80	84			
6161	C₅₄H₈₈O₁₇	Caulosaponin (Leontin)	1008.7	255			
6163	C₅₆H₇₄N₄O₁₂S	Psychotrine sulfate..	1026 7	217			
6164	C₅₆H₈₈O₉	Caulophyllosapogenin	904.68	315			
6165	C₅₇H₁₀₄O₆	Glycerol trielaidate	884 80	32			
6166	C₅₇H₁₀₄O₆	Glycerol trioleate	884 80	−17	240¹⁰	0 915	
6167	C₅₇H₁₀₄O₉	Glycerol triricinoleate	932.80			0.959	
6168	C₅₇H₁₁₀N₂O₁₅	Pyosin... .	1062.9	238		0 862⁴⁰	1115
6169	C₅₇H₁₁₀O₆	Tristearin.	890 85	54.5; 70 8			
6170	C₅₈H₄₄O₂₃	Fustin...	1110 4	219			
6172	C₆₆H₁₀₄O₁₇	Caullophyllosaponin	1168.8	260			
6173	C₆₈H₉₆N₂O₂₆S	Aconitine sulfate	1388.8				1333
6175	C₇₂H₈₈N₆O₂₀	Quinine citrate	1356.7	183 5			

REFRACTIVE INDEX

A. LIQUIDS

Serial No.	Gen. index No.	Refractive index n_D^{20}	Dispersion $H_\beta - H_\alpha$
1	586	1 306	0 0045
2	60	1 329	0 0054
3	208	1 3416	0 0061
4	141	1 3419	0 0051
5	213	1 341	0 0060
6	168	1 3474	0 0058
7	793	1 3526	0 0061
8	514	1 3531	0 0058
9	1072	1 355	0 0062
10	1073	1 3564	0 0040
11	1049	1 3574	0 0056
12	794 1	1 3576	
13	794	1 3579	0 0062
14	448	1 3591	0 0068
15	451	1 3597	0 0063
16	489	1 3613	0 0079
17	262	1 361	0 0061
18	452	1 3619	0 0062
19	396	1 363	0 0070
20	447	1 3636	0 0067
21	233	1 3639	0 0062
22	395	1 3664	0 0060
23	1716	1 369	0 0064
24	1086	1 3695	0 0063
25	37	1 3714	0 0072
26	212	1 3719	0 0066
27	1715	1 372	0 0065
28	773	1 3723	0 0078
29	725	1 3727	0 0064
30	718	1 3730	0 0070
31	984	1 3758	0 0080
32	1713	1 376	0 0065
33	665	1 3767	0 0051
34	1714	1 377	0 0065
35	727	1 3771	0 0066
36	726	1 3779	0 0065
37	506	1 378	0 0065
38	1712	1 3783	0 0064
39	823	1 3786	0 0070
40	719	1 3791	0 0071
41	1741	1 3807	0 0066
42	1746	1 3819	0 0065
43	48	1 382	0 0080
44	1610	1 3821	
45	2387	1 3825	
46	116	1 3828	
47	667	1 383	
48	1015	1 384	
49	1019	1 3840	
50	717	1 3843	0 0071
51	1017	1 3841	0 0066
52	1020	1 3844	0 0067
53	1739	1 3849	
54	247	1 385	
55	2380	1 385	0 0091
56	1063	1 3851	
57	1026	1 3852	0 0003
58	1016	1 3858	0 0008
59	505	1 386	0 0086
60	749	1 386	
61	2382	1 3861	0 0064
62	1007	1 3862	0 0070
63	450	1 3808	0 0068
64	792	1 387	0 0067
65	824	1 387	0 0075
66	269	1 3874	
67	1064	1 3874	0 0071
68	1018	1 3879	0 0066
69	1001	1 3881	0 0131
70	1004	1 3882	0 0072
71	468	1 3886	0 0065
72	524	1 389	0 0074
73	1653	1 389	
74	1014	1 3891	0 0008
75	1006	1 3895	0 0071
76	154	1 3898	0 0084
77	2393	1 390	0 0068
78	809	1 3902	0 0060
79	1002	1 3902	0 0080
80	1655	1 3903	0 0070
81	626	1 3904	0 0069
82	972	1 3900	
83	1649	1.391	
84	242	1 392	
85	625	1.3927	0 0079
86	1005	1 3927	0 0070
87	2933	1 3929	0 0080
88	724	1 3930	0 0070
89	2392	1 393	0 0068
90	3369	1 393	0 0062
91	1654	1 3932	0 0068
92	1659	1 3945	
93	822	1 394	0 0074
94	2926	1 3947	0 0066
95	1651	1 3951	0 0068
96	1639	1 3959	0 0071
97	2362	1 3959	
98	747	1 3960	
99	790	1 396	0 0068
100	2354 1	1 3960	
101	598	1 3962	
102	686	1 3962	
103	2937	1 3964	
104	791	1 397	0 0068
105	495	1 3972	0 0081
106	1045 1	1 3973	
107	228	1 3974	0 0073
108	2359	1 3975	
109	723	1 3979	0 0070
110	718	1 398	
111	821	1 398	0 0074
112	2911	1 3980	0 0069
113	624	1 3984	0 0086
114	2910	1 398	0 0070
115	1640	1 399	
116	780	1 3993	0 0069
117	671	1 3996	
118	1652	1 3997	0 0068
119	356	1 3998	0 0127
120	2005	1 3999	
121	917	1 4001	0 0096
122	2354	1 4005	0 0069
123	2361	1 4005	0 0069
124	1636	1 4006	0 0071
125	3365	1 4008	
126	2357	1 4009	0 0070
127	1534	1 4010	0 0098
128	1617	1 401	0 0090
129	1764	1 401	0 0081
130	2353	1 4012	
131	820	1 401	0 0075
132	746	1 4015	0 0071
133	2901 1	1 4015	
134	2038	1 4016	
135	2912	1 402	0 0070
136	487	1.4022	
137	775	1 4026	0.0080
138	2935	1 4026	
139	2009 1	1 4030	
140	2904	1 4035	
141	2912	1 4036	
142	1560	1 4038	0 0071
143	3347	1 404	
144	3349	1 4040	
145	1013	1 4043	0 0071
146	937	1.4045	0 0085
147	3353	1 4047	
148	2903	1 4049	
149	1760	1 405	0 0075
150	1768	1.405	
151	3354	1 405	
152	378	1 4051	0 0080
153	1010	1 4051	0 0071
154	1084 1	1 4053	
155	1045	1 4056	0 0084
156	2936	1 4058	
157	1046	1 4060	
158	1081	1 406	0 0070
159	1803	1 406	0 0069
160	2275	1 406	0 0087
161	3330 1	1 4060	
162	3334	1 4060	0 0071
163	2901	1 4065	
164	3333 1	1 4070	
165	1084	1 4072	0 0070
166	1079	1.4075	0 0071
167	3331	1.4076	
168	652	1.4079	0.0080
169	1767	1.4079	0.0069
170	3351	1.406	0.0071
171	3995	1 408	0 0072
172	4007	1 408	0 0068
173	2344	1 4082	
174	3998	1 408	0 0072
175	1012	1.4086	0 0072
176	1100	1.4088	0 0074
177	420	1 4093	
178	2934	1 4095	
179	1080	1 410	0 0070
180	2985	1 410	0 0076
181	1044	1 4103	
182	1570	1 4104	0 0074
183	1730	1 411	
184	3329	1 4110	
185	3994	1 411	
186	2331	1 4114	
187	2910	1 4111	
188	1602	1 4115	
189	1000	1 4116	
190	657	1 4118	
191	1013	1 4119	0 0073
192	2326	1 412	0 0090
193	651	1 4121	0 0081
194	3335	1 4122	
195	3311	1 4123	0 0071
197	3999	1 4126	
198	3986	1 4127	0 0072
199	1619	1 4128	
200	1070	1 4129	0 0118
201	1645	1 4130	0 0073
202	2343	1 4131	0 0073
203	2846	1 4131	0 0073
204	446	1 4134	0 0094
205	1730 1	1 4135	
206	948	1.4136	0 0051
207	1643	1.4138	0 0074
208	2309	1 4138	0 0072
209	3338	1 414	
210	4001	1 414	
211	1726 1	1 4141	
212	587	1 4144	
213	3982	1 4145	
214	1733 1	1 4146	
215	2411	1 4149	
216	1571	1 415	
217	1644	1 4150	0 0073
219	2873	1 415	0 0090
220	3993	1 415	0 0075
221	3336	1 4153	0 0073
222	375	1 4154	0 0100
223	966	1.4156	0 0081
224	2396	1 4159	
225	2896	1 4161	0 0075
226	66	1.4164	0 0076
227	189	1 4166	0 0080
228	2397	1 4172	
229	3936	1 4174	0 0194
230	3372	1 4176	0 0084
231	1736	1 4178	
232	911	1 4179	0 0044
233	2944	1.4184	
234	4178	1.4184	
235	968	1 4185	0 0075
236	969	1 4185	0 0105
237	479	1 4186	0 0101
238	1695	1 4194	0 0073
239	2302	1 419	
240	2320	1 4195	
241	943	1 4196	0 0091
242	1734	1.4196	0 0071
243	1561	1 4198	0 0081
244	1662	1.4198	0 0081
245	1732	1 420	
246	2847	1 420	0 0071
247	2955	1 4201	
248	2970	1 4203	
249	2971	1 4204	
250	3989	1 4204	0 0074
251	2400.1	1 4206	
252	896	1.4207	0 0087
253	2407 1	1 4209	
254	2954	1.4209	
255	4411	1.4209	
256	2399	1 421	0 0073
257	2869	1.421	0.0076
258	3088	1 421	
259	2407	1.4213	
260	569	1.4216	
261	2892	1.4217	0 0113
262	1067.1	1.4219	0 0071
263	2301	1 4223	0 0076
264	358	1 4224	
265	2100	1 4226	0 0070
266	2405	1 4226	
267	658	1 4227	0 0075
268	4412	1 4228	
269	2351	1.423	0 0075
270	2409	1 423	0 0052
271	3357	1 423	
272	2330	1 4235	0 0075
273	28	1 4237	0 0080
274	2965	1 4238	
275	220	1 4239	0 0003
276	711	1 4240	
277	999	1 4240	
278	2419	1 424	0 0078
279	2967	1 424	
280	3325	1 424	0 0193
281	4012	1 424	0 0078
282	4161	1.424	0 0073
283	1557	1 4242	0 0106
284	3308	1 4242	
285	657 1	1 4247	
286	3309	1 4248	
287	2403	1 425	0 0074
288	2868	1 425	
289	465	1 4251	0 0093
290	616	1 4254	0 0071
291	2406	1 4254	
292	2987	1 4254	
293	3314	1 4259	
294	4419	1 426	0 0081
295	928	1 4263	
296	2899	1 4268	0 0076
297	2962	1 427	0 0073
298	2963	1 4270	
299	4585	1 427	0 0075
300	4586	1.427	0 0074
301	949	1 4271	
302	3962	1.4271	
303	721	1 4272	
304	1612	1 4273	0 0075
305	264	1 4274	0 0072
306	3939	1 4275	
307	3975	1.4275	
308	2964	1 4278	
309	744	1 428	0 0095
310	3310	1 4284	
311	2386 1	1 4288	
312	4172	1.4280	0 0077
313	1027	1 429	
314	4162	1 4293	
315	449	1 4295	
316	4153	1 4299	0 0075
317	2867	1 430	0 0076
318	2966	1 430	0 0075
319	2086	1 430	0 0076
320	3356	1.430	0 0074
321	1629	1 4302	
322	2953	1.4303	
323	273	1 4306	0 0102
324	355	1 4306	0 0094
325	925	1 4306	0 0094
326	3289	1 4306	
327	3937	1.4309	0 0077
329	3361	1 4310	
330	3363	1 431	
331	3940	1 4311	
332	4843	1 4312	
333	620	1.4314	0 0114
334	2412	1 4314	0 0073
335	3355	1 4317	
336	736	1 432	
337	4852	1 4321	
338	3358	1 4322	0.0076
339	2952	1 433	0 0073
340	3328	1 4330	0 0076
341	4166	1 4333	
342	4169	1 4334	
343	712	1.4335	

Serial No	Gen. index No.	Refractive index n_β	Dispersion $H_\beta - H_\alpha$
344	3364	1.4338	
345	2318	1.434	
346	464	1.4341	0.0092
347	743	1.4344	
348	3362	1.4345	
349	192	1.4346	0.0089
350	158	1.4349	
351	5010	1.4350	
352	742	1.436	0.0092
353	924	1.436	0.0080
354	471	1.4362	
355	2849 1	1.4362	
356	258	1.4364	0.0126
357	2968	1.437	0.0074
358	3961	1.437	0.0078
359	5260	1.437	0.0076
360	3303	1.4371	
361	614	1.4373	0.0149
362	1253	1.4375	0.0126
363	3895	1.4376	
364	17	1.438	
365	762	1.438	0.0096
366	3944	1.4380	
367	604	1.4386	0.0082
368	811	1.4386	0.0097
369	3285	1.4388	0.0092
370	927	1.4390	0.0131
371	470	1.4392	
372	741	1.4398	0.0089
373	1506	1.4404	
374	4179	1.4404	
375	2813	1.4407	0.0098
376	1089	1.441	
377	2812	1.4410	0.0112
378	1041	1.4412	0.0083
379	1098	1.4412	0.0091
380	1366	1.4413	0.0122
381	457	1.4414	0.0077
382	1500	1.4415	0.0103
383	941	1.4416	
384	1252	1.4417	0.0131
385	2281	1.4419	
386	655	1.442	0.0084
387	3960	1.4420	
388	5156	1.442	0.0084
389	1042	1.4421	
390	814	1.4425	0.0099
391	1576	1.4425	
392	5888	1.4427	
393	764	1.4428	0.0098
394	2284	1.443	
395	648	1.4433	
396	1096	1.4437	
397	2825	1.4438	
398	2827	1.444	
399	3295	1.4441	
400	190	1.4443	0.0084
401	1040	1.4444	0.0089
402	4387	1.4445	
403	1056	1.4450	0.0094
404	1537	1.4451	0.0095
405	2327	1.4452	
406	2835	1.4453	
407	1055	1.4454	0.0094
408	2283	1.4454	
409	4381	1.4455	0.0083
410	3968	1.4456	
411	619	1.4457	0.0129
412	4856	1.4459	
413	1769	1.446	
414	4376	1.4460	
415	148	1.4462	
416	1699	1.4464	0.0120
417	19	1.4467	0.0080
418	4388	1.4468	
419	963	1.4469	
420	3827	1.4471	
421	2850	1.4476	
422	1692	1.4478	0.0088
423	3892	1.4481	0.0092
424	921.1	1.4482	0.0083
425	5940	1.4482	
426	2831	1.4486	0.0082
427	5013	1.4486	
428	1551	1.449	0.0169
429	1724	1.4490	
430	926	1.4495	
431	2282	1.4496	
432	1090	1.4499	
433	307	1.4500	0.0090
434	2890	1.4503	
435	3808	1.4505	
436	618 3	1.4506	
437	585	1.4507	0.0087
438	648 2	1.451	0.0092
439	929	1.4512	0.0176
440	3826	1.4515	
441	3917	1.4521	
442	2294	1.4524	0.0121
443	4010	1.4524	0.0095
444	1054	1.4530	0.0089
445	3805	1.4532	
446	285	1.4539	0.0055
447	2888	1.4540	
448	3893	1.4540	
449	5853	1.4543	
450	648 1	1.4550	
451	1595	1.455	0.0084
452	364	1.4554	
453	4144	1.4556	
454	1308 4	1.4556	0.0107
455	107	1.4557	0.0094
456	5356	1.4557	
457	5813	1.4558	
458	222	1.4562	0.0110
459	3889	1.4567	
460	648 4	1.4570	
461	696	1.457	
462	3933	1.457	0.0081
463	2889	1.4574	
464	3069	1.4579	
465	5482	1.4580	
466	2340	1.4580	
467	2311	1.4590	
468	2886	1.459	0.0082
469	2383	1.4594	
470	11	1.4595	0.0079
471	1478	1.4597	
472	5371	1.4602	0.0084
473	3971 1	1.4603	
474	3902	1.4608	
475	3992	1.4606	
476	12	1.4607	0.0097
477	3804	1.4609	
478	2439	1.461	
479	3296	1.4614	
480	3915	1.4623	
481	5605	1.4626	
482	1105	1.463	0.0088
483	4372	1.4630	
484	5606	1.4636	
485	3917	1.4642	
486	3273	1.4643	
487	1328	1.4646	0.0145
488	3918	1.4649	
489	366	1.4655	0.0132
490	136	1.4659	
491	4148	1.4659	
492	2814	1.4660	0.0151
493	4374	1.4660	
494	103	1.4666	0.0107
495	1756	1.467	
496	2882	1.467	0.0084
497	2796	1.4675	
498	2240	1.4680	
499	3854	1.4690	
500	2058	1.4691	0.0144
501	176	1.4697	0.0112
502	2059	1.470	0.0112
503	3811	1.4700	
504	3891	1.4701	0.0153
505	2057	1.4704	0.0094
506	159	1.4711	
507	3858	1.4715	0.0111
508	863	1.4717	
509	3913 1	1.4723	
510	3810	1.4727	0.0114
511	3952	1.4727	
512	515	1.4729	0.0078
513	3913	1.4729	
514	4115	1.473	
515	3806	1.473	0.0118
516	4371	1.4739	
517	3879	1.4741	
518	3864	1.4743	
519	1466	1.4744	0.0131
520	2745*	1.4745	
521	3924	1.4747	0.0105
522	368	1.4753	0.0094
523	582	1.4755	
524	2239	1.4763	
525	106	1.4777	
526	3927	1.4785	
527	3816	1.4788	
528	139	1.479	
529	2797	1.4792	0.0116
530	1370	1.4792	
531	3908	1.4798	
532	422	1.4801	0.0110
533	3026	1.4803	
534	887	1.4805	
535	5164	1.4806	0.0102
536	3682	1.482	
537	3624	1.4823	
538	3922	1.4827	0.0096
539	3800	1.4828	
540	5480	1.484	
541	3823	1.4816	
542	3761	1.4848	
543	1506	1.4867	
544	3865	1.4870	0.0147
545	1131	1.4872	0.0140
546	3860	1.488	
547	3886	1.488	
548	5001	1.4881	
549	2927	1.489	0.0120
550	3725	1.4890	
551	3765	1.4895	
552	2262	1.4903	0.0132
553	3857	1.4911	
554	3724	1.4914	
555	221	1.4915	
556	3229	1.4920	0.0147
557	4097	1.4922	
558	4344	1.4922	
559	3728	1.4925	0.0114
560	1697	1.4920	0.0125
561	3223	1.4930	0.0146
562	3736	1.493	0.0140
563	4097 1	1.4930	
564	3882	1.4935	
565	4367	1.4939	
566	4312	1.494	
567	110	1.4912	
568	3226	1.4943	0.0160
569	4980	1.4946	
569 1	3731	1.495	0.0144
570	5078	1.4951	
571	4098	1.4954	0.0133
572	1051	1.4955	0.0131
573	2088	1.4956	0.0158
574	4983	1.4956	
575	1588	1.4959	0.0104
577	2083	1.4959	0.0152
578	755	1.4960	0.0137
579	2112	1.4962	0.0160
580	3228	1.4967	0.0113
581	3856	1.4967	
582	3726	1.4969	
583	4366	1.4972	
584	2685	1.4973	0.0158
585	3225	1.4975	0.0152
586	3780	1.4976	
588	600	1.498	0.0117
589	3677	1.498	0.0137
590	4075	1.4981	
591	4978	1.4984	
593	3741	1.4986	
594	3286	1.4993	0.0116
595	3681.1	1.4995	
596	4974	1.4996	
597	476	1.4997	0.0056
598	3277	1.4998	0.0213
599	3152	1.500	
600	754	1.5001	0.0140
601	3727	1.5003	0.0146
602	4977	1.5005	
603	4976	1.5007	
604	1443	1.500	0.0160
605	4561	1.5011	
606	1365	1.5014	0.0167
607	4324	1.5019	0.0117
608	2810	1.5023	0.0245
609	4828	1.5025	
610	2590	1.503	0.0149
611	4999	1.5030	
612	589	1.5035	0.0151
613	3883	1.5036	
614	111	1.504	0.0104
615	3224	1.5042	0.0153
616	3761	1.5042	
617	4081	1.5042	
618	666	1.5046	
619	2710	1.505	0.0150
620	3763	1.5050	
621	475	1.5051	0.0148
622	3230	1.5051	0.0158
623	90	1.5055	0.0137
624	3679	1.5057	0.0163
625	4971	1.5057	
626	2684	1.5058	0.0161
627	2720	1.506	0.0161
628	3154	1.506	0.0161
629	3678	1.506	0.0162
630	815	1.5063	0.0130
631	4972	1.5065	
632	689	1.507	
633	2722	1.507	0.0164
634	4350	1.507	
635	4318	1.508	
636	3080	1.5081	0.0169
637	1827	1.5083	0.0140
638	4515	1.5085	
639	603	1.509	0.0127
640	2586	1.509	0.0188
641	870	1.509	0.0163
643	2775	1.5105	
644	244	1.512	0.0163
645	331	1.512	
646	2721	1.512	0.0160
647	183	1.5128	0.0132
648	3244	1.513	0.0171
649	3786	1.5131	0.0163
650	3227	1.5132	0.0157
651	404	1.5134	0.0168
652	1330	1.514	0.0169
653	4102	1.514	
654	3119	1.5143	
655	5141	1.516	0.0143
656	2589	1.5164	0.0132
657	5000	1.5164	
658	3754 2	1.5168	
659	2163	1.517	0.0173
660	3235 1	1.5174	
661	4091 1	1.5175	
662	3740	1.5187	0.0157
663	3788	1.5201	0.0117
664	412	1.5203	0.0131
665	4318	1.5207	
666	2041	1.521	0.0164
667	4500	1.521	
668	2713 1	1.5211	
669	3755	1.5218	
670	3170	1.5226	0.0206
671	413	1.523	0.0124
672	2040	1.523	0.0165
673	3149	1.5232	
674	3757	1.5234	
675	3096	1.523	
676	3655	1.5236	
677	2714	1.5240	
678	3752	1.524	0.0157
679	2503	1.5242	
680	3088	1.5249	0.0196
681	1307	1.525	0.0172
682	3258	1.525	
683	1090 1	1.5250	
684	3057	1.526	
685	859	1.5261	0.0270
686	2111	1.5261	0.0198
687	594	1.5206	0.0173
688	1250	1.5267	0.0232
689	3132	1.527	0.0183
690	3664	1.5271	0.0189
691	2039	1.528	0.0166
692	3034	1.5282	
693	576	1.5285	0.0173
694	4353	1.5285	
695	3747	1.5286	0.0160
696	45	1.5297	0.0221
697	2	1.5300	0.0117
698	3656	1.5301	0.0204
699	3265	1.5305	
700	1249	1.5311	0.0232
701	699	1.5313	0.0122
702	2084	1.5323	
703	3134	1.5335	
704	4064	1.5338	0.0202
705	2571	1.5339	0.0217

Serial No	Gen. index No	Refractive index n_β'	Dispersion $H_\beta - H_\alpha$	Serial No	Gen index No	Refractive index n_D	Dispersion $H_\beta - H_\alpha$	Serial No	Gen. index No	Refractive index n_D	Dispersion $H_\beta - H_\alpha$	Serial No.	Gen. index No.	Refractive index n_D	Dispersion $H_\beta - H_\alpha$
706	3237	1.5457	0.0168	731	1229	1.549	0.0176	756	2757	1.570	0.0217	781	102	1.6062	
707	1890	1.546	0.0216	732	2032	1.5490		757	2203	1.5714	0.0249	782	601	1.6077	
708	2618	1.5369	0.0222	733	3259	1.5492	0.0229	758	2204	1.5728	0.0230	783	1205	1.608	0.0217
709	2725	1.547	0.0180	734	2031	1.551		759	2004	1.5735	0.0315	784	1483	1.6081	0.0256
710	184	1.5379	0.0140	735	2639	1.551	0.0189	760	3642	1.5749		785	2061	1.609	0.0234
711	2038	1.549	0.0175	736	1347	1.5529	0.0252	761	2771	1.575	0.0162	786	2492	1.6094	
712	3606	1.5491	0.0210	737	1859	1.5547	0.0221	762	4930	1.576		787	1204	1.611	
713	2159	1.5309	0.0173	738	2030	1.555		763	4757	1.5761		788	3548	1.6149	0.0296
714	2161	1.540	0.0181	739	2763	1.5559	0.0225	764	1200 2	1.577		789	3549	1.616	0.0296
715	2162	1.540	0.0184	740	2633	1.556	0.0182	765	1200 1	1.5814		790	4038	1.618	0.0303
716	1388	1.5407	0.0213	741	1441	1.5562	0.0375	766	2255	1.583	0.0248	791	3069	1.6195	0.0424
716 1	1944	1.541		742	2762	1.558	0.0214	767	372	1.584		792	1333	1.621	0.0253
717	3789	1.5421	0.0220	743	964	1.559		768	1887	1.5861	0.0286	793	1369	1.6260	0.0265
718	2677	1.5125		744	2758	1.559	0.0217	769	1442	1.5863	0.0249	794	127	1.6277	0.0189
719	123	1.5137	0.0165	745	2578	1.5597	0.0270	770	2491	1.588		795	3455	1.633	0.0309
720	2195	1.5440	0.0175	746	1062	1.5598	0.0283	771	2756	1.5887	0.0248	796	128	1.638	0.0183
721	10	1.5442	0.0219	747	1291	1.590	0.0193	772	18	1.589	0.0176	797	428	1.642	
722	1389	1.5455	0.0202	748	2760	1.561	0.0214	773	151	1.5890	0.0162	798	1918	1.6509	0.0349
723	1230	1.516	0.0178	749	2098	1.5620	0.0227	774	1375	1.5895	0.0240	799	3453	1.658	0.0325
724	2081	1.5102		750	2767	1.5649	0.0230	775	4723	1.5921	0.0195	800	4263	1.6913	0.0356
725	2001	1.5464	0.0232	751	1857	1.5650	0.0209	776	1376	1.5931	0.0243				
726	3260	1.5469		752	619	1.567		777	1202	1.5979	0.0161				
727	2160	1.547	0.0185	753	1856	1.567	0.0230	778	101	1.5992	0.0193				
728	236	1.5472	0.0204	754	1176	1.5671	0.0207	779	4296	1.602	0.0290				
729	2082	1.5475		755	2123	1.5692	0.0214	780	126	1.603	0.0162				
730	3787	1.5481	0.0224												

Serial No	Gen index No	Temperature $t°C$	Refractive index n_D	Dispersion $H_\beta - H_\alpha$	Serial No	Gen index No	Temperature $t°C$	Refractive index n_D	Dispersion $H_\beta - H_\alpha$	Serial No	Gen index No.	Temperature $t°C$	Refractive index n_D	Dispersion $H_\beta - H_\alpha$
801	683	0	1.3752		857	1572	15	1.4644	0.0090	912	3955	17	1.4385	0.0090
802	310	0	1.4538	0.0058	858	4147	15	1.4708		913	568	17	1.4467	
803	209	7	1.3597		859	3912	15	1.4801		914	3819	17	1.4674	0.0109
804	1327	7	1.6053		860	3863	15	1.4849		915	3821	17	1.4784	
805	930	7.5	1.4341	0.0094	861	3859	15	1.4871	0.0130	916	4993	17	1.5332	
806	3054	8 2	1.571	0.0234	862	4979	15	1.4921		917	3649	17	1.5671	
807	969 1	8 4	1.417		863	117	15	1.4982	0.0233	918	4404	17.1	1.4435	0.0072
808	4339	9 5	1.5301	0.0171	864	118	15	1.4998	0.0227	919	3820	17 1	1.4774	0.0116
809	22	10	1.2675	0.0052	865	4986	15	1.5018		920	3849	17.1	1.4895	0.0157
810	4304	10 8	1.6265	0.0337	866	988	15	1.5094	0.0071	921	982	17.2	1.3817	0.0085
811	807	11	1.4198	0.0077	867	100	15	1.5219	0.0148	922	2267	17 2	1.4511	0.0111
812	3591	11	1.5425	0.0188	868	3589	15	1.5632		923	3928	17 2	1.4638	0.0085
813	2832	11 9	1.4519	0.0084	869	3590	15	1.5736		924	339	17 4	1.5337	
814	2570	11 9	1.5503	0.0229	870	29	15	1.7425		925	340	17.4	1.5369	
815	2276	12	1.4468		871	4306	15.1	1.6477	0.0404	926	2830	17.5	1.4771	0.0104
816	2337	12	1.467		872	558	15 2	1.4735	0.0103	927	609	17 6	1.4588	0.0157
817	4323	12	1.5703	0.0253	873	359	15 3	1.4302		928	3245	17 6	1.5058	0.0157
818	2824	12 5	1.1208	0.0080	874	1541	15 3	1.4526	0.0084	929	5359	17 7	1.463	0.0092
819	1535	12 5	1.4559	0.0167	875	525	15 4	1.3770	0.0071	930	3638	17 8	1.4804	0.0085
820	2453	12 5	1.5524	0.0338	876	1546	15 4	1.4213	0.0092	931	3637	17.8	1.5451	0.0169
821	2580	12 7	1.5764	0.0298	877	3128	15.5	1.5647		932	920	18	1.4079	
822	89	12 9	1.4340	0.0101	878	3122	15 7	1.5717	0.0236	933	1000	18	1.4282	0.0094
823	1078	13	1.411		879	3061	15.8	1.5196	0.0274	934	4375 1	18	1.4565	
824	3818	13	1.479		880	983	16	1.378		935	3125	18	1.5441	0.0180
825	3851	13	1.4971	0.0135	881	1613	16	1.4013	0.0090	936	3667	18	1.5680	0.0251
826	5	13	1.5831		882	942	16	1.4083	0.0076	937	4813	18	1.5933	0.0280
827	3861	13 6	1.4540	0.0083	883	737	16	1.4156		938	545	18 1	1.5004	0.0168
828	608	13 7	1.4786	0.0128	884	3874	16	1.438		939	1022	18 2	1.4513	
829	1518	13 7	1.4993	0.0141	885	1555	16	1.4506	0.0123	940	3753	18 2	1.4999	0.0136
831	4041	13 9	1.6232	0.0312	886	3304	16	1.452		941	3037	18 2	1.6283	0.0312
832	2880	14	1.458		887	2884	16	1.455		942	1568	18 3	1.4198	
833	2342	14	1.462		888	2883	16	1.458		943	916	18 3	1.4221	0.0148
834	2878	14	1.463		889	2887	16	1.458		944	400	18 4	1.4058	0.0070
835	3812	14	1.4883	0.0172	890	3923	16	1.4762		945	2855	18 4	1.4607	0.0090
836	2579	14	1.5566	0.0248	891	5003	16	1.480		946	2818	18 4	1.4904	0.0124
837	4707	14	1.610		892	908	16	1.4888	0.0149	947	1341	18 5	1.5389	0.0211
838	2336	14 4	1.4397	0.0092	893	3651	16	1.5514		948	4260	18 5	1.635	
839	3852	14 5	1.4647	0.0084	894	84	16	1.580		949	935	18 8	1.4357	0.0096
840	3919	14 5	1.4787		895	379	16 1	1.4397	0.0079	950	773 1	18 9	1.4200	
841	3666	14 5	1.5439	0.0189	896	2279	16 3	1.4554	0.0150	951	4560	18 9	1.5198	0.0195
842	2280 1	14 6	1.4505	0.0083	897	3847	16 3	1.4846	0.0126	952	170	19	1.4117	
843	979	14 7	1.4098	0.0071	898	608 1	16 3	1.4971	0.0133	953	1554	19	1.4375	
844	3574	14 7	1.5740	0.0222	899	1548	16 4	1.4458	0.0136	954	2929	19	1.4435	0.0087
845	3762	14 8	1.5104	0.0201	900	4279	16 4	1.6157	0.0296	955	3807	19	1.4724	
846	4967	14 8	1.5128	0.0153	901	918	16.5	1.4402		956	3850	19	1.4900	
847	3283	14 9	1.4463	0.0103	902	3324	16.5	1.4632	0.0090	957	4987	19	1.4992	
848	1616	15	1.4065	0.0090	903	880	16 6	1.4470	0.0129	958	4088	19	1.5092	
849	622	15	1.4257		904	934	16 6	1.4527	0.0127	959	4994	19	1.5289	0.0111
850	713	15	1.4413		905	2816	16 6	1.4561	0.0104	960	2568	19	1.5485	0.0227
851	4004	15	1.4372		906	2570	16 6	1.5469	0.0230	961	4150	19.1	1.4714	0.0134
852	1533	15	1.4421		907	2538	16.6	1.5485	0.0240	962	4023	19 3	1.6546	0.0409
853	132	15	1.4490	0.0116	908	4587	16 8	1.4419		963	2298	19 5	1.4310	0.0102
854	133	15	1.4519	0.0101	909	1519	16 8	1.5077	0.0147	964	2299	19.5	1.4355	0.0105
855	5007	15	1.4628		910	2328	16 9	1.425	0.0076	965	3959	21	1.447	
856	4834	15	1.4638		911	313	17	1.3870	0.0104	966	3639	21	1.5390	

Serial No	Gen. index No.	Temperature t°C	Refractive index n_D	Dispersion H_β - H_α
967	4998	21.3	1.4979	
968	2759	21.3	1.5591	
969	4307	21.3	1.6544	0 0408
970	3121	21.4	1.5370	0 0168
971	2569	21.4	1.5407	0 0223
972	2071	21.4	1 5637	0 0247
973	3600	21 4	1 5766	0 0311
974	1496	21 6	1 4351	0 0114
975	2859	21 6	1.4766	0 0089
976	4789	21 6	1 5743	0 0193
977	4814	21 6	1 6321	0.0400
978	2028	21.9	1 4512	
979	3297	22	1 4380	
980	5765.1	22	1 4538	
981	3916	22	1 4604	
982	3822	22	1.4754	
983	3815	22	1 4770	0 0085
984	3813	22	1 4959	
985	5005	22 2	1 4600	0 0081
986	3703	22 2	1 5604	
987	301	22.3	1 4075	0 0093
988	4559	22 3	1 4984	0 0140
989	2205	22 4	1 5711	
990	2199	22 5	1 5021	
991	1357	22 5	1 5642	0 0242
992	2493	22 5	1 5990	
993	3958	22 6	1 4484	
994	4373	22.6	1 4623	0 0083
995	46	22 7	1 4453	0 0113
996	893	22.7	1.4852	0 0166
997	2468	22 7	1 5645	0 0231
998	2134	22.7	1 5760	
999	3601	22 9	1 5494	0 0268
1000	2384	23	1 4531	
1001	4563	23	1.5300	0 0264
1002	1430	23	1 5861	0 0231
1003	3547	23	1 6141	0 0298
1004	2505	23 1	1 5272	
1005	3701	23 1	1 5802	0 0244
1006	3702	23 1	1 5898	
1007	886	23 2	1 4365	0 0147
1008	1628	23 3	1.4329	0 0094
1009	314	23 4	1 4597	0 0102
1010	4375	23 4	1 4619	0 0082
1011	4156	23 4	1 4624	
1012	3191	23 4	1 5798	
1013	3192	23.4	1 5933	0 0302
1014	4448	23 4	1 6060	0 0278
1015	561	23 5	1 5231	0 0170
1016	1700	23 6	1 4464	
1017	1482	23 6	1 4992	0 0175
1018	1444	24	1 5043	
1019	4241	24	1 5826	
1020	1701	24 3	1 4463	
1021	2289 3	24 4	1 4432	0 0083
1022	3728 1	24 5	1 4877	0 0139
1023	4385	25	1 4555	0 0080
1024	5875	25	1 1875	
1025	3687	25	1 5252	
1026	3036	25 1	1.6223	0 0302
1027	2289 2	25 2	1 4431	0 0082
1028	1885	25 5	1 5257	0 0191
1029	2338	26	1 4558	
1030	4490	26	1 575	0 0205
1031	4226	26	1 6644	

Serial No	Gen index No	Temperature t°C	Refractive index n_D	Dispersion H_β - H_α
1032	300	26 1	1 4540	0 0005
1033	994	26 4	1 4951	0 0147
1034	1587	26 8	1 4877	
1035	816	27 5	1 4769	0 0140
1036	5603	30	1 4559	0 0126
1037	3804	30	1 474	
1038	3981	31	1 4308	
1039	3126	33	1 5758	0 0295
1040	2293	33 8	1 4561	0 0082
1041	5380	33 9	1 4358	0 0077
1042	316	34 2	1 4146	
1043	5381	34 3	1 4147	0 0076
1044	3648	34 4	1 5537	0 0249
1045	5186	34 6	1 490	0 0076
1046	5852	35	1 4587	
1047	5391	35 2	1 4319	0 0075
1048	4530 3	35 2	1 5526	0 0292
1049	2490	36	1 6332	0 0293
1050	1011	36 5	1 3931	0 0070
1051	1627	37	1 4606	0 0078
1052	177	37 2	1 5258	0 0181
1053	2096	38 6	1 5703	
1054	6056	40	1 4446	
1055	1553	40	1 4467	0 0118
1056	3272	40	1 4511	0 0150
1057	5360	40	1 4533	
1058	1314	40	1 5473	
1059	1315	40	1 5565	
1060	1316	40	1 5579	
1061	4060 1	40	1 5726	0 0327
1062	4039	40	1 6026	0 0289
1063	860	40 3	1 5238	
1064	1413	41	1 5125	0 0189
1065	5610	42 9	1 434	0 0075
1066	4174	45 2	1 4294	0 0076
1067	5694	45 3	1 4346	0 0076
1068	3587	46	1 5836	
1069	931	46 7	1 4434	0 0123
1070	239	47	1 415	0 0098
1071	4297	47 3	1 5932	0 0281
1072	993	48	1 4126	0 0079
1073	30	48	1 4418	0 0085
1074	3802	48	1 4621	
1075	2464	48	1 6231	0 0343
1076	3412	48 5	1 6338	0 0305
1077	56	48 6	1 4616	0 0149
1078	5876	50	1 4663	
1079	5805	50	1 4689	
1080	3550	51 2	1 6703	0 0424
1081	4305	53 2	1 6443	0 0439
1082	4447	53 5	1 5975	0 0268
1083	1331	56	1 5010	0 0173
1084	1251	56	1 5150	0 0225
1085	5763	57 1	1 448	0 0084
1086	1480	57 7	1 6339	0 0305
1087	2206	59 1	1 5532	
1088	4851	60	1 4308	
1089	6147	60	1 4429	
1090	2263	60	1 4787	0 0228
1091	563	61	1 4953	
1092	1858	61	1 5553	0 0246
1093	1961	61 5	1 5557	
1094	1962	61 5	1 5577	
1095	1963	61 5	1 5647	
1096	2083	62 5	1 5346	

Serial No	Gen index No	Temperature t°C	Refractive index n_D	Dispersion H_β - H_α
1097	560	63 1	1 4165	
1098	288	63 9	1 4152	
1099	156	65	1 4297	
1100	3071	66	1 5377	0 0169
1101	1231	69 9	1.5266	0.0171
1102	3456	70 7	1 6079	0.0295
1103	2172	74	1 5425	0 0187
1104	3414	76	1 6228	0 0303
1105	4219	77 1	1 588	0.0205
1106	3503	77 8	1 5678	0 0375
1107	238	78 3	1 4274	0 0098
1108	5168	78 9	1 4283	0.0075
1109	2356	79	1 3732	0.0064
1110	6018	79 1	1 4331	0 0077
1111	5814	79 5	1 4283	0 0076
1112	617	79 7	1 4228	0 0126
1113	5150	79 8	1 4273	0 0075
1114	6157	80	1 4381	
1115	6169	80	1 4300	
1116	3801	80	1 4402	0.0089
1117	5379	80 2	1 4299	0 0076
1118	4756	80 6	1 539	0 0187
1119	525N	80 7	1 4175	0 0073
1120	5816	80 8	1 4236	0 0075
1121	936	81	1 4342	0.0123
1122	631	82 1	1 379	0.0067
1123	4406	82 1	1 4183	0.0074
1124	2386	83 9	1 421	0 0083
1125	6026	93 5	1 4297	0.0076
1126	3507	98 7	1 0206	0.0324
1127	4218	98 8	1 6048	0.0293
1128	5402	90	1 5839	0.0219
1129	2548	99 2	1 5522	0 0242
1130	5063	99 2	1 6762	0 0556
1131	921.2	99 3	1 4657	0 0121
1132	1206	99 3	1 5743	0.0204
1133	4024	99 4	1 6211	0.0387
1134	1807	99 4	1 6803	0 0541
1135	3584	99 4	1 6828	
1136	4800	99 4	1.6959	0 0591
1137	3583	90 4	1 7083	0.0515
1138	3291	99 5	1 4760	0 0094
1139	5223	99 5	1 5021	0 0133
1140	4640	99 5	1.6959	0 0561
1141	2810	99.6	1 4621	0.0094
1142	5224	99 6	1 5022	0.0134
1143	3494	99 6	1.5827	0.0287
1144	6145	100	1 4347	
1145	6144	100	1.4366	
1146	2864	100	1 4811	0.0085
1147	4947	100	1 5080	0.0060
1148	3144	100	1.5345	0 0177
1149	3417	100	1 6092	0.0291
1150	3418	100	1 6235	0.0313
1151	1480	100 4	1.4188	0.0065
1152	4119	107 2	1 480	0.0145
1153	482	107 8	1 4161	0.0090
1154	3282 1	109 4	1.4482	0.0085
1155	3307	110 6	1.4303	0.0077
1156	782	113	1.446	0.0097
1157	2585	114 6	1 512	0.0187
1158	4652	129	1 6567	
1159	5340	130 4	1.480	0.0133
1160	2007	131 9	1 504	0 0191
1161	3938	133 3	1 422	0.0073

B. SOLIDS

I. Mean Values

Serial No	Gen. index No	Refractive index n_β	Serial No	Gen index No	Refractive index n_β	Serial No	Gen. index No	Refractive index n_β	Serial No	Gen. index No.	Refractive index n_β
1162	481	1 4156	1164	1578 1	1 54	1165	5664	1 635	1166	414	1.755
1163	1070 1	1 525									

II. Uniaxial Group

Serial No.	Gen. index No	ω	ε	Serial No	Gen. index No	ω	ε	Serial No	Gen. index No	ω	ε	Serial No	Gen. index No	ω	ε
1167	55	1 484	1 602	1173	238*	1 51	1 46	1179	2174	1 569	1 666	1184	1416	1 633	1 626
1168	3973	1 497	1 476	1174	808	1 514	1 521	1180	6075	1 579	1 540	1185	2454	1 646	1 642
1169	535	1 499	1 49	1175	5002	1 545	1 548	1181	4043.1	1 581	1 493	1186	4672	1 6588	1.6784
1170	3756	1 525	1 609	1176	5142.1	1 545	1 548	1182	1769 1	1 590	1 650	1187	1625	1.700	1 640
1171	2373	1 529	1 513	1177	697 1	1 554	1 515	1183	4272	1 600	1.649	1188	4727	1 717	1 563
1172	2915	1 530	1 430	1178	1093	1 550	1 548					1189	21	1 800	1 750

* Stable modification

III. Biaxial Group

Serial No	Gen index No	α	β	γ	Serial No	Gen index No	α	β	γ	Serial No	Gen index No	α	β	γ
1190	679 1	1 367	1 409	1 536	1235	1688	1 545	1 546	1 837	1280	4330 1	1 564	1 6.8	
1191	361	1 4162	1 4603	1 5592	1236	786		1 547		1281	4752	1 621	1 629	1 661
1192	4184	1 402	1 463	1 617	1237	1530 1		1 548		1282	4943	1 590	1 639	1 640
1193	4218	1 407	1 468	1 620	1238	2916 1		1 550		1283	5317	1 620	1 6:0	1 650
1194	147	1.440	1 475	1 625	1239	853 1	1 459	1 555	1 582	1284	306		1 653	
1195	4397*		1 478		1240	988 1	1 546	1 559		1285	788		1 635	
1196	4368 3‡	1 471	1 479	1 519	1241	778	1 519	1 561	1 591	1286	5317*	1 543	1 636	1 684
1197	2920		1 484		1242	4396	1 5376	1 5651	1 5705	1287	3585		1 657	
1198	238†	1 370	1 485	1 585	1243	1042	1 551	1 567	1 571	1288	5319	1 607	1 642	1 675
1199	5066 1		1 488		1244	3961		1 570		1289	5067 1	1 621	1 613	1 618
1200	2234 1		1 496		1245	1472	1 56	1 57	1 60	1290	3087	1 505	1 645	1 655
1201	4368 3‡	1 479	1 496	1 524	1246	3716	1 54	1 571	1 59	1291	4750	1 587	1 616	1 769
1202	1507*	1 494	1 498	1 509	1247	5313 1	1 544	1 572		1292	1111 1	1 626	1 616	1 712
1203	2608 1	1 487	1 499	1 566	1248	1044	1 555	1 573	1 577	1293	5082 4	1 612	1 647	1 662
1204	2260 1	1.488	1 501	1 527	1249	193 1	1 515	1 575	1 586	1294	5213 1		1 650	
1205	776		1 503		1250	3199	1 560	1 576	1 617	1295	5304	1.463	1 653	1 780
1206	270	1.445	1 505	1 540	1251	5177	1 510	1 578	1 618	1296	4748	1.621	1 654	1 691
1207	666		1 509		1252	3778	1 5545	1 5787	1 5912	1297	1985	1.412	1 662	1 756
1208	994 1		1 510	1 607	1253	835	1 55	1 581		1298	5561	1.580	1 665	1 690
1209	3742		1 512		1254	708	1 549	1 583	1 625	1299	1749	1.586	1 668	1 680
1210	4008	1 505	1 512	1 524	1255	3194	1 556	1 587	1 700	1300	1987	1 479	1 669	1 734
1211	5028 1	1 511	1 512	1 836	1256	3111	1 535	1 592	1 760	1301	5428 1	1 529	1 670	1 716
1212	2260 2	1 495	1 513	1 672	1257	5228	1 522	1 594	1 616	1302	1149	1 640	1 670	1 819
1213	947 1	1 500	1 515	1 535	1258	161	1 538	1 600	1 602	1303	3539	1 493	1 675	1 739
1214	3344		1 520		1259	3222	1 550	1 600	1 680	1304	5442	1.570	1 685	1 690
1215	975 1	1 413	1 520	1 589	1260	5648	1.560	1 600	1 610	1305	1111 2	1.619	1 688	1 696
1216	5961		1 524	1 566	1261	976		1 6015	1.6187	1306	2566 2	1 597	1 692	1 806
1217	2373 1	1 528	1 529	1 537	1262	4530 2		1 602		1307	1058	1 5697	1 6935	1 7324
1218	1070 2	1 510	1 530	1 566	1263	1960		1 602		1308	84.1	1 631	1 698	1 713
1219	1672	1 524	1 531	1 534	1264	5320	1 571	1 602	1 647	1309	3103	1 479	1 710	1 810
1220	629	1 450	1 534	1 610	1265	1936 1	1 526	1 603		1310	4322	1 583	1 73	
1221	1705	1 525	1 535	1 560	1266	977	1 490	1 605	1 620	1311	445	1.490	1 743	1 872
1222	639	1 4955	1 5352	1 6045	1267	669 1	1 530	1 605	1 658	1312	4739	1 464	1 718	1 916
1223	67 1	1 1227	1 5358	1 5545	1268	3234	1 538	1 609	1 754	1313	1197	>1 56	1 75	>1 95
1224	638	1 495	1 536	1 605	1269	3208	1 600	1 610	1 675	1314	1200	1 650	1 760	1 870
1225	484	1 515	1 540	1 575	1270	1977	1 609	1 612	1 616	1315	1142	1 763	1 787	1 857
1226	5336	1 520	1 540	1 580	1271	3540	1 460	1 614	1 697	1316	87	1 710	1 847	1 863
1227	2367 1	1 536	1 540	1 541	1272	1414	1 601	1 614	1 731	1317	5818	1 524	1 867	1 873
1228	1035	1 532	1 541	1 549	1273	3732		1 615		1318	1112	1 508	1 870	1 907
1229	4391*	1 517	1 542	1 555	1274	241	1.495	1 615	1 650	1319	3060	1 535	1 873	1 893
1230	2372		1 513		1275	1115	1 578	1 620	1 627	1320	1364	1 54	>1 95	1 505
1231	1037	1 517	1 514	1 546	1276	3196	1 495	1 625	1 807					
1232	4318 1		1 515		1277	5202	1 580	1 625	1 645					
1233	303	1 4386	1 5157	1 5042	1278	5141	1 610	1 625	1 675					
1234	61 1	1 507	1 546	1 546	1279	5562	1 620	1 625	1 630					

Miscellaneous

Serial No	Gen index No	α	β	γ	Serial No	Gen index No	α	β	γ	Serial No	Gen index No	α	β	γ
1321	5135 1		1 52? (red)		1426	5221	1.49		1 58	1331	5511	1 625		1 690
1322	5244 1	1 529	1 533 (red)		1427	1069 1	1.495		1 565	1332	5424	1 565		1 768
1323	835 1		1 564 (red)		1428	610	1.570		1 660	1333	Bolland, 57, 31: 390, 10, approximate data only			
1324	868	1 385		1 530	1429	4500	1 583		1 747					
1325	3873*	1 480		1 522	1430	2135	1 602		1 627					

*Hydrated form   †Metastable modification   ‡Stable modification

INDEX TO C TABLE

Pyroracemic alcohol, 449
Pyrotartaric acid, 947
Pyrotritaric acid, 2180
Pyrrole, 589
Pyrrole-2-aldehyde, 874
Pyrrole-2-carboxylic acid, 877
Pyrrolidine, 759
α-Pyrrolidone, 672
Pyrroline, 670
Pyrrolylene, 596
Pyrrone, 3068
Pyruvic acid, 369
Pyruvic nitrile, 332 1
Quassin, 6034
Quebrachine, 5670
Quebrachine hydrochloride, 5672
Quebrachite, 2375
Quebrachol, 5698
Queroetagetin, 4891
Quercetin, 4889
d-Quercitol, 1670
l-Quercitol, 1671
Quercitrin, 5631
Quinaldine ethiodide, 4312
Quinaldine methiodide, 4056
Quinaldinic acid, 3479
Quinalgan, 5289
Quinamine, 5469
Quinazoline, 2463
Quinone, 5545
Quinhydrone, 4255
Quinic acid, 2313
Quinic amide, 2324
Quinic lactone, 2270
Quinicine, 5560
Quinide, 2270
Quinidine, 5561
Quinidine hydrochloride, 5567
Quinine, 5562, 5563, 5564
Quinine acetate, 5728
Quinine acetylsalicylate, 5971
Quinine butyrate, 5637
Quinine carbonate, 6127
Quinine citrate, 6175
Quinine dibromosalicylate, 5921
Quinine dichloride, 5555
Quinine diguaiacolsulfonate, 6003
Quinine disulfate, 5575
Quinine ethyl carbonate, 5801
Quinine formate, 5671
Quinine glycerophosphate, 6137
Quinine hydrate, 5587
Quinine hydrobromide, 5566
Quinine hydrochloride, 5568
Quinine hypophosphite, 5581
Quinine lactate, 5796, 5797, 5798
Quinine malate, 6139
Quinine β-naphtholsulfonate, 6122
Quinine nitrate, 5574
Quinine phenolsulfonate, 5807
Quinine propionate, 5795
Quinine salicylate, 5925
Quinine succinate, 6138
Quinine sulfate, 6124
Quinine tartrate, 6141
Quininic acid, 4036
Quinizarin, 4633
Quinoline, 3037
Quinoline-2-carboxylic acid, 3179
Quinoline-3-carboxylic acid, 3180
Quinoline-4-carboxylic acid, 3178
Quinoline-6-carboxylic acid, 3181
Quinoline-7-carboxylic acid, 3182
Quinoline-8-carboxylic acid, 3183
Quinoline-2, 3-dicarboxylic acid, 4021
Quinoline-2, 4-dicarboxylic acid, 4022
Quinoline ethiodide, 4057
Quinoline methiodide, 3406
Quinoline tartrate, 6136
Quinolinic acid, 1900
Quinone, 1287
Quinonedioxime, 1399

Quinoneoxime, 1348
Quinotoxine, 5560
β-Quinovin, 6001
Quinoxaline, 2464
Raffinose, 5350
Rapic acid, 5361
Ratanhine, 3720
β-Resalgin, 5198
Resorcinol, 1415
Resorcinol acetate, 2619
Resorcinol antipyrine, 5198
Resorcinol diacetate, 3620
Resorcinol diethyl ether, 3769
Resorcinol dimethyl ether, 2741
Resorcinol ethyl ether, 2742
Resorcinol methyl ether, 2175
Retamine, 1995
Retene, 5306
Rhamnitol, 1749
β-Rhamnose, 1672
Rhamnose phenylhydrazone, 4367 1
Rheadine, 5636
Rhein, 4886
Rhizochohc acid, 1016
Rhodamic acid, 24
Rhodeoretin, 6046
Rhodeose, 1673
Rhodinol, 3908
Rhodinyl acetate, 4374
Rhodinyl butyrate, 4835
d-Ribose, 1036
Ricinelaidic acid, 5368
Ricinic acid, 5369
Ricinine, 2554
Ricinmic acid, 1980
Ricinoleic acid, 5370
Robigenin, 4884
Robinin, 6049
Rochelle salt, 3184
Rosindon, 5696
Rosinduline, 5698
Rosindulon, 5696
Rosolic acid, 5518
Rubazonic acid, 5519
Ruberythric acid, 5895
Rubicine, 5882
Rubidium acid chloroacetate, 3234
Rubidium acid oxalate, 3225
Rubidium acid phthalate, 3228
Rubidium citrate, 3231
Rubidium cobalt malonate, 3263
Rubidium lithium d-tartrate, 3276
Rubidium sodium mesotartrate, 3277
Rubidium tartrate, 3229
Rubijervine, 5907
Rumicin, 4878
Rutin, 5928
Sabadinine, 5977
Sabadine, 5982
Sabinane, 3895
Sabinene, 3819
Sabinol, 3860
Saccharin, 1896
d-Saccharine, 1579
Saccharose, 4496
Safrol, 3591
Sakuranetin, 5076
Sakuranin, 5706
Saleetol, 3622
Salein, 4547
Salein benzoate, 5549
Saleitrin, 5620
Salicylaldehyde, 2004
Salicylaldehyde methylphenylhydrazone, 1769
Salicylamine, 2220
Salicylamide, 4485
Salicyl chloride, 1863
Salicylic acid, 2013
Salicylic nitrile, 1890
Salicylosalicylic acid, 4692

Salicyl-p-phenetidin, 4928
Salicyl sulfonic acid, 2027
Salicyluric acid, 3114
Saliformin, 4540
Saligenin, 2166
Salinigrin, 4535
Salipyrine, 5308
Salocol, 5210
Salol, 4467
Salophen, 4911
Saloquinine, 5925
Salvarsan, 4311
Samarium acetate, 2051
Samarium ethyl sulfate, 2054
Samarium formate, 2050
Samarium propionate, 2052, 2053
Sambunigrin, 4819
Sanguinarine, 5509
α-Santalene, 4979
β-Santalene, 4980
γ-Santalene, 4981
Santalic acid, 5987
Santalin, 5987
α-Santalol, 4987
β-Santalol, 4988
Santalyl chloride, 4970
Santalyl salicylate, 5730
Santene, 3273
dl-Santenol, 3302
Santinic acid, 4936
Santonic acid, 4956
Santonin, 1913
dl-Santonnine acid, 4955
Santyl, 5730
Saponarin, 5661
Sapotin, 5983
Sarcosine, 186
Sarsapic acid, 1289
Sarsasapogenin, 5905
Sarsasaponin, 6143
Scatole, 3100
α-Scatolecarboxylic acid, 3560
Scillitin, 5247
Scoparin, 5536
Scopolamine, 5222
Scopoletin, 5530
Scopolin, 5836
Scopoline, 2822
Scutellarein, 4887
Scutellarin, 5623
Sebacic acid, 3938
Sebanic acid, 3954
Sekisanine, 6061
Selenophenol, 1431
α-Selinene, 4982
Semicarbazide, 69
Semicarbazide hydrochloride, 76
Semicarbazide nitrate, 81
Seminose, 1682
Senecifolidine, 4990
Senecifoline, 5337
Senecifoline hydrochloride, 5338
Septentrinaline, 6051
dl-Serine, 193
d-Serine, 493.1
Sidonal, 5352
Silicon tetraethyl, 3418
Silicon tetraphenyl, 3424
Silver acetate, 1107
Silver lactate, 1108
Silver oxalate, 1106
Silver tartrate, 1109, 1110
Sinalbin, 5095
Sinapic acid, 4073
Sinapine sulfate, 6038
Sinapine thiocyanate, 5242
Sinapoline, 2285
Sinomenine, 5457
Sitosterol, 5935
Smilacin, 5906
Sodium acetate, 2756
Sodium acid malonate, 2758

Sodium acid tartrate, 2759
Sodium citrate, 2766
Sodium diacetate, 2760
Sodium elaidate, 2762
Sodium ethanedisulfonate, 2768
Sodium formate, 2754
Sodium glutamate, 2772
Sodium lithium tartrate, 2912
Sodium naphthalene-1, 5-disulfonate, 2767
Sodium 1, 4-naphthylaminesulfonate, 2775
Sodium oleate, 2763
Sodium palmitate, 2761
Sodium thallium tartrate, 2782
Sodium sulfanilate, 2774
Sodium tartrate, 2764
Sodium uranyl acetate, 2835
Solangustine, 6053
Solanidine, 6079
Solanine, 6158
Solanine hydrochloride, 6159
Sophoretin, 4889
Sophorin, 5927
Sorbic acid, 1498
Sorbic chloride, 1441
d-Sorbitol, 1752
d(l)-Sorbose, 1684
dl-Sorbose, 1685
Sordidin, 4471
Sparassol, 3695
Sparteine, 4994
Sparteine hydrochloride, 5011
Sparteine hydroiodide, 5012
Sparteine sulfate, 6009
Sphingosine, 5259
Spinacene, 5978
Spirosal, 3184
Stachydrine, 2323
Stannous acetate, 2519
Staphisagroine, 5556
Starch, 1578 1
Stearic acid, 5379
Stearic aldehyde, 5378
Stearic amide, 5390
Stearic anhydride, 6106
Stearic amilide, 5845
Stearolic acid, 5347
Stearone, 6090
Stearonitrile, 5374
Stearoxylic acid, 5349
Stearyl chloride, 5373
Stercorin, 5939
Stigmasterol, 6005
Stilbene, 4708
Stovaine, 4829
Strontium acetate, 2467
Strontium antimony tartrate, 2471
Strontium calcium propionate, 2493
Strontium ethanedisulfonate, 2468
Strontium ethylsulfate, 2469
Strontium formate, 2464
Strontium nitrotetronate, 2470
Strophantidin, 5929
Strophantin, 5998, 6125
Struxine, 5677
Stryacol, 5070
Strychnidine, 5656
Strychnine, 5642
Strychnine methylarsinate, 5719
Strychnine nitrate, 5653
Strychnine salicylate, 5948
Strychnine sulfate, 6129
Strychnine d-tartrate, 6148
Stylopine, 5425
Styphnic acid, 1198
Styracin, 5292
Styrene, 2538
Styrene dibromide, 2455
Styrolene alcohol, 2736
Styryl alcohol, 3126
Styrylamine, 3190

PROPERTY-SUBSTANCE TABLES

[All index numbers refer to the ℂ-Tables (p. 176) except those preceded by ℬ, which refer to the ℬ-Table (p. 106)]

I. MELTING POINTS

−207: ℬ337. −189.9: 500, 409, ℬ345, 54. −172: 252, 180, ℬ1810, ℬ465, ℬ95. −160: ℬ204, 1072, ℬ404, 145. −150: ℬ485.5, 1532, ℬ405, 2328, 337 −145: 31, 781.2, ℬ349, 1615, ℬ126, 982, 224, 17.1, 263, 148, 374, 375, 1712. −135: 781.1, 985, ℬ433, ℬ125, ℬ366, ℬ195, ℬ346, 1073, 747. −130: 598, 686, 446, ℬ1813, 505, 408, 2330. −125: 3890, 365, 526, 986, 208, 793, 746, 468, ℬ365, 1741, 63, 273. −120: 915, 916, 366, 220, 742, ℬ364, 9, 262, 1079, ℬ347, 469, 364, 793. −114.8: 1764, ℬ263, 667, 741, ℬ13, ℬ1811, 154, ℬ207, 518, 209, ℬ9, 2058. −110: ℬ53, 464, ℬ1752, ℬ466, 234, 790, ℬ339, ℬ1814, 310, ℬ467. −105: 1102, 513, 517, 338, 822, ℬ1816, 756, 1537, 754, ℬ97, 814, ℬ98, 247, 3229, 475, 525. −100: ℬ374, 594, 213, 837, 391, 2357, 717, 1652, 1610, 1716, 452. −98: 1466, 60, 41, 615, 3810, 28, 189, ℬ1597, 149, 282, 1016, 2361, 2112. −95: 920, 1763, 448, 1713, ℬ371, 378, ℬ348, 755, 3372, 979, 1654. −93: ℬ134, 40, 727, 2683, 65, 1020, 35, 1007, 395, 2933, 176 −90: ℬ42, ℬ472, 652, 1011, 2389, 789, 465, 651, 1655 −88: ℬ282, 356, 726, 106, 719, ℬ17, 2869, 506, 403, 821. −85: 2868, 1019, 1639, 725, 100, 272, ℬ4, ℬ205, 524, 1587, ℬ11. −82: 332, 115, 447, 2905, 283, 383, 133, 451. −80: ℬ57, ℬ62, ℬ72, ℬ105, ℬ265, 13, 2981, 1078, ℬ55, ℬ107, 1006, ℬ102. −77: ℬ113, ℬ350, ℬ372, 1651, 2415, ℬ6, 1659. −75: 614, 2840, 3728.1, 1021, 626, 2382, 1045, ℬ38, 1017, 2359, 693. −70: ℬ352, 2985, 1443, 227, ℬ206, 823, 2834, ℬ356, 45, 718. −65: 539, ℬ130, ℬ68, 11, 19, 2763, 948, 2955, 3230, 3258 −60: ℬ238, 38, 1012, 1548, 107, 2867, 91, 215, 459, 2203, 1638, 2941. −56: 2001, ℬ20, ℬ141, 412, 2741, ℬ2701, ℬ52, 1099, 3817, ℬ59. −54: 367, 2685, 2841, ℬ43, 26, 3995, 3228, 1630, 1728, 3684, 3149, 1603. −51: ℬ3, 176, 1002, 3354, 4450, ℬ26, 820, ℬ64, 88, 132, 683, 824, 2309, ℬ468. −48: ℬ28, ℬ84, 2040, ℬ19, 724, 2340, ℬ63, 2927, 1307. −45: ℬ1815, 169, 1560, 2343, 3893, 3984, 377, 4175, 140, 2491, ℬ46. −43: ℬ27, 1026, ℬ109, 870, 1005, 1328, 168, 3337, 1570. −40: ℬ121, ℬ142, ℬ214, ℬ267, ℬ797, 400, 576, 2031, 1760. −39: ℬ115, ℬ378, 622, 2038, ℬ47, 859, 2968, 3080. −38: ℬ110, 1013, ℬ60, 2163, 159, ℬ29, 190, 2030. −35: ℬ114, ℬ232, 987, 1644, 3219, 3731, 1388, 2403, 3154, 95, 413, 3789. −34: ℬ112, 186, 1606, ℬ360, 384, 2344, 3993, 260, 2205, 4400, 1333. −31: ℬ192, 1294, 1571, ℬ488, 2722, ℬ56, ℬ237, ℬ450, 3638, 48. −29: 111, 886, 749, 2030, ℬ209, ℬ798, ℬ619, 2684, 233, 4178, 151, 568, 809, 4789, 379. −25: ℬ48, ℬ1575, 827, 2393, 5688, 1230, ℬ12, 2204, 2503, 1253, 929. −23: 12, 2419, 911, 90, 4095, 450, 609, 1734, 4038, 4419, 2896, 1200.2. −21: ℬ1397, 2, 121, 1918, 2847, ℬ7, ℬ90, ℬ147, ℬ493, 205, 690,

714, 1074, 2961, 85, 3037. −19: 671, 1762.1, 4283, ℬ110.1, ℬ136, ℬ266, 4176, ℬ15, ℬ11, 1229, 329, 264, 1964. −17: 355, 449, 4398, 6166, 181, 2204, 2966, 2853, 112.1, 3133, ℬ10, 2159. −16: ℬ530, 388, 735, 22, 1646, 1725, 1885, ℬ186, 428, 1200.1, 2589. −12: 68, 89, 290.1, 2327, 4411, 4849, 1081, ℬ18, 1642, 4852, 2081, 1376, 1694. −10: 2509, 2351, 3667, 1643, ℬ96, 1200.5, ℬ914, 1054, 1551, 3573, 3978.1, 2618. −8: ℬ155, 561, 2518, 3129, 723, 4513, ℬ1866, 2493, 3069, 2004, 1205, 1442, 4587. −6: 7, 606, 826, 1366, 3863, 3929, 404, 1249, 994, ℬ156. −5: ℬ235, 46, 2514, 2766, 2949, 3364, 1870, 1314, 2081. −4: 615, 1812, 2029, 2711, 3740, 4169, 1856, 4493, 2928. −2: ℬ149, 3054, 3835, 1390, ℬ2, ℬ1515, ℬ353, 3902, 4847, 5153, 1859, 141. 0: ℬ1, ℬ40, 127, 1314, 1375, 1844, 2423, 3276, 128, 2633, ℬ2953, 458, 3752, 2812. 1: ℬ494, 1478, 4846, ℬ210, 3170, ℬ103, 2756, 1250, 1204. 2: ℬ384, 2669, 1200.4, 2788, ℬ1672, 2580, ℬ287, 407, 1003, 1880, 5690, ℬ2612, 3978. 4: 3640, 4399, 5156, 84, ℬ23, ℬ100, ℬ367, 24, 313, 3453, 4584, 29, 2216, 1365, 1733, 1949, 2725, 4296, 4856, 1296, 1347. 6: 102, 3852, 1612, 4062, ℬ79, 376, ℬ284, 863, 1314, 1959, 2492, 4002, 4761, 18, 2041, 3688. 8: ℬ34, ℬ381, 2882, 4442, 5940, ℬ76, 300, 37, 2676, 5480, ℬ46. 9: 319, 1105, 4157, 4402, 4707, 4794, 2098, ℬ5. 10: 136, 184, 307, 934, 2053, 2161, 2639, 5018, 5, ℬ45, 1662, 3944, 5160, ℬ135. 11: ℬ163, 345, 2886, 3035, 3591, 3662, ℬ121, 3328, 4172, 358, 3769. 13: ℬ218, 17, 1691, 3776, 4828, 2686, ℬ3219, 699, 1332, 2153, 359, ℬ146, 1330, ℬ35.1. 14: ℬ385, 5359, 5361, 2865, 1318, 619, 4170. 15: 25, 146, 721, 1482, 2134, 2190, 5851, ℬ2669, 2082, 2759, 3574. 16: 620, 656, 795, 2128, 2899, 1329, 655, 2294, 4180, 212, ℬ39. 17: ℬ1646, 372, 558, 621, 1176, 2398, 2850, 4011, 4542, 4843, 5370, 160, 4758, 515. 18: ℬ2843, 457, 646, 1506, 1857, 1863, 3933, 5377, 5380, 1369, 4733. 19: 602, 4179, 4697, 4848, 5017, 1568, 1483, 2571. 20: ℬ80, ℬ856, ℬ1039, ℬ2443, 2498, 3157, 3192, 4529, 5167, 5682, 126. 21: 841, 2632, 3132, 4505, 1251, 1958. 22: ℬ644, ℬ2698, ℬ2961, 562, 1117, 2120, 3160, 5164, 5481, 5812, 6105, ℬ191, 2737, 3648, 4392, 5260. 23: 1954, 2322, 3036, 4173, 5373, 1258, 120, 1627. 24: 1320, 2002, 4367.4, 4415, 2060, 5381, ℬ2673, 2664, 4158. 25: ℬ71, ℬ2450, ℬ2688, 672, 3078, ℬ1234, 792, ℬ2514, 368, ℬ1287, ℬ2813. 26: 1458, 2118, 2119, 2126, 2248, 2338, 2525, 2706, 3240, 4523, 5353, 5735, ℬ470, ℬ975, 1251, 4448, 4241. 27: ℬ1134, ℬ1345, 456, 833, 1392, 1432, 2086, 2500, 3717, 4490, 4590, 5484, ℬ35. 28: ℬ158, ℬ2739, 765, 1163, 2032, 2174, 2427, 2738, 3277, 4417, 5391, 1331. 29: 51, 1267, 1966, 2515, 4373, 5155, 5760, 7174, 2507, 4322, 5258, ℬ2622, 2160, ℬ2237. 30: ℬ99, ℬ123, ℬ219, ℬ331, ℬ2666, 267, 312, ℐ822, 1945, 2120.1, 2123, 2667, 3173, 4414, 4440,

5200, **5354, 6044, 4321, 1888, 2464, 4589, 5852. 31:** ℬ491, 1023, 1948, **2293, 3981, 4318, 4484, 6056,** 851, 1368. **32:** 627, 647, 1553, **2114, 2490, 3003, 4579, 5486,** 5738, 6165, 316, ℬ2693, 860, ℬ268, 1315. **33:** ℬ36, ℬ3084, 163, 1170, 1297, 3126, 3156, 3255, 3304, 3924, 5151, 941, 4019, 5764, 5608, 2162, 4914. **34:** ℬ111, ℬ886, 1807, 2136, 2434, 2715, 3002, 3689, 4125, 4205, 1268, 5068, 5087, 5360, 5389, 1259, 2544.2, ℬ1149, ℬ2724, 5767. ℬ1286. **35:** ℬ50, ℬ200, ℬ1367, ℬ2133, ℬ3313, 177, 1591, 2063, 2071, 2578, 3588, 3922, 4250, 4300, 4741, 5371.1, ℬ2752, 4039, 1011, 2247, 2688, 3974, 5184. **36:** ℬ244, 1207, 1255, 2116, 3028, 3600, 3757, 3972, 4226, 4288, 5255, 5355, 5805, 1163, ℬ768, 2096, 5020, 229, 3092 **37:** ℬ607, 1316, 1957, 2474, 3081, 3412, 3628, 4212, 4512, 4771, 4859, 5850, 1163, ℬ1962, 203, 4060.1, 4297, 1893. **38:** ℬ234, ℬ386, 56, 1270, 1743, 1794, 2099, 2638, 3000, 3703, 4362, 4383, 4858, 5485, 5610, ℬ2241, 1955, 3079, 1163, 1219. **39:** ℬ451, 602, 603, 1323, 1772, 1950, 4368, 4508, 4743, ℬ1147.5, 5072, 5150, 5966, 2900, 4530.3, 3317, ℬ2260. **40:** ℬ701, ℬ702, 193, 546, 961, 974, 1335, 1550, 2067, 2178, 3793, 3923, 4109, 4327, 6112, ℬ1147, 1334, 5694, 1893. **41:** ℬ252, ℬ1147, 118, 852, 1413, 1871, 1946, 2424, 2522.1, 2553, 2801, 2819, 2854, 3004, 3070, 4103, 4104, 4452, 4823, 5224, 5374, 97, ℬ78. **42:** ℬ37, ℬ2245, ℬ2718, ℬ3242, ℬ203, 66, 1715, 1792, 2743, 3587, 3905, 4588, 5332, 5609, 5739, 5848, 730, 1873, 3973, ℬ2259, 3802, 4220. **43:** ℬ1817, 1166, 1173, 1201, 2066, 2192, 3272, 3710, 4018, 4367.9, 4467, 4676, 2978, 4551, 2206, 156. **44:** ℬ2611, 30, 293, 654, 1157, 1283, 1811, 1956, 2223, 3220, 3692, 3870, 4053, 4330, 4503, 4984, 5292, ℬ766, 5014, 1216, 5772, 1962, 4405, 4247. **45:** ℬ49, ℬ2927, ℬ3223, ℬ3261, 346, 931, 933, 1233, 1339, 1349, 1552, 1850, 2501, 2723, 2781, 3005, 3766, 4287, 4449, 4831, 4837, 5044, 5310, 6114, ℬ767, 393, 3685, 4266, 4431, 4788, 5637, ℬ1343, 5876. **46:** 559, 1111, 1395, 1423, 1874, 1941, 1969, 2037, 2712, 2802, 3252, 4305, 4308, 4998, 5019, 5343.1, 5856, 6157, 553, 5092, 6120.1 **47:** ℬ1640, ℬ2655, ℬ3272, 239, 874, 1439, 1978, 3115, 3144, 3130, 3251, 3774, 3906, 4297.1, 5097, 5148, 161, 1135, 1858, 2073, 3670, 5816. **48:** ℬ2696, 125, 1581, 1937, 2463, 3172, 3299, 3617, 3416, 4328, 4406, 4520, 4815, 5285, 5347, 6145, 1331, 6, 4447, 3147. **49:** ℬ1980, 492, 1961, 2046, 2047, 2263, 2707, 2761, 3291, 3907, 4106, 4787, 4947, 5262, 5243, 5769, 5770, 5168, 1254, 5283. **50:** ℬ14, ℬ3498, ℬ1144, ℬ2792, 27, 150, 255, 389, 844, 1268, 1322, 1428, 2494, 2881, 3159, 3550, 3801, 4780, 4908, 5604, 5218, 156, 1325, 1986, 5348, 5859, ℬ2678. **51:** ℬ1509, 343, 1136, 1793, 1935, 2003, 2432, 3663, 2594, 3669, 3836, 4558, 4582, 4800, 4853, 5860, ℬ2258, 4203, 2083, 3756, 5357. **52:** 3, 244, 1175, 1303, 2807, 3221, 3250, 3658, 4065, 4094, 4318.1, 4435, 5557, 6086, 2504, 4756, 4850, 1231. **53:** ℬ387, 152.1, 563, 771, 1082, 2049, 2143, 2176, 2785, 4119, 4270, 5368, 5429, 5993, 956, 1165, 1178, 701. **54:** 361, 498, 631, 1066, 1851, 2100, 2577, 3619, 4473, 5016, 5521, 5861, 5881, 6047, 560, 1171, 3709, 5244, 5815, 6169. **55:** ℬ290, 2730, 663, 1968, 2051, 2544.1, 2591, 4049, 4333, 4443, 4736, 4763, 4765, 5043, 5222, 5729, 6147, 2575, 2386. **56:** ℬ216, ℬ1822, 15, 179, 348, 607, 882, 1131, 1202, 1208, 2095, 2256, 2470, 2559, 2739, 3456, 3704, 4238, 4323, 4368 7, 4369.2, 5142, 5176, 5625, 5917, 156, ℬ2010, ℬ2073, 1306, 1391, ℬ1552 **57:** ℬ61, 105, 548, 897, 1133, 1214.1, 1232, 2314, 3076, 3116, 3137, 4503, 4816, 5587, 5689, ℬ2672, 86, 109, 3652, 5761, 2218. **58:** ℬ70, ℬ508, ℬ1769, ℬ2757, ℬ1269, 497, 902, 921.2, 1154, 1234, 1719, 1810, 1894, 1942, 2650, 2815, 3015, 3175, 3671, 4045, 4851, 5646, 3056, 3101, 5158, 5392, 3473. **59:** ℬ1714, 288, 1167, 1852, 2497, 3454, 3470, 3708, 3921, 4273, 4460, 4770, 4996, 5063, 5183, 5201, 5431, 5440, 1984, ℬ845, 1150, 2115, 2717, 5301, 5942, 1988, 5257. **60:** ℬ1165, ℬ1667, ℬ2110, ℬ2720, ℬ2931, ℬ2932, ℬ2934, ℬ3142, ℬ3343, 96, 583, 960, 1168, 1821, 2508, 2678, 2857, 3099, 3218, 3509, 4223, 4445, 4557, 4650, 4724, 4727, 5147, 5256, 5279, 5560, 5719, 5918, 6065, 6151, 1875, 4024, 4516, 5116, 5213.1. **61:** ℬ220, ℬ1450, ℬ2439, ℬ2855, 315, 342, 369, 382, 481, 581, 611, 1191, 1192, 1947, 1960, 1987, 2258, 3038, 3071, 3197, 3241, 3413, 3593, 3742, 3779, 4046, 4213, 4261, 4740, 5261, ℬ376, 5437, 5915,

5919, 156, 1581, 5520, 5763, 1277. **62:** ℬ3222, 1152, 1169, 1393, 1663, 2034, 2103, 2249, 2783, 3417, 3776, 3912, 4782, 4838, 4893, 4902, 5054, 5216, 5810, ℬ120, ℬ1867, 1480. **63:** 214, 675, 838, 1177, 1213, 1321, 2036, 2864, 3183, 3257, 4320, 4655, 4802, 5066.1, 5093, 5846, 6010, ℬ2075, ℬ3271, 1298, 1704, 5378, 5985. **64:** ℬ455, ℬ732, 32, 343, 442, 461, 883, 936, 2033, 2251, 2672, 3099, 3700, 4340, 5159, 6045, 6144, ℬ1272, ℬ1907, 426, 432, 1116, ℬ2812 **65:** ℬ108, ℬ119, ℬ942, ℬ2607, ℬ2903, 257, 324, 1865, 2172, 2473, 2708, 2921, 3597, 1047, 1108, 4210, 4746, 5053, 5057, 5436, 5367, 5532, 5967, 6157, 1174, 3195. **66:** 333, 439, 580, 1854, 2048, 2137, 2171, 2478, 2697, 2740, 3269, 3398, 5483, 5963, 1370, 3584, 4020, 4228, ℬ2027. **67:** 830, 845, 1067, 1235, 1319, 2167, 2992, 3016, 3630, 4225, 4352, 5149, 1200.3, 1245, 4208, 5358. **68:** ℬ1081, 865, 1134, 1179, 1236, 1237, 1262, 1343, 1484, 2221, 2709, 2736, 3074, 3143, 3414, 3467, 3498, 3614, 3650, 3695, 3953, 4368 41, 4507, 4528, 4530 1, 4953, 5161, 5759, 6028, 1929, 1248. **69:** 401, 538, 1138, 2000, 2054, 2185, 2805, 3201, 3238, 3594, 4219, 4393, 5344, 5602, 5691, 5814, 5862, 1993, 5379, 238, 2091, 2913, 6058, 1985, 4204. **70:** ℬ908, ℬ1077, ℬ2638, ℬ3270, 320, 351, 802, 932, 1220, 1463, 1501, 2272, 2519, 3065, 3457, 3791, 3925, 4458, 4495, 4772, 5193, 5526, 5553, 5893, 6014, ℬ2056, 2044, 3991, 4242, ℬ120, 6160. **71:** 1377, 1809, 1963, 2009, 2670, 3198, 3462, 3622, 4549, 4764, 5061, 5611, 1324, 1396, 2068, 4734, ℬ2164. **72:** ℬ116, ℬ275, ℬ1141, ℬ1971, ℬ2481, ℬ3149, 617, 1261, 1437, 1808, 1832, 1970, 2042, 2532, 2879, 3010, 3242, 3768, 4042, 4786, 5854, 5857, 5959, 6106, ℬ2076, 1172, 2543, 311. **73:** ℬ168, ℬ915, ℬ1878, ℬ2614, 2469, 2473, 2592, 4912, 4919, ℬ286, ℬ2713, 962, 2455, ℬ202. **74:** ℬ647, ℬ1948, ℬ1988, 195, 380, 491, 674, 1703, 2085, 2726, 3558, 3595, 3686, 4393.1, 4480, 4742, 4769, 5154, 5692, 5773, 5991, 6013, 6081, 1666, 6091, 142. **75:** ℬ239, ℬ389, ℬ643, ℬ866, ℬ1626, ℬ2044, ℬ2848, 455, 584, 738, 842, 1371, 1806, 1919, 2561, 2682, 2705, 2749, 3193, 3775, 4233, 4285, 5313, 5364, 5422, 5847, 6048, 6003, 1151, ℬ679, ℬ1796, 784, 4207, 5173. **76:** ℬ270, ℬ3204, 292, 441, 907, 957, 1665, 2306, 2452, 2844, 2980, 3044, 3468, 3697, 3712, 4072, 4950, 5366, 5375, 5410, 5941, ℬ2055, 570, 1210, 5823, 6080, 6107, 2584. **77:** ℬ932, ℬ2033, 2540.2, 3254, 3544, 3745, 4389, 5067.1, 5405, 5607, 5638, ℬ2231, 1247, 2147, 3583, 5837, 5878, ℬ2498. **78:** ℬ2757, ℬ2976, 664, 673, 975, 2055, 2425, 3389, 3746, 3778, 3792, 4034, 4239, 4433, 5000, 5280, 5387, 5639, 5879, 5962, 710, 79, 2537, 4187, 2117. **79:** ℬ293, ℬ909, ℬ2733, 214, 482, 731, 1269, 1522, 2131, 2212, 3233, 3290, 3474, 3713, 4067, 4330.1, 4792, 5284, 5386, 5411, 5824, 5835, 1302. **80:** ℬ139, ℬ144, ℬ181, ℬ655, ℬ683, ℬ827, ℬ1535, ℬ1857, ℬ3071, 152, 1209, 2120, 2179, 3705, 3722, 3732, 3777, 3875, 4319, 4647, 4798, 4890, 5220, 5492, 5920, 6113, 3494, 1305, ℬ2240, 3394, 3631, 1924. **81:** ℬ2020, 238, 1257, 1664, 1690, 2596, 3387, 3428, 3543, 4026, 4466, 4903, 5073, 5369, 5501, 5855, 5223, 5385. **82:** ℬ245, ℬ249, 93, 167, 197, 394, 1450, 1513, 1621, 1872, 1967, 2207, 2558, 2694, 2727, 2843, 2916, 4271, 4640, 4702, 5174, 5545, 5808, 6109, 6110, 3091, 3603, 5968, 5916. **83:** 1127, 1181, 1263, 1286, 1600, 2056, 2125, 2950, 3080, 3165, 3970, 4432, 5178, 5365, 5766, 6026, ℬ2057, 1217, 3392, ℬ2000, 5840. **84:** 1187, 1304, 1418, 3216, 3410, 3749, 3750, 4425, 4506, 1985, 5263, 5407, 5592, 5765, 5768, 6160, ℬ649. **85:** 325, 398, 1153, 1352, 1580, 1766, 2510, 2695, 2993, 3315, 3370, 3519, 3627, 4833, 4907, 5383, 5384, 5799, ℬ458, 2997, 4332, 5858, 1996. **86:** ℬ1449, ℬ3269, 217, 1021, 1119, 1138, 1753, 2045, 2166, 2573, 2662, 2799, 2808.1, 3596, 4476, 4892, 4897, 4924, 5135.1, 5349, 6008, ℬ2735, 1162, 1206, 5984. **87:** 864, 1036, 1164, 1218, 1378, 1567, 1574, 2682.1, 2918, 3107, 4079, 4152, 4195, 4390, 4997, 5408, 5593, ℬ1974, 1156. **88:** 201, 240, 290, 1184, 1301, 2065, 2542, 3391, 3418, 4368.3, 4519, 4536, 4698, 5025, 5163, 5219, 5436, 6015, 6090, ℬ2032, 2253. **89:** ℬ133, ℬ375, 175, 677, 1039, 1111.2, 1132, 1855, 2811, 3013, 3537, 4206, 4209, 4522, 5085, 5119, 5244.1, 5286, 5583, 5673, 5779, 5922, ℬ2078, 2471, 3396, 6043, 1272. **90:** ℬ157, ℬ983, ℬ1002, ℬ1627, ℬ1708, 129, 350,

1111.1, 1185, 1440, 1449, 1582, 2533, 3401, 3524, 4258, 4521, 4927, 5341, 5418, 5762, 5951, 6100, 5745. **91:** ℬ253, ℬ1468, ℬ1910, ℬ1960, ℬ2069, 843, 898, 1300, 2094, 2462, 3163, 3213, 3400, 4515, 4958, 5142.1, 5230, 5801, 6011, 6054, ℬ648, ℬ2019, 2142. **92:** ℬ171, 504, 1200 6, 3388, 3706, 4703, 4915, 4928, 5308, 5412, 5438, 5533, 5633, 5830, 5955, ℬ2058, 1159, 1211, 3783, 5402, ℬ1908, 6024. **93:** ℬ391, 1989, 2087, 2555, 2564, 2995, 3124, 4127, 4184, 4436, 4530 2, 5002, 5153, 4709, 4918, 1964, 5044, 5579, 5618, 6027, 6156, ℬ1882, 543, 846, 5845. **94:** ℬ1082, 1336, 1381, 1769 1, 2222, 2286, 3060, 3460, 4969, 5781, 5802, 5938, 2540.1, 3471, 2089. **95:** ℬ368, ℬ650, ℬ727, ℬ2149, ℬ2609, 39, 281, 952, 2139, 2208, 2582, 3100, 3427, 4218, 4360, 4710, 4795, 5162, 5179, 5287, 5617, 5872, 6057, 1990, 4672, ℬ2034, 1372, 4921 **96:** ℬ2968, 69, 1160, 1196, 1216, 1340, 1350, 1566, 2139, 2385, 2520, 3123, 3507, 3580, 3711, 3744, 4275, 4509, 5231, 5601, 5675, 2647, ℬ291, ℬ1802. **97:** ℬ2021, ℬ2625, 211, 284, 662, 1241, 1922, 1991, 2653, 3390, 3522, 4082, 4868, 4906, 5065, 5363, 5555, 5667, 5904, ℬ1865, 946, 5128. **98:** ℬ1648, ℬ1972, 157, 955, 1433, 1564, 1890, 2614, 2829, 3302, 4113, 5221, 5640, 5974, 2145, ℬ2001, 1123, 5306, 6007. **99:** 141, 579, 996, 1118, 1356, 1995, 2250, 2539, 2617, 3231, 3523, 3867, 4325, 4615, 4896, 5954, ℬ1812, 4652. **100:** ℬ22, ℬ241, ℬ534, ℬ563, ℬ564, ℬ651, ℬ784, ℬ786, ℬ792, ℬ796, ℬ842, ℬ851, ℬ860, ℬ878, ℬ879, ℬ910, ℬ914 1, ℬ1108, ℬ1168, ℬ1197, ℬ1398, ℬ1454, ℬ1540, ℬ1613, ℬ1713, ℬ1716, ℬ1943, ℬ2297, ℬ2629, ℬ2729, ℬ2791, ℬ2793, ℬ2920, ℬ3155, ℬ3312, 635, 1199, 1248, 1526, 1913, 2285, 2448, 2615, 2998, 3784, 4013, 4401, 4797, 5124, 5133, 5215, 5250, 5432, 5725, 5880, 2475, 2515, 3751, 4451. **101:** 259, 540, 1688, 2562, 3008, 3207, 3422, 3466, 3621, 4854, 5622, 270, 1149, 2304, ℬ1975. **102:** ℬ230, ℬ1113, 1094, 1499, 3199, 3287, 3301, 3525, 3641, 4236, 4410, 5077, 5965, 2585, 502, 3282 1, 5038. **103:** ℬ927, ℬ1240, ℬ3054, 894, 1095, 1313, 1525, 2105, 2308, 3393, 3435, 3489, 4517, 4641, 4752, 4855, 5175, 1179. **104:** ℬ122, ℬ162, ℬ652, ℬ3040, 271, 1264, 1276, 1197, 1674, 1926, 1928, 2104, 2169, 2373, 2943, 2996, 3082, 3200, 3772, 3829, 4437, 4688, 5194, 5937, 6016, 3979. **105:** ℬ128, ℬ988, ℬ2949, 72, 1035, 1242, 1308, 1382, 1414, 2140, 2183, 2252, 2735, 2991, 3179, 3500, 3520, 3867.1, 4368.2, 4713, 4951, 4960, 5082.4, 5409, 5500, 5664, 5939, 6012, ℬ2035, 782, 1994. **106:** ℬ431, ℬ1162, 702, 1057, 1077, 1155, 1317, 1932, 2005, 2648, 2931, 3665, 4302, 5165, 5236, 5524, 5570, 5687, 5734, 5891, 1895, 3307. **107:** ℬ460, ℬ653, ℬ3265, 871, 922, 1436, 1931, 2305, 3077, 3115, 3535, 3930, 4196, 4237, 4807, 5115, 5394, 2782 **108:** ℬ2022, ℬ2114, 182, 483, 1282, 2011, 2130, 2173, 2227, 2863, 3123, 4051, 4087, 4257, 4434, 4475, 4537, 4730, 5055, 5710, ℬ2002, 250, 2574. **109:** ℬ1242, ℬ3251, 298, 1829, 2114, 2367.1, 2583, 2622, 3287, 4058, 4080, 4243, 4368.5, 4913, 4922, 5390, 5616, 3399, 3477, ℬ1078, 2456. **110:** ℬ30, ℬ32, ℬ106, ℬ221, ℬ459, ℬ533, ℬ726, ℬ925, ℬ941, ℬ978, ℬ1078, ℬ1089, ℬ1172, ℬ1673, ℬ1707, ℬ1728, ℬ2070, ℬ3203, ℬ3333, 143, 187, 700, 785, 958, 1115, 1709, 1752, 2149, 2168, 2820, 2822, 2919, 3062, 3194, 4272, 4866, 5177, 5499, 5507, 5693, 5732, 5787, 3551, ℬ1976, 2461, 2586. **111:** ℬ3334, 947, 1161, 1225, 1295, 1403, 1923, 2300, 2598, 3029, 3164, 3800, 4281, 4455, 4552, 4894, 5094, 5249, 5423, 5795, 945, 4935, 1275, 1397. **112:** ℬ654, ℬ2990, 322, 537, 740, 1212, 1386, 1841, 1921, 2133, 2368, 2665, 3499, 3607, 5185, 5233, 5400, 5674, 5771. **113:** 905, 1351, 1495, 1795, 1853, 3145, 3395, 3403, 4058, 4214, 5138, 5246, 5671, 6101. **114:** ℬ250, 289, 641, 1193, 2681, 2920, 3063, 3112, 3421, 4580, 4700, 4729, 5129, 2649, 3974, 4465. **115:** ℬ295, ℬ985, ℬ1032, ℬ1131, ℬ1133, ℬ1644, ℬ1704, 678, 1122.1, 1299, 1312, 1353, 2531, 2560, 2744, 3158, 3125, 3464, 3472, 3604, 4105, 4186, 5039, 5198, 5538, 5715, 6041, 3643.1, 5235, 1287. **116:** ℬ3347, 321, 349, 760, 1128, 1186, 1971, 2006, 2110, 2315, 2472, 2595, 3141, 3411, 4077, 4384, 4439, 4453, 4648, 4801, 4901, 5034, 5584, 1271, 3668. **117:** ℬ183, ℬ989, 866, 953, 1122.2, 2148, 2466, 2994, 3458, 3608, 3795, 4043.1, 4753, 5325, 4008, 2576, 4774,

1155.1, 1565, 1992, 4293, 5071, 5208, 5903. **118:** ℬ2132, 261, 1183, 1195, 1214, 1224, 1309, 1383, 1830, 2090, 2187, 2613, 2930, 3211, 3629, 3794, 4112, 5123. **119:** 21, 1213, 1718, 1788, 3136, 3718, 4469, 4570, 5099, 5126, 5350, ℬ1083, 188, 3884, 571, 1158, **120:** ℬ166, ℬ179, ℬ274, ℬ566, ℬ709, ℬ820, ℬ905, ℬ992, ℬ1135, ℬ1715, ℬ2117, ℬ2440, ℬ2504, ℬ2956, ℬ3108, 245, 352, 1125, 1194, 1221, 1265, 1292, 1488, 1930, 2027, 2511, 2527, 2597, 2679, 2746, 3024, 3419, 3447, 3534, 3581, 4110, 4955, 5237, 5252, 5325 1, 5406, 5589, 5665, 5714, ℬ379. **121:** ℬ3029, 178, 363, 1130, 1192, 1749, 2307, 2530, 3007, 4762, 4783, 4839, 4900, 5074, 5111, 5454, 5571, 1827, 2007, ℬ922, 1197. **122:** 313.1, 1174.1, 1824, 1877, 2236, 2458, 3503, 3508, 4459, 4818, 5116, 5554, 5980, 8, 2502. **123:** ℬ242, ℬ979, 81, 947.1, 954, 1122, 1447, 1487, 1833, 1997, 3169, 3343, 3404, 3461, 5461, 5466, ℬ251, 4245. **124:** ℬ2025, 330, 592, 1686, 1878, 2566, 1504.1, 3429, 3618, 4192, 4361, 1477, 4708, 4739, 5062.1, 5465, 5829, 5869, 5947. **125:** ℬ172, ℬ656, ℬ705, ℬ716, ℬ904.2, ℬ3048, 836, 1311, 1502, 2077, 2170, 2535, 2593, 2731, 3166, 3429.1, 3448, 3510, 3536, 3782, 4123, 4128, 4749, 5117, 5598, 5804.1, 5841, 6096, 6136, 5613. **126:** ℬ531, 521, 697.1, 808, 1348, 1672, 1689, 2877, 3139, 4063, 4274, 4461, 4805, 5114, 5728, 5930, 1279, 861, 3465, 5281. **127:** ℬ152, ℬ706, ℬ1658, ℬ2116, 1203, 1222, 1358, 1999, 2224, 2661, 2750, 2916.1, 3217, 3504, 3585, 3938, 4135, 4539, 4677, 4796, 4799, 5128.1, 5497, 5505, 1190, 2141. **128:** ℬ1665, 291, 463, 1108, 1710, 2211, 2416, 2693, 3463, 3485, 3839, 3881, 4009, 4355, 4681, 5712, 4286. **129:** ℬ2606, 636, 761, 872, 1387, 1783, 2093, 2220, 2601, 2674, 3059, 3167, 3431, 3437, 3934, 4249, 4301, 4420, 4744, 4806, 4961, 5096, 5340, 5621, 1260, 5204, 4216. **130:** ℬ24, ℬ178, ℬ848, ℬ1181, ℬ2303, ℬ2676, 104, 634, 1607, 1826, 1842, 1910, 1933, 1972, 2070, 2072, 2109, 2839, 3045, 3374, 4021, 4695, 5070, 5101, 6035, 6063, 574, 680, 2566.1, 3095, 2431. **131:** 1038, 1244, 2000, 2493.1, 3026, 3397, 3602, 4289, 4755, 5102, 5415, 5794, 5864, 6051, 2699, 3159. **132:** ℬ140, ℬ646, ℬ1255, 906, 1682, 2176.1, 2324, 2563, 3030, 3058, 3140, 3424, 3838, 4369, 5534, 5741, 5886, 4923, 4936, 55. **133:** ℬ369, 326, 867, 975.1, 1226, 1681, 1683, 2612, 2673, 3012, 3072, 3075, 3256, 3496, 4234, 4334, 4468, 4623, 4728, 4811, 5157, 3087, 5551. **134:** 899, 1180, 1278, 1354, 1119, 1504, 2860, 3073, 4454, 4699, 5515, 5539, 5908, 6126, 1498, 1825, 1952, 4651. **135:** ℬ986, 630, 904, 1070.2, 1461, 1462, 2180, 2215, 2371, 3027, 3420, 3449, 3528, 3716, 4485, 4497, 5275, 5456, 5612, 5615, 360. **136:** 862, 1144, 1400, 1891, 2999, 3282.2, 4052, 4687, 4826, 5188, 5225, 5527, 5700, 5723, 5890, 5934, 5388, 5105. **137:** 1058, 1344, 1360, 1669, 2600, 2730, 3014, 3031, 3161, 3501, 3797, 4313, 4335, 4925, 5098, 5403, 5445, 6094, 1925, 2228. **138:** 454, 1137, 1361, 1711, 2146, 2652, 3442, 3513, 4704, 5952, 5997, 6025, 5995. **139:** 1071, 1823, 3176, 4781, 4808, 4904, 5596, 5746, 1481. **140:** ℬ583, ℬ728, ℬ907, ℬ987, ℬ1074, ℬ1106, ℬ1136, ℬ1396, ℬ3201, 638, 1147, 2078, 2264, 2536, 2845, 2875, 3380, 3515, 3616, 4358, 4920, 5058, 5078, 5417, 5597, 5935, 5996, 6005, 1420, 5130, 5774, 1860. **141:** 138, 551, 572, 1355, 1951, 2024, 2025, 3178, 3377, 3672, 3711, 4054, 4367.3, 4568, 4952, 5088, 5095, 5114, 1898, 3102. **142:** ℬ539, ℬ803, ℬ1254, ℬ2988, 98, 1563, 3032, 3344, 3719, 4015, 4426, 4479, 5288, 5339, 5558, 6020, 5027, 1982. **143:** ℬ931, ℬ1142, ℬ1451, ℬ1868, 1401, 2268, 2680, 3373, 3502, 3505, 3771, 4229, 4276, 4654, 4675, 4748, 4959, ℬ496, 5309, 1818, ℬ444. **144:** ℬ471, ℬ981, ℬ1864, 781, 1274, 1673, 1676, 1907, 2529, 2541, 3113, 3138, 3696, 3887, 4462, 4766, 5319, 5479, 5873, 2214, 2689. **145:** 336, 889, 1238, 1668, 2074, 2088, 2876, 3408, 3436, 3492, 3841, 4422, 4811, 5069, 5118, 5414, 5548, 6075. **146:** ℬ269, 308.1, 869, 1677, 2948, 3275, 3281, 4200, 4644, 5697, 5708, 14, 1979, 4472, ℬ170. **147:** ℬ2135, 1223, 1799, 1831, 2135, 2565, 2626, 3490, 4133, 4817, 5399, 5541, 5595, 5868, 1897, 2692, 5628. **148:** 633, 1145, 1280, 1398, 1406, 1845, 2602, 3187, 3500, 3831, 4518, 4548, 4692, 4732, 4865, 5112, 5427, 5933, 873, **149:** ℬ1536, 2517, 2656, 4259, 4605, 4942, 5104, 5634, 6069, 6074, 3674, 57. **150:** ℬ159, ℬ161, ℬ454,

ℬ540, ℬ541, ℬ885, ℬ965, ℬ1090, ℬ1183, ℬ1625, ℬ1942, ℬ1961, ℬ2009, ℬ2026, ℬ2043, ℬ3032, ℬ3111, 53, 114, 305, 542, 1503, 1678, 1796, 1802, 1939, 2516, 2728, 2932, 3011, 3210, 3248, 3611, 3624, 3886.1, 4074, 4382, 4538, 4738, 4784, 5026, 5076, 5990, 6031, 6078, 2556. **151:** ℬ137, 1562, 2734, 3426, 4141. **152:** ℬ247, 70, 219, 575, 1451, 1524, 1846, 2138, 2308.1, 3475, 3559, 3675, 3888, 4199, 4227, 4486, 4607, 4671, 4819, 5145, 5239, 5899, 2607. **153:** ℬ899, 437, 1037, 1379, 1507, 1748, 1798, 2623, 2789, 3089, 3196, 3623, 4055, 4086, 4183, 4668, 4830, 5494. **154:** 433, 1362, 1589, 1684, 1685, 1841, 2012, 2150, 3061, 3531, 4504, 4754, 5247, 5620, 5902, 1817, 1973, 5676, 1861. **155:** 34, 425, 1530, 1927, 2418, 2651, 4309, 4525, 4566, 4731, 4954, 5028.1, 5317, 5523. **156:** 853, 1189, 1452, 1938, 2749, 3117, 3440, 3479, 4146, 4189, 4237, 4446, 4494, 4564, 4565, 4934, 5473, 5512, 5655, 5831, 5973, 6017, 2226, 3876. **157:** ℬ912, 884, 1061, 1310, 1384, 3086, 3282, 3840, 4057, 5139, 5187, 5217, 5525, 5971, 5999. **158:** 362, 1435, 1464, 1585, 1696, 3491, 3980, 4310, 4750, 4785, 5302, 6046. **159:** 577, 819, 1363, 2013, 2654, 3104, 3516, 4142, 4367.1, 4811, 5084, 5650, 5909, 296, 1032. **160:** ℬ85, ℬ1080, ℬ1127, ℬ1132, ℬ1395, ℬ1612, ℬ1947, ℬ3302, 36, 309, 637, 1508, 1786, 3025, 3068, 3114, 3186, 3203, 3578, 3646, 4029, 4075, 4201, 4311, 4331, 1367 2, 4562, 4867, 5568, 5575, 5906, 5950, 5960, 5977, 6000, 6088, 5103, 2691, 3379, 5491. **161:** 64.1, 308, 777, 900, 1129, 1124, 1460, 1579, 2209, 2265, 2321, 2370, 2428, 2790, 4071, 4214, 4351, 4456, 4478, 4596, 4705, 5457, 5477, 6059, 3880. **162:** ℬ980, ℬ2992, 499, 923, 1142, 1881, 2080, 2700, 3205, 3433, 3439, 3843, 4553, 4610, 4678, 4745, 4995, 5238, 5421, 5726, 2675, 5404. **163:** 82, 1784, 1785, 1838, 2313, 2698, 2729, 3103, 3118, 3553, 3949, 4222, 4656, 4751, 5251, 6068, 5100. **164:** 868, 1240, 1528, 2010, 2141, 3177, 3487, 4037, 4185.1, 4747, 4773, 5086, 5186, 5213, 5514, 5874, 6084, 1816, ℬ480, 1034, 5573, 5613. **165:** ℬ81, ℬ735, ℬ926, 353, 434, 1141, 1364, 1404, 1425, 1717, 1849, 1998, 2360, 2540, 2621, 3142, 3288, 3560, 3885, 4164, 4240, 4303, 4714, 4803, 5240, 5426, 5867, 5932, 6021, 5796. **166:** 1505, 1815, 3148, 3632, 4365, 5212, 5927, 1751. **167:** ℬ292, ℬ3146, 1913, 2019, 2106, 2193, 2603, 4481, 5498, 5542, 5577, ℬ2980, 1864, 4716, 5901. **168:** ℬ1221, ℬ2949, 1115, 1485, 1675, 1675.1, 2372, 2657, 2828, 3093, 3486, 4496, 5329, 5561, 1143. **169:** ℬ774, 1290, 1605, 1797, 1914, 2188, 2549, 3605, 3609, 4028, 4474, 5081, 5144, 5540, 6127. **170:** ℬ175, ℬ248, ℬ288, ℬ359, ℬ929, ℬ1079, ℬ1220, ℬ1610, 436, 639, 776, 1200, 1239, 1446, 1486, 1977, 2467, 3141, 3521, 3533, 3612, 3954, 4646, 4653, 4667, 4767, 4943, 5171, 5172, 5196, 5330, 5659, 5790, 5818, 6022, 6042, 1116, 2079. **171:** 297, 438, 523, 1583, 2489, 2755, 3488, 3647, 4188, 4255, 4574, 4872, 5352, 5651, 5718, 5798, 6032, 1256. **172:** 2235, 2609, 4230, 4809, 5469, 5648, 5652, 5668, 1273, 5508, 5578, 2604, ℬ226, 2541. **173:** 76, 3232, 4143, 4190, 4291, 6030, ℬ3011, 5271. **174:** ℬ1219, 1394, 1671, 1776, 1840, 2075, 2213, 3707, 3845, 4127, 4182, 4498, 5180, 5207, 5298. **175:** ℬ88, ℬ678, ℬ1198, ℬ1209, ℬ1313, ℬ1618, 501, 903, 2210, 2266, 2378, 2476, 3206, 3409, 3527, 3625, 3830, 4356, 4368.9, 4369.1, 4534, 4546, 4820, 4829, 4873, 5082 2, 5489, 5511, 5562, 5657, 5704, 5747, 5797. **176:** 650, 2369, 3106, 3376, 3511, 3842, 5242, 5267, 5744, 5912, 1981, 2587. **177:** ℬ1346, 1836, 2659, 2752, 5572, 5777, 5291, 6139, 2611. **178:** ℬ164, 200, 1124, 1148, 1285, 1577, 1911, 2605, 3514, 3541, 3832, 4027, 4265, 5269, 5278, 5535, 5547, 5645, 5711, 5783, 5785, 6104, 1953, 5724. **179:** 549, 567, 1835, 2660, 3846, 4656 1, 4877, 4910, 4956, 5826, 6125. **180:** ℬ504, ℬ544, ℬ565, ℬ645, ℬ906, ℬ916, ℬ950, ℬ1126, ℬ1166, ℬ1195, ℬ1554, ℬ3143, ℬ3288, 703, 1198, 2260.1, 2422.2, 2512, 3023, 3378, 3386, 3896, 4215, 4717, 5311, 5549, 5658, 5660, 5666, 5945, 6076. **181:** 1188, 2433, 2185, 2599, 3105, 3181, 4014, 4554, 4614, 5320, 5395, 5519, 5546, 5581, 5885, 6137, 2534, 5245. **182:** ℬ519, ℬ1182, 58, 503, 1879, 3495, 3579, 3837, 4878, 5042, 5090, 5530, 5649, 5946, 5949, 6131. **183:** 71, 2132, 2547, 2606, 3430, 5248, 5430, 5627, 5905, 5928, 6175, 1974. **184:** ℬ849, 679.1, 767, 1182, 1448, 2546, 2794, 4556, 4929, 5803,

6082, 2616. **185:** ℬ370, 92, 629, 1586, 1661, 1882, 1915, 2426, 3108, 4030, 4134, 4825, 5107, 5137, 5463, 5559, 5631, 5998, 6002, 6029, 1266. **186:** ℬ397, ℬ3304, 550, 1777, 2184, 2526, 2990, 3626, 3644, 4396, 5889, 6050, 6122, 1975, 5175. **187:** 94, 840, 1781, 2076, 2270, 2374, 3869, 4428, 4711, 5109, 5228, 5925, 6018, 6085, 3111, 3483, 4598, 5742, 5493. **188:** ℬ923, ℬ1201, 1120, 1281, 1359, 1750, 2186, 2417, 4111, 4198, 4911, 5232, 5591, 5778, 5784, 6119, 1820, 299. **189:** 147, 3049, 3064, 3526, 3698, 4424, 4501, 5046, 5331, 5428, 5467, 5516, 5564, 5788, 5793, 5898, 3407, 3582. **190:** ℬ256, ℬ639, ℬ720, ℬ721, ℬ1589, ℬ1604, ℬ3023, ℬ3206, ℬ3209, 818, 1121, 1624, 1789, 1900, 1940, 2435, 2459, 2989, 3020, 3518, 4056, 4078, 4217, 4277, 4423, 4541, 5136, 5495, 5842, 6019. **191:** ℬ1869, ℬ3291, 1129, 2375, 2421, 2479, 2808, 3084, 3645, 3871, 4073, 4593, 4631, 4722, 5125, 877. **192:** ℬ850, 707, 2630, 5234, 5290, 5792, 5931, 6138, 6106. **193:** ℬ930, 251, 265, 768, 944, 988 1, 1060, 1892, 3042, 4107, 4202, 4875, 5268, 5303, 5435, 6038, 5563. **194:** ℬ1649, ℬ1797, ℬ1895, 1779, 2373.1, 3375, 3476, 4430, 4659, 4679, 1821, 5226, 5337, 5413, 6070. **195:** ℬ1180, ℬ1555, ℬ3234, 709, 876, 2917, 3088, 3162, 3384, 3445, 4357, 4121, 4526, 4535, 4633, 5050, 5113, 5276, 5626, 5844, 5929, 5957, 6003, 6049, 6067, 6135, ℬ3298. **196:** 1780, 3873, 1221, 5189, 4136, ℬ1137, ℬ1253, 535, 2287, 2528, 3051, 4463, 5420. **198:** ℬ131, 976, 994 1, 1159, 1472, 1778, 1884, 2225, 3293, 3446, 3770, 4193, 5315, 5372, 5839, 5921, 5265, 6116. **199:** ℬ686, ℬ1194, ℬ1218, 2020, 2376, 4059, 4601, 5419, 5424, 5624, 5681, 5698. **200:** ℬ185, ℬ945, ℬ958, ℬ984, ℬ1192, ℬ1237, ℬ1605, ℬ1860, 280, 461, 1228, 2018, 2022, 2026, 2550, 2923, 3039, 3067, 4367.5, 4669, 4694, 4706, 4937, 5312, 5458, 5476, 5510, 5566, 5580, 5623, 5782, 5989, 6061, 6120, 6129. **201:** ℬ1196, 1775, 1834, 2438, 2554, 3010, 5326, 5614, 2014, 1983, 4547, 4394. **202:** 901, 1402, 1837, 2465, 3868, 4944, 5211, 5425, 5447, 6141. **203:** 1787, 2260 2, 2631, 2655, 3046, 3385, 4810, 4932, 5273, 5304, 6066. **204:** 2016, 2460, 3520, 3530, 4555, 4600, 4612, 4905, 5127, 5433, 5496, 6098, 1819, 6073, 4948. **205:** ℬ169, ℬ772, ℬ1125, ℬ3211, ℬ3290, 599, 1839, 3554, 3635, 3866, 4231, 4368.1, 4487, 4719, 4938, 5036, 5270, 5274, 5328, 5594, 5821, 5884, 6039. **206:** ℬ716, ℬ2937, 640, 676, 1405, 1584, 2017, 3085, 4191, 4489, 4871, 4936.1, 5294, 5296, 5586, 5727, 5737, 5838, 5849, 6071. **207:** 207, 2610, 2628, 3271, 4597, 4622, 4682, 4808, 5048, 5305, 5396, 5529, 5892, ℬ3293, 1722, 5443. **208:** ℬ1188, 440, 977, 2608, 3083, 4611, 4680, 5091, 5214, 5327, 5733, 5786, 6004, 6108, ℬ3024, 3901, 4619, 5035, 5887. **210:** ℬ118, ℬ1184, ℬ2707, ℬ2948, ℬ3303, 174, 486, 704, 1337, 2323, 3094, 3828, 4083, 4145, 4397, 4464, 4471, 5197, 5289, 5351, 5434, 5441, 5446, 5528, 5865, 5953, 5958, 3900, 4311, ℬ770. **211:** ℬ1677, 544, 2945, 3033, 4604, 5082.1, 6034, 6102. **212:** ℬ685, ℬ1217, ℬ3232, 1373, 2259, 3294, 3693, 3909, 4290, 4621, 4990, 5131, 5588, 5706, 5743, 6099, 6159. **213:** ℬ1208, ℬ3213, ℬ3326, 1866, 2015, 2733, 2947, 3556, 3964, 4295, 5509, 5707, 5069. **214:** ℬ89, ℬ1189, ℬ1624, 1801, 2274, 2437, 2557, 5316, 5809, 2243, 5641. **215:** ℬ775, ℬ990, ℬ1216, ℬ3103, 1281, 1609, 2377, 2624, 2625, 2747, 2806, 3673, 4232, 5080, 5871, 6023, 6079, 6153, ℬ490. **216:** 3021, 3438, 3517, 3910, 4114, 4663, 4684, 5683, 5699, 5775, 5970. **217:** ℬ2529, ℬ3301, 435, 831, 854, 1062, 2436, 2798, 2946, 4254, 4881, 5132, 5629, 6037, 6087, 6163, 1976. **218:** ℬ320, ℬ1105, ℬ1186, 3047, 3405, 4441, 4649, 4879, 5209, 5452, 5503, 5590, 5647, 5705, 5836. **219:** ℬ2527, 1421, 4031, 4457, 4625, 5536, 6170. **220:** ℬ153, ℬ968, ℬ1028, ℬ1205, ℬ1213, ℬ1215, ℬ2726, 354, 1288, 1426, 1531, 1623, 1848, 1906, 2023, 2101, 2440, 2551, 3402, 4170, 4606, 4673, 4718, 4775, 4804, 4957, 5082.3, 6142. **221:** ℬ959, ℬ1803, 2732, 3773, 4310, 4617, 5544, 5883, ℬ1490, 1126, 2429, 4252, 5191, 5290, 5979. **223:** 1070.1, 5040, 5080, 6032.1, ℬ2686, 554, 1680, 2273, 3041. **225:** ℬ317, ℬ928, ℬ3210, ℬ3214, 1114, 3202, 4081, 4530, 4691, 4840, 4944.1, 5181, 5398, 5487, 5684, 5888, ℬ516. **226:** ℬ517, 74, 708, 1110, 1903, 3007, 3432, 5079, 5630, 5987. **227:** ℬ2762, 317, 2021, 2450, 3017, 3538,

4483, 5192, 5780, ℬ638. **228:** ℬ1190, ℬ1212, 1896, 1909, 2244, 2482, 2566.2, 3234, 3610, 4685, 5182, 5297, 5439, 5926, 6052, 6148, 1474, 1493, 4876, 6083. **230:** ℬ165, ℬ299, ℬ316, ℬ895, ℬ1060, ℬ3296, 527, 878, 1800, 1828, 1912, 2430, 2754, 3443, 3853, 4613, 4890, 5462, 5468, 5654, 5685, 5736, 5791, 6092. **231:** 59, 3555, 4768, 5062, ℬ1680, 1139, 1140, 1345, 4599, 5636, 5661, 5910, 5956. **233:** ℬ424, 241, 1492, 4016, 4194, 5056, 5703, 5713, 1670, 3552, 3557, 4312, 4945, 5914, 5986. **235:** ℬ1211, ℬ2502, ℬ2763, ℬ3299, 260, 610, 1076, 1598, 1782, 1908, 2449, 3270, 3444, 4085, 4500, 5089, 5333, 6001, 6053, 6124. **236:** ℬ1214, ℬ2608, 4657, 5552, 5599, 5907, ℬ804, ℬ894, ℬ1200, 1075, 1902, 2701, 3451, 3743, 4224, 4499, 4660, 4712, 5293, 5716, ℬ254, 5041. **238:** ℬ758, 1338, 2107, 2439, 3043, 3150, 4151, 4524, 1696, 4812, 4910, 5994, 6168, 2245, 2186, 4256, 5227, 5663 **240:** ℬ1033, ℬ1804, 839, 1399, 1578, 3050, 3066, 3204, 3723, 4429, 4608, 4664, 4666, 5033, 5272, 5455, 5776, 5914, 5982, 5983 **241:** 3274, 3613, 4035, 4294, 5416, 5900, 1862, ℬ2984, 494, 5151, 5600, 5669, 6115, 1899, 706, 1514, 4084, 5190. **244:** 2183, 4616, 5259, 6077, 1211, ℬ184, ℬ1187, 565, 2443, 3185, 5471, 5632, 5804, 5896. **246:** 493, 705, 3298, 3636, 4022, 5060, ℬ487, 1679, 3406, 5203, 5343, 5822. **248:** ℬ1674, ℬ2674, ℬ3297, 904.3, 1901, 3572, 5670, 6143, 3482. **250:** ℬ380, ℬ884, ℬ916 4, ℬ1130, ℬ1202, ℬ1207, ℬ1210, ℬ1711, ℬ3059, ℬ3102, 541, 2122, 2446, 2177, 2627, 3109, 3965, 4040, 4181, 4569, 4715, 4880, 4946, 5051, 5075, 5206, 5677, 5679, 5701, 5820, 6040, 6062, 5656. **251:** ℬ2134, 1539, 1817, 4571, 5264, ℬ1073, ℬ2689, ℬ1199, ℬ2935, 3110, 3720, 4364, 4683, 4693, 5709, 6006. **253:** ℬ2754, 1093, 1374, 1529, 5047, 5964, ℬ173, ℬ1129, 1146, 1771, 3178, 6089, 6158 **255:** ℬ2621, 557, 1508, 2422.1, 2444, 4618, 4926, 6161, ℬ872, ℬ1203, 547, 995, 1523, 2658, 3018, 4527, 4638, 4642, 4737, 4931. **257:** 2420, 2792, 3292, 4502, 5807, 6117, 609.1, 1625, 1904, 3512, 1124, 5045, 5277. **259:** ℬ93, ℬ808, ℬ1204, 1920, 3381, 5567, ℬ891, ℬ1186, ℬ3311, 682, 1575, 3946, 4315, 4822, 4869, 5300, 5323, 5338, 5702, 5843, 5895, 6172. **261:** 2451, 5460, 5504, 5696, 4167, 4634, 4933, 5972, 3452, 3862, 5324, 5488, 5506, 5442. **265:** ℬ1896, 875, 2552, 3209, 5322, 5517, 5913, ℬ1193, 1883, 2791, 4602, 4690, 4909. **268:** 4251, 4963, 5642, 6037, 3280, 3434. **270:** ℬ722, ℬ904.1, ℬ982, ℬ1173, ℬ2761, 679, 2445, 2817, 3019, 3382, 3796, 4197, 4674, 5401, 5518. **271:** ℬ2704, 4949, 5513, 2151, 4246, 4629, 5295, 5911, 6154, ℬ3282. **273:** ℬ1475, 1068, 2521, 4884, 5513, 5537, 5866. **275:** ℬ1675, ℬ1729, ℬ2687, 532, 1708, 3480, 4025, 4874, 4882, 5032, 5556, 5619, 6155. **276:** ℬ221, ℬ1678, ℬ883, ℬ1679, ℬ1191, 3691, 4314, 6097, 2690. **280:** ℬ91, ℬ642, ℬ828, ℬ1880, ℬ2532, ℬ2906, ℬ3193, 1380, 1707, 2102, 2108, 3721, 4036, 4628, 4870, 5307. **282:** ℬ1312, 4603, 6036, 3208, 3481, ℬ2115, 4488. **285:** 766, 1521, 2620, 4620, 4662, 4862, 4888, 5585, 5863, 5936. **286:** 1471 1, 1191, 3052, 4182, 5448, 5464, ℬ319, 573, ℬ2971, 1455, 1494, 1773, 2488, 4861, 4060, 4615, ℬ1128. **290:** ℬ223, ℬ228, ℬ896, ℬ954, ℬ2590, 1109, 1706, 2158, 3090, 3294.1, 4620, 4689, 5672, 5748, 5811, 6055, 1059. **292:** ℬ283, ℬ2983, 3481, 4630, 1113, 2793, 1860, 185, 552, 780, 1705, 2823, 3053, 3222, 3848. **297:** ℬ2958, 5318, ℬ225, 1069, 1980, 5806, 2152. **300:** ℬ25, ℬ229, ℬ272, ℬ549, ℬ550, ℬ691, ℬ794, ℬ823, ℬ1056, ℬ1749, ℬ1915, ℬ2682, ℬ2706, ℬ3194, ℬ3195, 1106, 1407, 2442, 4887, 5023, 5502, 5550, 5686, 5721, 5834, 5943, 5981. **302:** ℬ882, 4624, 4643, 5064, ℬ1879, 1289, 4863, 1107, 1475, 2487, 5682, ℬ271, 5976, ℬ1993, ℬ2705, 5817. **310:** ℬ227, ℬ1818, ℬ3221, 1385, 4185, 4253, 4635, 4889, 5975, ℬ2952, 4886. **315:** ℬ617, ℬ3309, 1069.1, 6133, 6164, 1346, 4891, ℬ2668. **320:** ℬ315, ℬ495, ℬ1111, ℬ2815, 1127, 2702, 3586, 4665, 4885, 5030. **321:** ℬ1474, ℬ2170, 1905, 2417, ℬ2756, ℬ1746, 6060. **330:** 2480, 4594, 4595, 4627, 4636, 5022, 5029, 5722, 5082, ℬ715, 888, 2153, ℬ711, ℬ2960, 564. **340:** ℬ66, ℬ710, ℬ869, 2157, 2703, 4661, 4686, 5024, 5031. **350:** ℬ561, ℬ844, ℬ1245, ℬ2790, ℬ3106, 1112, 3022, 5819, ℬ684, 4438, ℬ3359, ℬ182, ℬ3051. **360:** ℬ1859, 4883, 5695, ℬ749, 879,

ℬ1836, ℬ2922, ℬ2925, 1411, ℬ543. **380:** ℬ92, ℬ2917, 2704, ℬ2679, ℬ1785, ℬ832, ℬ2623, 2155, ℬ2811, 5028, ℬ752, ℬ3101. **400:** ℬ657, ℬ2923, ℬ2939, ℬ2959, ℬ3015, ℬ3190, ℬ3279, 1409, 2156, ℬ548, ℬ2446, 2154, ℬ713, ℬ2503, ℬ3310, ℬ310, 199 **420:** ℬ3289, ℬ939, ℬ3238, ℬ1777, ℬ696, ℬ3202, ℬ1062, ℬ1837, ℬ742, ℬ1058, ℬ322, ℬ703, ℬ2103, ℬ3022, ℬ3237, ℬ1075, ℬ2685, ℬ753, ℬ2615, ℬ704. **450:** ℬ82, ℬ2602, ℬ1779, ℬ867, ℬ1059, ℬ2947, ℬ700, ℬ3300, ℬ2744, ℬ562, ℬ2933, ℬ1140, ℬ1835, ℬ1036. **480:** ℬ1757, ℬ1086, ℬ2675, ℬ1061, ℬ2104, ℬ948, ℬ940. **500:** ℬ94, ℬ529, ℬ616, ℬ1244, ℬ1710, ℬ3130, ℬ699, ℬ2505, ℬ2105, ℬ174, ℬ3175, ℬ535, ℬ300, ℬ2610. **550:** ℬ296, ℬ859, ℬ2788, ℬ1064, ℬ3117, ℬ2928, ℬ2257, ℬ2836, ℬ2773, ℬ193, ℬ1778, ℬ825, ℬ880, ℬ2458, ℬ328. **575:** ℬ2244, ℬ1163, ℬ2929, ℬ829, ℬ1088, ℬ2077, ℬ303, ℬ2531, ℬ3168. **600:** ℬ861, ℬ3280, ℬ542, ℬ3006, ℬ302, ℬ951, ℬ2973, ℬ301, ℬ2821, ℬ2605, ℬ2711, ℬ2634, ℬ3292. **625:** ℬ326, ℬ2063, ℬ304, ℬ1984, ℬ707, ℬ3287, ℬ2373, ℬ3167, ℬ3205, ℬ2442, ℬ881, ℬ3284. **650:** ℬ1268, ℬ1963, ℬ2680, ℬ1068, ℬ2841, ℬ279, ℬ2911. **675:** ℬ2233, ℬ2496, ℬ2080, ℬ2601, ℬ2831, ℬ3200, ℬ2833, ℬ324, ℬ2039, ℬ1017, ℬ2820, ℬ536, ℬ2824. **700:** ℬ1275, ℬ2136, ℬ2822, ℬ2832, ℬ2829, ℬ327, ℬ2131, ℬ2162, ℬ3197, ℬ665, ℬ2908, ℬ1773. **725:** ℬ2599, ℬ2924, ℬ2238, ℬ757, ℬ2513, ℬ2909, ℬ2847, ℬ2666. **750:** ℬ2907, ℬ3158, ℬ2677, ℬ3172, ℬ692, ℬ1154, ℬ3196, ℬ2239, ℬ663, ℬ2921, ℬ1873, ℬ788, ℬ2236, ℬ1153, ℬ2926. **775:** ℬ1042, ℬ2748, ℬ503, ℬ2849, ℬ2024, ℬ3161, ℬ1543, ℬ1775, ℬ1642, ℬ1541. **800:** ℬ567, ℬ810, ℬ1247, ℬ1440, ℬ1744, ℬ2837, ℬ3171, ℬ576, ℬ2671, ℬ2965, ℬ2974, ℬ2893, ℬ2008, ℬ307, ℬ669, ℬ3349, ℬ568. **825:** ℬ1066, ℬ1004, ℬ1018, ℬ2628, ℬ3224, ℬ2654, ℬ1631, ℬ2509, ℬ1979, ℬ1087, ℬ1772. **850:** ℬ1041, ℬ528, ℬ1265, ℬ2745, ℬ309, ℬ572, ℬ2616, ℬ501, ℬ579, ℬ2584, ℬ2838, ℬ2604, ℬ747, ℬ3131, ℬ2438, ℬ2777. **875:** ℬ1838, ℬ1839, ℬ1243, ℬ499, ℬ1070, ℬ2918, ℬ2692, ℬ524, ℬ2975. **900:** ℬ780, ℬ857, ℬ2253, ℬ2656, ℬ1959, ℬ937, ℬ560, ℬ3115, ℬ3116, ℬ2487, ℬ1939, ℬ571, ℬ3129. **950:** ℬ1246, ℬ1669, ℬ1774, ℬ1072, ℬ2161, ℬ2441, ℬ2500, ℬ2846, ℬ1564, ℬ2002.1, ℬ2262, ℬ3100, ℬ3013, ℬ557, ℬ2670, ℬ3017, ℬ570, ℬ3267, ℬ2716, ℬ1385, ℬ1567. **1000:** ℬ836, ℬ2598, ℬ2852, ℬ2588, ℬ2507, ℬ3305, ℬ569, ℬ789, ℬ1384, ℬ2645, ℬ3132, ℬ1862, ℬ843, ℬ3014, ℬ577. **1050:** ℬ2863, ℬ2587, ℬ3215, ℬ558, ℬ2938, ℬ1668, ℬ2776, ℬ2003, ℬ2967. **1100:** ℬ824, ℬ956, ℬ1223, ℬ1593, ℬ1870, ℬ2360, ℬ2379, ℬ2486, ℬ2865, ℬ2174, ℬ1374, ℬ1561, ℬ970, ℬ552, ℬ1694, ℬ957, ℬ2488, ℬ587, ℬ2334. **1150:** ℬ3138, ℬ573, ℬ1571, ℬ1851, ℬ553, ℬ1572, ℬ3139, ℬ2035.1, ℬ1976,1. ℬ1651, ℬ2141, ℬ2437, ℬ1348. **1200:** ℬ2313, ℬ2644, ℬ2354, ℬ1850, ℬ2263, ℬ765, ℬ876, ℬ877, ℬ3283, ℬ1407, ℬ2646, ℬ2275, ℬ1317, ℬ1518, ℬ1314, ℬ2499, ℬ1372. **1300:** ℬ1319, ℬ2380, ℬ1519, ℬ1581, ℬ2589, ℬ1316, ℬ2597, ℬ1318, ℬ1978, ℬ1520, ℬ1845, ℬ2712, ℬ2966, ℬ947, ℬ1957.1. **1351:** ℬ1846, ℬ2235, ℬ2431, ℬ2663, ℬ2659, ℬ2427, ℬ1517, ℬ2130. **1400:** ℬ2660, ℬ1325, ℬ2125, ℬ1333, ℬ2559, ℬ2355, ℬ811, ℬ2248, ℬ2561, ℬ2394, ℬ1424, ℬ2430, ℬ1801, ℬ1671, ℬ3223, ℬ2426. **1500:** ℬ858, ℬ812, ℬ1795, ℬ2860, ℬ2274, ℬ2392, ℬ1337, ℬ2270, ℬ2315, ℬ1406, ℬ2404, ℬ2400, ℬ2175, ℬ1334, ℬ2451, ℬ2472, ℬ2521, ℬ2538, ℬ2410. **1600:** ℬ2391, ℬ2557, ℬ2267, ℬ2273, ℬ447, ℬ1258, ℬ2850, ℬ1621, ℬ343, ℬ2266, ℬ2537. **1700:** ℬ2600, ℬ340, ℬ2393, ℬ2544, ℬ2328. **1800:** ℬ1904, ℬ1393, ℬ755, ℬ1619, ℬ2177. **1900:** ℬ1590, ℬ2494, ℬ1743, ℬ1977, ℬ1763, ℬ1858, ℬ2318, ℬ2222, ℬ1877. **2200:** ℬ2109, ℬ1724, ℬ2283, ℬ1821, ℬ1663. **2400:** ℬ1725, ℬ2100, ℬ1945, ℬ2434, ℬ1662, ℬ483, ℬ2232. **2700:** ℬ473, ℬ1689, ℬ1767, ℬ2099, ℬ2128, ℬ456, ℬ1690. **3000:** ℬ461.

II. BOILING POINTS

−192: ℬ337, 54, ℬ95, ℬ345, 180, ℬ1813. **−95:** ℬ204, ℬ465, ℬ97, 252, ℬ195, ℬ9, 115, ℬ351, 44. **−75:** ℬ205, ℬ17, ℬ350,

B126, B41, B125, 145, B263, B265, B485.5, 112. **—48:** 17.1, 409, 500, B68, B67, B206, B73, B26, B87, 408, B102, 232, 337. **—30:** B53, B364, 338, 263, 41, 31, 35, 95. **—18:** 686, B282, B349, B346, 13, 153, 73, 49, 781.2. **—10:** B38, B213, 65, 684, B130, 596, B141, 99, B84. **0.6:** 781.1, 685, B371, 64, 474, B98, 526, 40. **5:** B131, 687, B339, 310, 282, 63, 508, B365, 9, 1074 **10:** B40, 209, 224, B1814, 683, 42, 7, B149, 148, 283, B1811, 243, 598, 595, B4, B1672. **20.1:** 985, 208, 162, 52, 980, B96, 374, 1.1, 294, 295, 22, 1072, 597, B466, 921. **31:** B404, 569, 213, B142, 113, 983, B366, 525, 916, 615, 793, 273. **35:** B1646, 716, 278, 279, 373, 272, 1073, 982, 469, B105, 220, 986, 794, 613. **40:** 920, 28, 984, 539, 915, 530, 981, 1611, 286, B101, 313, 880, 45, 172, 913, 823. **44:** 402, 513, 918, B39, 375, 490, B797, 461, 715, 468. **47:** B99, B1149, 132, 365, 524, 447, 979, 1716. **50:** 881, 146, 749, 914, B1509, 154, 356, 520. **53:** B347, 479, 276, 529, 1050, 1087, 451. **55:** 218, 164, 237, 824, 917, 448, B111. **57:** 489, 452, 189, B352, 1712, B55, 1618, 228, 169, 465. **60:** 397, 519, 1534, 1715, 364, 133, 318, 718, 19, 1086, 3, 586 **62:** 27, 612, 1049, 357, 822, 89, 301, 1714, 3776, 1610, 891, 801, 4, 60 **65:** 50, 323, 800, 1616, B372, 517, 773, 813, 1613, 1001. **68:** 287, 518, 748, 821, 1742, 747, 1713, B59, 1533, 420. **70:** 2277, 1617, 1088, 464, 366, 728, 155, 1532, B469, 234, 112.1, 1615, 1619, 744, B207, 10. **74:** 231, 158, B42, 772, 1003, 118, 717, 83, 323.1, 820, 1048, 149, 12, 725. **78:** B513, 746, 1102, 262, 1468, B1515, 2394, B58, 332, 396, 719, 1365, 726. **80:** B348, B376, 165, 378, 752, 925, 1738, 2333, 624, 1466, 1469, 531, 2393. **81:** 130, 758, 727, 1612, B8, 168, 277, 1535, 506, 792. **83:** 623, 1252, 1537, 190, 2387, 47, 1536, 1761, 2325, 1632, 798, 117. **85:** 576, 812, 825, 1366, 1470, 1045, B109, 1328, B468. **87:** 222, 417, 669, 1008, 2388, B1810, 77, 106, 625, 811, 759, 1739, 247, 1253. **89:** 226, 1021, 1091, 1741, 462, 476, 1547, 1764, 458, 2390, 134, B1815. **91:** 670, 1101, 1740, 743, 1737, 742, 814, 17.4, 68, 122, 290.1, 652, 2391, 1002, 1019. **93:** 171, 210, 332.1, 837, 1007, 1065, 2276, 2392, 1496, 1631, 1544. **95:** 383, 1100, 2334, B390, B918, 418, B34, B35, 446, 1015, 395, 505, 2331, 26. **98:** 161, 203, 751, 810, 2326, 107, 1016, 2389, 173, 2329. **99:** 643, 1064, 2332, 1017, 1044, 791, 938. **100:** B1, B72, B455, B2010, B2044, 33, 195, 1261, 1418, 1690, 4013, 4333, 37, 2330. **101:** 642, 302, 741, 1020, 1005, 1006, 1081, 48. **102:** 496, 651, 1656, 1746, 2281, 1018, 475, 2866. **103:** 390, 2284, 2328, 391, 233, 1004, 181, B377, 614, 1099, 121, 85. **105:** B128, B530, B870, 135, 937, 1046, 1047, 2283, 1043, 1054. **106:** B215, 192, 588, 2240, 2278, 1630, 254, 1014, 2943, 568, 1699, 2279, B210, 790. **108:** 108, 137, 166, 668, 695, 816, 2238, 965, 735, 1042, 2933, 2935. **110:** B619, B1822, 183, 284, 522, 2510, 2239, 123, B467, 495, 2112, 2275, 1760, 2944. **111:** 2282, 1660, 1655, 1629, 2413, 1694, 1653, 11, 967, 1650, 2280, B103, B382, 159, 100. **114:** 714, 1082, 1085, 2057, 2241, 2934, 2938, B488, 414, 1637, 999, 242. **115:** 239, 355, 663, 1193, 2942.1, 870, 908, 1080. **116:** B37, 67, 110, 205, 590, 660, 2058, 2937, 587, 998, 926, 2936, 1658, 334. **117:** 300, 379, 2059, 2414, 2871, 756, 789, 1640. **118:** B233, B798, B1714, 274, 563, 667, 307, 711, 1084.1, 1097, 2111, 2940, 4386, 212, 2327, 1652, 2942. **119:** 177, 533, 1639, 1700, 2194, 2382, 895, 1084. **120:** B1825, 347, 488, 827, 1733.1, 2452, 2891, 1755, 2873, 755, 2825, 2869, 939, 90, 2364. **121:** 377, 1041, 1635, 1733, 2237, 2975, 1654, 1634. **122:** B266, B426, 1725, 2974, 1734, 775. **123:** 387, 2874, 3048, 1659, 1649, 2345, 2868, 1763. **124:** 562, 927, 1636, 1662, 2827, 712, 713, 512, 2941, 2355. **125:** B232, B436, 236, 419, 578, 896, 1053, 2824, 2872, 17, 1026. **126:** 340, 470, 1096, 1701, 266, 1651. **127:** B250, 367, 608, 754, 1638, 1657, 91, 1040. **128:** 966, 1083, 1443, 1727, 1762, 1769, 2363, 2384, 2870, 745, 227. **129:** 1556, 1633, 1702, 973, 2867, 1098, 1546. **130:** B1893, 78, 253, 411, 970, 1212, 2397, 2796, 5350, 1648, 139, 924. **131:** 589, 1028.1, 1520, 2354.1, 1558, 1079, 384, 487, 184, 1730. **132:** B845, 404, 1691, 2359.1, 2398, 3350, 1307, 176, 3295, 3347, B1147, 2296, 471, 1695, 1732, 2826. **135:** B23,

B164, 119, 571, 1548, 1723, 1726, 2357, 2795, 339, 799, 2362, 534, 2797, 3349. **136:** 38, 774, 2347, 481, B450, B1552. **137:** B360, 385, 628, 929, 1509, 1604, 2976, 3348, 1545, 1511, 2683, B218, 2686, 1078. **138:** B57, B460, 410, 817, 1510, 1730.1, 2359, 1587, 2985, 969.1, 3272. **139:** B353, B693, 992, 1557, 2685, 3297, 3351, 626. **140:** B120, B177, B378, B459, 304, 412, 635, 803, 2100, 2110, 2843, 2973. **141:** 972, 907, 1756, 2409, 2411, 450, 3353, 358. **142:** 721, 1754, 2298, 2412, 2979, 3273, 3808, 2396, 341, 1595, 2351, 2926 **143:** 910, 2197, 2295, 2453, 3321, 1445, 3352, 1444, 2344, 2902, 2361. **144:** 319, 431, 942, 2349, 2684, 3805, 658, 403, 737. **145:** 657.1, 697, 989, 2833, 2356, 2415. **146:** 191, 275, 449, 644, 782, 968, 1512, 2538, 140. **147:** 2929, 3800, 2961, 2353, 3827, 421, 1736. **148:** 386, 429, 466, 659, 850, 969, 1052, 1070, 1434, 2401, 3365, 849, 2316, B1752, 2905, 2199. **149:** 327, 693, B284, 2360, 3296, 3323, 2977, 3322. **150:** B237, B915, 121, 636, 2304, 2348, 2402, 2108, 2915, 3320, 3807, 221, 18, 2865, 3354, 594. **151:** 1667, 2288, 2957, 3826, 460, B62. **152:** 704, 2299, 2404, B2, 1735, 160, 2964, 258. **153:** B60, 1596, 3223, 815, B362, B367, 681, 2358, 329. **154:** B452, 406, 1028, 2951, 2971, 3817, 724. **155:** 186, 757, 1482, 2343, 2517, 2836, 2898, 3119, 3212, 3331, 3621, 2419, 2399, 1603, 1728, 2163, 2912. **156:** 388, 1051, 1597, 1692, 3372, 3803, 1204, 1767, 2202, 1541, 2904. **157:** 753, 2196, 2906, 3727, 3895, 3229, 3963. **158:** 691, 833, 2407, 2853, 604, 3937, 3962. **159:** B919, 648.2, 2772, 3369, 3741, 3802, 3994, 2039, 1770, 2381. **160:** B799, 43, 256, 433, 497, 498, 545, 733, 945, 963, 1165, 1687, 3801, 3888, 3995, 3998, 2901, 2952, 2963, 3997. **161:** 593, 2289, 2084, 3956, 389, 807, 1627, 2856, 2962, 859, 4001. **162:** B208, 111, 453, 978, 1022, 1718, 1876, 2953, 3224, 3226, 3996, 2040, 142, 969, 2041, 2297, 2405, 3225. **163:** 620, 622, 1602, 1765, 2818, 2972, 361, 631, B472, 723, 2894, 1011. **164:** 101, 1593, 2958, 2970, 2987, 3809, 3818, 5213, 2198, 3228, 3894. **165:** B219, 163, 359, 1517, 1563, 1724, 2776, 2983, 3819, 2200, 582, 1588, 1880, 762. **166:** 424, 688, 1542, 1543, 1550, 1641, 1757, 1886, 2201, 2878, 2955, 3362, 3368, 4000, 662, 2903, 2928, 2907, 3366, 3999. **167:** B47, 413, 2859, 3333, 3811, 2406, 848, 2911. **168:** 585, 1056, 2778, 2893, 2959, 3717, 3892, 2365, 3960, 1055, 2338, 3334, 3726. **169:** 779, 1519, 1572, 2339, 2366, 2854, 2880, 2960, 3958, 2909, 3893, 1430, 3976, 3230. **170:** B974, B1658, 661, 773.1, 826, 943, 951, 1693, 2881, 3340, 3961, 893, 1029. **171:** 285, 1518, 2982, 3358, 3816, 3959, 2719, 3363, 2164, 619. **172:** B383, 2, 84, 328, 1572.1, 2722, 2779, 2855, 2858, 2965, 3318, 3327, 3369, 3055, 2910, 4007, 2831, 2896, 1743, B214. **173:** 847, 1230, 1231, 1314, 1559, 2008, 2925, 3367, 346, 950. **174:** 103, 422, 689, 1010, 2303, 2342, 2379, 2718, 2774, 2834, 3993, 3357, 3736, 2885. **175:** B1610, 988, 2341, 2468, 3120, 3361, 3725, 3728, 3815, 3821, 3825, 783, 3345, 2403. **176:** B1597, 1067.1, 2340, 2890, 3728.1, 3806, B76, 3902, 2721, 3121, 3227, 1013, 2908. **177:** 235, 244, 797, 2837, 3810, 3820, 2720, 3247, 230. **178:** 696, 901, 1105, 2773, 2884, 3810, 3330, 2380. **179:** 196, 1229, 1569, 1592, 2883, 5317, 2038, 2001, 2290, 2956. **180:** B209, B508, B1648, B2932, 29, 492, 666, 933, 1561, 1697, 2066, 2895, 2986, 3355, 3724, 4210, 4516, 5052, 5081, 5329. **181:** B515, 185, 1341, 2337, 2951, 948, 2030. **182:** 734, 1413, 1608, 2981, 3326, 3730, 3823, 3054, 621, 1888, 2841, B1864. **183:** B81, 170, 423, 2320, 2838, 3731, 3733, 1431, 2032, 2031. **184:** 400, 796, 921.1, 1622, 2195, 3858, 1442, 3729, 2924, 3258, 3329. **185:** 92, 617, 931, 2888, 3055, 3734, 3824, 5637, 2335, 3891, 3339, 3332, B470. **186:** 202, 381, 674, 1704.1, 2317, 1570, 1389, 2302. **187:** B1673, 993, 1012, 1417, 1646, 1933, 2809, 3335, 2770, 2889, 3324. **188:** B2112, 351, 928, 1331, 2777, 4165, 151, 1333, 269. **189:** 480, 510, 930, 1390, 1804, 2799, 3639, 2887, 4116, 6, 156, 2572, 2641, 605, 3245. **190:** 116, 348, 1265, 1489, 1642, 2645, 2988, 3576, 3814, 4006, 4012, 3342, 1571, 3246, 1885, 2160. **191:** 371, 2642, 3336, 4407, 4419. **192:** B385, 558, 1499, 1601, 2316, 2318, 2319, 3127, 3244, 3303, 1568, 3337. **193:** B216, 46, 528, 1805, 4166, 3890, 136, 1647, 246.

194: B1894, 600, 1551, 2567, 2756, 2966, 3356, 3986, 4118, 2190.
195: 932, 1296, 2189, 2192, 2386, 2512, 2861, 2862, 3360, 3371, 3732, 3852, 3916, 3961, 109, 4156, 2191, 3189, 2568, 2569, 2590.
196: 1200.2, 1560, 1772, 2050, 2643, 2851, 3302, 3311, 3758, 2203, 3786, 1200 3, 720, 2004, 3304. **197:** 806, 1599, 1645, 2724, 3739, 4178, 5083, 5977, 1859, 264. **198:** 331, 690, 702, 858, 2310, 3760, 3950, 4906, 2771, 2840, 3915, 936, 3277, 2496, 2309. **199:** 606, 2029, 2495, 2640, 3319, 3897, 3914, 2589. **200:** B361, B1812, 556, 501, 740, 771, 1515, 2003, 2314, 2614, 2781, 2852, 3299, 3744, 3844, 3861, 3949, 4117, 5212, 4897, 2206, 2204. **201:** 370, 3905, 3906, 2162, 2846, B491, 548, 971, 1613, 2644, 2900, 3305, 3912, 5326, 4115, 2571, 2161. **203:** B1148, 570, 1500, 3575, 2205, 2588, 368, 3982, 2383, 382, 608 2, 882, 1857, 2062, 2505, 2570, 2637, 2897, 2922, 3653, 3740, 3911, 3983, 3904. **205:** B235, 763, 1250, 1251, 1681, 1720, 1856, 2713, 3301, 3306, 3788, 3921, 1385, 4412, 2174, 2301, 1600, 3735, 2639, 3882, 2159, 3984. **206:** 729, 911, 1067, 1283, 1803, 2635, 2737, 2768, 2769, 1137, 2763. **207:** 892, 2636, 2767, 3638, 3883, 3918, 3978 4, 4122, 3637, 3881, 3972, 2710, 1644, 3341. **208:** 150, 561, 809, 1554, 2434, 2775, 3150, 3903, 3985, 4365.1, 2311, 3992, 1177, 2687, 4121, 4416, **209:** 1009, 1552, 3188, 3276, 3860, 3907, 3975, 4120, 3846. **210:** B1817, 87, 1233, 2185, 2917, 3316, 3917, 3921, 3977, 3989, 4413, 3759, 1375, 2267, 1317. **211:** 828, 2064, 2814, 2848, 3157, 3978, 4005, 2063, 2700, 3260, 120, 1234, 1089. **212:** B211, B1827, 731, 2261, 2707, 3151, 3235, 3574, 3927, 3928, 3973, 3974, 4155, 3152, 3259, 3470, 2739, 4404. **213:** B356, 473, 182, 1176, 2581, 3128, 3249, 3268, 4130, 3154, 3149, 3901. **214:** 511, 1315, 1858, 1961, 1905, 1906, 2035, 2506, 2711, 3261, 3926, 3987, 4418, 1349, 2633, 1249. **215:** B2931, 975, 1703, 1887, 2711, 2765, 3316, 3361, 3935, 3936, 3947, 2634. **216:** 760, 852, 1768, 2758, 2764, 3129, 3655, 4010, 4129, 4411, 4132, 2350, 2766, 2847, 3469, 3789, 3133, 2760. **217:** B679, 405, 1205, 1316, 1553, 2507, 2716, 2717, 2718, 2759, 2849, 3156, 3263, 3661, 4157, 860, 2632, 3923, 3494. **218:** 2292, 2575, 2705, 3132, 3834, 3855, 4011, 4366, 4367, 4934, 3265, 1918, 2812, 2819, 3081. **219:** B1796, 415, 416, 1175, 1206, 1844, 2127, 2165, 2712, 3236, 3922, 3966, 4175, 2709. **220:** B381, B800, 97, 126, 315, 316, 761, 922, 1235, 1149, 1704, 2047, 2269, 2821, 3155, 3238, 3676, 3765, 3782, 4254, 1376, 4378, 1480, 5846, B286, 1870, 2638. **221:** B268, 342, 399, 1201, 2490, 2492, 2714, 2762, 2949, 4585, 3678, 3968, 2725. **222:** 238, 372, 2081, 3131, 3262, 3267, 3967, 311, 2577, 2784, 4161, 2618, 2715, 673, 2351, 2757. **224:** 428, 987, 2216, 2257, 3219, 3763, 3856, 3859, 4176, 6608, 1327. **225:** 553, 1794, 2045, 2046, 2131, 2813, 3071, 3131, 3257, 3753, 4089, 4131, 5815, 2708, 3920, 478, 2668, 2927, 188. **226:** 2464, 2761, 2930, 3243, 3654, 3677, 3755, 4100, 4171, 4373, 4170, 3679, 3239, 4367.7. **227:** 86, 2084, 2294, 1388, B1677, 2255, 3285, 863, 866. **228:** 1498, 2051, 3240, 3680, 3683, 4097, 4138, 4172, 3313, 1868, 4177. **229:** 2122, 2178, 3286, 3757, 3849, 3908, 3942, 4139, 2786, B1802, 1376. **230:** B451, 314, 315, 555, 603, 795, 1092, 1600, 1916, 2042, 2123, 2322, 2579, 3857, 4119, 4147, 5853, 3660, 3283, B386. **231:** 1377, 2082, 2688, 3170, 1002, 3598, 5859, 3684, 2454, 3756. **232:** 125, 1058, 1872, 1917, 2498, 3685, 4750, 3787, 3662, 784, 3241, 4581. **233:** 1744, 1810, 2293, 2864, 3191, 3264, 3663, 3847, 1237, 2503. **234:** 1458, 1869, 1871, 2043, 3266, 3863, 4180, 4587, 3591, 157. **235:** B384, 629, 708, 2011, 2033, 2044, 2048, 2049, 2271, 2501, 2522 1, 3173, 3242, 3209, 3656, 3764, 5200, 3769, 3648, 1216. **236:** 3659, 4159, 1297, 2931, 1200, 1795, 2525, 2803, 3125, 3190, 3312, 3657, 3784, 3929, 4090, 4099, 5021, 3237, 2899, 4367.6, 3037, 3752 **238:** 265, 583, 1298, 1758, 1812, 1957, 2083, 3221, 3284, 2120, 1207, 1791, 2523, 3682, 2120.1, 1000, 4088, 2671. **240:** B368, B519, B1032, B1180, B1513, 138, 1919, 1924, 2005, 2064, 2115, 2117, 2119, 2457, 2804, 3171, 3690, 3885, 4096, 4547, 5033, 5102, B1797, 1875, 3785. **241:** 2125, 2544.1, 2738, 3251, 4368 8, 2052, 1217, 1576, 1955, 2243, 2578, 3174, 3790, 4374, 3098, 4337. **243:** 1432, 1790, 2124, 2176, 2163, 2522, 2822, 2863, 3036, 3589, 4038,

4346, 5860, 3941, 1483. **244:** 1477, 3070, 4091, 2175, 3592, 1179, B2113, 104, 994, 1321, 1414, 1472, 2121, 2126, 2218, 2524, 2541, 4039, 4326, 4408, 5066, 3701. **246:** 941, 1136, 1137, 1949, 2669, 3291, 3767, 4849, 3754, 1318, 1688, 1792, 1873, 2497, 2580, 2740, 2742, 3783, 5896, 2745. **248:** 312, 802, 1811, 2472, 2787, 3590, 3884, 4092, 4098, 4968, 2007, 2244. **250:** 805, 901, 1464, 1745, 1956, 2829, 3124, 3748, 4828, 4900, 5259, 4095, 4338, 672, 2666, 3943. **251:** 1322, 1368, 1369, 2217, 3069, 3192, 4283, 5718, 4160, 883, 1320, 1479, 2172, 2646, 4856, 4979, 3548. **253:** 1178, 1291, 1793, 2849.1, 3081, 3135.1, 3666, 1236, 1135, 2504, 3328, 4339, 4827, 2788, 4140, 3199, 4219. **255:** B387, B1828, 1502, 2249, 2544.2, 3035, 3220, 3547, 3689, 3793, 3933, 4127, 4363, 3001, 3250. **256:** 350, 974, 1173, 1203, 2508, 3005, 3172, 3874, 4048, 4978, 1202, B254, 1527, 2118, 3931, 3953, 2245, 3028. **258:** 1169, 1326, 2099, 2667, 2890, 3455, 3979, 4041, 4282, 4843, 3126, 2723, 2425, 3289, 3709, 4049, 4211, 4415, 4973, 5568, 2585, 2247, 3600. **260:** 178, 565, 730, 861, 1325, 1367, 1822, 2100, 2116, 3214, 3746, 3791, 3836, 3930, 4491, 4500, 4974, 4857, 5005, 5347, 3002, 1959. **261:** 571, 1201, 4045, 4308, B253, 3004, 3546, 3587, 3593, 3833, 3835, 4315, 4490, 4567, 2801, 1247, 4832, 607. **263:** 1077, 1317, 1626, 1958, 2474, 2586, 3688, 4380, 4976, 1304, 4044, 4093, 4975, 3456, 4279. **265:** B1129, 2253, 2658, 3076, 3146, 3606, 3980, 4046, 4347, 4351, 4365, 4391, 4590, 3650, 2584, 3651. **266:** B1676, 1171, 1506, 4280, 3100, 3044, 560, 1481, 3077, 3558, 3643, 4128, 4106, 4493, 3667, 5166. **268:** B1649, B1865, 2677, 3003, 3139, 3140, 4205, 3981, 309, 2518, 4852. **270:** B2114, B2606, 502, 765, 1246, 1970, 2078, 2622, 3773, 4032, 4043, 4064, 4983, 4033, 4543, 5018, 1979. **271:** B770, 2073, 2367, 3771, 4062, 4542, 5142, 1324, 3078, 4757, 4758, 3099. **273:** 3671, 4349, 2670, 1574, 2251, 2676, 3778, 4042, 5156, 2736, 3747. **275:** 2587, 3000, 4034, 4184, 4238, 4972, 3248, B1674, 1157, 2424, 3652, 4267, 4304, 1296, 1415, 4359, 2423, 3702. **277:** 889, 1119, 2098, 4492, 4707, 5003, 5141, 4218, 1158, 1807, 3620, 3792, 4778, 3703, 2519, 2731, 3147. **280:** 769, 1255, 1581, 2207, 2250, 2616, 2729, 2850, 3130, 3507, 4173, 4278, 4327, 4368 2. **281:** 871, 4760, 3153, 1580, 3412, 3454, 3615, 4163, 4207, 4297, 4544, 4318. **283:** 2619, 2578, 3710, 4319, 2651, 4756, 2431, 1259, 1260, 2254, 2596, 2617, 2697, 3420, 3542, 4533, 5153, 4317. **286:** 1397, 2780, 3417, 3508, 4268, 4299, 1416, 1258, 897, 1180, 3183, 3290, 4269, 4508, 886, 4759, 5167, 3414. **288:** B74, 592, 1133, 3006, 4779, 1384, 3543, 3595, 4195, 4761, 5244, 1256, 3113, 3497, 5002. **290:** B46, B379, 515, 573, 1254, 2072, 2173, 2473, 2647, 3198, 3670, 3771, 4531, 4635, 4784, 4802, 5286, 5490, 5681. **291:** B255, 1926, 4047, 1215, 4023, 5169, 4532, 4996, 3642, 3692, 4050, 4260, 4845, 1513. **294:** 4322, 4381, 4997, B230, 1478, 2648, 4158, 4439, 4762, 5160, 5017. **296:** 1154, 3194, 4352, 4324, 3498, 4019, 5110, 4325, 4793, 4225, 4017, 3115, 4362, 4777, 4939, 5340, 4507, 1273, 4266, 4930. **300:** B363, B695, 1213, 1985, 2167, 2416, 3010, 3075, 3094, 3158, 3197, 3526, 3630, 3696, 4029, 4109, 4473, 4512, 4529, 4650, 4789, 4838, 4867, 4987. **301:** 1817, 3550, 5020, 3016, 1272, 1921, 3596, 4262, 4270, 4466, 4967, 3195, 3945, 5260, 2649. **304:** B883, 946, 3473, 4323, 3499, 3216, 4020, 4305, 4441, 4447. **306:** 4018, 4448, 4794, 3551, 3196, 4708, 3466, 4240, 4846, 1419, 4988. **310:** B896, 4261, 5010, 5100, 1120, 4329, 1505, 1925, 4790, B444, 4458, 4697. **315:** B1342, B1895, 1163, 1165, 1166, 1982, 4451, 4916, 4725, 4734, 4726. **317:** B66, 1573, 1981, 2462, 5391, 4442, 1271, 4204, 4917. **320:** B1804, 1149, 4212, 4263, 4715, 4791, 5816, 2844, B894, 4506, 4724, B88, 5861. **325:** 4393, 4687, 5037, 5072, 4994, 1110, 4243, 5059, 4727, B1679, 4220, 5607, B251, 1123. **330:** B2116, 831, 4431, 5074, 5486, 4792, 4914, B2115, B1678, 808, 1923. **335:** 1922, 1925, 4915, 5044, 1525, 4203, 5689, B91, 2421. **340:** B496, B1624, 1526, 4214, 4242, 4271, 4460, 5043, 5135, 5262, 4652, B2117, 4425, 4244, 4649, 4455, 4728, 5168, 5335. **345:** 2212, 5038, 4434, B1675, 4672, 4902, 5521. **350:** 4515, 5183, 5184, 5747, 4427, 4435, 4436, B898, 4285, 4211, 4465, 5520, 5402. **360:** 1991, 3042, 3307, 4137, 4622,

4676, 4912, 4913, 5193, 5491, 5746, 5616, 4287, 4892, 5281, 4012. 371: 4215, ℬ471, 4514, 5053, 4249, 4620, 6010, ℬ1869, 5379, 5887, ℬ882, 4907, 5173, 5306, 5055. 400: ℬ958, ℬ1798, ℬ2959, 4690, ℬ292, 4286, 5395, ℬ226, ℬ2608, ℬ89, ℬ480, 5494. 421: ℬ92, 5883, 5274, ℬ716, 4626, 5863, 4722, ℬ1075, 5172, ℬ316, 5264. 452: 5493, ℬ320, 4637, 4636, 5508, ℬ1749, ℬ170, ℬ223. 500: ℬ322, ℬ769, 5817, ℬ224, ℬ228, 5695, ℬ227, ℬ678, ℬ271, ℬ2105. 600: ℬ193, ℬ1879, ℬ490, ℬ487, ℬ753, ℬ752, ℬ881. 707: ℬ272, ℬ832, ℬ495, ℬ749, ℬ696, ℬ700, ℬ703, ℬ2936. 916: ℬ543, ℬ548, ℬ529, ℬ829, ℬ825, ℬ940, ℬ779, ℬ1268, ℬ2613. 1230: ℬ499, ℬ3283, ℬ2610, ℬ528, ℬ951, ℬ3284, ℬ2680, ℬ3205, ℬ3287, ℬ2917, ℬ2926, ℬ3200, ℬ947, ℬ2605, ℬ939, ℬ2924, ℬ2668, ℬ2677, ℬ3197. 1400: ℬ2499, ℬ3196, ℬ2131, ℬ2671, ℬ2921, ℬ2769, ℬ2918, ℬ1337, ℬ1059, ℬ1870, ℬ1334, ℬ2500. 1670: ℬ2604, ℬ2670, ℬ1858, ℬ341. 3800: ℬ1619, ℬ1721, ℬ1799, ℬ481, ℬ1805, ℬ1689, ℬ1690.

III. DENSITY

A. Liquids

0.415: 54, 409, ℬ102, 1072, 1073, ℬ406, 1716, 1715, 980, 1713, 1714. 0.670: 2392, 2394, 915, 916, 2387, 1610, ℬ407, 2389, 917, 2301, 1534. 0.692: 1613, 2933, 525, 823, 918, 1617, 22, ℬ110, 914, 2939, 824. 0.712: 1619, ℬ409, ℬ414, 822, 1535, 2331, 3354, 524, 2334, 2936, 1761, 2940, 3995. 0.724: 2873, ℬ125, 1764, 2279, ℬ412, 1086, 4000, 3994, 794.1, 3999, 821, 794, 1760. 0.740: 820, ℬ415, 3351, 4178, 396, ℬ416, 2985, 1711, 3993, 3957, 1101 0.750: 1615, 1100, 1738, 1737, 979, 1739, 2975, 4112, 3372, 4587. 0.760: 669, 4586, 479, 2974, 4165, 2211, 2328, 2330, 2413, 1001, 4856, ℬ418, 1099, 1762.1, 3323, 4012, 4111, 2869, 2973. 0.771: 2868, 2987, 5018, 3365, ℬ420, 4006, 4849, 5167, 913, 1632, 2419, 4418, 5260, ℬ421, 1612, 2867. 0.781: ℬ422, 208, ℬ423, 168, 395, 506, 3320, 1049, 262, 792, 5156. 0.790: 3960, 3297, 5377, 60, 1003, 3961, 301, 667, 718, 448, 2825, 2284, 3812. 0.800: 790, 1769, 2281, 972, 1603, 2827, 973, 3811, 1639, 3295, 505, ℬ411, 2382. 0.805: 719, 880, 1366, 1544, 2283, 2345, 417, 791, 2955, 1081, 1084, 1602, 2282. 0.810: 789, 1537, 1084.1, 1630, 2327, 2898, 2965, 1754, 3895, 3959, 1640, 1730.1, 2320, 2317, 313, 1083, 2396, 2397, 2872. 0.817: 717, 1078, 2403, 2897, 2960, 1005, 1085.1, 1636, 1699, 2896, 1085, 1726, 1733.1, 2407, 2407.1, 2408, 2968. 0.820: 1728, 2399, 5169, 2892, 2970, 3827, 2967, 1725, 2400.1, 2409, 2796, 2954, 2962, 3356, 1727, 1734, 3978.1 0.825: 2971, 3978, 4005, 4170, 4172, 4848, 800, 2240, 2963, 2966, 2956, 3364, 1736, 2400, 3361, 4002. 0.830: 1469, 2797, 3826, 1547, 1732, 1746, 2929, 4415, 237, 587, 3362, 925, 2410, 3326, 4179, 4836, 998, 1633, 3355, 3821. 0.835: 1098, 1629, 2239, 810, 811, 3358, 517, 814, 837, 999, 1628, 1000, 2952, 3889, 2865, 3893. 0.840: 273, 749, 2412, 3808, 3822, 356, 1466, 3810, 3809, 3815, 4010, 2928, 1546, 1468, 1470, 272. 0.850: 2343, 1572, 3816, 1063, 711, 993, 2890, 3333, 3334, 446, 1048, 3823, 3824. 0.856: 927, 3894, 3903, 5606, 1096, 2288, 3728.1, 1545, 3727, 5380, 3331, 5978. 0.860: 469, 1054, 2834, 3333.1, 3725, 3728, 3730, 3992, 4115, 2686, 3734, 3805, 3969, 4168, 513, 3226, 3228, 3721, 4408, 3229.1. 0.863: 1548, 2835, 2912, 3820, 4367, 2909 1, 3223, 2685, 3731, 3806, 5853, 2359.1. 0.866: 801, 2112, 2357, 2901.1, 3729, 4175, 3225, 3726, 4365.1, 2354.1, 3229, 3740 1, 3330.1, 3807, 3899, 3988, 2359. 0.870: 926, 1046, 1653, 4992, 5813, 2901, 748, 1649, 1652, 1655, 1064, 1695, 2855, 2903, 3891, 798, 2355, 2683, 747. 0.875: 2354, 3915, 3230, 4576, 2356, 3987, 533, 2858, 3733, 3817, 1654, 2353, 2684, 2953, 4117. 0.880: 1365, 5003, 1658, 3908, 3920, 1015, 1651, 3224, 4366, 1016, 1043, 1659, 3329, 4991. 0.884: 746, 4144, 4118, 4370, 1020, 3337, 4827, 1496, 2111, 3850, 4828. 0.890: 468, 1017, 1019, 3119, 1044, 3897, 4980, 1047, 3227, 4376, 5001, 3303, 3918, 5141, 2415, 3917, 397, 1018, 3890, 5362, 713, 725, 3974.1. 0.901: 727, 3639, 3740, 3902, 4385, 4835, 5253, 2538, 5152, 5346, 451, 4842, 4974, 2884, 3328, 4158, 5015, 3324,

4977, 1056, 4148. 0.910: 670, 2899, 3961, 4368.8, 908, 2888, 3913.1, 4841, 642, 2883, 2777, 3861, 1055, 2340, 4982, 5342, 5605. 0.915: ℬ429, ℬ1824, 2831, 3786, 3813, 3913, 6166, 891, 2337, 3788, 4156, 726, 3369, 2298, 4578, 4972, 1557, 3923, 3924, 4388. 0.920: 4131, 3854, 3928, 2351, 764, 2339, 2341, 3575, 938, 2299, 3341, 5482. 0.925: 1558, 1644, 2289, 3847, 3927, 4971, 452, 937, 1617, 4130, 1643, 2882, 3258, 3926, 3935, 4975. 0.930: 2153, 2859, 4976, 4978, 3931, 671, 4843, 965, 2830, 3936, 3735, 3764, 3789. 0.935: 489, 799, 1519, 2861, 2201, 2810, 3922, 4157, 4981, 569, 3260, 3787, 3859, 375, 4371, 3263, 4561. 0.94: 2979, 3790, 3882, 3883, 1010, 3259, 3947, 4990, 763, 1012, 2204, 3858, 762, 978, 2386 1, 3860, 3852, 4560. 0.945: 909, 3857, 997, 2818, 589, 623, 3948, 724, 1541, 3214, 3267, 5005. 0.950: 1443, 2199, 2811, 3265, 783, 924, 1478, 1444, 3319, 3762, 3865, 3904, 4132, 1326, 5940. 0.955: 2775, 624, 1415, 2756, 4378, 752, 2335, 3765, 723, 1555, 2200, 6167 0.960: 3753, 1554, 307, 2763, 3204, 2914, 1553, 2722, 3121, 3655, 2778, 4089, 2365, 3246, 2840. 0.970: 1551, 2721, 3933, 3637, 355, 2762, 4823.1, 1595, 2758, 213, 625, 2766, 3638, 4091 1. 0.975: 929, 1511, 3752, 3856, 4067, ℬ432, 2767, 3751 2, 5009, 3656, 1026, 2760. 0.980: 1089, 2195, 1067.1, 2719, 870, 3654, 1314, 2764, 3878, 930, 3661, 3763, 4579. 0.985: 1372, 4573, 2203, 3648, 935, 2718, 3662, 3761, 4941, 5000, 5688, 1312. 0.990: 931, 1182, 4161, 681, 3235.1, 400, 450, 2757, 162, 815, 3664, 4315, 1090, 1509, 1662, 2163, 3235. 0.995: 3311, 103, 1070, 1510, 3236, 3573, 2204, 3243, 3574, ℬ1, 2058, 4761. 1.000: 1095, 4097.1, 66, 3128, 4513, 5140, 5334, 258, 797, 896, 3134, 3054, 4490, 4757, 4930, 3237, 773.1, 3747, 4147. 1.010: 594, 2713, 3132, 5110, ℬ197, 1560, 590, 2713.1, 620, 2535, 4098, 3780, 4096, 4097, 1279, 652, 928, 2846, 2848, 2302, 2569. 1.020: 608 1, 795, 2570, 3701, 285, 608.2, 1442, 5371, 2322, 4904, 1328, 1561, 3312, 4038, 1789. 1.026: 2571, 3680, 4090.1, 619, 2567, 3681.1, 3684, 5010, ℬ426, 651, 1022, 3133, 3679, 3703. 1.03: ℬ104, 1028, 3677, 3125, 3678, 218, 4939, 2161, 496, 2706, 3676, 2568. 1.040: 2255, 2745, 4545, 4970, ℬ440, 2847, 5678, 3285, 266, 274, 2001, 2159, 720, 3154, 3286, 212, 3069, 4062. 1.050: 593, 3152, 3284, 358, 2812, 4350, 511, 4153, 2300, 4348, 2318, 2748, 3192, 3872, 4093, 4383, 2180, 3149, 309. 1.061: 2788, 4296, 1029, 3283, 911, 4353, 616, 3135, 3191, 378, 576, 989, 1441, 3601, 3547, 2813, 176, 1606, 458. 1.071: 2041, 2040, 3548, 3549, 1430, 2572, 5944, 807, 913, 969.1, 2310, 2590. 1.080: 737, 1570, 2039, 3667, 2588, 449, 626, 609, 3546, 968, 621, 2008, 4726, 1572.1. 1.090: 3649, 4102, 578, 1092, 1559, 2468, 2725, 420, 665, 2814, 3037, 2589, 1889, 3591, 1357, 1483, 3642, 3036. 1.100: 4723, 3169.1, 4917, 471, 722, 2038, 154, 170, 1571, 4670, 247, 3688, 4368.4, 561, 1307, 2087, 1117. 1.11: 492, 2267, 2071, 657, 233, 969, 4733, 264, 470, 4297.1, 672, 736, 2579, 2269. 1.121: 1568, 2134, 4064, 4324, 275, 2580, 5164, 520, 2509, 1341, 2669, 2849.1. 1.131: 3170, 805, 2578, 893, 4381, 3171, 46, 48, 383, 3945, 146, 3253, 3886, 4023. 1.150: 1756, 1388, 2127, 1390, ℬ439, 948, 1917, 994, 2284.1, 3606, 658, 859. 1.160: 2084, 3289, ℬ438, 1253, 453, 2004, 460, 2499, 1252, 1692, 189, 949, 2696. 1.180: 3694, 887, ℬ798, 379, 2618, 5282, 655, 659, 2498, 1042, 3455, 334. 1.200: 1031, 2850, 1347, 1859, 227, 696, 1375, 858, 1041, 1376, 279, 632, 710. 1.220: 37, 384, 744, 1040, 2316, ℬ514, 1576, 4442, 4441, ℬ435, 803, 1314, 1857, 863, 921.1, 1916. 1.252: 190, 1856, 515, 742, 67, 359, 2098, 741, 604, 3937, 1230, ℬ442, 1959, 1229. 1.310: ℬ1575, 465, 192, 1327, 1506, 472, 473, ℬ441, 1251, 1250, 604.1, 1540, 421, 1588, 158, 28, 1249, 2053. 1.340: ℬ366, 464, 423, 2639, 230, 365, 2637, 422, 2633, 1326, ℬ42, 585, 963, 276, 558, 582, 366. 1.400: 497, 2491, 2423, 545, 2031, 605, 2030, 2492, 2493, 304, ℬ634, 2020, 1697, ℬ2, 159, ℬ96, ℬ635, 220, ℬ1397. 1.460: ℬ11, 648, 5350, 1672, 225, 3453, 106, ℬ10, 61, ℬ636, ℬ352, 19, 329, 648.3, 1053, 1294, 2119. 1.500: 1578.1, ℬ632, ℬ637, 43, 1052, 1822, 107, 648.1, 137, 1051, 2454, 648.4, ℬ629. 1.526: 141, ℬ633, 467, 136, 1844, 1367, ℬ207, 645, 139, ℬ630, 12, 756. 1.600: 140, 367, 755, 754, 90, 1601, ℬ521, ℬ232,

B58, 2494, B129, B512, B628, B59, 757, B210, B57, B100, 221, 2061, 2062. **1.700**: 368, 555, 476, 987, 694, 475, B62, 693, B13, 414, B622, 690. **1.800**: 2064, 689, 1949, 688, 1759, 1333, B523, B45, 390, B1597, B60, 38, B1808, 116, B621. **1.901**: B163, 600, B39, 412, 341, 234, 1205, 413, B619, 83, 339, 340, 183, B218, B522. **2.110**: 415, 122, 184, 649, 186, B188, 123, B236, 45, 522, 370, B378, B76, B919, 4, 427. **2.529**: 601, 20, 151, B1815, B63, B142, 345, B64, 101, 5, 127, 18, 235, 128. **3.022**: B204, B918, B497, B381, 20, B34, B206, 87, B205. **4.49**.

B. Solids

0.760: 846, 5881, 5918, 5967, 5985, 6014, 6080, B2916, 5214, 2266, B2601, 1502, 936, 4406, 6010. **0.919**: B2667, 548, 3016, B1812, 3257, 4805, 1058, 239, 3756, 481, 3302. **1.008**: 607, 5343.1, 3901, B2791, 761, 2573, 4322, 1057, 1652, 3307, 760, 2801, 5902, 482, 1077, 2206, 831. **1.051**: 2160, 5817, 5933, 1771, 3140, 289, 571, B2643, 3853, 3550, 502, 2116, 3494, 5214.1. **1.150**: 5213.1, 238, 4270, 2166, 3498, 4352, 832, B431, B430, B2623, 5887, 4943, 5404, 5284, 4894, 2595. **1.203**: 4225, B2626, 259, 5818, 3886 1, B2908, 504, 298, 3867.1, 5428.1, B5, B1896, 2701, 4480, 2308 1, 4226. **1.250**: 4167, 4956, 503, 5573, 1705, B2624, 5435, 2032, 5202, B2306, 1287, 1992, 308.1, 1581, 55, 5541, 5028.1, 1990, 1414. **1.35**: 6104, 4739, 5647, B111, 5028, 4656.1, 802, 3697, B173, 3111, 5704, B2655, 5522. **1.40**: 498, 2175, 58, 4622, 1929, 947, B134, B2170, B2347, 1398, 6148, 1397, 5659, B2300, 4620, 2013, 1349, B3086, 3778. **1.45**: B2757, 808, 3178, 1419, B2171, 630, B2807, 1231, B2636, 976, B2119, B2693, 1351. **1.47**: B2990, 204, 1464, 1991, 2682.1, B2814, 1172, 1350, B1400, B1809, B201, B2855. **5.0**: B502, B1328, B1350, B1426, B1428, B1844, B1904, B289, B1969, B1260, B1375, B2282, B1712, B2202, B1539, B499. **5.10**: B311, B1130, B2017, B734, B1334, B994, B2035.1, B3329, B1021, B2030, B2513, B456, B507, B554, B1258, B1441, B3061, B829. **5.2**: B280, B1096, B1337, B1682, B1711, B1063, B1371, B1590, B1686, B2518, B1990, B1992, B2516, B618, B462. **5.3**: B600, B677, B716, B724, B1154, B1634, B313, B595, B1423, B593, B1049, B1236, B1103, B1767, B883, B1457, B862, B608, B715, B864, B473, B1095. **5.50**: B592, B1630, B1671, B1852, B1542, B1065, B544, B723, B956, B1059, B708. **5.6**: B306, B306.1, B1304, B1710, B1726,

B744, B601, B603, B951, B971, B1636, B1763, B1123, B279, B670, B1064, B1996, B1440, B1455. **5.7**: B320, B322, B1372, B1418, B1614, B2339, B714, B2494, B473.1, B1421, B546, B2338, B1632, B1098, B1723, B957, B582, B2599. **5.8**: B568, B596, B1117, B1685, B1978, B1391, B2048, B529, B574, B2571, B2049, B1163, B541. **5.9**: B602, B1118, B1652, B1703, B907, B1071, B565, B2507, B597, B2538, B1736, B1562. **6.0**: B401, B936, B1050, B1506, B1781, B1227, B540, B2059, B894, B2366, B1442, B1105. **6.1**: B594, B1022, B1101, B1402, B1666, B1784, B402, B658, B657, B548, B1655, B501, B606, B2483, B1327. **6.2**: B553, B614, B1124, B1390, B1617, B863, B539, B1800, B898, B1116, B897, B1055. **6.3**: B604, B607, B1100, B1119, B1517, B1570, B1631, B1366, B2580, B1722, B559, B1086. **6.4**: B335, B605, B667, B934, B935, B995, B1834, B1025, B905, B575, B616, B889, B834, B672, B1051, B1062, B503, B833, B663, B1121. **6.5**: B609, B660, B1102, B1501, B1958, B1629, B3118, B659, B509, B598. **6.6**: B611, B617, B1573, B2827, B1285, B824, B1698, B543, B996, B1143, B1619. **6.7**: B1405, B2007, B2006, B545, B666, B1374, B1620, B1024, B719, B1502. **6.8**: B573, B671, B327, B336, B551, B576, B581, B1776, B2005, B712, B1700, B1306. **6.9**: B610, B661, B1040, B1103, B1681, B1688, B1840, B2834, B557, B612, B1621, B484, B1235. **7.0**: B485, B578, B588, B613, B696, B1386, B1104, B1854, B599, B2041, B1807, B536, B584. **7.1**: B586, B589, B1565, B585, B725, B3188, B587, B334, B590, B882, B1171, B1842, B681, B1734, B2828. **7.2**: B1233, B1697, B535, B2023, B1847, B615, B2826, B2830, B577, B1247, B1977, B893, B1705, B1067, B1066, B910, B325. **7.4**: B1128, B1385, B1393, B1843, B1849, B2062, B2060, B2037, B1057, B1528. **7.5**: B305, B314, B330, B552, B900, B1833, B1041, B700, B904, B538, B1170, B1464, B324. **7.7**: B328, B896, B318, B902, B2079, B1384, B1848, B1146, B323, B891, B676. **8.0**: B525, B704, B1004, B1070, B1732, B1850, B580, B321, B558, B901, B821, B560, B822. **8.2**: B308, B1695, B528, B1326, B888, B890, B1662, B1701, B1550, B888, B1017, B309, B1072, B1684, B1780. **8.64**: B2082, B887, B880, B895, B1137, B1806, B1169, B307, B1663, B881, B675. **9.04**: B1139, B527, B892, B2087, B526, B524, B2099, B668, B879, B1152, B1702, B1179, B1855, B1693. **11.1**: B878, B1725, B1724, B1224, B1225, B1689, B1690. **16.06**.

LIQUID CRYSTALS

H. W. FOOTE

The term "transition temperature" refers in the tables to the temperature at which the solid and crystalline-liquid phases are in equilibrium at a pressure of one atmosphere; by "melting point," is meant the corresponding temperature at which the crystalline-liquid and isotropic liquid phases are in equilibrium. In some cases, more than one stable liquid crystal phase exists, giving an additional transition temperature for each additional liquid crystal phase. These transition temperatures between two liquid crystal phases are indicated by *. In most cases, they are only approximate. Melting points which are quite uncertain, usually due to partial decomposition, have "d " written after the value. No attempt has been made to estimate the accuracy of values obtained by a single investigator, as the methods of determination are the same in nearly every case and the result obviously depends on the skill of the investigator and the purity of the compounds.

A series of apparently good determinations by different observers is apt to vary by considerably more than one degree, and it seems unlikely that any transition temperature or melting point of liquid crystals is known with an accuracy much better than one degree.

For this reason, the weighted average of a number of different determinations is usually given to the nearest whole degree. When the number of determinations is sufficient, the weighted average deviation, usually to the nearest whole degree, is given also.

The melting points of unstable liquid crystals, in monotropic systems, are not included in the tables, and transition temperatures, in the ordinary sense, do not exist in this case. Many observations on monotropic compounds will be found in nearly all the Halle dissertations and in the publications by Vorländer, which are listed at the end of the tables.

For the effect of pressure on the transition temperature and melting point of liquid crystals, see G. Hulett, 7, **28**: 629; 99. For approximate data on liquid crystals of alkali salts of higher fatty acids (chiefly) see Vorländer, **25**, **43**: 3120; 10. For similar data regarding compounds which are optically active, see H. Stoltzenberg, Diss., Halle (1911). For qualitative data regarding liquid crystals, see E. Wolferts, Diss., Halle (09), R. Wilke, Diss., Halle (09); K. Mattenklodt, Diss., Halle (11); and Vorländer, **25**, **40**: 1415, 1966; 07.

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
C$_{10}$H$_{10}$O$_3$	CH$_3$OC$_6$H$_4$CH:CHCOOH	p-Methoxycinnamic acid .	170 ± 1	186 ± 1	(7, 11, 30, 33, 34, 42, 43, 45)
C$_{11}$H$_{12}$O$_3$	C$_2$H$_5$OC$_6$H$_4$CH:CHCOOH	p-Ethoxycinnamic acid .	192	197	(43)
C$_{12}$H$_{14}$O$_3$	C$_2$H$_5$OC$_6$H$_4$CCH$_3$:CHCOOH	p-Ethoxy-β-methylcinnamic acid	122 5	159	(37)
C$_{14}$H$_{10}$BrNO$_3$	BrC$_6$H$_4$CH:NC$_6$H$_4$COOH	p-Bromobenzal-p-aminobenzoic acid	272	274	(12)
C$_{14}$H$_{10}$ClNO$_3$	ClC$_6$H$_4$CH:NC$_6$H$_4$COOH	p-Chlorobenzal-p-aminobenzoic acid	260	263	(12)
C$_{14}$H$_{10}$INO$_3$	IC$_6$H$_4$CH:NC$_6$H$_4$COOH	p-Iodobenzal-p-aminobenzoic acid	279	287	(12)
C$_{14}$H$_{10}$O$_6$	HOC$_6$H$_4$COOC$_6$H$_4$COOH	p-(p-Hydroxybenzoxy)-benzoic acid	258	266 ±	(45)
C$_{14}$H$_{11}$NO$_2$	C$_6$H$_5$CH:NC$_6$H$_4$COOH	Benzal-p-aminobenzoic acid	183	191	(26)
C$_{14}$H$_{12}$N$_2$O$_3$	O$_2$NC$_6$H$_4$CH:NC$_6$H$_4$OCH$_3$	p-Nitrobenzalanisidine	135		(26)
C$_{14}$H$_{14}$N$_2$O$_3$	CH$_3$OC$_6$H$_4$NONC$_6$H$_4$OCH$_3$	p-Azoxyanisol .	116 ± 1	135 ± 1	(1, 3, 6, 7, 9, 11, 14, 19, 23, 30, 32, 35, 36, 42, 45)
C$_{14}$H$_{15}$N$_3$	CH$_3$NHC$_6$H$_4$CH:NNHC$_6$H$_5$	p-Methylaminobenzalphenylhydrazone	170	190	(34)
C$_{15}$H$_{10}$N$_2$O$_2$	CNC$_6$H$_4$CH:NC$_6$H$_4$COOH	p-(p-Cyanobenzalamino)-benzoic acid	247	>320	(17)
C$_{15}$H$_{12}$N$_2$O	CNC$_6$H$_4$CH:NC$_6$H$_4$OCH$_3$	p-Cyanobenzalanisidine	115	125	(17)
C$_{15}$H$_{12}$N$_2$O	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$CN	Anisal-p-cyanoaniline	103	113 5	(12)
C$_{15}$H$_{13}$N$_3$O$_4$	CH$_3$COOC$_6$H$_4$N:NC$_6$H$_4$COOH	p-Acetoxyazobenzoic acid	254	d.	(31)
C$_{15}$H$_{12}$O$_2$	C$_6$H$_5$C$_6$H$_4$CH:CHCOOH	p-Phenylcinnamic acid	221	236	(2)
C$_{15}$H$_{12}$O$_5$	CH$_3$OC$_6$H$_4$COOC$_6$H$_4$COOH	p-(p-Methoxybenzoxy)-benzoic acid	223	272	(45)
C$_{15}$H$_{13}$NO$_2$	CH$_3$C$_6$H$_4$CH:NC$_6$H$_4$COOH	p-(p-Methylbenzalamino)-benzoic acid	220	243	(26)
C$_{15}$H$_{13}$NO$_3$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$COOH	p-(Anisalamino)-benzoic acid	197	298 d.	(15, 46)
C$_{15}$H$_{14}$N$_2$O$_3$	O$_2$NC$_6$H$_4$CH:NC$_6$H$_4$OC$_2$H$_5$	p-Nitrobenzalphenetidine	124		(26)
C$_{15}$H$_{16}$N$_2$O$_3$	CH$_3$OC$_6$H$_4$NONC$_6$H$_4$OC$_2$H$_5$	p-Anisylazoxyphenetol	94 ± 1	149 ± 1	(4, 7, 32)
C$_{15}$H$_{17}$N$_3$	C$_2$H$_5$NHC$_6$H$_4$CH:NNHC$_6$H$_5$	p-Ethylaminobenzalphenylhydrazone	160	182	(34)
C$_{16}$H$_{12}$O$_6$	CH$_3$COOC$_6$H$_4$COOC$_6$H$_4$COOH	p-Hydroxybenzoic acid p-acetoxybenzoate	228 d.	>250	(45)
C$_{16}$H$_{12}$O$_7$	CH$_3$OCOOC$_6$H$_4$COOC$_6$H$_4$COOH	p-Hydroxybenzoic acid p-carbomethoxyoxybenzoate	218 d.	d.	(45)
C$_{16}$H$_{14}$N$_2$O	CNC$_6$H$_4$CH:NC$_6$H$_4$OC$_2$H$_5$	p-Cyanobenzalphenetidine	115	132	(17)
C$_{16}$H$_{14}$N$_2$O	C$_2$H$_5$OC$_6$H$_4$CH:NC$_6$H$_4$CN	p-Ethoxybenzal-p-cyanoaniline	105	124	(12)
C$_{16}$H$_{14}$N$_2$O$_2$	O$_2$NC$_6$H$_4$CH:CHCH:NC$_6$H$_4$CH$_3$	p-Nitrocinnamal-p-toluidine	130	141	(26)
C$_{16}$H$_{14}$N$_2$O$_3$	O$_2$NC$_6$H$_4$CH:CHCH:NC$_6$H$_4$OCH$_3$	p-Nitrocinnamalanisidine	155	160	(26)
C$_{16}$H$_{15}$NO$_2$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$COCH$_3$	Anisal-p-aminoacetophenone	121 5	135	(15)
C$_{16}$H$_{15}$NO$_3$	CH$_3$COOC$_6$H$_4$CH:NC$_6$H$_4$OCH$_3$	p-Acetoxybenzalanisidine	112	128	(15)
C$_{16}$H$_{15}$NO$_3$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$OCOCH$_3$	p-(Anisalamino)-phenol acetate	81 5	108	(15)
C$_{16}$H$_{16}$N$_2$O$_2$	CH$_3$COC$_6$H$_4$N:NC$_6$H$_4$OC$_2$H$_5$	p-Acetophenoneazophenetol	130		(47)
C$_{16}$H$_{16}$N$_2$O$_3$	CH$_3$OC$_6$H$_4$CH:NN:CHC$_6$H$_4$OCH$_3$	Anisaldazine	165 ± 3	180 ± 1	(5, 6, 7, 19)
C$_{16}$H$_{16}$N$_2$O$_3$	C$_2$H$_5$OC$_6$H$_4$N:NC$_6$H$_4$OCOCH$_3$	p-Phenetolazophenetidine	121	138	(46, 47)
C$_{16}$H$_{16}$N$_2$O$_4$	CH$_3$OC$_6$H$_4$N:NC$_6$H$_4$OCOOC$_2$H$_5$	p-Anisylazocarbethoxyphenol .	90	114	(46, 47)
C$_{16}$H$_{18}$N$_2$O$_3$	C$_2$H$_5$OC$_6$H$_4$NONC$_6$H$_4$OC$_2$H$_5$	p-Azoxyphenetol..	137 ± 1	167 ± 1	(3, 14, 19, 23, 30, 32, 35, 42, 45)
C$_{16}$H$_{20}$N$_2$	C$_2$H$_5$NHC$_6$H$_4$C$_6$H$_4$NHC$_2$H$_5$	Diethylbenzidine . .	115.5	120 5	(34)
C$_{17}$H$_{15}$NO$_3$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$CH:CHCOOH	p-(Anisalamino)-cinnamic acid	208	d.	(15)
C$_{17}$H$_{16}$N$_2$O$_3$	O$_2$NC$_6$H$_4$CH:CHCH:NC$_6$H$_4$OC$_2$H$_5$	p-Nitrocinnamalphenetidine .	134	137	(26)
C$_{17}$H$_{16}$N$_2$O$_4$	CH$_3$COC$_6$H$_4$N:NC$_6$H$_4$OCOOC$_2$H$_5$	p-Acetophenoneazocarbethoxyphenol	120	126	(47)
C$_{17}$H$_{16}$N$_2$O$_4$	CH$_3$COOC$_6$H$_4$N:NC$_6$H$_4$COOC$_2$H$_5$	Ethyl p-acetoxyazobenzoate .	99	102	(31)
C$_{17}$H$_{17}$NO$_3$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$CH$_2$CH$_2$COOH	p-(Anisalamino)-hydrocinnamic acid	136	162	(45)
C$_{17}$H$_{18}$N$_2$O$_4$	C$_2$H$_5$OC$_6$H$_4$N:NC$_6$H$_4$OCOOC$_2$H$_5$	p-Phenetolazocarbethoxyphenol	96	137	(47)
C$_{18}$H$_{15}$ClO$_4$	CH$_3$COOC$_6$H$_4$CH:CClC$_6$H$_4$OCOCH$_3$	p-Dihydroxychlorostilbene diacetate	125	138	(11, 29)
C$_{18}$H$_{16}$N$_2$O$_4$	CH$_3$COOC$_6$H$_4$CH:NN:CHC$_6$H$_4$OCOCH$_3$	Di-(p-acetoxybenzalazine)	185	192	(16, 40)
C$_{18}$H$_{17}$NO$_3$	CH$_3$OC$_6$H$_4$CH:NC$_6$H$_4$CH:CHCOOCH$_3$	Methyl anisal-p-aminocinnamate	156	176	(43, 47)
C$_{17}$H$_{17}$N$_2$O$_3$	CH$_3$OC$_6$H$_4$N:NC$_6$H$_4$CH:CHCOOC$_2$H$_5$	Ethyl p-anisylazocinnamate. . . .	116, 123*	143	(46, 47)
C$_{18}$H$_{18}$N$_2$O$_5$	C$_2$H$_5$OCOC$_6$H$_4$NONC$_6$H$_4$COOC$_2$H$_5$	p-Azoxyethyl benzoate.	114 ± 0.6	121 ± 0.5	(7, 11, 19, 27, 40, 42, 45)
C$_{18}$H$_{18}$N$_2$O$_6$	C$_2$H$_5$OCOOC$_6$H$_4$N:NC$_6$H$_4$OCOOC$_2$H$_5$	p-Azocarbethoxyphenol	97	118	(15)

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
C₁₈H₁₈N₂O₇	C₂H₅OCOOC₆H₄NONC₆H₄OCOOC₂H₅	p-Azoxycarbethoxyphenol	95	130	(15)
C₁₈H₁₈O₃	CH₃OC₆H₄CH:CHCH.CHC₆H₄OCH₃	Di-(p-anisylbutadiene)	225	238	(34)
C₁₈H₂₀N₂O₂	C₂H₅OC₆H₄.CH.NN.CHC₆H₄OC₂H₅	Di-(p-ethoxybenzalazine)	172	195	(13, 24, 45)
C₁₈H₂₀N₂O₂	CH₃OC₆H₄C((CH₃):NN:C((CH₃)C₆H₄-OCH₃	Di-(p-methoxyacetophenoneazine)..	195	202	(16)
C₁₈H₂₀N₂O₄	HOC₂H₄OC₆H₄CH·NN:CHC₆H₄-OC₂H₄OH	Di-(hydroxyethoxybenzalazine) ...	184	207	(13)
C₁₈H₂₂N₂O₂	C₃H₇OC₆H₄NONC₆H₄OC₃H₇	Di-(p-n-propoxyazoxybenzene)	116	122	(4, 40)
C₁₉H₁₆N₂O₂	CNC₆H₄CH:NC₆H₄CH:CHCOOC₂H₅	Ethyl p-cyanobenzal-p-aminocinnamate	131	179	(17)
C₁₉H₁₈N₂O₄	CH₃COOC₆H₄N:NC₆H₄CH:CHCOO-C₂H₅	Ethyl p-acetoxyphenylazocinnamate.	132	152	(47)
C₁₉H₁₉NO₂	CH₃C₆H₄CH:NC₆H₄CH.CHCOOC₂H₅	Ethyl p-(p-methylbenzalamino)-cinnamate	96, 107*	118	(46, 47)
C₁₉H₁₉NO₃	C₂H₅OC₆H₄CH.NC₆H₄CH:CCH₃-COOH	p-(p-Ethoxybenzalamino)-α-methyl-cinnamic acid	180	265	(20)
C₁₉H₁₉NO₃	CH₃OC₆H₄.CH.NC₆H₄CH:CH-COOC₂H₅	Ethyl (p-anisalamino)-cinnamate.	100, 108*, 117*	138	(9, 43, 46, 47)
C₁₉H₁₉NO₃	C₂H₅OC₆H₄CH.NC₆H₄CH:CH-COOCH₃	Methyl p-(p-ethoxybenzalamino)-cinnamate	132	187	(43, 47)
C₁₉H₂₂N₂O₃	C₂H₅OC₆H₄N:NC₆H₄OCOC₄H₉	p-Phenetolazophenol n-valerate	78 83	125	(47)
C₂₀H₁₃N₃O₂	CNC₆H₄N:NC₆H₄OCOC₆H₅	p-Cyanobenzeneazophenol benzoate	181	226	(12)
C₂₀H₁₄Br₂N₂	BrC₆H₄N.CHC₆H₄CH:NC₆H₄Br	p-Phthalal-di-(p-bromoaniline)	208	288	(17)
C₂₀H₁₄Cl₂N₂	ClC₆H₄N.CHC₆H₄CH:NC₆H₄Cl	p-Phthalal-di-(p-chloroaniline)	176	282	(17)
C₂₀H₁₄I₂N₂	IC₆H₄N.CHC₆H₄CH.NC₆H₄I	p-Phthalal-di-(p-iodoaniline)..	262	268	(12)
C₂₀H₁₄N₄O₄	O₂NC₆H₄CH:NC₆H₄N.CHC₆H₄NO₂	(Di-p-nitrobenzal)-p-phenylenedia-mine	242	315	(46)
C₂₀H₁₆N₂O₃	CH₃OC₆H₄N NC₆H₄OCOC₆H₅	p-Anisylazophenol benzoate	159 163	178	(47)
C₂₀H₁₇NO	CH₃OC₆H₄CH:NC₆H₄C₆H₅	Anisal-p-aminodiphenyl	161	177	(12, 46)
C₂₀H₁₇N₃O	CH₃OC₆H₄CH:NC₆H₄N:NC₆H₅	Anisal-p-aminoazobenzene	151	182	(15, 39, 46)
C₂₀H₁₈N₂O₅	CH₃OCOCH:CHC₆H₄NONC₆H₄CH-CHCOOCH₃	Methyl azoxycinnamate	221	257	(40)
C₂₀H₂₀N₂O₂	CH₃OC₆H₄CH:CHCH:NN:CHCH:-CHC₆H₄OCH₃	Di-p-methoxycinnamicaldazine	210	218	(34)
C₂₀H₂₀N₂O₄	C₂H₅COOC₆H₄CH:NN:CHC₆H₄OCO-C₂H₅	Di-p-propionylhydroxybenzalazine	160	187	(16)
C₂₀H₂₀N₂O₅	C₂H₅OCOOC₆H₄N:NC₆H₄CH CHCO-OC₂H₅	Ethyl p-carbethoxyphenolazocin-namate	114	152	(47)
C₂₀H₂₁NO₃	C₂H₅OC₆H₄CH:NC₆H₄CH:CHCOO-C₂H₅	Ethyl p-(p-ethoxybenzalamino)-cin-namate.	69, 113,* 152*	159	(43, 45, 46, 47)
C₂₀H₂₁NO₃	CH₃OC₆H₄CH·H₄CH:NC₆CCH₃COO-C₂H₅	Ethyl p-(anisalamino)-α-methylcin-namate	90	93	(20, 43)
C₂₀H₂₁NO₃	C₂H₅OC₆H₄CH:NC₆H₄CH:CCH₃CO-OCH₃	Methyl p-(p-ethoxybenzalamino)-α-methylcinnamate	105	147	(20, 43)
C₂₀H₂₄N₂O₂	C₂H₅OC₆H₄CCH₃:NN:CCH₄C₆H₄O-C₂H₅	Di-p-ethoxyacetophenoneazine	142	163	(16)
C₂₁H₁₄O₇	HOC₆H₄COOC₆H₄COOC₆H₄COOH	p-Hydroxybenzoic acid p-(p-hydroxy-benzoxy) benzoate	283	d.	(45)
C₂₁H₁₆N₂O₃	CH₃COC₆H₄N:NC₆H₄OCOC₆H₅	p-Acetophenoneazophenol benzoate.	211 d.		(47)
C₂₁H₁₇NO	C₆H₅C₆H₄CH:NC₆H₄COCH₃	p-(p-Phenylbenzalamino)-a c e t o-phenone	187 5		(2)
C₂₁H₁₈N₂O₃	C₂H₅OC₆H₄N:NC₆H₄OCOC₆H₅	p-Phenetolazophenol benzoate ...	173	193	(46, 47)
C₂₁H₁₉NO	C₂H₅OC₆H₄CH:NC₆H₄C₆H₅	p-(p-Ethoxybenzalamino) diphenyl	145	184	(12)
C₂₁H₁₉NO	C₆H₅C₆H₄CH:NC₆H₄OC₂H₅	p-Phenylbenzal-p-phenetidine	164	189 5	(2)
C₂₁H₁₉N₃O	C₂H₅OC₆H₄CH:NC₆H₄N NC₆H₅	p-(p-Ethoxybenzalamino)-a z o b e n-zene	131 5	199	(2)
C₂₁H₂₁NO₅	C₂H₅OCOOC₆H₄CH:NC₆H₄CH:CH-COOC₂H₅	Ethyl p-[(p-carbethoxyoxybenzal)-amino] cinnamate	80	151	(47)
C₂₁H₂₃NO₃	CH₃OC₆H₄CH:NC₆H₄CH CH-COOC₄H₉	n-Butyl anisal-p-aminocinnamate.	58	76	(43)
C₂₁H₂₃NO₃	C₂H₅OC₆H₄CH.NC₆H₄CH:CCH₃CO-OC₂H₅	Ethyl p-(p-ethoxybenzalamino)-α-methylcinnamate.	95	122 ± 2	(9, 19, 20, 39, 43, 46)

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
$C_{21}H_{23}NO_3$	$CH_3OC_6H_4CH:NC_6H_4CH:CCH_3$-$COOC_3H_7$	n-Propyl p-(anisalamino)-α-methyl-cinnamate	50	85	(20, 43)
$C_{22}H_{14}H_4$	$CNC_6H_4N:CHC_6H_4CH:NC_6H_4CN$	p-Phthalal-di-(p-cyanoaniline)	164	209	(12)
$C_{22}H_{17}NO_4$	$C_6H_5CH:NC_6H_4COOC_6H_4COOCH_3$	Methyl benzal-p-aminobenzoyl-p-hydroxybenzoate	174	177	(48)
$C_{22}H_{19}NO_2$	$C_6H_5C_6H_4CH:NC_6H_4COOC_2H_5$	Ethyl p-(p-phenylbenzalamino)-benzoate	121 5	128 5	(2)
$C_{22}H_{20}N_2$	$CH_3C_6H_4CH:NC_6H_4N:CHC_6H_4CH_3$	Di-(p-tolual)-p-phenylenediamine	194	266	(46)
$C_{22}H_{20}N_2$	$CH_3C_6H_4N:CHC_6H_4CH:NC_6H_4CH_3$	p-Phthalal-di-(p-toluidine)	186	238	(17)
$C_{22}H_{20}N_2O_2$	$CH_3OC_6H_4CH:NC_6H_4N.CHC_6H_4OCH_3$	Dianisal-p-phenylenediamine	210	338	(46)
$C_{22}H_{22}N_2O_2$	$CNC_6H_4C:HNC_6H_4CH:CHCOOC_5H_{11}$	act-Amyl p-(p-cyanobenzalamino)-cinnamate	95	107	(17, 38, 46)
$C_{22}H_{22}N_2O_4$	$C_2H_5OCOCH:CHC_6H_4N:NC_6H_4CH.$-$CHCOOC_2H_5$	Ethyl p-azocinnamate	155	230	(15, 43)
$C_{22}H_{22}N_2O_5$	$C_2H_5OCOCH:CHC_6H_4NONC_6H_4$-$CH:CHCOOC_2H_5$	Ethyl p-azoxycinnamate	140 ± 1	240 ± 1	(7, 15, 25, 40, 43, 45)
$C_{22}H_{24}O_3$	$CH_3OC_6H_4CH:C_5H_6O:CHC_6H_4OCH_3$	Dianisalcyclohexanone	159	170	(2, 38, 44)
$C_{22}H_{24}N_2O_4$	$C_3H_7COOC_6H_4CH:NN:CHC_6H_4O$-$COC_3H_7$	Di-p-butyryloxybenzalazine	116	181	(16)
$C_{23}H_{25}NO_3$	$CH_3OC_6H_4CH:NC_6H_4CH:CH$-$COOC_5H_{11}$	act-Amyl anisal-p-aminocinnamate	49	90	(43)
$C_{23}H_{25}NO_3$	$CH_3OC_6H_4CH:NC_6H_4CH:CH$-$COOC_5H_{11}$	iso-Amyl anisal-p-aminocinnamate	52	90	(43)
$C_{23}H_{25}NO_3$	$C_2H_5OC_6H_4CH:NC_6H_4CH:CHCOO$-$C_4H_9$	n-Butyl p-(p-ethoxybenzalamino)-cinnamate	68, 88*	125	(43)
$C_{23}H_{25}NO_4$	$C_2H_5OC_6H_4CH:NC_6H_4CHCOH_3COO$-$C_3H_7$	n-Propyl p-(p-ethoxybenzalamino)-α-methylcinnamate	88	121	(20, 43)
$C_{23}H_{18}O_8$	$CH_3COOC_6H_4COOC_6H_4COO$-$C_6H_4COOH$	p-Hydroxybenzoic acid p-(p-acetoxy-benzoxy)-benzoate	248	d.	(48)
$C_{23}H_{19}NO_2$	$C_6H_5C_6H_4CH:NC_6H_4CH:CHCOOCH_3$	Methyl p-(p-phenylbenzalamino)-cinnamate	208, 216*	247	(2)
$C_{23}H_{19}NO_5$	$CH_3OC_6H_4CH:NC_6H_4COOC_6H_4COO$-$CH_3$	Methyl p-(anisalamino)-benzoyl-p-hydroxybenzoate	217	300	(48)
$C_{23}H_{21}NO_4$	$CH_3OC_6H_4CH:NC_6H_4CH_2OC_6H_4CO$-$OCH_3$	Methyl p-(anisalamino)benzyl-p-hydroxybenzoate	157	165	(48)
$C_{23}H_{24}O_3$	$C_2H_5OC_6H_4CH:C_5H_6O:CHC_6H_4OC_2H_5$	Di-(p-ethoxybenzal)-cyclopentanone	189, 194*	200	(44)
$C_{23}H_{27}NO_3$	$C_2H_5OC_6H_4CH:NC_6H_4CH:CHCOO$-$C_5H_{11}$	act-Amyl p-(p-ethoxybenzalamino)-cinnamate	68, 114*	121	(43)
$C_{23}H_{27}NO_3$	$C_2H_5OC_6H_4CH:NC_6H_4CH:CHCOO$-$C_5H_{11}$	iso-Amyl p-(p-ethoxybenzalamino)-cinnamate	81	137	(43)
$C_{23}H_{27}NO_3$	$C_2H_5OC_6H_4CH:NC_6H_4CH:CCH_3CO$-$OC_4H_9$	n-Butyl p-(p-ethoxybenzalamino)-α-methylcinnamate	55, 65*	82	(20, 43)
$C_{23}H_{27}NO_3$	$CH_3OC_6H_4CH:NC_6H_5CH:CCH_3COO$-$C_5H_{11}$	act-Amyl p-(anisalamino)-α-methyl-cinnamate	62	69	(46)
$C_{24}H_{18}O_8$	$C_2H_5OCOOC_6H_4COOC_6H_4COOC_6H_4$-$COOH$	p-Hydroxybenzoic acid p-(p-carbeth-oxyoxybenzoxy) benzoate	215	d.	(48)
$C_{24}H_{20}N_2O_4$	$C_6H_5COOC_6H_4N:NC_6H_4CH:CHCOO$-$C_2H_5$	Ethyl p-benzoyloxyphenylazocin-namate..	135	212	(47)
$C_{24}H_{21}NO_2$	$C_6H_5C_6H_4CH:NC_6H_4CH:CH$-$COOC_2H_5$	Ethyl p-(p-phenylbenzalamino)-cin-namate.	145, 180,* 205,* 210*	219	(2, 39, 43, 46)
$C_{24}H_{22}N_2O_4$	$CH_3OC_6H_4CH:NC_6H_4CONHC_6H_4$-$COOC_2H_5$	Ethyl p-(anisalamino)-benzoyl-p-aminobenzoate. .	212, 220*	247	(45, 46)
$C_{24}H_{24}Br_2N_2O_5$	$C_2H_5OCOCCH_3:CBrC_6H_4NONC_6H_4$-$CBr:CCH_3COOC_2H_5$	Ethyl p-azoxy-α-methyl-β-bromcin-namate	110, 132*	138	(20)
$C_{24}H_{24}N_2O_2$	$C_2H_5OC_6H_4CH:NC_6H_4N:CHC_6H_4O$-$C_2H_5$	Di-(p-ethoxybenzal)-p-phenylenedi-amine	200		(2)
$C_{24}H_{24}N_2O_2$	$C_2H_5OC_6H_4N:CHC_6H_4CH:NC_6H_4O$-$C_2H_5$	p-Phthalal-di-(p-phenetidine)	197	324	(17)
$C_{24}H_{24}N_2O_3$	$C_3H_5OCOCH:CHC_6H_4NONC_6H_4$-$CH:CHCOOC_3H_5$	Allyl p-azoxycinnamate.	124	235	(40)
$C_{24}H_{26}N_2O_5$	$C_2H_5OCOCCH_3:CHC_6H_4NONC_6H_4$-$CH:CCH_3COOC_2H_5$	Ethyl p-azoxy-α-methylcinnamate.	109, 134*	140	(20, 21)

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
C₂₄H₂₆N₂O₄	C₃H₇OCOCH.CHC₆H₄NONC₆H₄- CH:CHCOOC₃H₇	iso-Propyl p-azoxycinnamate. . . .	150	184	(40)
C₂₄H₂₆N₂O₄	C₃H₇OCOCH CHC₆H₄NONC₆H₄- CH CHCOOC₃H₇	n-Propyl p-azoxycinnamate . . .	123	243	(40)
C₂₄H₂₆O₄	C₂H₅OC₆H₄CH C₆H₄O·CHC₆H₄- OC₂H₅	Di-(p-ethoxybenzal)-cyclohexanone.	146	176	(44)
C₂₄H₂₈N₂O₄	C₄H₉COOC₆H₄CH.NN:CHC₆H₄- OCOC₄H₉	Di-(p-valerylhydroxy)-benzalazine	145	160	(16)
C₂₄H₂₈N₂O₄	C₄H₉COOC₆H₄CH.NN CHC₆H₄- OCOC₄H₉	Di-(p-isovalerylhydroxy)-benzalazine	131	156	(16)
C₂₄H₂₉NO₃	C₂H₅OC₆H₄CH NC₆H₄CH·CCH₃- COOC₅H₁₁	act-Amyl p-(p-ethoxybenzalamino)- α-methylcinnamate	86	100	(20, 43)
C₂₄H₂₉NO₃	C₂H₅OC₆H₄CH NC₆H₄CH CCH₃- COOC₅H₁₁	iso-Amyl p-(p-ethoxybenzalamino)- α-methylcinnamate	83	90	(20, 43)
C₂₅H₁₉N₂O₂	C₆H₅C₆H₄N·NC₆H₄OCOC₆H₅	p-Diphenylazophenol benzoate	194	240	(12)
C₂₅H₁₉N₃	C₆H₅C₆H₄CH NC₆H₄N NC₆H₅	p-(p-Phenylbenzalamino)-azobenzene	207	252	(2)
C₂₅H₂₀O₈	CH.COOC₆H₄COOC₆H₄COOC₆H₄- COOC₂H₅	Ethyl p-hydroxybenzoate p-(p-acet- oxybenzoxy) benzoate	112	282	(45)
C₂₅H₂₃NO₄	C₆H₅COOC₆H₄CH NC₆H₄CH·- CHCOOC₂H₅	Ethyl p-(p-benzoxybenzalamino)- cinnamate	125	217	(47)
C₂₅H₂₃NO₂	C₆H₅C₆H₄CH:NC₆H₄CH CCH₃- COOC₂H₅	Ethyl p-(p-phenylbenzalamino)-α- methylcinnamate	120, 148*	175	(20, 43)
C₂₅H₂₈N₂O₅	C₃H₇OCOCCH CHC₆H₄NONC₆H₄- CH.CHCOOC₃H₇	n-Propyl p-azoxy-α-methylcinnamate	70, 125*?	128	(20)
C₂₆H₁₈Br₂N₂	BrC₆H₄CH NC₆H₄C₆H₄N CHC₆H₄Br	Di-(p-bromobenzal)-benzidine	285	312	(12)
C₂₆H₁₈Cl₂N₂	ClC₆H₄CH·NC₆H₄C₆H₄N CHC₆H₄Cl	Di-(p-chlorobenzal)-benzidine	265	318	(12)
C₂₆H₁₈Cl₂N₂O	ClC₆H₄N.CHC₆H₄NONC₆H₄CH. NC₆H₄Cl	p-Azoxybenzaldi-m-chloramine	174, 181,* 198*	213	(46)
C₂₆H₁₈I₂N₂	IC₆H₄CH.NC₆H₄C₆H₄N.CHC₆H₄I	Di-(p-iodobenzal)-benzidine	>300		(12)
C₂₆H₁₈N₂O₂	C₆H₅COOC₆H₄N.NC₆H₄OCOC₆H₅	p-Dibenzoylazophenol	208	250	(15, 39)
C₂₆H₁₈N₂O₃	C₆H₅COOC₆H₄NONC₆H₄OCOC₆H₅	p-Dibenzoylazoxyphenol	192	280	(15)
C₂₆H₁₈N₄O₆	O₂NC₆H₄CONHC₆H₄C₆H₄NHCO- C₆H₄NO₂	Di-(p-nitrobenzoyl)-benzidine	365	d.	(45)
C₂₆H₁₈O₄	C₆H₅OCOC₆H₄C₆H₄COOC₆H₅	Diphenyl p, p'-diphenylcarboxylate	213	245	(45)
C₂₆H₂₀N₂	C₆H₅CH:NC₆H₄C₆H₄N.CHC₆H₅	Dibenzalbenzidene	234	260	(6, 24)
C₂₆H₂₀N₂	C₆H₅C₆H₄CH:NN.CHC₆H₄C₆H₅	Di-p-phenylbenzalazine	245	271	(2)
C₂₆H₂₂N₂	CH₃C₆H₄CH·NC₁₀H₆N CHC₆H₄CH₃	Di-p-tolual-1, 5-naphthylenediamine	210	230	(46)
C₂₆H₂₂N₂O₂	CH₃OC₆H₄CH:NC₁₀H₆N:- CHC₆H₄OCH₃	Dianisal-1, 5-naphthylenediamine	206	313	(46)
C₂₆H₂₂N₄O₂	H₂NC₆H₄CONHC₆H₄C₆H₄NHCO- C₆H₄NH₂	Di-(p-aminobenzoyl)-benzidine	312	d.	(45)
C₂₆H₂₄N₂O₄	C₆H₄(CH·NC₆H₄COOC₂H₅)₂	Ethyl p-phthalal-di-(p-aminobenzo- ate) ..	189	230	(17)
C₂₆H₂₅NO₂	C₆H₅C₆H₄CH:NC₆H₄CH:- CHCOOC₄H₉	n-Butyl p-phenylbenzal-p-aminocin- namate .	167	203	(43)
C₂₆H₂₆N₂O₄	C₂H₅OCOCCH₃.CHC₆H₄NONC₆H₄- CH:CCH₃COOC₂H₅	Allyl p-azoxy-α-methylcinnamate.	75	115	(20)
C₂₆H₂₆N₂O₉	C₂H₅OCOCH₂OCOCH CHC₆H₄- NONC₆H₄.CH·CHCOOCH₂- COOC₂H₅	p-Azoxycinnamic acid ethyl glyco- late ester	148	235	(40)
C₂₆H₃₀N₂O₅	C₄H₉OCOCH.CHC₆H₄NONC₆H₄- CH:CHCOOC₄H₉	n-Butyl p-azoxycinnamate . .	111	214	(40)
C₂₇H₂₇NO₂	C₆H₅C₆H₄CH:NC₆H₄CH:- CHCOOC₅H₁₁	act-Amyl p-(p-phenylbenzalamino)- cinnamate	115, 153*	180	(43)
C₂₇H₂₇NO₂	C₆H₅C₆H₄CH:NC₆H₄CH:- CHCOOC₅H₁₁	iso-Amyl p-(p-phenylbenzalamino)- cinnamate .	164, 188*	197	(43)
C₂₇H₂₇NO₂	C₆H₅C₆H₄CH:NC₆H₄CH:- CCH₃COOC₄H₉	n-Butyl p-(p-phenylbenzalamino)-α- methylcinnamate	99, 137*	149	(20, 43, 46)
C₂₇H₂₇NO₂	C₆H₅C₆O₄CH:NC₆H₄CH.- CC₂H₅COOC₃H₇	n-Propyl p-(p-phenylbenzalamino)- α-ethylcinnamate	119	135	(20, 21, 43)
C₂₈H₁₈O₆	C₆H₅COOC₆H₄C:CC₆H₄OCOC₆H₅	Di-p-oxytolanedibenzoate	214	254	(41)
C₂₈H₂₀N₂O₄	C₆H₅COOC₆H₄CH:NN:CHC₆H₄- OCOC₆H₅	Di-p-benzoxybenzalazine.	227	290	(16, 40)

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
C₃₄H₃₀O₆	C₆H₄COOC₆H₄CH:CHC₆H₄OCOC₆H₄	Di-p-hydroxystilbene dibenzoate	224	285 d.	(41)
C₂₈H₂₄N₂	(C₆H₄N:CHC₆H₄CH₃)₂	Di-(p-tolual)-benzidine	231	>300	(6, 34)
C₂₈H₂₄N₂O₂	(C₆H₄N:CHC₆H₄OCH₃)₂	Dianisalbenzidene	258		(46)
C₂₈H₃₃N₂O₄	C₆H₅COOC₆H₄N:NC₆H₄CH:CCH₃-COOC₅H₁₁	act-Amyl p-benzoylazophenol-α-methylcinnamate	88	120	(20)
C₂₈H₃₄N₂O₄	C₄H₁₁OCOCH:CHC₆H₄NONC₆H₄-CH:CHCOOC₅H₁₁	iso-Amyl p-azoxycinnamate	144	186	(40)
C₂₈H₃₄N₂O₄	C₄H₉OCOCCH₃:CHC₆H₄NONC₆H₄-CH:CCH₃COOC₄H₉	iso-Butyl p-azoxy-α-methylcinnamate	86, 110*	125.5	(20)
C₂₈H₃₄N₂O₄	C₄H₉OCOCCH₃:CHC₆H₄NONC₆H₄-CH:CCH₃COOC₄H₉	n-Butyl p-azoxy-α-methylcinnamate	60	100	(20)
C₃₀H₂₂N₂O₃	C₆H₅COCH:CHC₆H₄NONC₆H₄CH:-CHCOC₆H₅	p-Azoxybenzalacetophenone	213		(47)
C₃₀H₂₈N₂O₂	(C₆H₄N:CHC₆H₄OC₂H₅)₂	Di-(p-ethoxybenzal)-benzidine	248	>300	(13)
C₃₀H₂₈N₂O₂	(C₆H₄N:CHC₆H₄CH₂OCH₃)₂	Di-(p-methoxy-o-methylbenzal)-benzidine	171	>300	(13)
C₃₀H₃₀N₂O₄	C₆H₄(CH:NC₆H₄CH:CHCOOC₂H₅)₂	Ethyl p-phthalal-di-(p-aminocinnamate)	174, 270*	310	(17)
C₃₀H₅₀O₂	C₂H₅COOC₂₇H₄₅	Cholesterol propionate	97 ± 2	112 ± 2	(6, 10, 18, 30)
C₃₀H₅₀O₃	C₂H₅OCOOC₂₇H₄₅	Cholesterol ethyl carbonate	83	103 5	(8)
C₃₁H₅₂O₂	C₃H₇COOC₂₇H₄₅	Cholesterol n-butyrate	96 4	107 3	(18)
C₃₁H₅₄O₃	C₃H₇OCOOC₂₇H₄₅	Cholesterol n-propyl carbonate	99	101	(8)
C₃₂H₂₄N₂	C₆H₄(N:CHC₆H₄C₆H₅)₂	Di-(p-phenylbenzal-p-phenylenediamine	284	>300	(2)
C₃₂H₂₄N₂O₄	C₆H₅CH:CHCOOC₆H₄CH:NN:CH-C₆H₄OCOCH:CHC₆H₅	Di-(p-cinnamylhydroxy)-benzalazine	206	245	(16)
C₃₂H₃₄O₁₀	CH₃COOC₆H₄COOC₆H₄COOC₆H₄-COOC₆H₄COOC₂H₅	Ethyl p-hydroxybenzoate p-[p-(p-acetoxybenzoxy)benzoxy]benzoate.	187 d	d.	(45)
C₃₂H₂₆O	C₆H₅C₆H₄CH:C₆H₅O CHC₆H₄C₆H₅	Di-(p-phenylbenzal)-cyclohexanone	236 5	237 5	(2)
C₃₃H₃₃N₂O₂	C₂H₅OCH₂C₆H₂CH:NC₆H₄C₆H₄N:C-HC₆H₄CH₂OC₂H₅	Di-(-p-ethoxy-o-methylbenzal)-benzidine	167	>300	(13)
C₃₃H₅₆O₂	C₄H₉COOC₂₇H₄₅	Cholesterol valerate	91 8	99.2	(18)
C₃₃H₅₆O₃	C₄H₉OCOOC₂₇H₄₅	Cholesterol n-butyl carbonate	78	90	(8)
C₃₃H₂₆O₅	C₆H₅COOC₆H₄CH:C₅H₄O:CHC₆H₄O-COC₆H₅	Di-(p-benzoxybenzal)-cyclopentanone	234	236	(44)
C₃₃H₅₆O₂	C₅H₁₁COOC₂₇H₄₅	Cholesterol capronate	91 2	100	(18)
C₃₄H₃₂N₂O₇	C₆H₅COCH₂OCOCH.CHC₆H₄NON-C₆H₄CH:CHCOOCH₂COC₆H₅	Phenacyl p-azoxycinnamate	231	238	(40)
C₃₄H₄₆N₂O₄	C₈H₁₇OCOCH:CHC₆H₄NONC₆H₄-CH:CHCOOC₈H₁₇	n-Octyl p-azoxycinnamate	91	175	(40)
C₃₄H₅₀O₂	C₆H₅COOC₂₇H₄₅	Cholesterol benzoate	146 ± 1	178.5 ± 0.3	(18, 22, 30, 35, 42, 45)
C₃₆H₄₀N₂O₄	C₆H₄(CH:NC₆H₄CH:CHCOOC₅H₁₁)₂	act-Amyl p-phthalal-di-(p-aminocinnamate)	133, 195*	268	(17)
C₃₆H₅₀N₂O₅	C₈H₁₇OCOCCH₃:CHC₆H₄NONC₆H₄-CH:CCH₃COOC₈H₁₇	n-Octyl p-azoxy-α-methylcinnamate	41, 62*	85	(20)
C₃₇H₆₄O₂	C₉H₁₉COOC₂₇H₄₅	Cholesterol caprinate ...	82 2	90.6	(18)
C₃₈H₄₄N₂O₄	C₆H₄(CH:NC₆H₄CH:CCH₄COO-C₆H₁₁)₂	act-Amyl p-phthalal-di-(p-amino-α-methylcinnamate)	144, 211*	248	(17)
C₄₀H₂₈N₆O₆	(C₆H₄NHCOC₆H₄N:CHC₆H₄NO₂)₂	Di-(m-nitrobenzal-p-aminobenzoyl)-benzidine.	>370	d.	(45)
C₄₀H₃₄N₄	C₆H₅CH:NC₆H₄CH₂NHC₆H₄C₆H₄N-HCH₂C₆H₄N:CHC₆H₅	Di-p-(benzalaminobenzyl)-benzidine.	217	246 d.	(46)
C₄₂H₃₈N₄O₂	(C₆H₄NHCH₂C₄H₄N:CHC₆H₄OCH₃)₂	Di-p-(anisalaminobenzyl)-benzidine	202 d.	d.	(45)
C₅₀H₇₈N₂O₄	C₁₆H₃₃OCOCH:CHC₆H₄NONC₆H₄-CH:CHCOOC₁₆H₃₃	n-Cetyl p-azoxycinnamate..	105	141	(40)
C₅₂H₈₂N₂O₅	C₁₆H₃₃OCOCCH₃:CHC₆H₄NONC₆H₄-CH:CCH₃COOC₁₆H₃₃	n-Cetyl p-azoxy-α-methylcinnamate	77	84	(20)
C₅₅H₉₀O₃	C₂₇H₄₅OCOOC₂₇H₄₅	Cholesterol carbonate... ...	177	235	(8)
C₁₄H₁₂ClHgNO	CH₃OC₆H₄CH:NC₆H₄HgCl	p-Anisalaminophenylmercury chloride....................	274	d.	(46)

Index formula	Formula	Name	Trans. temp.	M. P.	Lit.
$C_{15}H_{13}ClHgN$	$C_6H_5CH:CHCH:NC_6H_4HgCl$	p-Cinnamalaminophenylmercury chloride......	255	265	(46)
$C_{16}H_{15}HgNO_3$	$CH_3OC_6H_4CH:NC_6H_4HgOCOCH_3$	p-Anisalaminophenylmercury acetate	177	180	(46)
$C_{26}H_{18}HgN_4O_4$	$O_2NC_6H_4CH:NC_6H_4HgC_6H_4N:CHC_6-H_4NO_2$	Mercury di-(p-nitrobenzalamino-phenyl)	236	241	(46)
$C_{26}H_{20}HgN_2$	$C_6H_5CH:NC_6H_4HgC_6H_4N.CHC_6H_5$	Mercury di-(benzalaminophenyl)	180	184	(46)
$C_{28}H_{24}HgN_2$	$Hg(C_6H_4N:CHC_6H_4CH_3)_2$	Mercury di-(p-toluialaminophenyl)	217	229	(46)
$C_{28}H_{24}HgN_2O_2$	$Hg(C_6H_4N:CHC_6H_4OCH_3)_2$	Mercury di-(anisalaminophenyl)	209	285	(46)
$C_{30}H_{24}HgN_2$	$Hg(C_6H_4N:CHCH CHC_6H_5)_2$	Mercury di-(cinnamalaminophenyl)	208	269	(46)
$C_{30}H_{28}HgN_2O_2$	$Hg(C_6H_4N:CHC_6H_4OC_2H_5)_2$	Mercury di-(p-ethoxybenzalamino-phenyl)	204	272	(46)

LITERATURE

(For a key to the periodicals see end of volume)

(1) Auwers, 7, **32**: 30, 00.   (2) Bertleff, Diss, Halle, 08.   (3) Bogojawlensky and Winogradow, 7, **60**: 433, 07.   (4) Bogojawlensky and Winogradow, 7, **64**: 229, 08.   (5) Bose and Conrat, 65, **9**: 169, 08.   (6) Bredig and v Schukowsky, **25**, **37**: 3419, 04.   (7) Böhner, Diss, Marburg, 06.   (8) Däumer, Diss, Halle, 12.   (9) Dickenschied, Diss, Halle, 08.   (10) Dorn, 03, **11**: 777, 10.   (11) Eichwald, Diss, Marburg, 05.   (12) Fröhlich, Diss, Halle, 10.   (13) Gatterman, 14, **357**: 313, 07.   (14) Gatterman and Ritschke, **25**, **33**: 1738, 90.   (15) Hansen, Diss Halle, 07.   (16) Hulme, Diss, Halle, 07.   (17) Huth, Diss, Halle, 10.   (18) Jaeger, 64P, **9**: 78, 06.   (19) Jaeger, 64, **101**: 1, 17.

(20) Kasten, Diss, Halle, 09.   (21) E Lehmann, Diss, Halle, 10.   (22) Lehmann, 7, **4**: 462, 89.   (23) Lehmann, 8, **40**: 401, 90.   (24) Lehmann, 8, **2**: 649, 00.   (25) Lehmann, 8, **19**: 22, 06.   (26) Meye, Diss, Halle, 08.   (27) Meyer and Dahlem, 13, **326**: 331, 03.   (28) Müller, **25**, **54**: 1481, 21.   (29) Münch, Diss, Marburg, 03.   (30) Prins, 7, **67**: 689, 09.   (31) Reichardt, Diss, Halle, 09.   (32) Rising, **25**, **37**: 13, 01.   (33) v Romburgh, 64V, **9**: 9, 01.   (34) Rotarski, **25**, **41**: 1994, 08.   (35) Schenk, 7, **25**: 337, 98.   (36) Schenk, 7, **28**: 280, 99.   (37) Schroeter, **25**, **41**: 5, 08.   (38) Stumpf, 65, **11**: 780, 10.   (39) Sultze, Diss, Halle, 08.

(40) Vorländer, **25**, **39**: 803, 06.   (41) Vorländer, **25**, **40**: 4527, 07.   (42) Vorländer, 196, **12**: 321, 07.   (43) Vorländer, **25**, **41**: 2033, 08.   (44) Vorländer, **25**, **54**: 2261, 21.   (45) Vorländer, 7, **105**, 211, 23.   (46) D Vorländer, Chem Kristallog der Flüssigkeiten, 1924.   (47) Wilke, Diss, Halle, 09.

CRYSTALLOGRAPHY OF COMPOUNDS OF CARBON

GEORGE L. KEENAN AND RAYMOND M. HANN

Standard arrangement. For abbreviations, see p. 100. Literature, p. 338

ℬ-TABLE

Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
16 See **C**-Table								
18 SiC₂₄H₂₀N₄	Silico tetraphenylamide	M	Bi	··	17° 40′		Ax pl b (010); X∧c = 27½° in obtuse ∠β	(G)
SiC₂₁H₁₆	Tetra-p-tolylsilicane	M.	Bi	–		83° 30′	Ax. pl ⊥b(010)	(G)
SnC₁₄H₁₀N₂Cl₂	p-Toluidine tin chloride	M.	Bi.	↑	77°		Ax. pl ⊥b(010); Z∧c = 19° in obtuse ∠β	(G)
23 PbC₂H₂O₄	Lead formate	R	Bi	–	70° 34′		Ax. pl. b(010), X‖c	(G)
PbC₄H₆O₄.3H₂O	Lead acetate	M	Bi	+	83° 55′		Ax. pl b(010), Z∧c = 55° 18′ in obtuse ∠β	(G)
PbC₁₀H₁₄O₁₀S₂.6H₂O	Lead sulfocamphylate	R	Bi	–		78° 17′	Ax. pl b(010), X‖c	(G)
27 TlC₂HO₄	Thallium acid oxalate	M	Bi	+	74° 5′ (red)		Ax. pl ⊥b(010)	(G)
TlC₂HO₄.½H₂O	Thallium acid oxalate	M	Bi	+	106° 5′ (red)		Ax pl b(010), Z∧c = 79° 36′ (red) in obtuse ∠β	(G)
Tl₂C₄H₄O₆	Thallium mesotartrate	Tri	Bi	+	73° 54′			(G)
Tl₂C₄H₄O₆.½H₂O	Thallium tartrate	R. (?)	Bi.	–		69°	Ax pl b(010), X‖c	(G)
TlC₅H₂O₇N₃	Thallium picrate	M	Bi.				Ax pl. b(010)	(G)
Tl₂C₄H₄O₆	Thallium dl-tartrate	M	Bi	+	88° 22′		Ax pl b(010), Z∧c = 84° 44′ in obtuse ∠β	(G)
Tl₂C₄H₄O₆	Thallium tartrate	Trig	Un	+				(G)
TlC₄H₄O:Sb.H₂O	Thallium antimonyl tartrate	R	Bi	–		20°-25°		(G)
28 ZnC₄H₆O₄.3H₂O	Zinc acetate	M	Bi.	+	84° 30′		Ax pl b(010), Z∧c = 54 75° in acute ∠β	(G)
ZnC₈H₁₄O₄	Zinc butyrate	M.	Bi.	+	Large			(37)
ZnC₈H₁₈O₄	Zinc methylethylvalerate	?	Bi.					(37)
ZnC₃H₂O₄Br.8H₂O	Zinc bromomesaconate	M	Bi	–	71° 21′	118° 15′	Ax. pl ⊥b(010); X∧c = 14° in obtuse ∠β	(G)
ZnC₁₀H₆O₆S₂.6H₂O	Zinc naphthalene-1, 5-disulfonate	M.	Bi		58° 16′		Ax. pl.‖(010); η_α∧c = 74°	(41)
ZnC₂₄H₃₀N₂I₄	Phenyldimethylethylammonium zinc iodide	M	Bi.	+	80° 52′		Ax. pl ⊥b(010); Z∧c = 43° in acute ∠β	(G)
ZnC₆H₁₃ON₃Cl₄.3H₂O	Triacetonediamine hydrochloride zinc chloride	M.	Bi.	+	36° 14′	58° 20′	Ax pl ⊥b(001); Z∧c = 49° in obtuse ∠β	(G)
30 HgC₂H₈NI₃	1, 1-Dimethylammonium mercuric iodide	M.	Bi	–	Large			(16)
HgC₃H₁₀NI₃	1, 1-Trimethylammonium mercuric iodide	R.	Bi.	–	Large			(16)
HgC₄H₁₂NI₃	1, 1-Diethylammonium mercuric chloride	R.	Bi.	+	Very large			(16)
CuC₂H₂O₄.4H₂O	Cupric formate	M.	Bi.	–	34° 54′	55° 6′	Ax. pl. b(010); X∧c = 23° 35′ in obtuse ∠β	(G)
CuC₁₀H₆O₆S₂.6H₂O	Copper naphthalene-1, 5-disulfonate	M.	Bi.				Ax. pl ‖(010), η_α∧c = 75°	(14)

Ag	Al	As	Au		B	Ba	Be	Bi	Br		C	Ca	Cb	Cd	Ce		Cl	Co	Cr	Cs	Cu		Dy	Er	Eu	F	Fe		Ga	Gd	Ge	Gl	H		Hf	Hg	Ho	I	In		Ir	K	La	Li	Lu
47	13	33		64	79	75	15	5		16	77	51	29	59		4	44	46	85	31		67	60	64	3	43		25	65	20	75	2		73	30	68	6	26		36	53	56	51	72

	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
32	AgC₄O₂N₄	Ethylene dicyanide silver nitrate	R	Bi	–	42° 36 5'		Ax. pl. c(001); X‖b	(G)
	AgC₄H₄O₁₂N₄	Ethylene dicyanide silver nitrate	R	Bi	–	42° 41'		Ax. pl. c(001); X‖a	(G)
	AuC₁₄H₁₄SCl	Gold dibenzylsulfine chloride (meta-stable form)	Tet	Un					(G)
	AuC₅H₁₂NCl₄	Piperidine chloroaurate	R	Bi	+		70° 40'	Ax pl. b(010); Z‖c	(G)
	AuC₅H₁₀O₂NCl₄.H₂O	4-Aminovaleric acid chloroaurate	M	Bi	–		70° (apprx)	Ax pl ⊥b(010); X∧c = 91 5° in obtuse ∠β	(G)
	AuC₉H₁₆NCl₄	3, 4, 5, 6-Tetramethyl-1, 2-dihydro-pyridine hydrochloride chloroaurate	M	Bi	+		91° (apprx)	Ax pl ⊥b(010)	(G)
	K₃IrC₂O₄Cl₄.H₂O	Iridium tetrachloro tripotassium oxalate	R	Bi	–		94° 40'	Ax pl (010), Bxₐ⊥(001)	(11)
37	PtC₂H₁₂N₂Cl₆	Methylammonium chloroplatinate	C						(21)
	PtC₁₀H₁₄N₂Cl₆	Pyridine chloroplatinate	Tri	Bi	·		59° 54'	Ax pl nearly ⊥c-axis	(G)
	PtC₁₀H₁₆O₂N₂Cl₆	Choline chloroplatinate	M	Bi	+		25° 52'	Ax pl ⊥b(010); Z∧c = 75° 12' in acute ∠β	(G)
	PtC₁₂H₁₄N₂Cl₆	α-Picoline chloroplatinate	M	Bi	–		93° 13 5'	Ax pl b(010)	(G)
	PtC₁₈H₂₀N₆Cl₆	1-Phenyl-3-imino-5-methyl triazoline chloroplatinate	M	Bi				Ax pl b(010); Z nearly ⊥c(001)	(G)
	PtC₁₂H₁₄O₄N₂Cl₆ 2H₂O	Pipecolinic acid chloroplatinate	M	Bi			66° 56'	Ax pl b (010)	(G)
	PtC₁₂H₁₄O₄N₂Cl₆	α-Homobetaine chloroplatinate	M	Bi	+	88° 12'		Ax pl b(010), Z∧c = 89° in obtuse ∠β	(G)
	PtC₁₄H₁₈N₂Cl₆	Ethyl pyridine chloride chloroplatinate	R	Bi	–		44°	Ax pl a(100), X‖b	(G)
	PtC₁₄H₁₈N₂Cl₆	Dipropyl carbinol amine chloroplatinate	M	Bi	–		72° 40'	Ax pl ⊥b(010); X nearly ⊥c (001)	(G)
	PtC₁₆H₁₈O₂N₂Cl₆	Tropanine chloroplatinate	M	Bi		52° 12'		Ax pl ⊥b(010)	(G)
	PtC₁₀H₁₈N₂Cl₆	Tropidine chloromethylate chloroplatinate	R	Bi.	+		70°	Ax pl b(010); Z‖c	(G)
	PtC₁₀H₂₄N₂Cl₆	Ethyldipropyl ammonium chloroplatinate	R	Bi.			61° 26'	Ax pl c(001); Z‖a	(G)
	PtC₂₀H₄₀N₂Cl₆	Anhydrolupinin chloroplatinate (stable mod)	M	Bi			38° (apprx)	Ax pl ⊥b(010)	(G)
	PtC₂₂H₂₈N₂Cl₆	Diethyl-p-toluidine chloroplatinate	R	Bi	+	63° 0'		Ax pl a(100), Z‖b	(G)
39	RuN₃H₁₂O₃Cl₃	Ruthenium ammonium chloral hydrate	M	Bi		56° 20'			(L.-B)
	MnC₁₂H₄O₁₄N₆ 5H₂O	Manganese picrate	R	Bi	–		15° 30'	Ax pl. b(010); X‖c	(G)
43	FeC₁₂H₄O₁₄N₆ 5H₂O	Ferrous picrate	R	Bi.	·		24° 48'	Ax pl. a(100); X‖c	(G)
	FeC₁₅H₂₁O₆	Ferriacetylacetone	R	Bi	·		50° (apprx)	Ax pl. a(100); X‖c	(G)
	FeC₂₀H₁₄O₆S₂.6H₂O	Ferrous naphthalene-β-sulfonate		Bi	+				(1)
44	CoC₄H₆O₄.4H₂O	Cobalt acetate	M	Bi.	–	30° 13'	48° 12'	Ax pl b(010); X∧c = 53 5° in acute ∠β	(G)
	CoC₆H₂₄N₆I₂.H₂O	d-Luteo triethylenediamine cobalt iodide	R	Bi	+		Small	Ax pl (001); Bxₐ = b-axis	(15)
	CoC₆H₂₄N₆I₂.H₂O	dl-Luteo triethylenediamine cobalt iodide	R	Bi			Small	Ax pl (010), Bxₐ = c-axis	(15)
	CoC₁₀H₆O₆S₂.6H₂O	Cobalt naphthalene-1, 5-disulfonate	M	Bi		61° 40'		Ax pl ‖(010), ηₐ∧c = 72° 0.5'	(41)
	NiC₁₀H₆O₆S₂ 6H₂O	Nickel naphthalene-1, 5-disulfonate	M	Bi		50° 56'		Ax pl ‖(010), ηₐ∧c = 74°	(41)
49	UC₂H₁₂O₃N	Ammonium uranyl acetate	Tet	Un					(G)
	UCdC₁₂H₁₈O₁₀ 6H₂O	Cadmium uranylacetate	R	Bi	–		57° 54' (red)	Ax. pl. a(100)	(G)
	UMnC₁₂H₁₈O₁₆.6H₂O	Manganese uranyl acetate	R	Bi	–		31°	Ax pl. a(100)	(G)
	(UO₂)₃CoC₁₂H₁₈O₁₂ 7H₂O	Cobalt diuranyl acetate	R	Bi	–		103° 30'	Ax. pl. c(001)	(G)
55	Al₂C₁₂O₁₂ 18H₂O	Mellite	Tet	Un.					(25)
	YtC₁₂H₃₀O₁₂S₄ 18H₂O	Yttrium ethyl sulfate	H	Un					(24)
	YC₁₂H₁₇O₁₅N₃S₃ 7H₂O	Yttrium m-nitrobenzenesulfonate	M	Bi.	+			Ax pl b(010); Z∧c = 85° in obtuse ∠β	(G)
58	LaC₁₂H₃₀O₂₄S₄ 18H₂O	Lanthanum ethyl sulfate	H	Un					(24)
	CeC₁₂H₃₀O₂₄S₄ 18H₂O	Cerium ethyl sulfate	H	Un					(24)
60	PrC₁₂H₃₀O₂₄S₄ 18H₂O	Praseodymium ethyl sulfate	H	Un					(24)
	NdC₁₂H₃₀O₂₄S₄ 18H₂O	Neodymium ethyl sulfate	H	Un					(24)
63	SaC₁₂H₃₀O₂₄S₄ 18H₂O	Samarium ethyl sulfate	H	Un					(24)
	EuC₁₂H₃₀O₂₄S₄ 18H₂O	Europium ethyl sulfate	H	Un					(24)
	GdC₁₂H₃₀O₂₄S₄.18H₂O	Gadolinium ethyl sulfate	H	Un					(24)
67	DyC₁₂H₃₀O₂₄S₄ 18H₂O	Dysprosium ethyl sulfate	H	Un					(24)
	ErC₁₂H₃₀O₂₄S₄ 18H₂O	Erbium ethyl sulfate	H	Un					(24)
	TmC₁₂H₃₀O₂₄S₄ 18H₂O	Thulium ethyl sulfate	H	Un					(24)
	YbC₁₂H₃₀O₂₄S₄ 18H₂O	Neoytterbium ethyl sulfate	H	Un					(24)
75	BeC₄H₈O₄N₂	Ammonium beryllium oxalate	M.	Bi			27° 47'	Ax. pl. b(010), Z∧c = 37.5° in obtuse ∠β	(G)
	Be₃C₄H₁₀O₈S₂ 4H₂O	Diethyl beryllium sulfate (basic)	Tet	Un					(24)
	MgC₄H₆O₄ 4H₂O	Magnesium acetate	M	Bi	–	56° 34'	89° 54'	Ax. pl b(010); X∧c = 48.25° in acute ∠β	(G)
	MgC₆H₁₀O₆.2.5H₂O	Magnesium dilactate.	M	Bi	+		79° (apprx)	Ax. pl. b(010)	(G)
	MgC₄H₄O₆ 6H₂O	Magnesium dl-tartrate	M	Bi.	–		102°	Bxₐ∧c = 30° in acute ∠β	(17)
	MgC₁₀H₆O₆S₂ 6H₂O	Magnesium naphthalene-1, 5-disulfonate	M.	Bi		52° 20'		Ax pl.‖(010), ηₐ∧c = 73° 0.5'	(41)
77	CaC₂O₄.H₂O	Calcium oxalate.	M.	Bi	+	80°		Ax pl. b(010); Z∧c = 64 25° in acute ∠β	(G)
	CaC₂H₂O₄	Calcium formate	R.	Bi	+	26° 47'	41° 2'	Ax. pl. b(010); Z‖a	(G)
	CaC₃H₂O₄.2H₂O(?)	Calcium malonate	?	Bi.	+		moderate		(27)
	CaC₄H₂O₄.2H₂O	Calcium fumarate	R.	Bi.	–	22° 24'	37° (apprx)	X = a, Y = b, Z = c	(28)
	CaC₄H₂O₄ H₂O	Calcium maleate	R	Bi	–	77° 36' (calc.)	164° (calc)	X = c, Y = a, Z = b	(28)

Mg Mn Mo N — Na Nb Nd Ni O — Os P Pb Pd — Pr Pt Ra Rb — Rh Ru S Sa — Sb Sc Se Si Sn — Sr Ta Tb Te Th — Ti Tl Tm U V — W Y Yb Zn Zr
75 43 47 11 — 82 51 61 45 1 — 35 12 23 41 — 60 37 80 84 — 40 39 5 63 — 14 56 9 13 22 — 78 52 66 10 24 — 19 27 70 49 50 — 43 57 71 28 31

Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
CaC₄H₄O₅.3H₂O	Calcium malate	R	Bi.	+			Ax. pl. b(010); Z∥a	(37)
CaC₄H₄O₄.3H₂O	Calcium succinate	?	Bi			Very large		(37)
CaC₄H₄O₆.3H₂O	Calcium mesotartrate	M	Bi.	– (?)		Very large	Ax. pl b(010)	(G, 37)
CaC₄H₁₀O₄	Calcium crotonate	(?)	Bi.	–				(37)
CaC₄H₁₀O₁₀.6H₂O	Calcium acid malate	R	Bi	+		109° 6' (red)	Ax. pl. a(100); Z∥c	(G)
Ca₂C₁₂H₆O₁₂	Calcium aconitate	?	Bi			100° (apprx.)		(37)
Ca₃C₁₂H₁₀O₁₄.4H₂O	Calcium citrate	?	Bi					(37)
CaC₄H₄O₁₀N₂.(7)H₂O	Calcium nitrotetronate(?)	M	Bi.		32° 26'		Ax pl ⊥b(010); Z nearly ⊥a (100)	(G)
Ca₃PbC₁₅H₁₀O₁₅	Dicalcium lead propionate	Tet	Un	+				(G)
Ca₄PbC₂₀H₁₀₀O₃₄.12H₂O	Tetracalcium butyrate pentalead propionate	C						(G)
CaCuC₄H₁₂O₄.6H₂O	Calcium cupric acetate	Tet	Un					(G)
78 SrC₂H₂O₄	Strontium formate	R	Bi	+	74° 14'	143° 36'	Ax pl. a(100); Z∥b	(G)
SrC₂H₂O₄.2H₂O	Strontium formate	R	Bi	–	66° 59.33'	114° 8'	Ax. pl. b(010), X∥c	(G)
SrC₂H₄O₆S₂.H₂O	Strontium disulfonate	M	Bi			Large	Ax pl ⊥(010)	(8)
SrC₄H₁₀O₈S₂.2H₂O	Strontium ethyl sulfate	M.	Bi		75° 4'		Ax. pl. ⊥b(010); Z∧c = 70° in acute ∠β	(G)
SrC₄H₄O₁₀N₂.(?)H₂O	Strontium nitrotetronate	M	Bi		30° 23'		Ax pl b(010); X⊥a(100)	(G)
SrC₈H₄O₁₂Sb₂	Strontium antimonyl tartrate	H	Un	–				(G)
Sr₂CuC₄H₄O₈.8H₂O	Cupric strontium formate	Tri	Bi		72° 4'			(L-B)
SrCa₂C₁₅H₁₀O₁₂	Dicalcium strontium propionate	Tet	Un	+				(G)
79 BaC₂H₂O₄	Barium formate	R	Bi	+	77° 54 33'		Ax. pl. b(010); Z∥a	(G)
BaC₄H₄O₆.5H₂O	Barium dl-tartrate	M	Bi	+	93° 1'		Ax. pl. ⊥b(010)	(G)
BaC₄H₄O₄.H₂O	Barium acetate	Tri	Bi					(1a)
BaC₆H₁₀O₄.H₂O	Barium propionate	R	Bi	–	81° 36'		Ax. pl. a(100); X∥b	(G)
BaC₆H₁₂O₇.(?)H₂O	Barium d-galactonate	M.	Bi.			77° 37'	Ax. pl. ⊥b(001); Z∥b	(G)
BaC₁₀H₁₆O₄.4H₂O	Barium methyluvinate	R.	Bi.		88° 12'		Ax. pl. a(100), Z∥b	(G)
BaC₆H₄O₈S₂.2H₂O	Barium m-benzenedisulfonate	R	Bi.		62° 19' (red)		Ax pl. a(100); Z∥c	(G)
BaC₆H₄O₈S₂.4H₂O	Barium phenol-2, 4-disulfonate	M	Bi	–	61° 58'		Ax pl ∥a(100), X∧c = 5° 20' in acute ∠β	(G)
BaC₂H₂N₈.3.5H₂O	Barium tetrazole	R	Bi.			40° (apprx)	Ax pl a(100); Z∥c	(G)
BaC₆H₂O₅N₂S 3.5H₂O	Barium dinitrophenol sulfonate	M	Bi			72° 13'	Ax pl b(010); X∧c = 77° in acute ∠β	(G)
BaC₄H₄O₄N₂.2H₂O	Barium methyloxamate	M.	Bi.			40° (apprx)	Ax. pl. b(010), Z∧c = 8° in obtuse ∠β	(G)
BaC₁₀H₁₀O₄N₄ 1 5H₂O	Barium methylpyrazole carbonate	Tri	Bi		56° 42'		Ax pl ⊥b(010)(apprx)	(G)
BaC₁₀H₁₀O₈P₂ 2H₂O	Barium diacetonephosphonate	R.	Bi	+		122° 44'	Ax pl b(010), Z∥c	(G)
BaC₂₆H₁₆O₆N₂S₂	Barium p-amidobenzophenone-p-sulfonate	M.					Ax pl ∥(010)	(8)
BaCdC₄H₄O₄.2H₂O	Barium cadmium formate	M.	Bi	+	67° 36'	117°	Ax pl ⊥b(010), Z∧c = 46° 23' in acute ∠β	(G)
Ba₂CuC₄H₄O₁₂	Barium copper formate	R.	Bi	+		79°	Ax pl b(010)	(G)
BaCa₂C₁₅H₁₀O₁₂	Dicalcium barium propionate	C.						(G)
81 LiC₄H₄O₅ 5H₂O	Monolithium malate	M.	Bi	–		100°	Ax pl b(010)	(G)
Li₂C₁₀H₆O₆S₂ 2H₂O	Lithium naphthalene-1, 5-disulfonate	M.	Bi		23°		Ax. pl. ⊥(010)	(41)
LiC₄H₄O₆N H₂O	Ammonium lithium tartrate	R.	Bi	+	87° 6'			(G)
LiC₄H₄O₆N H₂O	Lithium ammonium dl-tartrate	M.	Bi	+	81° 42'		Ax. pl. b(010); Z∧c = 76.5° in obtuse ∠β	(G)
LiTlC₄H₄O₆.H₂O	Lithium thallium tartrate	R	Bi.	+		24° 40' (red)	Ax. pl. c(001)(red); Z∥b	(G)
Li₂Cr₂C₁₂O₁₄ 18(?)H₂O	Lithium chromic oxalate	R	Bi	–		95° 26'	Ax pl b(010); X∥c	(G)
LiUO₂C₄H₆O₆ 5H₂O	Lithium uranyl acetate	M	Bi	–		65° 14'	Ax pl b(010); X∧c = 12° in obtuse ∠β	(G)
Li₃Al₂C₁₂O₂₄.12H₂O	Lithium aluminum oxalate	Tri	Bi	–		100° 30'	Ax pl ⊥b(010)	(G)
82 NaC₂H₃O₂ 3H₂O	Sodium acetate	M	Bi	–	62° 50'		Ax pl ⊥b(010); X∧c = 44° in acute ∠β	(G)
NaC₃H₃O₄ H₂O	Sodium acid malonate	R	Bi	–	39° 20'	55° 21'	Ax. pl. a(100), X∥c	(G)
NaC₄H₅O₆.H₂O	Sodium dl-tartrate	R	Bi	+	51° 31' (red)	83° 34' (red)	Ax. pl. a(100); Z∥c	(G)
NaC₄H₇O₄	Sodium diacetate	C						(G)
NaC₅H₅O₄	Sodium citraconate	M	Bi.	–	53° 25' (red)		Ax. pl. b(010)	(G)
NaC₈H₅O₄	Sodium acid phthalate.	R	Bi		30° (apprx.)		Ax. pl. c(001)	(G)
NaC₁₅H₁₉O₄ 3 5H₂O	Sodium santonate	R	Bi.	–		51° 46'	Ax. pl. a(100); X∥b	(G)
NaC₁₅H₂₁O₄.3H₂O	Sodium hydrosantonate	R	Bi.	+		37° 24' (red)	Ax. pl. a(100); Z∥c	(G)
NaC₆H₅O₄S 2H₂O	Sodium p-phenolsulfonate	M	Bi.	+	69° 58'	125° 47'	Ax pl. b(010); Z∧c = 9° in obtuse ∠β	(G)
NaC₇H₅O₅S 2H₂O	Sodium m-sulfobenzoate	Tri	Bi.	–		86° 7'	X⊥b(010)	(G)
NaC₇H₇O₃S	Sodium p-xylenesulfonate	R	Bi.	–		27° 46'	Ax. pl. c(001); X∥b	(G)
Na₂C₂H₄O₆S₂ 2H₂O	Sodium ethane disulfonate	M	Bi.			Large	Ax. pl (010)	(8)
Na₂C₁₀H₆O₈S₂ 2H₂O	Sodium naphthalene-1, 5-disulfonate	M	Bi.	–	24° 0 5'		Ax pl ⊥(010)	(41)
Na₂CH₂O₆N₄	Sodium diisonitramidomethane	M	Bi.	–	89° 20'		Ax. pl. b(010); X∧c = 43.66° in acute ∠β	(G)

	Ac Al As Au	B Ba Be Bi Br	C Ca Cb Cd Ce	Cl Co Cr Cs Cu	Dy Er Eu F Fe	Ga Gd Ge Gl H	Hf Hg Ho I In	Ir K La Li Lu
	28 55 13 33	54 79 75 15 5	16 77 51 29 59	4 44 46 85 31	67 69 64 3 43	25 65 20 75 2	73 30 68 6 26	36 53 56 81 72

Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
NaC₄H₅O₄N.H₂O	Sodium aspartate	M	B₁			31° 30′	Ax pl. b(010); Z∧c = 51° in acute ∠β	(G)
NaC₄H₄O₆N.H₂O	Sodium ammonium dl-tartrate	M	B₁	−	44° 20′		Ax. pl. ⊥b(010)	(G)
NaC₄H₄O₆N.4H₂O	Sodium ammonium tartrate	R	B₁	−	59° 52′	96° 30′	Ax. pl. a(100); X‖c	(G)
NaTlC₄H₄O₆.4H₂O	Sodium thallium tartrate	R	B₁.	−		75° 49′- 76° 47′ (red)	Ax pl. a(100); X‖c	(G)
NaC₅H₈O₄N	Sodium acid glutamate	M	B₁		63° 3 5′		Ax. pl. ⊥b(010), Z⊥γ(102)	(G)
NaC₆H₆O₃NS 2H₂O	Sodium sulfamlate	R	B₁	+	65° 21′	115° 24′	Ax pl b(010), Z‖c	(O)
NaC₁₀H₇O₃NS 4H₂O	Sodium naphthalenesulfonate (stable)	M	B₁	+	69° 10′		Ax pl b(010), Z∧c = 3° 35′ in acute ∠β	(G)
NaTl₃C₄H₄O₁₁	Sodium trithallium tartrate	R	B₁	+		75° 40′	Ax pl c(001); Z‖b	(G)
NaCuC₁₅H₂₇O₁₈ 9H₂O	Sodium cupric triuranyl acetate	M	B₁	+		90° 50′	Ax pl ⊥b(010)	(O)
Na₃Fe₂C₁₂O₃₆ 10H₂O	Sodium ferric oxalate	M	B₁	−	30° 0′	46° 53′	Ax pl b(010), X∧c = 12° in obtuse ∠β	(G)
Na₃Cr₂C₁₂H₁₂O₂₄N₂ 7H₂O	Sodium ammonium chromic oxalate	M	B₁	−		98° 20′	Ax pl. ⊥(010)	(G)
NaUC₂H₃O₄	Sodium uranyl acetate	C						(G)
NaU₃MnC₁₈H₂₇O₃₆ 9H₂O	Sodium manganese triuranyl acetate	M	B₁	.		105° 30′	Ax pl ⊥b(010), X∧c = 70 5° in obtuse ∠β	(G)
Na₃AlₑC₆H₁₂O₂₄N₃ 7H₂O	Sodium ammonium aluminium oxalate	M	B₁	.		134°	Ax pl ⊥b(010); X∧c = 76° in obtuse ∠β	(G)
Na₃Al₂C₁₂H₁₂O₂₄N₃ 7H₂O	Sodium ammonium aluminium oxalate	M	B₁					(21)
Na₃Al₂C₁₂O₂₄ 10H₂O	Sodium aluminium oxalate	M	B₁	−		83° 30′	Ax pl b(010); X∧c = 7.5° in obtuse ∠β	(G)
Na₃₄Al₁₂C₇₂H₃₆₆O₃₆N₁₂	Ammonium sodium aluminium oxalate	Tr₁	B₁	−		138°	Ax pl ⊥(001); Bx₁⊥(001)	(21)
NaLiC₄H₄O₆ 2H₂O	Sodium lithium dl-tartrate	M	B₁	−	68° 57′ (red)		Ax pl b(010); X∧c = 34.5° in obtuse ∠β	(G)
83 K₂C₂O₄ H₂O	Potassium oxalate	M	B₁		82°	156°	Ax pl b(010), X∧c = 40° 45′ in obtuse ∠β	(G)
KC₂HO₄	Potassium acid oxalate	M	B₁	.	40°	64°	Ax. pl. ⊥b(010), X ⊥c(100)	(G)
KC₂HO₄ H₂O	Potassium acid oxalate	R	B₁	−		75° 40′	Ax pl c(001); X‖b	(G)
KC₄H₅O₄	Potassium acid succinate	M	B₁			113°	Ax pl ⊥b(010)	(G)
KC₄H₅O₄.2H₂O	Potassium acid succinate	R	B₁				Ax pl c(001), Z‖a	(G)
KC₄H₅O₆	Potassium acid tartrate	R	B₁			161° 40′	Ax pl c(001); X‖b	(G)
KC₈H₁₁O₈	Potassium acid disuccinate	M	B₁	.		122° 50′	Ax pl ⊥b(010); X∧c = 44° in obtuse ∠β	(G)
K₂C₄H₄O₆ ½H₂O	Potassium tartrate	M	B₁	.	62°	102° 16′ (red)	Ax pl ⊥b(010)	(G)
K₂C₄H₄O₆ 2H₂O	Potassium dl-tartrate	M	B₁			130° 2′ (red)		(G)
K₄C₄H₂O₁₂ 2H₂O	Potassium tetraoxalate	R	B₁				Bx₁⊥(001)	(13)
K₆C₁₂O₁₂ 9H₂O	Potassium mellitate	R	B₁	−		73° 30′	Ax pl b(010); X‖c	(G)
KCH₂O₄S	Potassium formaldehyde sulfite	M	B₁			98° 18′	Ax pl. b(010)	(G)
KC₆H₅O₄S	Potassium phenolsulfonate	R	B₁	+	69° 4′ (apprx)		Ax pl c(001); Z‖b	(G)
KC₆H₅O₄S 2H₂O	Potassium phenolsulfonate	R	B₁	+			Ax pl a(100); Z‖c	(G)
KC₆H₅O₄S	Potassium phenylsulfate	R	B₁	+		87° 58′	Ax pl. b(010), Z‖c	(G)
KC₇H₇O₃S H₂O	Potassium p-toluenesulfonate	R	B₁	−	67° 4′		Ax pl a(100), X‖b	(G)
K₂CH₂O₆S₂	Potassium methanedisulfonate	M	B₁		72°		Ax pl ⊥b(010), Z∧c = 41° in obtuse ∠β	(G)
K₂C₆H₄O₆S₂ H₂O	Potassium m-benzenedisulfonate	M	B₁.			96° (apprx)	Ax pl ⊥b(010)	(G)
K₂C₆H₄O₇S₂ H₂O	Potassium phenoldisulfonate	R	B₁	−	65° 35′		Ax. pl. b(010); X‖a	(G)
KC₆H₄O₃SCl	Potassium p-chlorobenzenesulfonate	M	B₁		81° 25′ (red)		Z‖b	(G)
K₂C₁₀H₆O₆S₂ 2H₂O	Potassium napthalene-1, 5-disulfonate	M	B₁		38° 50′		Ax. pl. ⊥(010); η₍α₎∧c = 78°	(41)
KC₈H₄O₃N	Potassium phthalaminate.	R	B₁	−		21° 2′	Ax pl b(010); X‖a	(G)
KC₇H₃O₆N₂	Potassium 3, 5-dinitrobenzoate	M	B₁	−		55° 25′	Ax pl. b(010); X∧c = 65° in acute ∠β	(G)
KC₆H₂O₇N₃	Potassium picrate	R	B₁	−	33° 34′	67° 39′	Ax pl a(100), X‖c	(G)
KC₅H₃N₄O₄	Potassium acid uroxanate		B₁					(21)
KC₄H₄O₇Sb.H₂O	Potassium antimonyl tartrate	R	B₁	−	42° 34′	72° 50′	Ax pl. c(001), X‖b	(G)
K₂IrC₄O₄Cl₂ H₂O	Potassium iridium chloroxalate	M	B₁	+	76° 23′		Ax. pl b(010); Z∧c = 13° 53′ in obtuse ∠β	(G)
K₂PtC₂O₂N₂ H₂O	Potassium platino nitrito oxalate	M	B₁		89° 40′		Ax pl ⊥b(010)	(O)
K₄Fe₂C₁₂O₂₄ 6H₂O	Potassium ferric oxalate	M	B₁	−	80° 4′ (red)		Ax pl b(010), X∧c = 1.25° in obtuse ∠β	(G)
K₂NiC₄O₈S₄	Potassium nickel dithioxalate	M	B₁					(27)
KCaC₄H₄O₁₂Sb₂N H₂O	Calcium antimonyl tartrate potassium nitrate	R	B₁			64° 1′	Ax. pl. a(100); Z‖b	(G)
KLiC₂H₄O₆S₂ H₂O	Lithium potassium ethanedisulfonate	M	B₁		82°		Ax pl. (010); Bx₁ ⊥(001) = 41° in obtuse ∠β	(6)
KLiC₄H₄O₆ H₂O	Lithium potassium tartrate	R	B₁	−	73° 58′		Ax pl b(010); X‖a	(G)
KNaC₄H₄O₆ 4H₂O	Sodium potassium tartrate	R	B₁	+	69° 40′	117° 2′	Ax pl. b(010); Z‖a	(G)
KNaC₄H₄O₁₂Sb N H₂O	Potassium antimonyl tartrate sodium nitrate	R	B₁	−		90° 45′	Ax pl c(001), X‖a	(G)
KNaC₁₂H₁₂O₂₄SbN 2H₂O	Potassium antimonyl tartrate sodium nitrate	R	B₁.	−		88° 37′	Ax pl b(010); X‖c	(G)
K₂NaIrC₂O₄Cl₂ 2H₂O	Potassium sodium iridium chloronitrito oxalate	R	B₁.	+		63° 24′	Ax. pl. a(100); Z‖b	(G)

Mg Mn Mo N   Na Nb Nd Ni O   Os P Pb Pd   Pₑ Pt Ra Rb   Rh Ru S Sa   Sb Se Si Sn   Sr Ta Tb Te Th   Ti Tl Tm U V   W Y Yb Zn Zr
76 42 47 11   82 51 61 45 1   35 12 23 41   60 37 80 84   40 39 8 63   14 56 9 18 22   78 52 66 10 24   19 27 70 49 50   48 57 71 28 21

	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit
84	Rb₂C₄H₄O₆.2H₂O	Rubidium dl-tartrate	M	Bi	−	56° 6′		Ax. pl. b(010); X ∧ c = 82° 18′ in acute ∠β	(G)
	Rb₂C₄H₄O₆ H₂O	Rubidium mesotartrate	Tri	Bi	−	75° 18′		Ax. pl. 19° with c-axis	(G)
	Rb₂Al₂C₁₂O₂₄ 6H₂O	Rubidium aluminium oxalate	M	Bi	−	80° 22′		Ax. pl. (010)	(G)
	RbLiC₄H₄O₆ H₂O	Lithium rubidium tartrate	R	Bi	−	57° 10′ (red)		Ax. pl. c(001); X∥a	(G)
	Rb₂Na₂Cr₂C₁₂O₂₄ 7H₂O	Sodium rubidium chromic oxalate	M	Bi	−		56°	Ax pl. b(010); X ⊥c(001)	(G)
	Rb₁₂Na₁₂Al₂C₆O₄₈ 23H₂O	Sodium rubidium aluminium oxalate	M	Bi	−		24° 30′	Ax. pl. b(010); X ⊥(001)	(G)

ℭ-TABLE

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit
21	CHI₃	Iodoform	H	Un	−				(G)
55	CH₄ON₂	Urea	Tet	Un					(G)
58	CH₄N₂S	Thiourea	R	Bi	−		69° 54′– 70° 59′	Ax pl a(001); X∥b	(G)
64.1	CH₅O₃As	Methyl arsenate	M	Bi	−	14° 24′		Ax. pl. ⊥b(010); X ∧ c = 53° 20′ in acute ∠β	(G)
70	CH₄O₄N₂	Urea nitrate	M	Bi.	−		23° 10′	Ax pl. b(010), X ⊥c(001)	(G)
	CH₁₀O₄N₂S	Ammonium methanedisulfonate	M	Bi	−	79° 34′		Ax. pl. ⊥b(010), X ∧ c = 39° in obtuse ∠β	(G)
84.1	C₂Cl₄Br₂	1, 2-Dibromo-1, 1, 2, 2-tetrachloroethane	R	Bi	−		87° 45′	Ax. pl. a(100); X∥c	(G)
87	C₂Br₆	Hexabromoethane	R	Bi	−		79° 30′	Ax. pl. a(100); X∥c	(G)
92	C₂Cl₆	Hexachloroethane	R	Bi	−		66° 28′	Ax. pl. a(100)	(G)
	C₂O₂N₃I₃	Diiodofuroxane	R	Bi		63° 38′		Ax pl. c(001); Z∥b	(G)
147	C₂H₂O₄	Oxalic acid	R	Bi	+			Ax. pl. c(001); Z∥b	(G)
	C₂H₂O₄ 2H₂O	Oxalic acid	M	Bi	−	68°		Ax. pl. ⊥b(010), X∥b	(G)
161	C₂H₃O₂Cl₃	Chloral hydrate	M	Bi	−	20° 48′	35° (apprx)	Ax. pl b(010); X ∧ c = 58° 45′ in obtuse ∠β	(G)
238	C₂H₅ON	Acetamide (Unst mod)	?	Bi.			120° (apprx)		(37)
238	C₂H₅ON	Acetamide (St mod)	Trig	Un	−				(G)
248	C₂H₅O₄N H₂O	Ammonium hydrogen oxalate	R	Bi	−		22° 32′	Ax. pl. a(100); X∥c	(G)
	C₂H₆O₂NCl	Glycocoll hydrochloride	R	Bi	−		63° 50′	Ax. pl. a(100); X∥b	(G)
303	C₂O₄H₈N₂.H₂O	Ammonium oxalate	R	Bi	−	61° 44′	110° 8′	Ax. pl. a(100); X∥c	(G)
306	C₂H₁₀N₂Cl₂	Ethylenediamine hydrochloride	M	Bi.	−	81° 4′		Ax. pl b(010); X ∧ c = 6° in acute ∠β	(G)
308.1	C₃N₃Cl₃	Cyanuric trichloride	M	Bi			28°	Ax pl. ⊥b(010)	(G)
313.1	C₃H₂ON₂Br₂	Dibromocyanacetamide	M	Bi	+		29° 52′	Ax pl ⊥b(010); Z ∧ c = 31° in obtuse ∠β	(G)
	C₃H₃N₂Cl	4-Chloropyrazole	R	Bi	+		100° (apprx)	Ax. pl. a(100)	(G)
	C₃H₄O₃Br₂ H₂O	Dibromopyroracemic acid	M	Bi	+	34° 9′		Ax pl ⊥b(010)	(G)
	C₃H₄ON₂S	Pseudothiohydantoin	R	Bi	−		81° 30′	Ax pl a(100), X∥b	(G)
	C₃H₄O₃N₂S	Pyrazol-4-sulfonic acid	Tet	Un.					(L-B)
436	C₃H₆O₂N₂	Malonamide (metast mod)	Tet	Un.	−				(G)
444	C₃H₄O₃N₄	Ammonium fulminurate	M	Bi					(G)
	C₃H₇O₂N	β-Alanine	R	Bi	−		70° (apprx)	Ax. pl. c(001); X∥b	(G)
	C₃H₁₀NBr	Trimethyl ammonium bromide	M	Bi	+		50° (apprx)	Ax. pl. (010)	(G)
	C₃H₁₀NI	Trimethyl ammonium iodide	M	Bi	+		53° (apprx)	Ax. pl. (010)	(G)
535	C₃H₁₂O₃N₆	Guanidine carbonate	Tet	Un					(G)
	C₄H₃O₂NBr₂	Dibromosuccinimide	M	Bi	+		20° 50′	Ax. pl. b(010); Z ∧ c = 8° in obtuse ∠β	(G)
579.1	C₄H₅O₄N 2H₂O	Nitrotetrolic acid	M	Bi				Ax pl. b(010)	(G)
	C₄H₄O₂Br₂	trans-α-β-Dibromocrotonic acid	M	Bi		56° 1′		Ax. pl. ⊥b(010)	(G)
	C₄H₄O₂N₂	Mesotartaric acid nitrile	M	Bi	+	50° (apprx.)		Ax pl ⊥b(010), Z ∧ c = 35° in obtuse ∠β	(G)
	C₄H₅O₂Cl	α-Chlorocrotonic acid	M	Bi.	+	68° 17′			(G)
592	C₄H₅O₂N (St mod)	Succinimide	R	Bi.			90°	Ax. pl. (010); Bx_a ⊥(010)	(28)
602	C₄H₆Br₄	Butadiene tetrabromide	R	Bi.	+		57° (apprx)	Ax pl a(100); Z∥c	(G)
	C₄H₆O₂NCl₃	Ammonium trichloroisobutyrate	R	Bi	+		96°	Ax pl. c(001)	(G)
	C₄H₆O₃N₂S	3-Methylpyrazole-4-sulfonic acid	M	Bi		53°	92°	Ax. pl. ⊥b(010); Z∥b	(G)
610	C₄H₆O₃N₄	Allantoin	H	Un					(21)
	C₄H₆O₄Se	Selenodiglycolic acid	M	Bi		78° 30′		Ax. pl b(010); Z ∧ c = 41° in obtuse ∠β	(G)
640	C₄H₆O₆ H₂O	dl-Tartaric acid	Tri	Bi		67° 10′		Ax. pl. ∥p(110)	(G)
	C₄H₇O₄N	dl-Aspartic acid	M	Bi		81° 44′		Ax. pl. ⊥b(010)	(G)
	C₄H₇O₃N	Acetamide oxalate	R	Bi	−		25°	Ax. pl. a(100); X∥c	(G)
697.1	C₄H₈O₂Cl₂	Dichlorobutylene glycol	Trig	Un					(G)
	C₄H₈O₇NSb H₂O	Ammonium antimonyl tartrate	R.	Bi.	−		130° 46′	Ax. pl c(001); X∥b	(G)
708	C₄H₈O₃N₂ H₂O	Asparagine	R.	Bi.	+	l. 86° 40′ d. 87° 16′		Ax pl b(010); Z∥c	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
709	$C_4H_7O_4N_3$	Tartramide	R	Bi	−	.	43° (apprx.)	Ax. pl. b(010); X‖a	(G)
	$C_4H_9O_4N$	Ethylamine dioxalate	M	Bi	−		89° 20'	Ax. pl. b(010)	(G)
776	$C_4H_9O_6N$	Ammonium hydrogen malate	R	Bi	−	47° 54'	75° 24'	Ax. pl. b(010); X‖c	(G)
778	$C_4H_9O_6N$	Ammonium hydrogen tartrate	R	Bi	-	79° 54'		Ax. pl. c(001); X‖b	(G)
786	$C_4H_9N_3O_3$	Guanidine lactate	R	Bi	+	79° 12'		Ax. pl. a(100), Z‖b	(G)
788	$C_4H_{10}N_2S_2$	Ethylenediamine thiocyanate	M	Bi.	−	51°	89° 20'	Ax. pl. b(010), X∧c = 64° 30' in obtuse ∠β	(G)
808	$C_4H_{10}O_4$	ι-Erythrite	Tet	Un					(G)
	$C_4H_{12}NI$	Diethyl ammonium iodide	R	Bi	+		52° 15' (apprx.)	Ax. pl. (001); Z‖a	(G)
	$C_4H_{11}O_5N_1$	Ammonium malate	R	Bi		47° 34' (red)			(L-B)
835	$C_4H_{11}O_6N_1$	Ammonium tartrate	M	Bi.	−	39° 36'	64° 46'	Ax. pl. b(010), X∧c = 18° 41' in obtuse ∠β	(G)
835.1	$C_4H_{11}O_6N_1$	Ammonium racemate	M	Bi	+	60° 54		Ax. pl. b(010)	(G)
	$C_5H_5O_3Cl$	Chlorocitraconic acid	R	Bi	+	46° 24' (blue)	75° 5' (blue)	Ax pl. b(010), Z‖c	(G)
	$C_5H_4O_4N_2.H_2O$	Pyrazole dicarboxylic acid	M	Bi		77°		Ax. pl. ⊥b(010); Z apprx ⊥a(403)	(G)
868	$C_5H_4O_6$	Aconic acid	R	Bi	−			Ax. pl. a(100); X‖b	(G)
877	$C_5H_5O_2N$	Pyrrole-2-carboxylic acid	M	Bi	+	62° 7'		Ax. pl. b(010); Z∧c = 23° 45' in obtuse ∠β	(G)
	$C_5H_5O_5N_3$	Urimidosuccinic acid	R	Bi	+	78° 14'		Ax. pl. a(100); Z‖c	(G)
900	$C_5H_6O_4$	Itaconic acid	R	Bi	+		97° 40' (red)	Ax. pl. b(010); Z‖a	(G)
	$C_5H_7O_5Br$	Citrabromopyrotartaric acid	M	Bi		76°		Ax pl. ⊥b(010), Z∧c = 62° in acute ∠β	(G)
	$C_5H_7O_3N_3$	Urimidosuccinic acid amide	M	Bi		79° 35'		Ax pl. b(010)	(G)
947.1	$C_5H_6O_4$	Methyltetronic acid lactone	R	Bi	+		120° 10'		(14)
957	$C_5H_8O_6.H_2O$	Methyl hydrogen d-tartrate	R	Bi		60° (apprx)		Ax pl. a(100); Z‖c	(G)
	$C_5H_9O_7Br$	Bromohydrotiglic acid	M	Bi			150°		(G)
	$C_5H_9O_2N$	Hydroxypiperidone	M	Bi	+		92° 33'	Ax pl. ⊥b(010); Z nearly ⊥a(100)	(G)
975.1	$C_5H_9O_3N$	α-Acetylaminopropionic acid	R	Bi	−	36° 9'		Ax pl. a(100); X‖c	(G)
977	$C_5H_9O_4N$	d(l)-Glutamine acid	R	Bi	−	40° 27'	66° 35'	Ax. pl. b(010); X‖a	(G)
988.1	$C_5H_9O_4NCl$	d(l)-Glutamine acid hydrochloride	R	Bi	+	70° 44'		Ax. pl. a(100); Z‖b	(G)
994.1	$C_5H_{10}O_2N_2$	Dimethylmalonamide	R	Bi	+		58° 27'	Ax. pl. b(001); Z‖c	(G)
996	$C_5H_{10}O_4N_2$	Amylene nitrosate	M	Bi	+	62° 55'	103° 53'	Ax. pl. ⊥b(010); Z∧c = 7° in obtuse ∠β	(G)
1035	$C_5H_{10}O_5$	d-Lyxose	M	Bi	−			Ax pl. b(010)	(G)
1070.2	$C_5H_{10}O_4N$	Methyltetronamide	Not det	Bi	+		Large		(14)
	$C_5H_{12}NBr$	Piperidine hydrobromide	R	Bi		35° (apprx)		Ax pl. b(010); Z‖a	(G)
1075	$C_5H_{12}NCl$	Piperidine hydrochloride	R	Bi	−	52° 56'		Ax pl. c(001); X‖a	(G)
1093	$C_5H_{12}O_4$	Pentaerythritol	Ditet	Un					(G)
	$C_5H_{13}NBr_2$	Trimethyl-bromoethyl ammonium bromide	M	Bi	+		40° 2'	Ax pl. ⊥(010); Z∧c = 39° 30' in acute ∠β	(G)
	$C_6O_2N_2Br_4$	1, 2, 3, 5-Tetrabromodinitrobenzene	M	Bi	−		45° 54'	Ax pl b(010); X⊥r(201)	(G)
	C_6OCl_8	β-Octochlorocyclohexenone	R	Bi	+			Ax pl. b(010); Z‖a	(G)
	C_6OCl_8	γ-Octochlorocyclohexenone	M	Bi	.	37° 38'	65° 59'	Ax. pl. b(010); X∧c = about 93° in obtuse ∠β	(G)
1120	C_6HCl_5O	Pentachlorophenol (β-mod).	M	Bi	+		65° 23.5'	Ax pl. ⊥b(010), Z∧c = 3° in acute ∠β	(G)
	$C_6H_2O_4N_2Br_2$	1, 3-Dinitro-4, 6-dibromobenzene (St mod)	R	Bi	+		56° 52'	Ax. pl. a(100); Z‖c	(G)
	$C_6H_2O_4N_2Br_2$	1, 3-Dinitro-4, 6-dibromobenzene (metast mod)	R	Bi	−		73° 5'	Ax pl. ⊥b(010); X⊥a(100)	(G)
	$C_6H_2O_4N_2Br_2$	1, 2-Dinitro-4, 5-dibromobenzene	R	Bi	−	2H −	88° 22'	Ax pl. a(100); X‖c	(G)
	$C_6H_2O_2NBr_3$	2, 4, 6-Tribromonitrobenzene	M	Bi	--		90° 13'	Ax pl ⊥b(010)	(G)
1142	$C_6H_3O_4N_2I_2$	1, 3-Dinitro-2, 4-diiodo-benzene	R	Bi	+	63° 26'		Ax. pl. b(010); Z‖c	(G)
1149	$C_6H_3O_4N_2Br$	3-Bromo-1, 2-dinitrobenzene	R	Bi	+	51° 30' (red)		Ax. pl. b(010); Z‖c	(G)
1155	$C_6H_3O_2NBr_2$	3, 5-Dibromonitrobenzene	M	Bi	−		72° 19'	X∧c = 29° in obtuse ∠β	(G)
1155.1	$C_6H_3O_3NBr_2$	Nitrodibromophenol	M	Bi		70° 73°		Ax pl. ⊥b(010)	(G)
1163	$C_6H_3O_4N_2Cl$	4-Chloro-1, 2-dinitrobenzene	M		−	45° 31'		Ax pl. ⊥b(010)	(G)
1165	$C_6H_3O_4N_2Cl$	α-4-Chloro-1, 3-dinitrobenzene (St mod)	R	Bi		102° 46' (red)		Ax. pl. b(010), Z‖c	(G)
1165	$C_6H_3O_4N_2Cl$	α-4-Chloro-1, 3-dinitrobenzene (metast mod)	R	Bi	+	94° 15'		Ax. pl. a(100); Z‖b	(G)
1174.1	$C_6H_3O_3NCl_2$	4, 6-Dichloro-2-nitrophenol	M	Bi	−		62° 29'		(G)
	$C_6H_3O_3NI_2$	2, 6-Diiodo-4-nitrophenol	Tri	Bi			55° 30'		(G)
1200	$C_6H_3O_8N_6$	Tetranitroaniline	M or Tri	Bi.	−		120° (at least)		(37)
1216	$C_6H_4O_2NCl$	m-Chloronitrobenzene	R	Bi.	−		91° 23'	Ax. pl. a(100); X‖a	(G)
	$C_6H_4O_4NSCl$	p-Nitrobenzenesulfonyl chloride	M	Bi.	−		65° (apprx)	Ax. pl. b(010); X∧c = 33° 36' in obtuse ∠β	(G)
1243	$C_6H_4O_4S_2Cl_2$	m-Benzenedisulfonyl chloride..	M	Bi.	−		80° 35'	Ax. pl. b(010); X∧c = 85° in obtuse ∠β	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
1274	C₆H₄O₅N₂	2, 3-Dinitrophenol	M.	Bi.			16°	Ax. pl. ⊥(010)	(29)
1277	C₆H₄O₅N₂	2, 6-Dinitrophenol	R.	Bi	+		95° 40′	Ax. pl. b(010); Z‖a	(G)
1278	C₆H₄O₅N₂	3, 4-Dinitrophenol	Tri.	Bi			65°		(29)
1377	C₆H₆NBr	p-Bromoaniline	R.	Bi	+		26° 57.5′	Ax pl. c(001); Z‖a	(G)
	C₆H₅O₂NCl	Nicotinic acid hydrochloride	R.	Bi	−		96° 22′	Ax. pl. a(100); X‖c	(G)
	C₆H₅O₂NCl	Picolinic acid hydrochloride	R	Bi	−	41° 16′	73° 52′	Ax. pl. b(010); X‖c	(G)
1384	C₆H₆Cl₆	α-trans-Benzenehexachloride	M	Bi	+		62° 2′	Ax. pl. b(010); Z∧c = 42° 25′ in obtuse ∠β	(G)
	C₆H₆ON₂	Picolinamide	M	Bi	+		73° 20′ (red)	Ax. pl. b(010)	(G)
	C₆H₆O₂N₂	2-Methylpyrazine-5-carboxylic acid	R	Bi			35° (apprx)	Ax. pl. a(100); Z‖c	(G)
	C₆H₆O₄N₂S	p-Nitrobenzenesulfamide	M	Bi		59°		Ax. pl. b(010); Z∧c = 70° in acute ∠β	(G)
1412	C₆H₆O₇N₄	Ammonium picrate	R	Bi	−		56°		(37)
1414	C₆H₆O₂	o-Dihydroxybenzene	M	Bi	+		58° (apprx)	Ax. pl. ⊥b(010); Z∧c = 6°-7°	(G)
1415	C₆H₆O₂	Resorcinol	R	Bi	−	46° 14′	76° 6′	Ax. pl. c(001); X‖a	(G)
1416	C₆H₆O₂	Hydroquinone	Trig	Un					(G)
	C₆H₆O₃ 2H₂O	Phloroglucinol	R	Bi	−		63° 49′	Ax pl c(001); X‖a	(G)
	C₆H₆O₃	α-Methyl-β-hydroxy-γ-pyrone (β-mod)	R	Bi			Small	Ax pl (001); Bx₀ = b-axis	(30)
1448	C₆H₅ON	p-Aminophenol	R	Bi	−		47° 37′	Ax pl c(001); X‖a	(G)
	C₆H₅O₂NS	Phenylsulfohydroxamic acid	R	Bi	+		43° 29′	Ax. pl. c(001); Z‖a	(G)
	C₆H₈NBr	Aniline hydrobromide	R	Bi	−		35°	Ax. pl a(100)	(G)
	C₆H₆O₂Br₄	Tetrabromocaproic acid	M	Bi.	+		21° 52′	Ax pl. ⊥b(010); Z∧c = 100° in obtuse ∠β	(G)
	C₆H₆O₄N₂Cl₂	1, 4-Dichloro-1, 4-dinitrosocyclohexane	M	Bi.	+	61° 58′ (blue)	100° 15′ (white)	Ax pl b(010); Z∧c = 40° 30′ in acute ∠β	(G)
	C₆H₈O₄NCl₂ 2H₂O	Ammonium trichlorodihydroxycyclopentane carboxylate	R	Bi			81° (apprx)	Ax pl. (100)	(4)
	C₆H₈N₂	2, 6-Dimethylpyrazine	M	Bi			86° (apprx)	Ax. pl. b(010); Z∧c = 20° in obtuse ∠β	(G)
1507	C₆H₈O₇ H₂O	Citric acid	R	Bi	+	65° 12′	108° 40′	Ax pl a(100), Z‖a	(G)
1523	C₆H₉O₃NS	Ammonium benzenesulfonate	R	Bi	+		33° 36′	Ax pl. a(100); Z‖c	(G)
	C₆H₉O₃N	Trimorpholine	M	Bi	+	80°		Ax pl b(010)	(G)
	C₆H₉O₃N	Acetamide dioxalate	Tri	Bi	−		69° 20′		(G)
	C₆H₁₀O₆Br₂	Inosite dibromhydrin	R	Bi	+	67° 30′		Ax. pl. b(010); Z‖a	(G)
	C₆H₁₀ClNO₃	Trimorpholine hydrochloride	M	Bi			50° 60′	Ax. pl ⊥b(010) (red)	(G)
1562	C₆H₁₀O₄	Adipic acid	M	Bi	−		47° 30′	Ax pl b(010)	(G)
1563	C₆H₁₀O₄	1, 1-Dimethylsuccinic acid	M	Bi		16° 12′	41° 28′	Bx₈ nearly ⊥(001), Ax pl (010)	(22)
	C₆H₁₀O₅	l-Glycosan (l-Glucose anhydride)	R	Bi			71° 45′	Ax pl a(100); X‖c	(G)
	C₆H₁₀O₅	dl-Dilactylic acid	R	Bi			65°	Ax pl ‖(010), Bx₈ ⊥(001)	(17)
	C₆H₁₀O₅	Dilactylic acid	R	Bi	−		65° (apprx)	Ax pl b(010); X‖c	(G)
	C₆H₁₀O₅	Isosaccharine	M.	Bi.	+		25° 19′	Ax. pl. ⊥b(010), Z∧c = 63° 15′ in obtuse ∠β	(G)
	C₆H₁₀O₇N	Acetamide ditartrate	M	Bi	−		70° 30′	Ax pl b(010), X∧c = 30° in acute ∠β	(G)
	C₆H₁₁O₃N₂	Pyrrolidine-α, α-dicarboxylic acid diamide	R	Bi	+		63° 30′ (apprx)	Ax. pl. b(010), Z‖c	(G)
	C₆H₁₁O₄N₂S₂ H₂O	Ammonium phenol-2, 4(?)-disulfonate.	M	Bi	+		113° 45′	Ax. pl b(010); Z∧c = 25° 21′ in obtuse ∠β	(G)
.	C₆H₁₂O₄	cis-o-Dihydroxyhexahydrobenzene	R.	Bi	+		53° 10′	Ax pl b(010); Z‖c	(G)
	C₆H₁₂O₅	α-Methylxyloside	M.	Bi.	−	35° 14′	54° 55′	Ax pl b(010); X∧c = 30° in acute ∠β	(G)
1670	C₆H₁₂O₅	d-Quercitol	M.	Bi.	+		58° 1′	Ax pl b(010); Z∧c = 11° 46′ in acute ∠β	(G)
1672	C₆H₁₂O₅ H₂O	β-Rhamnose	M	Bi		58° 5′		Ax pl b(010)	(G)
	C₆H₁₂O₆ 2H₂O	d(l)-Inosite	R	Bi	+		42° 30′	Ax. pl. a(100); Z‖c	(G)
	C₆H₁₂O₆ 2H₂O	Damboso ("meso")-inosite)	M	Bi	+		47° 20′	Ax. pl ⊥b(010); Z∧c = 17° in obtuse ∠β	(G)
	C₆H₁₂O₅N H₂O	Ammonium hydrogen ethoxysuccinate	R	Bi			20° (apprx)	Ax pl c(001); Z‖b	(G)
	C₆H₁₂ON₂	2-Propylantipyrine	M	Bi		52° 50′			(L-B)
	C₆H₁₂OS₂N₂Cl₂	Cystine hydrochloride	M	Bi	+		3° 16′	Ax pl ⊥b(010); Z⊥a(10̄1)	(G)
1750	C₆H₁₄O₆	Dulcitol	M.	Bi	−		151° 10′ (red)	Ax. pl ⊥b(010); X‖b	(G)
1751	C₆H₁₄O₆	d-Mannitol (α-mod)	R.	Bi	−		100° (apprx)	Ax pl c(001); X‖b	(G)
1751	C₆H₁₄O₆	d-Mannitol (β-mod)	R	Bi	−		71° 30′	Ax pl a(100); X‖b	(G)
1752.1	C₆H₁₄O₆ ½H₂O	Sorbitol	M.	Bi.	−		100° (apprx)	Ax pl b(010); Z nearly ⊥c(001)	(G)
1769 1	C₆H₁₅PS	Triethylphosphine sulfide	H.	Un					(G)
	C₆H₁₆N₂Br₂ H₂O	β-2, 5-Dimethylpiperazine hydrobromide	R	Bi.	+		72° (apprx)	Ax. pl. a(100); Z‖c	(G)
	C₆H₁₆NI	Dimethyl diethyl ammonium iodide	R	Bi.			82°	Z‖c	(G)
	C₇H₃O₃Cl₅	1-Methyl-1, 3, 3, 5, 5-pentachlorocyclohexan-2, 4, 6-trione	R	Bi	+		15° (apprx)	Ax pl a(100). Z‖c	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
1789	C₇H₃O₆N₃	2, 4, 6-Trinitrobenzoic acid	R	Bi	+		84° 36′	Ax. pl. c(001); Z∥b	(G)
	C₇H₄O₃Cl₂	3, 5-Dichlorosalicylic acid	R	Bi	+		29° 15′	Ax. pl. b(010); Z∥a	(G)
1835	C₇H₄O₆N₂	2, 4-Dinitrobenzoic acid	M	Bi	—		18°	Ax. pl. (010); Bx₂ nearly ⊥(101)	(11)
1837	C₇H₄O₆N₂	2, 6-Dinitrobenzoic acid	R	Bi	+		103°	Ax pl. (100), Bx₂ ⊥(010)	(11)
1839	C₇H₄O₆N₂	3, 5-Dinitrobenzoic acid	M	Bi	—		80° 16′	Ax. pl. b(010); X∧c = 48° in acute ∠β	(G)
	C₇H₄O₆	Chelidonic acid	M	Bi	—		40° (apprx.)	Ax. pl. ⊥b(010); X nearly ∥r(101)	(G)
1843	C₇H₄O₆ 3H₂O	Meconic acid	R	Bi	—		48° 55′	Ax pl. b(010); X∥c	(G)
1881	C₇H₅O₂I	o-Iodobenzoic acid	M	Bi			70° (apprx.)	Ax. pl. ⊥b(010); Bx₂ ∥c-axis	(G)
1903	C₇H₅O₄N 2H₂O	Dipicolinic acid	R	Bi			99°	Ax pl. (001); Bx ⊥(010)	(11)
1909	C₇H₅O₅N	5-Nitro-2-hydroxybenzoic acid	M	Bi	+		105° 38′		(G)
1977	C₇H₆N₂	Benzimidazol	R	Bi	+	86° 45′		Ax pl. c(001); Z∥b	(G)
1979	C₇H₆N₂	Indazole	M	Bi		50° (apprx.)		Ax pl. b(010); Z∧c = 18° in obtuse ∠β	(G)
1985	C₇H₆O₄N₂	2, 4-Dinitrotoluene	M	Bi				Ax pl. ⊥b(010); X∧c = 32° in acute ∠β	(G)
1987	C₇H₆O₄N₂	2, 6-Dinitrotoluene	R	Bi.	—			Ax pl. a(100); X∥c	(G)
1989	C₇H₆O₄N₂	3, 5-Dinitrotoluene	M	Bi	—		98° 4′	Ax pl. ⊥b(010)	(G)
	C₇H₆ON₄ H₂O	c-Phenylhydroxytetrazole	R	Bi		60° 70′		Ax pl. a(100); Z∥c	(G)
2074	C₇H₇O₂N	Anthranilic acid	R	Bi		78° 30′ (Hg. yellow)		Ax pl. c(001); Z∥a; Bx₂ ⊥(100)	(G)
	C₇H₇O₂N	Benzohydroxamic acid	R	Bi	+	50° 2′		Ax pl. a(100); Z∥b	(G)
	C₇H₇O₂N H₂O	Pyridinebetaine	M	Bi	—	25° 16′		Ax. pl. b(010); X∧c = 12° 45′ in obtuse ∠β	(G)
	C₇H₇O₄N₃	3, 5-Dinitro-p-toluidine	R	Bi					(3)
	C₇H₈ONCl	Isobenzaldoxime hydrochloride	R	Bi			100° (apprx.)	Ax pl. a(100); Z∥b	(G)
	C₇H₈O₂NCl	Pyridinebetaine hydrochloride	M	Bi	+	52° 3′	88° 8′	Ax pl. ⊥b(010); Z∧c = 9° 27′ in acute ∠β	(G)
	C₇H₈O₂N₃ H₂O	Benzenylamidine nitrite	M (?)	Bi	—		78° 55′	Ax pl. ∥d(010)	(G)
2174	C₇H₈O₂	Guaiacol	Trig	Un					(G)
2185	C₇H₈O₄	Hydrochelidonic anhydride	R	Bi	—		120° (apprx.)	Ax. pl. c(001); X∥a	(G)
	C₇H₈O₅Br	Bromo-shikimilactone	R	Un					(G)
	C₇H₉N₂Cl 2H₂O	Benzenylamidine hydrochloride	R	Bi			35° (apprx.)	Ax pl. a(100); Z∥c	(G)
	C₇H₉O₂Cl 2H₂O	α, α-Dimethyl-γ-pyrone hydrochloride	R	Bi	—		90° (apprx.)	Ax pl. a(100); X∥b	(G)
	C₇H₉ON	3-Amino-p-cresol	R	Bi	+		44° 46′	Ax pl. a(100); Z∥c	(G)
	C₇H₉ON 3H₂O	2, 6-Dimethyl-4-hydroxypyridine	M	Bi			110° 41′	Ax. pl. b(010)	(G)
2225	C₇H₉O₂N	Ammonium benzoate	R	Bi	+		67°	Ax pl. a(100); Z∥c	(G)
2233	C₇H₉O₃NS	p-Toluidine-2-sulfonic acid	M	Bi.	+		87° 54′	Ax pl. b(010); Z∧c = 8° in obtuse ∠β	(G)
2234.1	C₇H₉O₃NS	Ammonium o-sulfobenzoate	R	Bi	—	53° 29′	84° 39′	Ax pl. b(010); X∥a	(G)
	C₇H₁₀NBr	Toluidine hydrobromide	R	Bi	—	82° 37′		Ax pl. c(001); X∥b	(G)
	C₇H₁₀O₄Br₂	Dibromotrihydroxy tetrahydrobenzoic acid	R	Bi	+	76° 32′		Ax. pl. c(001)	(G)
2260.1	C₇H₁₀O₇N₂	Mono-ureindihydroxy dimethyl succinate	R	Bi		72° 15.5′		Ax. pl. b(010); Z∥c	(G)
2260.2	C₇H₁₀O₅N₄	Isohydroxydimethylurea	M	Bi	+	40° 9.5′	62° 34.25′	Ax. pl. ⊥b(010); Z∧c = 2° 15′ in acute ∠β	(G)
	C₇H₁₂O₄N₂S.2H₂O	2, 4-Toluylendiamine sulfate.	M	Bi			100° (apprx.)		(G)
	C₇H₁₂O₄	Trimethyl succinic acid	R	Bi		84° 11′		Ax pl. (100); Bx₂ ⊥(001)	(11)
	C₇H₁₄O₅	l-Methylrhamnoside	R	Bi	—	36° 11′	57° 8′	Ax pl. b(010), X∥c	(G)
	C₇H₁₄O₆	α-Methyl mannoside	R	Bi	+	46° 58′	75°	Ax pl. b(010), Z∥a	(G)
2372	C₇H₁₄O₆	α-Methyl glucoside	R	Bi	+	85° 18′		Ax. pl. b(010); Z∥c	(G)
2373	C₇H₁₄O₆	β-Methyl glucoside	Tet	Un					(G)
	C₇H₁₄O₆ H₂O	dl-α-Methyl galactoside	R	Bi	+	53° 5′	85° 45′	Ax pl. a(100); Z∥c	(G)
	C₈H₄O₄N₂Cl₂	2, 4, 6-Trichloro-3-nitrobenzoic acid methyl nitramide	M	Bi	—		42° (apprx.)	Ax. pl. ⊥b(010); X∧c = 69° in acute ∠β	(G)
	C₈H₅O₃N	Isatoic acid anhydride	M	Bi			90° (apprx.)	Ax. pl. ⊥b(010)	(G)
	C₈H₆O₂N	Phthaloxime	M	Bi					(16)
2452	C₈H₆NBr	Bromobenzyl cyanide	Trig	Un					(L-B)
	C₈H₆O₃N₃Br	1-Nitro-3-bromo-4-acetanilide (St. mod.)	M	Bi	—		124° 10′	Ax. pl. ⊥b(010)	(G)
	C₈H₆O₂Cl₄	Tetrachlorophloroglucinol dimethyl ether	R	Bi	+		90° (apprx.)	Ax. pl. a(100)	(G)
	C₈H₆O₃N₃Br	Nitrobromoacetanilide (α-mod.)	M	Bi.	—		124° 10′	Ax. pl. ⊥(010); Bx₂ nearly ⊥(001)	(3)
	C₈H₆ONCl₂	Dichloroacetanilide	M	Bi.	+	83° 35′		Ax pl. ⊥b(010); Z∧c = 61° in obtuse ∠β	(G)
2536	C₈H₇O₆N₃	2, 3, 6-Trinitro-p-xylene	M.	Bi.	—	64° 32′		Ax pl. b(010), X∧c = 28° in obtuse ∠β	(G)

Index No	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
	C_8H_9ONCl	Methylphenylurea chloride	R	Bi.			27° 41′	Ax. pl. c(001); Z∥b	(G)
	$C_9H_9ON_4$	Methoxyphenyltetrazole	Tri	Bi	−	74° 48′		Ax. pl. ⊥b-axis	(G)
2556	$C_8H_8O_3N_2$	m-Nitroacetanilide	M	Bi			80° (apprx.)	Ax. pl. ⊥b(010)	(G)
2564	$C_8H_8O_4N_2$	2, 3-Dinitro-p-xylene	M	Bi	+		105° 8′	Ax. pl. ⊥b(010)	(G)
	$C_8H_8O_4N_4$	9-Allyluric acid		Un					(21)
	$C_8H_8O_6$	Hematinic acid anhydride	R	Bi	+		53° (apprx)	Ax. pl. b(010); Z∥a	(G)
	$C_8H_8O_7$	Acetylcitric anhydride	R	Bi	−	71° 2′		Ax. pl. a(100); X∥c	(G)
	$C_8H_8N_4Cl\,H_2O$	Phenyliminotriazoline hydrochloride	M	Bi	+		120° 10′	Ax. pl ⊥b(010), Z∧c = 44° in acute ∠β	(G)
	$C_8H_9O_2SCl$	Chloromethyl p-tolyl sulfone	R	Bi	?		110° (apprx)	Ax. pl. b(010); Z∥c	(G)
2649	C_8H_9ON	Acetanilide	R	Bi	+	88° 36′		Ax. pl b(010); Z∥c	(G)
2657	$C_8H_9O_2N$	p-Acetaminophenol	M	Bi	−		90°	Ax. pl. ⊥b(010); X∥b	(G)
2681	$C_8H_9O_4N$	Biliverdic acid	M	Bi			31° (apprx)	Ax. pl ⊥b(010); X∧c = 55° in obtuse ∠β	(G)
	$C_8H_9O_4N_3$	2, 4-Dinitrodimethylamine	R	Bi	−		23° 30′	Ax. pl c(001); X∥a	(G)
	$C_8H_{10}O_2NCl$	Phenylglycocoll hydrochloride	R	Bi	−	18° 9′		Ax. pl b(010); X∥a	(G)
	$C_8H_{10}O_2$	p-Hydroxyphenylethyl alcohol (Tyrosol)	R	Bi			84° 30′	Ax pl ⊥b(010)	(*)
	$C_8H_{10}O_3$	Dimethylpyrogallol	M.	Bi	?		53° 47′	Ax. pl b(010)	(G)
	$C_8H_{12}NBr$	Xylidine hydrobromide	R	Bi			55° 19′	Ax. pl b(010); X∥a	(G)
	$C_8H_{18}O_2NBr$	Tetramethylsuccinic bromoimide.	R	Bi	−		62° 15′ (Hg, yellow)	Ax. pl (100); Bxa ⊥(001)	(24)
	$C_8H_{18}O_2NCl$	Tetramethylsuccinic chloroimide	R	Bi	−		47° 29′ (Hg, yellow)	Ax. pl (010), Bxa ⊥(001)	(24)
	$C_8H_{12}O_3NCl$	Vanillylamine hydrochloride	M	Bi	−		70°		(23)
	$C_8H_{12}NI$	Ethylaniline hydroiodide	R	Bi	−		65° (apprx)	Ax. pl. a(100); X∥c	(G)
2808 1	$C_8H_{14}O_4N_2$	Tetraacetylhydrazine	R	Bi	+	17° 5′	79° 33′	Ax. pl. c(001); Z∥b	(G)
	$C_8H_{14}O_4$	trans-Hexahydroterephthalic acid	M	Bi			65° (apprx)	Ax. pl ⊥b(010)	(G)
	$C_8H_{14}O_4$	Norpinic acid	M	Bi	+		7° (apprx)	Ax pl. ⊥b(010)	(G)
	$C_8H_{14}O_4$	Isopropylisoparaconic acid	M	Bi	+		51° 12′	Ax pl. ⊥b(010), Z∧c = 83° in obtuse ∠β	(G)
	$C_8H_{14}O_6N_2$	Lysidine d-ditartrate	M	Bi	−	80° 1′		Ax. pl b(010); X∧c = 30° in obtuse ∠β	(G)
	$C_8H_{14}O_6N_2Sb_2\,H_2O$	Ammonium antimonyl tartrate	R	Bi		68° 8′			(L-B)
2915	$C_8H_{16}O_4$	Metaldehyde	Tet	Un				Ax. pl ⊥b(010)	(G)
2916 1	$C_8H_{16}O_4$	bis-Methoxyacetol	M	Bi				Ax pl b(010); X∥a	(G)
2920	$C_8H_{16}O_6$	d, α-Ethyl glucoside	R	Bi	−	51° 14′		Ax pl b(010); X∧c = 21° in obtuse ∠β	(G)
	$C_8H_{17}N_2Cl$	1, 4-Dimethyl-5-isopropylpyrazoline hydrochloride	M	Bi	−	56°	94° 40′	Ax pl b(010); X∧c = 21° in obtuse ∠β	(G)
	$C_8H_{17}N_2Cl$	Isobutyraldazine hydrochloride	M	Bi	−	56°	94° 41′	Ax pl b(010); X∧c = 21° in obtuse ∠β	(G)
2945	$C_8H_{18}NBr$	d-Conine hydrobromide	R	Bi	+	45° 50′		Z∥c	(G)
2946	$C_8H_{18}NCl$	d-Conine hydrochloride	R	Bi	+	20° 0′		Ax pl c(001); Z∥b	(G)
2948	$C_8H_{18}NI$	d-Conine hydroiodide	M	Bi	−	107° 30′ (apprx)		Ax pl b(010)	(G)
	$C_8H_{20}PI$	Tetraethyl phosphonium iodide	Trig	Un					(G)
	$C_9H_6OBr_2$	Dibromohydrindone	R	Bi	−		36° 29′	Ax pl b(010); X∥a	(G)
	C_9H_5OBr	Phenyl-α-bromoacrolein	R	Bi	+		39°	Ax pl b(010); Z∥c	(G)
	C_9H_5OCl	Phenyl-α-chloroacrolein	R	Bi	?		22°	Ax pl a(100); Z∥c	(G)
	$C_9H_5O_2Br_2$	Phenyldibromopropionic acid	M	Bi	+		57° (apprx)	Ax pl ⊥b(010)	(G)
	$C_9H_8O_3Cl_2$	Ethyl dichlorosalicylate	R	Bi.	−			Ax pl. b(010); X∥c	(G)
3080	$C_9H_8N_2$	3-Aminoquinoline	R	Bi.	−		45°	Ax pl. c(001); X∥b	(G)
	$C_9H_8O_4$	Acetylsalicylic acid	Tri	Bi.	−	Small		Sections ⊥Bxa; elongation = Z	(42)
	$C_9H_8O_{12}N_4$	Pentaerythritol nitrate	Tet	Un.					(19)
	$C_9H_8O_6N_3Br$	Bromodinitromesitylene	M	Bi	−	42° 19′	88° 13′	Ax. pl. ⊥b(010); X∥b	(G)
	$C_9H_9Br_3$	Tribromomesitylene	Tri	Bi.			24° 3′		(G)
	$C_9H_9O_3Cl_3$	1, 3, 5-Trimethyl-1, 3, 5-trichlorocyclohexan-2, 4, 6-trione	M	Bi.	−		60° (apprx)	Ax. pl. b(010)	(G)
3103	C_9H_9ON	Hydrocarbostyril	R	Bi.	−	60° (apprx)		Ax pl a(100); X∥c	(G)
	$C_9H_9O_3N$	Benzoylacetohydroxamic acid	M.	Bi.	−	47° 10′		Ax. pl. ⊥b(010); X∧c = 66° in acute ∠β	(G)
3111	$C_9H_9O_3N$	Hippuric acid	R	Bi.	+	65° 49′		Ax pl c(001)	(G)
	$C_9H_9ON_2$	1-Phenyl-3-methylpyrrodiazoline	R.	Bi.	−		64° (red)	Ax. pl. b(010); X∥c	(G)
	$C_9H_{10}ON_2$	Isonitrosoanilacetone	R.	Bi.	−		41° 40′	Ax. pl. a(100); X∥c	(G)
	$C_9H_{10}O_4N_2$	Dinitromesitylene	R.	Bi.			50° (apprx.)	Ax. pl. a(100); X∥c	(G)
	$C_9H_{14}O$	D hydrodiacetyllevulinic acid hes	M.	Bi.	+	74° 45′		Ax. pl b(010); Z∧c = 5° in obtuse ∠β	(G)

Index No	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit
	$C_{10}H_9O_4N$	Phthalylethylhydroxylamine	R.	Bi.	−		91° 17′	Ax. pl. a(100); X‖c	(G)
	$C_{10}H_9O_4N$	Phthaloxime ethyl ether	R.	Bi.			70° (apprx.)	Bx$_a$ ⊥(001)	(26)
	$C_{10}H_9O_4N$	Dimethylnitroterephthalate	Tri.	Bi.	−		95° 30′	X ⊥b(010)	(G)
	$C_{10}H_9O_5N_3$	Nitrodinitrosomenthol peroxide	M.	Bi.		73° 48′		Ax. pl. b(010); Z∧c = 38° in acute ∠β	(G)
3585	$C_{10}H_{10}O_2N_2$	N-Phenyl-3-methylpyrazolone	M	Bi		72° 56′		Ax. pl. ⊥b(010); Z‖b	(G)
	$C_{10}H_{10}O_3N_3$	Disonitrosomenthol anhydride	M.	Bi				Ax. pl. ⊥b(010); Z∧c = 40° in acute ∠β	(G)
	$C_{10}H_{10}O_3$	Phenylisooxybutyrolactone	M	Bi				Ax. pl. b(010); Z∧c = 96° in obtuse ∠β	(G)
	$C_{10}H_{10}O_4$	2, 4-Dihydroxycinnamic acid	M.	Bi.	−		106° 20′ (red)	Ax. pl. ⊥b(010)	(G)
	$C_{10}H_{11}O_4N$·Cl	Dinitrochlorocymene	?	Bi.	+		120°		(37)
	$C_{10}H_{11}O_4N$·Cl	2-Chloro-5, 6-dinitrocymene	M.?	Bi.	−		70°		(37)
	$C_{10}H_{11}O_2N$	β-β-Dimethyl-α-indolinone	R.	Bi	−	46° 39′	81° 48′	Ax. pl. c(001); X‖a	(G)
	$C_{10}H_{11}O_2N$	β-Ethyl-α-indolinone	M.	Bi.	−		38° (apprx.)	Ax. pl. ⊥b(010)	(G)
	$C_{10}H_{11}O_4N$	Nitrocumic acid	M.	Bi.	−	36° 58′	64° 25′	Ax. pl. b(010); X∧c = 14° 11′ in acute ∠β	(G)
	$C_{10}H_{12}O_4N_2$	p-Aminophenaceturic acid	M.	Bi	−		102° 30′	Ax pl. ⊥b(010); X nearly ‖c	(G)
	$C_{10}H_{13}O_4N_3$	α-Disonitrosomenthol	M.	Bi	+		30° 45′	Ax. pl. ⊥b(010)	(G)
	$C_{10}H_{13}O_4N_3$	Ethyl N$^\omega$-phenyl allophonate		Bi					(85)
	$C_{10}H_{12}O_3$	p-Methoxyhydroatropic acid	M.	Bi	+	77° 58′		Ax pl b(010); Z∧c = 57° in acute ∠β	(G)
	$C_{10}H_{12}O_4$	Cantharidin	R.	Bi.		89° 7′		Ax. pl c(001); Z‖b	(G)
	$C_{10}H_{12}O_4S$	α-Phenylsulfonebutyric acid	R.	Bi	−	46° 45′		Ax pl. b(010); X‖a	(G)
	$C_{10}H_{12}O_4$	Methyl 4-hydroxy-3, 5-dimethoxybenzoate	M	Bi			63° (apprx)	Ax pl. b(010); X ⊥r(101)	(G)
	$C_{10}H_{13}Br_3$	Tribromocamphene	R	Bi		80° (apprx)		Ax. pl. c(001); X‖b	(G)
3700	$C_{10}H_{13}ON$	N-Ethylacetanilide	R	Bi.	+		103° 27′	Ax. pl. b(010); Z‖c	(G)
3716	$C_{10}H_{13}O_2N$	Phenacetin	M	Bi		62° 14′		Ax pl b(010)	(G)
	$C_{10}H_{13}O_2N$	p-Tolyl urethane	M.	Bi	−		59° 46′	Ax pl b(010); X∧c = 27° in acute ∠β	(G)
	$C_{10}H_{13}O_4N$	Vanillyl acetamide	M	Bi.	+		110° (115° calc)		(24)
3732	$C_{10}H_{14}$	1, 2, 4, 5-Tetramethylbenzene	M	Bi.	−	87° 22′		Ax. pl. b(010); X∧c = 0° 54′ in obtuse ∠β	(G)
	$C_{10}H_{14}O_2Br$	d-Bromopseudonitrocamphor	R.	Bi	+	79° (apprx)		Ax pl c(001); Z‖a	(G)
3742	$C_{10}H_{14}OBr_2$	d-α, α′-Dibromocamphor	R.	Bi	−	56° 5′	90° 38′	Ax pl a(100); X‖b	(G)
	$C_{10}H_{14}OBr_2$	d-α, β-Dibromocamphor	R	Bi	−	77° 51′		Ax pl b(010); X‖c	(G)
	$C_{10}H_{14}OCl_2$	d-α, π-Dichlorocamphor	R	Bi	+	62° 18′		Z‖c	(G)
	$C_{10}H_{14}O_3SCl_2$	d-α-Chloro-π-camphosulfonic chloride	R	Bi.		59° (apprx.)		Ax pl a(100); Z‖b	(G)
3756	$C_{10}H_{10}O_6N_2S_2$	Ammonium naphthalene-1, 5-disulfonate	M	Bi		49° 40′		Ax. pl. ⊥(010)	(41)
	$C_{10}H_{14}O$	Thymol	Trig	Un	+				(G)
	$C_{10}H_{14}O_3$	d(l)-Camphoric anhydride	R.	Bi	−		31° 20′ (red)	Ax pl a(100); X‖c	(G)
	$C_{10}H_{14}O_4$	Tetramethylapionol	R	Bi	+	49° 13′	80° 1′	Ax pl a(100); Z‖c	(G)
	$C_{10}H_{14}O_5$	Methyl α-anhydrocamphoronate	R.	Bi	−		120° (apprx)	Ax pl a(100); X‖b	(G)
	$C_{10}H_{14}O_5$	Methyl β-anhydrocamphoronate	R.	Bi.	−		33° (apprx)	Ax pl a(100); X‖b	(G)
	$C_{10}H_{14}O_4$	Dimethyl diacetylracemate	R	Bi	+	62° 36′	103° 29′	Ax. pl. c(001); Z‖b	(G)
3779	$C_{10}H_{15}OBr$	d-β-Bromocamphor	R.	Bi	+	76° (apprx)		Ax. pl. a(100); Z‖c	(G)
	$C_{10}H_{15}O_2N_2Br$	α-Bromopernitrosocamphor	R	Bi	+	90° 28′		Ax. pl. b(010); Z‖c	(G)
	$C_{10}H_{15}O_2N_2Br$	β-Isobromopernitrosocamphor	R	Bi	+	69° 20′		Ax pl a(100); Z‖c	(G)
	$C_{10}H_{15}O_2Br_3$	d(l)-Dihydrocarvone tribromide	R.	Bi	+	59° 45′		Ax. pl (100); Z‖c	(G)
	$C_{10}H_{15}O_3SBr$	d-π-Camphoresulfonyl bromide	R.	Bi	+	35°			(G)
	$C_{10}H_{15}O_3SCl$	d-π-Camphoresulfonyl chloride	R.	Bi	+	45° (apprx)			(G)
	$C_{10}H_{16}O_7N$	l-Ratanhin sulfate	R.	Bi		75° (apprx)		Ax pl c(001)	(G)
	$C_{10}H_{16}NBr$	Diethylaniline hydrobromide	M.	Bi.	−	77° 33′		Ax pl ⊥b(010); X∧c = 70° in obtuse ∠β	(G)
	$C_{10}H_{16}OBr_2$	Pinol dibromide	R.	Bi	−	131° 21′		Ax pl a(100), X‖c	(G)
	$C_{10}H_{16}NI$	p-Tolyltrimethylammonium iodide	R.	Bi	+	20° 36′		Ax pl b(010); Z‖c	(G)
3867.1	$C_{10}H_{16}O_3$	dl-Pinonic acid	M.	Bi		88° 32′		Ax pl b(010); Z∧c = 57° in acute ∠β	(G)
	$C_{10}H_{16}O_3$	d-α-Thugene ketonic acid	R.	Bi	+	74° 14′		Ax. pl. a(100); Z‖c	(G)
	$C_{10}H_{16}O_3$	Isoketocamphoric acid	M.	Bi	+	80° (apprx.)		Ax. pl. b(010); Z nearly ⊥c(001)	(G)
3873	$C_{10}H_{16}O_6·H_2O$	l-Cineolic acid	R.	Bi.	−	25° 30′		Ax. pl b(010); X‖c	(G)
3886.1	$C_{10}H_{17}O_2N$	dl-α-Pinonoxime	M.	Bi.	+		60°–70°	Ax pl. b(010); Z∧c = 10° in acute ∠β	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
	$C_{10}H_{16}O_2$	2-Hydroxy-Δ', 3-p-menthenone.	M	Bi.	—			X $\wedge$ c = 63° 6' in obtuse $\angle\beta$	(G)
	$C_{10}H_{18}O_5$	α, α'-Methylisopropyl-α, α'-dihydroxy-adipic acid	?	Bi.	—	75°			(87)
	$C_{10}H_{19}ON$	Δ^6, 8-Methylnonenyl amide		Bi	+	60°			(88)
3964	$C_{10}H_{19}ONCl$	Lupinine hydrochloride	R.	Bi.	+	50° 18'	102° 10'	Ax. pl. c(001), Z‖a	(G)
	$C_{10}H_{20}O_4N_2$ 3H$_2$O	α-2, 5-Dimethylpiperazine tartrate	M	Bi			80° (apprx)	Ax. pl. ⊥b(010)	(G)
	$C_{10}H_{20}NPS$	Triethylallylphosphothiourea	M.	Bi	—	72° 30'		Ax. pl. b(010), X$\wedge$c = 24° in acute $\angle\beta$	(G)
	$C_{10}H_{20}O_2$	cis-Terpine hydrate	R	Bi	+	77° 27'		Ax. pl. b(010), Z‖a	(G)
3980	$C_{10}H_{20}O_2$	trans-Terpine	M	Bi.	+		74° 15'	Ax. pl. ⊥b(001); Z$\wedge$c = 3° 6' in acute $\angle\beta$	(G)
	$C_{11}H_6O_{10}$ 5H$_2$O	Benzenepentacarboxylic acid .	R	Bi	—		57° 30'	Ax. pl. b(010), X‖c	(G)
	$C_{11}H_{12}N_4O_2$	9-Phenyluric acid .		Un					(8.8)
	$C_{11}H_7O_4Br$	Phenylbromoparaconic acid	R	Bi	—	56° 50'		Ax. pl. b(010); Z‖a	(G)
	$C_{11}H_9O_2N$	Citraconanil	M	Bi	+	14° 56'		Ax. pl. b(010)	(G)
	$C_{11}H_9O_3Cl_3$	Trichloromethyl-o-methoxy p h e n y l-carbinol acetic ether	M	Bi	—	75° 11'		Ax. pl. ⊥b(010)	(G)
	$C_{11}H_9O_2N$	Glutaric anilide	M	Bi		90°		Ax. pl. (010)	(88)
4043 1	$C_{11}H_{10}ON_3Br$	4-Bromoantipyrine	Ditrig	Un					(G)
	$C_{11}H_{11}O_3N$	β-Benzyl malimide	R	Bi	-	62° 66°		Ax. pl. b(010); X‖c	(G)
4053	$C_{11}H_{11}O_4N$	Ethyl o-nitrocinnamate	R	Bi	—		57° 40'	Ax. pl. c(001); X‖a	(G)
	$C_{11}H_{11}ON_3$	4-Iodoantipyrine	Trig	Un					(G)
	$C_{11}H_{10}O_2Br_2$	Ethyldibromocinnamate	M	Bi	--	88° (apprx)		Ax. pl. b(010); X$\wedge$c = 7° in acute $\angle\beta$	(G)
4058	$C_{11}H_{12}ON_2$	Antipyrine	?	Bi		54° 20'	103° 21'		(L-B)
	$C_{11}H_{12}O_2N_2$	4-Hydroxyantipyrine	M.	Bi			116° 23'	Ax. pl. b(010); Z⊥c(001)	(G)
	$C_{11}H_{13}O_3N$	Methyl phenaceturate	R	Bi				Ax. pl. b(010)	(G)
4086	$C_{11}H_{14}ON_2$	Cytisine	R	Bi	+	61° 36.5'		Ax. pl. a(100); Z‖c	(G)
	$C_{11}H_{14}O_2N_2$	Ethyl α-phenylhydrazine pyroracemate	M	Bi	—			Ax. pl. ⊥b(010); X$\wedge$c = 47° 4' in acute $\angle\beta$	(G)
	$C_{11}H_{14}O_4$	Methyl 3, 4, 5-methoxybenzoate	M	Bi.			113° 13' (white)	Ax. pl. ⊥b(010)	(G)
	$C_{11}H_{15}ON_3Br$ H$_2$O	Cytisine hydrobromide	M	Bi	-	87° (apprx)		Ax. pl. b(010)	(G)
	$C_{11}H_{15}O_4NCl$	Methyl 3, 4, 5-trimethoxy-2-aminoben-zoate	R	Bi	—		70° (apprx)	Ax. pl. c(001); X‖a	(G)
	$C_{11}H_{15}ON_2Cl$ H$_2$O	Cytisine hydrochloride	M.	Bi		72° (apprx)		Ax. pl. b(010); Z$\wedge$c = 55° in obtuse $\angle\beta$	(G)
	$C_{11}H_{15}O_3N$	Vanillyl propionamide	R	Bi	—	100° (98° calc)			(84)
	$C_{11}H_{15}O_3N$	Pyrocatechol carboxyl diethylamide	M.	Bi	+	7° 56'		Ax. pl. b(010); Z$\wedge$c = 55° in obtuse $\angle\beta$	(G)
	$C_{11}H_{16}O_7N$	s-Benzylhydroxylamine ditartrate	R	Bi		90° (apprx)		Ax. pl. a(100); Z‖b	(G)
	$C_{11}H_{16}O_2N_2$	Nitrosoamylene nitraniline	R.	Bi	+	82° 51'		Ax. pl. b(010); Z‖c	(G)
	$C_{11}H_{15}O_4N_3$ H$_2$O	Cytisine nitrate	M.	Bi	+	38° 49'		Ax. pl. b(010)	(G)
	$C_{11}H_{16}ON_2$	Amylene nitraniline	R	Bi	+	88° 21'		Ax. pl. a(100); Z‖c	(G)
	$C_{11}H_{18}O_4$	Dimethyl camphoronate	R.	Bi.	—		50° (apprx)	Ax. pl. b(010); X‖a	(G)
	$C_{11}H_{17}ON_2Cl$	Amylene nitraniline hydrochloride	M.	Bi	+	75° 41'		Ax. pl. ⊥b(010)	(G)
	$C_{11}H_{16}NBr$	Diethyl-p-toluidine hydrobromide	M.	Bi	+	69° 41.5'		Ax. pl. ⊥b(010)	(G)
	$C_{11}H_{18}O_4$	Ethyl camphoronate	M	Bi			56° (apprx)	Ax. pl. ⊥b(010)	(G)
	$C_{11}H_{18}O_5$	Triethyl desoxalate	M	Bi	—	61° 59'		Ax. pl. ⊥b(010)	(G)
	$C_{11}H_{20}ON_2$	Terpinene nitrolmethylamine	M	Bi		55° 20'	93° 56'	Ax. pl. ⊥b(010); Z$\wedge$c = 31° in obtuse $\angle\beta$	(G)
	$C_{11}H_{21}O_3N$	N-Methyl-2, 2, 6, 6-tetramethyl-4-hy-droxypiperidine carboxylic acid	R	Bi	--	82° 31'		Ax. pl. a(100); X‖b	(G)
4184	$C_{12}H_8$	Acenaphthylene	R	Bi	+	70° 16'	114° 46'	Ax. pl. a(100); Z‖b	(G)
4185 1	$C_{12}H_8Br_2$	p, p'-Dibromodiphenyl	M.	Bi		50° 60° (apprx)		Ax. pl. ⊥b(010)	(G)
4218	$C_{12}H_{10}$	Acenaphthene	R	Bi	+	70° 20'	115° 40'	Ax. pl. a(100); Z‖b	(G)
4221.1	$C_{12}H_{10}ICl$	Diphenyliodonium chloride	M.	Bi			Large	Ax. pl. b(010)	(G)
4225	$C_{12}H_{10}N_2$	Azobenzene	M.	Bi	+		59° 5'	Ax. pl. ⊥b(010); Z$\wedge$c = 62° in acute $\angle\beta$	(G)
	$C_{12}H_{10}ON_2$	α-Benzoylpyridine oxime	R.	Bi.		66°		Ax. pl. b(010); Z‖a	(G)
	$C_{12}H_{10}ON_2$	γ-Benzoylpyridine oxime	M	Bi		28°		Ax. pl. b(010); Z$\wedge$c = 62° in obtuse $\angle\beta$	(G)
	$C_{12}H_{10}O_4S_4$	Benzenesulfone trisulfide	Tet	Un.					(G)
4261	$C_{12}H_{10}S_2$	Diphenyl disulfide	R.	Bi.	—		85° (apprx)	Ax. pl. b(010); X‖c	(G)
	$C_{12}H_{11}O_3SBr$	Ethyl 1, 5-bromonaphthalene sulfonate	R.	Bi			29° 52'	Ax. pl. a(100); Z‖b	(G)
	$C_{12}H_{11}O_3SCl$	Ethyl 1, 5-chloronaphthalene sulfonate.	M.	Bi.		42° (apprx)		Ax. pl. b(010)	(G)
	$C_{12}H_{11}ON$	α-Phenylpyridyl carbinol	R.	Bi		65°		Ax. pl. c(001); Z‖a	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit
4272	$C_{13}H_{13}O_3NS$	Benzenesulfanilide	Tet	Un					(G)
	$C_{13}H_{17}O_3N$	Vanillyl n-butyramide	Tn	Bi.	+		Very large		(24)
	$C_{13}H_{17}O_3N$	Vanillyl isobutyramide	R.	Bi.	−		18° (17° 48′ calc.)		(24)
	$C_{13}H_{14}O_3$	Ethyl β-methylcoumarilate	R	Bi			72° 34′	Ax. pl. b(010); Z‖c	(G)
	$C_{13}H_{14}O_4$	cis-Dimethylsuccinic acid	R	Bi			124° 4′ (Hg, yellow)	Ax. pl. (010); Bx₀ ⊥(001)	(24)
	$C_{13}H_{13}O_3$	Acetotetrahydrocinchonine acid	R	Bi	−		12° 24′	X‖b	(G)
	$C_{12}H_{28}NI$	Tetrapropyl ammonium iodide	R	Bi	−		30° 1′	Ax. pl. (100); X‖b	(G)
	$C_{13}H_{16}NI$	1, 3, 3-Trimethyl-2-methylene indoline hydriodide	R	Bi		23° 48′ (red)	57° 16′ (red)	Ax. pl. c(110); X‖b	(G)
	$C_{12}H_{14}ON_2$	1-P h e n y l-3-methyl-4-dimethylpyrazolone	M	Bi		74° 2′		Ax. pl. ⊥b(010)	(G)
	$C_{12}H_{14}ON_2$	4-Methylantipyrine	M	Bi		86° (apprx)		Ax. pl. b(010); Z∧c = 47° in acute ∠β	(G)
4318.1	$C_{12}H_{14}O_4$	Ethyl p-methoxycinnamate	M	Bi			10° (apprx)	Ax. pl. b(010)	(G)
	$C_{12}H_{14}O_4$	Dimethyl phenylsuccinate	M	Bi	+			Ax. pl. ⊥b(010)	(G)
	$C_{12}H_{15}ON_2I$	1-P h e n y l-3-methyl-5-methoxypyrazole-2 methiodide	M	Bi.	−	72°		Ax. pl. b(010); X∧c = 73° in obtuse ∠β	(G)
	$C_{12}H_{15}ON_2I$	Antipyrine pseudomethiodide				75° 41′			(L-B)
	$C_{12}H_{15}ON_2I$	Antipyrine pseudoethiodide	M	Bi	+	74° 45′		Ax. pl. b(010); Z∧c = 84° 30′ in obtuse ∠β	(G)
	$C_{12}H_{15}ON$	7-Isopropylhydrocarbostyril	R	Bi		64° 51′		Ax. pl. b(010); Z‖a	(G)
4330.1	$C_{12}H_{15}O_3N$	Ethyl phenaceturate	R	Bi				Ax. pl. b(010)	(G)
	$C_{12}H_{15}O_3N$	Vanillyl crotonylamide	R	B.	+		Large		(24)
	$C_{12}H_{16}O_4$	2, 5-Dioxyacetophenone diethyl ether	Tn	Bi			85° (apprx)	Ax. pl. ⊥c(001)	(G)
	$C_{12}H_{17}O_2N_3$	Nitrosoamylenenitrol p-toluidine	R	Bi	+	77° 50′	167° 37′	Ax. pl. ⊥b(010); Z‖c	(G)
	$C_{12}H_{18}ON_2Cl$	Amylenenitrol p-toluidine hydrochloride	M	Bi	+	59° 26′	97° 30′	Ax. pl. ⊥b(010); Z∧c = 12° in obtuse ∠β	(G)
	$C_{12}H_{16}ON_2$	Amylenenitrol-p-toluidine	M	Bi	−		72° 40′	Ax. pl. b(010); X∧c = 35° in acute ∠β	(G)
	$C_{12}H_{18}O_4$	Dimethylanthantidin	R	Bi	+		116°	Ax. pl. b(010)	(G)
4368.3	$C_{12}H_{18}O_6$	Diethyl 1, 1-diacetosuccinate	M.	Bi	+	64° (apprx)		Ax. pl. b(010)	(G)
	$C_{12}H_{20}O$	Matico camphor	Trig	Un.					(G)
	$C_{12}H_{20}OS_2$	Methyl l-bornyl xanthate	R	Bi	−	33° 24′		Ax. pl. b(010); X‖a	(G)
	$C_{12}H_{22}ON_2$	Terpinene nitrolethylamine	M	Bi.		70° 53′	128° 32′	Ax. pl. ⊥b(010); Z∧c = 26° in obtuse ∠β	(G)
4394	$C_{12}H_{22}O_{11}.H_2O$	Lactose	M	Bi.	−		33° 35′	Ax. pl. ⊥b(010); X∧c = 10° 11′ in obtuse ∠β	(G)
4396	$C_{12}H_{22}O_{11}$	Saccharose	M	Bi	−	48° 0′	79° 7′	Ax. pl. b(010); X∧c = 67° 45′ in obtuse ∠β	(G)
4397	$C_{12}H_{22}O_{11}.2H_2O$	Trehalose	R	Bi	+	50° 16′	78° 56′	Ax. pl. b(010), Z‖b	(G)
	$C_{12}H_{12}O_6N.2H_2O$	d-Conine ditartrate	R	Bi	+		43° 33′	Ax. pl. a(100); Z‖c	(G)
	$C_{12}H_{12}O_{12}N_4.9H_2O$	Ammonium mellitate	R	Bi	−		17° (apprx)	Ax. pl. b(010) (red), X‖c	(G)
	$C_{13}H_8O_3Cl_2$	Phenyl 3, 5-dichlorosalicylate	R	Bi	−		70° 35′	Ax. pl. a(100); X‖c	(G)
4434	$C_{13}H_9N$	Acridine	R				117° (apprx)	Ax. pl. c(001); Z‖a	(G)
	$C_{13}H_{10}N_2$	Benzenyl-o-phenylenediamine	M	Bi	+		63°	Ax. pl. b(010); Z nearly ⊥c(001)	(G)
4454	$C_{13}H_{10}O_2$	p-Hydroxybenzophenone	R	Bi	−		96° 20′	Ax. pl. b(010); X‖a	(G)
	$C_{13}H_9O_2Br$	Phenyl m-bromobenzoate	R	Bi	+		41° 4′	Ax. pl. b(010), Z‖c	(G)
	$C_{13}H_{11}O_3NS$	p-Aminobenzophenone-p′-sulfonic acid	M	Bi				Ax. pl. ‖(010); Z = c	(8)
	$C_{13}H_{10}O_4Br_2$	Ethyl dibromohydroxydimethylisocoumarilate	M	Bi			80° (apprx)	Ax. pl. b(010); Z∧c = 30° in obtuse ∠β	(G)
	$C_{13}H_{12}O_4Cl_2$	Ethyl dichlorohydroxydimethylcoumarilate	M	Bi			75° (apprx)	Ax. pl. ⊥b(010); Z∧c = 30°-35° in obtuse ∠β	(G)
4500	$C_{13}H_{12}ON_2$	p-Hydroxy-p′-methylazobenzene	M.	Bi	−		52° 30′ (apprx)	Ax. pl. b(010); X∧c = 57° in obtuse ∠β	(G)
	$C_{13}H_{12}O_3N_4$	1, 3-Dimethyl-9-phenyluric acid		Bi			Large		(21)
	$C_{13}H_{12}O_3N_4$	1, 3-Dimethyl-9-phenylpseudouric acid		Bi			Large		(21)
4509	$C_{13}H_{12}O_3S$	Phenyl p-toluene sulfonate	R	Bi	−		84° 19′	Ax. pl. a(100); X‖b	(G)
	$C_{13}H_{13}O_5N$	Acetanilinopyrotartaric anhydride	M.	Bi			86° 2′	Ax. pl. ⊥b(010); Z⊥c(001)	(G)
	$C_{13}H_{14}O_4$	Ethyl hydroxydimethylisocoumarilate	R	Bi	+		65° (apprx)	Ax. pl. c(001); Z‖a	(G)
4530.1	$C_{13}H_{16}ON_2$	4-Ethylantipyrine	M.	Bi			30° (apprx)	Ax. pl. b(010); Z∧c = 40° in obtuse ∠β	(G)
4530.2	$C_{13}H_{16}ON_2$	1-Phenyl-2-propyl-3-methylpyrazolone	M.	Bi		52° 50′	79° 59′	Ax. pl. ⊥b(010); Z‖b	(G)
	$C_{13}H_{20}O_{10}$	Glycogallin	M	Bi.	−		55° (apprx.)	Ax. pl. b(010); X∧c = 16° in obtuse ∠β	(G)
	$C_{13}H_{17}ON_2I$	1-Phenyl-3-methyl-5-ethoxypyrazole-2-methiodide	M.	Bi.	−		88° (apprx.)	Ax. pl. ⊥b(010); X‖b	(G)

Index No	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
	$C_{13}H_{18}NCl$	2-Methyl-3, 3-diethyl 2, 3-dihydroindol hydrochloride	M	B₁	−	81° 51′			(G)
	$C_{13}H_{20}NI$	Methylethylallyl-p-t o l y l a m m o n i u m iodide	R	B₁			80° (apprx)	Ax pl c(001), Z∥c	(G)
	$C_{13}H_{20}O_8$	Pentaerythritol tetraacetate	Tet	U n					(18)
	$C_{13}H_{20}OS_2$	Ethyl dl-bornylxanthate	R	B₁	−	51° 16′		Ax pl b(010)	(G)
	$C_{13}H_7O_4N_2Cl_3$	Dinitrodichlorodiphenyltrichloroethane	M	B₁	−	58° (apprx)		Ax pl b(010); X ∧ c = 28° 30′ in obtuse ∠β	(G)
	$C_{14}H_8Cl_2Br_2$	1, 1-Di(bromophenyl)-2-dichloroethylene	R	B₁	+	34° 22′		Ax. pl. c(001), Z∥a	(G)
	$C_{14}H_8Cl_4$	1, 1-Di(chlorophenyl)-2-dichloroethylene	R	B₁	+	34° 26′		Ax pl b(010), Z∥a	(G)
	$C_{14}H_8Cl_3Br_2$	1, 1-Di(bromophenyl)-2-trichloroethane	R	B₁	+	62° 12′		Ax pl c(001), Z∥b	(G)
4650	$C_{14}H_{10}$	Diphenylacetylene	M	B₁		42° (red)		Ax pl ⊥b(010)	(G)
	$C_{14}H_{10}Cl_2$	1, 1-Diphenyl-2-dichloroethylene	M	B₁	−	30° 30′		Ax pl ⊥b(010)	(G)
4656 1	$C_{14}H_{10}O_2N_2$	Phthalylphenylhydrazine (orange-yellow)	M	B₁		85° (apprx)		Ax pl ⊥b(010)	(G)
4672	$C_{14}H_{10}O_2$	Benzil	Trig	U n					(G)
4681	$C_{14}H_{10}O_3$	Disalicylaldehyde	M	B₁					(G)
4688	$C_{14}H_{10}O_4$	Benzoyl peroxide	R	B₁				Ax pl a(100); Z∥b	(G)
	$C_{14}H_{11}Br_3$	Diphenyltribromoethane	M	B₁	+		110°	Ax pl b(010)	(G)
4705	$C_{14}H_{11}O_3N$	Dibenzohydroxamic acid	R	B₁	+	54° 45′ (red)		Ax pl a(100); Z∥b	(G)
4708	$C_{14}H_{12}$	Stilbene	M	B₁	+	91° 33′		Ax pl ⊥b(010), Z ∧ c = 60° in acute ∠β	(G)
	$C_{14}H_{12}N_4$	1, 5-Diphenyl-3-iminotriazoline	M	B₁				Ax pl b(010)	(G)
	$C_{14}H_{12}O$	Phenyl p-tolyl ketone	M	B₁	−	35° 15′		Ax pl ⊥b(010), X ∧ c = 36° 57′ in acute ∠β	(G)
	$C_{14}H_{12}N$	o-Iminodibenzyl	M	B₁		69° 58.5′		Ax pl ⊥b(010)	(G)
4748	$C_{14}H_{13}ON$	N-Benzoyl-o-toluidine	R	B₁	+	87° 33′		Ax pl a(100)	(G)
4749	$C_{14}H_{13}ON$	N-Benzoyl-m-toluidine	M	B₁			38° 10′	Ax pl ⊥b(010)	(G)
4750	$C_{14}H_{13}ON$	N-Benzoyl-p-toluidine	R	B₁		73° 43′		Ax pl c(001), Z∥b	(G)
4752	$C_{14}H_{13}ON$	N-Diphenylacetamide	R	B₁	+	52° 2′		Ax pl c(001), Z∥a	(G)
	$C_{14}H_{13}O_3N_2$	o-Nitrobenzyl-o-toluidine	R	B₁		49° (red)		Ax pl a(100), Z∥b	(G)
	$C_{14}H_{13}O_2N_3$	ω, ω'-Diphenylbiuret		B₁					(a.b)
	$C_{14}H_{14}ON_2$	Phenyl-o-phenetol	M	B₁	−	68°	154° (apprx)	Ax pl ⊥b(010); X ∧ c = 39° in acute ∠β	(G)
4783	$C_{14}H_{14}O_2$	Isohydrobenzoin	M	B₁	−	84° 59′		Ax pl ⊥b(010)	(G)
	$C_{14}H_{14}O_2$	1, 2-Dihydroxyphenylethane	R	B₁	+	122° 14′		Ax. pl. (100)	(*)
	$C_{14}H_{14}O_2$	o, o'-Dimethoxydiphenyl	R	B₁	−	5°		Ax pl (010); Bx₈ ⊥c(001)	(20)
	$C_{14}H_{14}O_3S_2$	Tolyl p-toluol thiosulfonate	M	B₁			19° 29′	Ax pl ⊥b(010); Z∥b	(G)
	$C_{14}H_{14}O_6S_3$	p-Toluenesulfone trisulfide	Tet	U n					(G)
4787	$C_{14}H_{14}S$	Dibenzyl sulfide	R	B₁		67° 38′		Ax pl. b(010); X∥c	(G)
	$C_{14}H_{16}NO_4Br H_2O$	Dipyridinebetaine hydrobromide	R	B₁	+	87° 30′		Ax pl c(001); Z∥b	(G)
	$C_{14}H_{16}O_4NCl H_2O$	Dipyridinebetaine hydrochloride	R	B₁	+	83° 52′		Ax pl c(001); Z∥b	(G)
	$C_{14}H_{16}ONCl$	Diphenylhydroxyethylamine h y d r o chloride	R	U n	−				(G)
	$C_{14}H_{18}O_6$	β-Methyltetramethoxycinnamic acid	M	B₁	+		102° 4′	Ax pl ⊥b(010); Z ⊥c(001)	(G)
	$C_{14}H_{18}O_2N$	Thalin tartrate.	R	B₁	+	78° 11′		Ax pl a(100)	(G)
	$C_{14}H_{20}O_2NI$	Ethyl tetrahydroquinoline-N-acetate methiodide	M	B₁		65° 70′		Ax. pl. ⊥b(010)	(G)
	$C_{15}H_{10}O_2$	Phenylcoumarin	M	B₁				Ax. pl b(010); Z ∧ c = 30° 15′ in acute ∠β	(G)
	$C_{15}H_{12}N_2$	3, 5-Diphenylpyrazole	M	B₁			43° 30′	Ax pl ⊥b(010); Z ∧ c = 44° in acute ∠β	(G)
	$C_{15}H_{13}O_3N$	syn-Benzoylbenzohydroxam m e t h y l ether	R	B₁	−	70° 10′		Ax pl. a(100); X∥c	(G)
	$C_{15}H_{12}O_3$	o-Hydroxydibenzoylmethane	M	B₁	+		75°	Ax. pl (010); Bx₈∥c-axis	(23)
4919	$C_{15}H_{13}O_3$	Methyl benzilate	M	B₁	−		74° 52′	Ax. pl. ⊥b(010)	(G)
	$C_{15}H_{13}O_3N$	Vanillyl benzoyl amide	R	B₁	−		85° (89° calc)		(24)
	$C_{15}H_{15}O_4NS H_2O$	p-Dimethylaminobenzophenone sulfonic acid	Tri	B₁			79° (apprx)	Ax pl. ∥m(110)	(G)
	$C_{15}H_{16}O_3$	2, 6, 2′, 5′-Tetrahydroxydiphenylmethyl ethyl ether	R	B₁		79° 11′		Ax pl a(100); Z∥b	(G)
4936 1	$C_{15}H_{16}O_6 H_2O(?)$	Pierotoxinin	R	B₁				Ax pl c(001)	(G)
	$C_{15}H_{16}O_2$	Hyposantonin	R	B₁			46° (apprx)	Ax pl b(010); Z∥b(?)	(G)
4943	$C_{15}H_{18}O_3$	Santonin	R	B₁	+	41° 17′ 43° 33′		Ax. pl. a(100); Z∥b	(27)
	$C_{15}H_{18}O_3$	Santonide	R	B₁	+	67° 1′ (red)		Ax. pl. a(100); Z∥c	(G)
	$C_{15}H_{18}O_3$	Parasantonide	R	B₁	−	59° 25′ (red)		Ax pl a(100); X∥c	(G)
	$C_{15}H_{18}O_6$	Triethyl trimesate	H	U n	−				(G)
	$C_{15}H_{19}O_2N_2Cl_3$	Butyl chloral antipyrine	Tri	B₁	−		110°		(G)
	$C_{15}H_{20}O_3$	Hydrosantonide	R	B₁	+	55° 10′ (red)	93° 43′ (red)	Ax. pl. a(100); Z∥c	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit
	$C_{15}H_{20}O_4$	Santonic acid	R.	Bi.		87° 40′		Ax. pl. a(100)	(G)
	$C_{15}H_{20}O_4$	Metasantonic acid	R.	Bi.	+		68° 25′ (red)	Ax. pl. a(100); Z‖c	(G)
	$C_{15}H_{20}O_4$	Parasantonic acid	R.	Bi	−	88° 13′ (red)		Ax. pl. a(100); X‖c	(G)
	$C_{15}H_{17}O_4N$	α-Isopropylglutaraminic acid	R	Bi.	+		117° 15′	Ax. pl. b(010); Z‖c	(G)
4960	$C_{15}H_{21}O_2N_3$	Physostigmine	R	Bi		77° 42′		Ax. pl. b(010); X‖c	(G)
	$C_{15}H_{20}O_4$	Hydrosantonic acid	R	Bi	+		100° (red)	Ax. pl. a(100); Z‖c	(G)
	$C_{15}H_{22}O_5$	Photosantonic acid	R.	Bi	−		107° 25′ (red)	Ax. pl. a(100); X‖c	(G)
	$C_{15}H_{23}O_3N$	Vanillyl n-heptoylamide	M.	Bi.	−		110° (107° calc.)		(24)
	$C_{15}H_{26}O(?)$	Juniperol	Tri (?)	Bi.	−	34° 46′		Ax. pl. nearly ‖b(010); X∧c = 72° in acute ∠β	(G)
	$C_{15}H_{25}O_3N$	Sesquiterpene nitrate	R.	Bi			18° 32′	Ax. pl. a(100) (red)	(G)
	$C_{15}H_{24}Cl_2$	l-Cadinene dihydrochloride	R.	Bi	+		50° (apprx)	Ax. pl. b(010); Z‖c	(37)
	$C_{15}H_{26}O$	Cypress camphor	R.	Bi	+		61° 30′	Ax. pl. b(010); Z‖a	(G)
4997	$C_{15}H_{26}O$	Cedrol	R.	Bi	+		64° 45′	Ax. pl. b(010); Z‖a	(G)
	$C_{15}H_{26}O_5$	Triacetone mannite	M	Bi	+	77° 4′	138° 13′	Ax. pl ⊥b(010); Z∧c = 26° 54′ in obtuse ∠β	(G)
5026.1	$C_{16}H_{10}O_3$	Diphenylmaleic anhydride	R.	Bi	+		Small	Ax. pl. a(100); Z‖c	(G)
	$C_{16}H_{11}O_3Br$	2, 3-Diphenyl-3-bromo-Δ'-crotono lactone	M	Bi.			55° (apprx)	Ax. pl. ⊥b(010)	(G)
	$C_{16}H_{12}O_3$	Diphenylsuccinic anhydride	R.	Bi			166° (Li) (apprx)	Ax. pl. b(010); Z‖a	(G)
	$C_{16}H_{12}N_4$	Di-p-dicyanobenzylamine	Tri.	Bi		69° 39′		Ax. pl. ‖c(001)	(G)
	$C_{16}H_{13}O_4N$	α-Benzoyl l-β-acetylbenzoylhydroxylamine	M.	Bi	+	75° 20′		Ax pl. ⊥b(010)	(G)
5066 1	$C_{16}H_{14}N_2$	1, 5-Diphenyl-3-methyl pyrazole	M	Bi		68° 22′		Ax pl b(010); Z∧c = 7° in obtuse ∠β	(G)
5067 1	$C_{16}H_{14}O$	Benzylidene-p-tolyl ketone	R.	Bi	+	36° 4′	61° 7′	Ax. pl c(001); Z‖b	(G)
	$C_{16}H_{14}Cl_3$	Di-p-tolyltrichloroethane	M.	Bi	+		85° 5′	Ax. pl. b(010); Z∧c = 4° in acute ∠β	(G)
	$C_{16}H_{15}O_3N$	Ethyl benzohydroxamic benzoate	R	Bi	+		94° 55′	Ax. pl. a(100); Z‖c	(G)
	$C_{16}H_{15}O_3N$	anti-Benzoyl benzohydroxamic ethyl ether	Tri.	Bi.	−		18° 30′ (apprx.)		(G)
	$C_{16}H_{15}O_4N$	Anisoyl p-toluohydroxamic acid	M	Bi.	+	63° 49′	113° 6′	Ax. pl b(010); Z⊥c(001)	(G)
	$C_{16}H_{15}O_4N$	p-Toluyl anisohydroxamic acid	M	Bi.	+	50° 10′	82° 52′	Ax pl b(010); Z∧c = 49° in acute ∠β	(G)
	$C_{16}H_{14}ON_2$	Phenyl styryl ketone	R. (?)	Bi				Ax. pl. b(010); Z⊥a(100)	(13)
	$C_{16}H_{16}N_2$	Acetophenone methylphenylhydrazone	M.	Bi.			Large	Ax. pl. b(010); Z⊥a(100)	(G)
5082.4	$C_{16}H_{16}O_2N_2$	Diacetylhydrazobenzene	R	Bi.	−	88° 45′		Ax. pl. b(010); X‖a	(G)
	$C_{16}H_{16}O_4N_2$	2-Phenyl-1-allylbenzimidazolium sulfate	M	Bi	+		56° 48′	Ax pl ⊥b(010); Z∧c = 33° 51′ in obtuse ∠β	(G)
	$C_{16}H_{16}O_8N_4$	2, 3-Dinitro-p-xylene + 2, 6-dinitro-p-xylene	R.	Bi	−		38° 36 5′	Ax. pl. a(100); X‖c	(G)
	$C_{16}H_{25}O_4N·4H_2O$	l-Benzoylecgonine tetrahydrate.	R.	Bi.			45° (apprx)	Ax pl. a(100); Z‖b	(G)
5131	$C_{16}H_{22}O_3NBr$	Homatropine hydrobromide	R.	Bi	−		69°–70°	Ax pl c(001); X‖b	(G)
	$C_{16}H_{23}O_3N_3$	Antipyrine isovalerianate	M.	Bi.		68° (apprx)		Ax pl c(001); Z∧c = 17° in obtuse ∠β	(G)
5135 1	$C_{16}H_{22}O_4$	Methyl santonate	R.	Bi		74° 24′ (red)	134° 12′ (red)	Ax pl. a(100); X‖c	(G)
	$C_{16}H_{22}O_4$	Methyl metasantonate	M.	Bi		90°		Ax. pl. ⊥b(010)	(G)
	$C_{16}H_{22}O_4$	Methyl parasantonate	R.	Bi	−		58° 25′ (red)	Ax pl. a(100); X‖c	(G)
	$C_{16}H_{24}O_4Br$	β-Bromoacetyltetramethylphloroglucinol	M.	Bi.	+		50° (apprx)	Ax. pl. ⊥b(010)	(G)
5142.1	$C_{16}H_{23}O_6N·H_2O$	l-Phenyl-α'-methylpiperidine d-tartrate	R	Bi.	−		55° 42′	Ax. pl. b(010); X‖c	(G)
	$C_{16}H_{26}O$	Guaiol (Champacol)	Trig	Un.					(G)
	$C_{17}H_{17}O_4N$	Ethyl anisohydroxamic benzoate	M.	Bi	+	71° 55′	66° 13′	Ax. pl ⊥b(010); Z‖b	(G)
	$C_{17}H_{17}O_4N$	syn-Anisoylbenzohydroxamic ethyl ether	M.	Bi	−		66° 13′	Ax. pl. ⊥b(010); X∧c = 55° 30′ in acute ∠β	(G)
	$C_{17}H_{17}O_4N$	anti-Benzoylanishydroxamic ethyl ether	M	Bi	−		63° 7′	Ax. pl. ⊥b(010)	(G)
5202	$C_{17}H_{19}O_3N·H_2O$	Morphine	R.	Bi	−		125° (apprx)	Ax. pl. ⊥ to elongation	(39)
	$C_{17}H_{20}NBr$	α-Benzylphenylallylmethylammonium bromide	R.	Bi		30°–40° (apprx)		Ax. pl. c(001); Z‖b	(G)
	$C_{17}H_{20}NCl$	α-Benzylphenylallylmethylammonium chloride	R.	Bi.			100° (apprx.)	Ax. pl. c(001); Z‖b	(G)
	$C_{17}H_{20}ON_2$	Oxymethylenecamphor phenylpyrazole	M.	Bi.	+	26° 40′		Ax. pl. ⊥b(010)	(G)
	$C_{17}H_{20}ON_2$	Pseudoephedrine phenylthiourea	R	Bi.	+	76° 15′		Ax. pl. c(001); Z‖b	(G)
	$C_{17}H_{20}ON_2S$	Ephedrine phenylthiourea	R.	Bi.	+	66° 25′	89° 43′	Ax. pl c(001); Z‖a	(G)
5213.1	$C_{17}H_{20}O_2$	(p-Dianisyl)dimethylmethane	R	Bi.	−	89° 54 5′			(G)
5226	$C_{17}H_{21}O_4NBr·3H_2O$	Hyoscine hydrobromide	R.	Bi	−		101° 12′	Ax pl b(010); X‖c	(G)
5228	$C_{17}H_{22}O_4NCl$	Cocaine hydrochloride		Bi.			Large (> 120°)	Ax. pl. (010)	(37)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
	C₁₇H₂₁O₃Br	Ethyl d(l)-bromosantonigate	R.	B₁	+	123° 26′	Ax pl a(100); Z‖c	(G)
	C₁₇H₂₃O₄N	Menthyl-o-nitrobenzoate	R.	B₁	−	30° 32′	47° 24′	Ax pl b(010), X‖c	(G)
	C₁₇H₂₂O₃N₂	2-Keto-6-methyl 4-(p-isopropyl phenyl)-1, 2, 3, 4-tetrahydropyrimidine-5-ethyl carboxylate.	M	B₁	+	44° (apprx.)	Ax pl b(010)	(G)
	C₁₇H₂₄ON₂	α-Dipentene nitrolbenzylamine	M	Bi	+	108° 14′	Ax pl b(010); Z∧c = 18° in acute ∠β	(G)
	C₁₇H₂₄ON₂	d(l)-Pinene nitrolbenzylamine	R	B₁	†	80° 0′	Ax pl c(001); Z‖a	(G)
	C₁₇H₂₄O₂	1, 1, 2-Trimethyl-2-phenylcyclopentane-3-ethyl carboxylate.	M	B₁	−	65° 20′	Ax pl b(010); X∧c = 50° in acute ∠β	(G)
5244	C₁₇H₂₄O₂	Menthyl benzoate	R	B₁		70° (apprx.)	Ax pl c(001); Z‖b	(G)
5244 1	C₁₇H₂₄O₄	Ethyl santoate	R	B₁	†	84° 0′ (red)	Ax pl a(100); Z‖c	(G)
	C₁₇H₂₄O₄	Ethyl parasantoate	R	B₁		35° 35′ (red)	Ax pl a(100); X‖c	(G)
	C₁₇H₂₆O₁₀	Ethyl tetraacetylquinate	R	B₁		79° 58′	Ax pl. a(100); X‖c	(G)
	C₁₈H₁₂O₁₂N₃S₃Bi 7H₂O	Bismuth m-nitrobenzene sulfonate	M.	B₁	†		Ax pl b(010); Z∧c = about 93° in obtuse ∠β	(G)
	C₁₈H₁₄O₃N₄	γ-Benzoylpyridine picrate	M	B₁		62°	Ax pl ⊥b(010), Z∧c = 65° in obtuse ∠β	(G)
	C₁₈H₁₆O₇N₄	α-Benzylpyridine picrate	M	B₁		10°	Ax pl. b(010)	(G)
	C₁₈H₁₆O₇N₄	γ-Benzylpyridine picrate	Tri	B₁		28°		(G)
	C₁₈H₁₆O₄	Diacetyl dihydroxy stilbene	M	B₁	−	81° 39′	Ax pl. ⊥b(010); X∧c = 13° in acute ∠β	(G)
5304	C₁₈H₁₆O₇	d(l)-Usnic acid	R	B₁.	†		Ax pl a(100); Z‖c	(G)
	C₁₈H₁₈O	Diethylanthrone	R	B₁.		60° (apprx.)	Ax pl c(001); Z‖a	(G)
	C₁₈H₁₆O₄	Hydrobenzoin diacetate	M	B₁		85° (apprx.)	Ax pl b(010); Z∧c = 12° in obtuse ∠β	(G)
	C₁₈H₁₆O₄	Isohydrobenzoin diacetate	R	B₁		80° 54′	Ax pl b(010); X‖c	(G)
	C₁₈H₂₀	sym-Tetramethylanthracene hydride	R	Bi	−	79° 83′	Ax pl b(010) (blue); X‖c	(G)
	C₁₈H₂₀	Tetramethyl-p-stilbene	M	B₁	+	24° (apprx.)	Ax pl b(010), Z∧c = 90° in obtuse ∠β	(G)
	C₁₈H₂₂O₂	Benzoyl-p-tert.-amyl phenol	R	B₁		58° 47′	Ax pl b(010); X‖a	(G)
5317	C₁₈H₂₁O₃N	Codeine	R	B₁.	†	125° (apprx.)		(39)
5317	C₁₈H₂₁O₃N H₂O	Codeine		B₁.	--	130° (apprx.)		(39)
5319	C₁₈H₂₁O₃N	Isocodeine	R	B₁	−		Ax. pl. b(010); X‖c	(G)
5320	C₁₈H₂₁O₃N	Pseudocodeine	M	Bi	+		Ax pl. ⊥b(010); Z∧c = 22° in acute ∠β	(G)
	C₁₈H₂₄O₄N₂	Tetraethyl-p-diaminopyromellitate	M	Bi.		85°–90°	Ax pl. b(010)	(G)
5336	C₁₈H₂₇O₃N	Capsaicin		B₁.				(21)
	C₁₈H₂₉O₃N	Hydrocapsaicin		B₁				(21)
	C₁₈H₂₉O₃N	Vanillyl n-decoylamide	R	B₁	+	23° (calc.)		(24)
5343.1	C₁₈H₃₂	Fichtelite (Retene perhydride)	M	B₁.	--		Ax pl b(010), X‖a-axis	(G)
	C₁₈H₃₂O₁₆ 2H₂O	Melezitose	R	B₁	--	85°	X = a, Y = b, Z = c	(39)
	C₁₈H₃₄O₅	Methyl pulvinate .	M	B₁			Ax. pl b(010); X‖c	(G)
	C₁₉H₁₄O₄NS	ms-Phenylacridonium hydrosulfate (green mod)	Tri	B₁.	--	42°		(G)
	C₁₉H₁₄O₄NS	ms-Phenylacridonium hydrosulfate (red mod)	M.	B₁.	+		Ax pl b(010); Z∧c = 78° 30′ in obtuse ∠β	(G)
5414	C₁₉H₁₇N₃	α-Triphenylguanidine	R	B₁	†	38° 3′	Ax. pl c(001); Z‖a	(G)
	C₁₉H₁₉N₃I	Phenyldiallylbenzimidazolium iodide	M.	B₁	+	85° 40.5′	Ax pl ⊥b(010); Z∧c = 38° 52′ in obtuse ∠β	(G)
5424	C₁₉H₁₉O₄N	Bulbocapnine	R	Bi	−		Ax. pl. a(100); X‖b	(G)
	C₁₉H₂₀N₂	Cinchene .. .	R	B₁		100° 56′	Ax. pl. c(001); Z‖b	(G)
	C₁₉H₂₀ON₂	Phenyldiallylbenzimidazolium hydroxide	M	B₁	†	60° 21′	Ax. pl. b(010); Z⊥c(001)	(G)
5428 1	C₁₉H₂₀N₂O₂	Cinchoninone	R	B₁	+	65° 20′	Ax. pl. c(001); Z‖b	(G)
	C₁₉H₂₁N₂Cl 2H₂O	Cinchonine chloride	R	B₁	†	13° (apprx.)	Ax. pl. a(100); Z‖c	(G)
544?	C₁₉H₂₂ON₂	Cinchonidine .	R	B₁.	+	100° ± 10°	Z = b	(40)
	C₁₉H₂₂ON₂ C₂H₆	Cinchonidine	R	B₁	†	Large		(40)
5442	C₁₉H₂₂ON₂	α-Cinchonine	M	B₁	--	38° ± 2°		(40)
5442	C₁₉H₂₂ON₂	α-Cinchonine	M	B₁	--	35° 52′	Ax pl ⊥b(010); X∧c = 57° in obtuse ∠β	(G)
	C₁₉H₂₂O	d-Cinnamalidene camphor	R	Bi	+	28° (apprx.)	Ax pl b(010); Z‖a	(G)
	C₁₉H₂₂ON₂Br H₂O	Cinchonine hydrobromide	R.	B₁.		150°		(G)
	C₁₉H₂₂ON₂Br.¼C₂H₆O	Cinchonine hydrobromide	R	B₁.		155°		(G)
	C₁₉H₂₂ON₂Br ¾(?)H₂O	Cinchonidine hydrobromide	R	B₁.	+	140°	Ax pl a(100); Z‖c	(G)
	C₁₉H₂₂ON₂Cl.2H₂O	Cinchonine hydrochloride	M	B₁	−	102°	Ax pl ⊥b(010); X∧c = 35° in obtuse ∠β	(G)
	C₁₉H₂₂ON₂Cl.¼C₂H₆O	Cinchonine hydrochloride	R	B₁	+	147°	Ax. pl. b(110); Z‖c	(G)
	C₁₉H₂₂ON₂I.1.5CH₆O	Cinchonine hydroiodide	R	B₁	+	147° 40′	Ax. pl. c(001); Z‖b	(39)
	C₁₉H₂₄O₂N₂.H₂O	Codethyline	R	Bi	+	About 125°		(G)
	C₁₉H₄O₄N₂S.5H₂O	Cinchonidine sulfate .	M.	Bi.	+	115° 36′	Ax pl ⊥b(010); Z∧c = 59° in obtuse ∠β	(G)

Index No	Formula	Name	System	Class	Sign	2V	2E	Orientation	Iıt
	$C_{19}H_{22}O_2N_2Se\ 5H_2O$	Cinchonidine selenate	M.	Bı	+		156° 40′	Ax. pl. ⊥b(010); Z∧c = 59° in obtuse ∠β	(G)
5477	$C_{19}H_{30}O_2$	Abietic acid	M	Bı.	−		65°	Ax pl. b(010); X∧c = 13° in acute ∠β	(G)
	$C_{19}H_{27}O_3N$	Vanillyl undecenoylamide	R.	Bı	−		Very large		(24)
	$C_{19}H_{29}O_3N$	Vanillyl n-undecoylamide	Trı	Bı	+		110° (106° calc)		(24)
	$C_{20}H_{14}$	Benzal fluorene	R	Bı.	+		13°	Ax pl. a(100); Z‖c	(G)
	$C_{20}H_{16}O_4$	2, 4-Dihydroxytriphenylacetic acid	M	Bı	−	77° 18′		Ax pl ⊥b(010); X∧c = 7° in obtuse ∠β	(G)
	$C_{20}H_{17}O_4NS$	α-Naphylamine naphthalene-α-sulfonate		Bı					(1)
	$C_{20}H_{17}O_4NS$	β-Napthylamine naphthalene-β-sulfonate		Bı					(1)
	$C_{20}H_{17}O_4NS$	α-Napthylamine naphthalene-β-sulfonate		Bı					(1)
	$C_{20}H_{17}O_4NS$	β-Napthylamine naphthalene-α-sulfonate		Bı	+		85° 5′		(1)
	$C_{20}H_{18}O_4$	Pulvinic acid ethyl alcoholate	R	Bı		114°	61° 6′	Ax. pl. a(100); Z‖b	(G)
	$C_{20}H_{18}O_5$	Atranoric acid	R	Bı.	+			Ax. pl. c(001); Z‖a	(G)
	$C_{20}H_{30}ON$	Benzoyl-β, β-diethylmethylindolenine	M	Bı	−		41° 25′	Ax. pl b(010); X∧c = 30° in acute ∠β	(G)
	$C_{20}H_{27}O_4N$	d(l)-Bulbocapnine methyl ether	Tet	Un					(G)
	$C_{20}H_{23}O_4N$	Corydin	Tet	Un					(G)
5561	$C_{20}H_{24}O_2N_2$	Quinidine	R	Bı	−		100° ± 10°		(40)
	$C_{20}H_{26}O_8N_4$	Diethyl dihydroxysuccinate γ osazone	R	Bı	+		143° 28′	Ax pl. a(100), Z‖b	(G)
	$C_{20}H_{24}O_2N_2\ C_2H_6O$	Quinidine	R	Bı	+		80° ± 5°		(40)
	$C_{20}H_{24}O_2N_2\ C_2H_4$	Quinidine	R	Bı	+		85° ± 2°		(40)
5567	$C_{20}H_{24}O_2N_2$	Quinine	R (?)	Bı					(40)
	$C_{20}H_{24}O_2N_2\ C_2H_6$	Quinine	R	Bı	+		Large		(40)
	$C_{20}H_{24}O_2N_2\ C_6H_6$	Quinine (1 nst mod)	R	Bı	−		110° ± 10°		(40)
	$C_{20}H_{24}ON_2Br\ H_2O$	Bromomethylenchonine	M	Bı			80° (apprx)	Ax. pl ⊥b(010)	(G)
	$C_{20}H_{26}O_6N_2S\ 7H_2O$	Quinine sulfate	R	Bı			19° 15′	Ax pl. a(100), X‖c	(G)
	$C_{20}H_{26}O_5N_2Se\ 7H_2O$	Quinine selenate	R	Bı	−		77° 15′	Ax pl. a(100), X‖c	(G)
	$C_{20}H_{27}O_3N_2Br$	Cinchonidine hydrobromide methyl alcoholate	R	Bı			142°		(G)
	$C_{20}H_{27}O_3N_2Br$	Cinchonine hydrobromide methyl alcoholate	R	Bı	+		40° 40′	Ax. pl. b(010); Z‖c	(G)
	$C_{20}H_{27}O_2N_2Cl$	Cinchonidine hydrochloride methyl alcoholate	R	Bı	+		140°	Ax. pl. a(100); Z‖c	(G)
	$C_{20}H_{27}O_2N_2Cl$	Cinchonine hydrochloride methyl alcoholate	R	Bı	+		157°	Ax. pl. b(010); Z‖c	(G)
	$C_{20}H_{27}O_3N_2I$	Cinchonine hydroiodide methyl alcoholate	R	Bı	+		126° 50′	Ax. pl. b(010); Z‖c	(G)
	$C_{20}H_{28}N_4$	Diethylamidine azyline	M	Bı			78° 36′	Ax pl a(100); Z‖c	(G)
5588	$C_{20}H_{30}O_2$	d-Pimaric acid	R	Bı	+		110° 22′	Ax pl a(100), Z‖c	(G)
	$C_{20}H_{30}O_2$	l-Pimaric acid	R	Bı	+ (?)	61° 45′	110° 22′	Ax pl a(100), Z‖b	(G)
	$C_{20}H_{30}O_3$	Camphorpinacone	R	Bı			126° 50′	Ax pl a(100)	(G)
	$C_{20}H_{27}O_3N_2Cl_2$	d(l)-α-Limonene nitrosochloride	M	Bı	+		99° 34′- 100° 15′	Ax pl b(010); Z∧c = 4° 50′ in acute ∠β	(G)
	$C_{20}H_{31}O_3N$	Vanillyl n-dodecoylamide	M	Bı.	+		100° (calc)		(24)
	$C_{20}H_{30}O_4N$	Methylcapsaicin	M.	Bı					(24)
	$C_{21}H_{16}O_3$	Benzil benzilate	M.	Bı	−	74° 10′	149° 46′	Ax pl. b(010); X∧c = 104° in obtuse ∠β	(G)
	$C_{21}H_{19}N_3Br$	Amarine hydrobromide	Trig	Un					(G)
	$C_{21}H_{19}N_3Cl$	Amarine hydrochloride	Trig.	Un.					(G)
	$C_{21}H_{20}$	Diphenyl-p-xylylmethane	M	Bı.	+	57° 43′			(G)
	$C_{21}H_{23}O_2N_2Br$	α-Bromostrychnine	R.	Bı.	−		58°	Ax pl a(100), X‖c	(G)
5642	$C_{21}H_{22}O_2N_2$	Strychnine	M. (?)	Bı					(37)
	$C_{21}H_{23}O_3N_3$	Tribenzylamine nitrate	R	Bı	−		45° 20′ (red)	Ax pl c(001), X‖a	(G)
5648	$C_{21}H_{22}O_5N$	Diacetylmorphine	R	Bı	−		110° (apprx)		(39)
	$C_{21}H_{24}O_7N_4$	β, β-Triethyl α methyleneindoline picrate	M	Bı	−		16° 7′		(G)
	$C_{21}H_{27}ON_2Br\ H_2O$	Cinchonine ethobromide	R	Bı		87° 50′		Ax pl. b(010), Z‖c	(G)
	$C_{21}H_{27}ON_3Cl_2$	Dichloromalene-p-tolyl-dipiperidide	M	Bı	+		44° 40′	Ax pl b(010)	(G)
	$C_{21}H_{27}O_2N_2I\ H_2O$	Cinchonidine hydroiodide ethiodide	M	Bı			90°	Ax pl ⊥b(010)	(G)
	$C_{21}H_{28}O_2N_2$	Quinidine methyl alcoholate	R	Bı	+		78°	Ax pl. a(100), Z‖c	(G)
	$C_{21}H_{29}O_2N_2I$	Cinchonine hydroiodide ethyl alcoholate	R	Bı	−		19°	Ax. pl. b(101), X‖c	(G)
	$C_{21}H_{28}O_2$	d-Bornyl methylene ether	R	Bı	+	75° 44′		Ax pl b(010), Z‖c	(G)
	$C_{21}H_{20}O_3$	p-Cresolphthalein	R	Bı	+	39°		Ax. pl. c(001); Z‖a	(G)
	$C_{21}H_{23}ON$	α, β-Dibenzoylcinnamenimide	R	Bı		82° 10′		Ax. pl b(010); Z‖a	(G)
	$C_{21}H_{19}O_4N$	Benzoyl benzohydroxamic ansate (α-mod)	M	Bı.	−		86° 30′	Ax pl. b(010)	(G)
	$C_{21}H_{19}O_4N$	Ansoyl benzohydroxamic p-toluate (β-mod)	M	Bı	+	56° 24′	100° 44′	Ax pl. b(010)	(G)
	$C_{21}H_{20}N_2$	1, 3, 4-Triphenyltetrahydropyrazine	R	Bı	+	66° 4′		Ax pl a(100); Z‖c	(G)
	$C_{21}H_{24}O_2N_4$	Bisantipyrine	M	Bı		60° 52′	98° 4′	Ax pl b(010); Z∧c = 37° in obtuse ∠β	(G)
5704	$C_{22}H_{23}O_7N$	Narcotine	R.	Bı.	−		50° (apprx.)	Ax. pl. a(100); X‖c	(G)

Index No.	Formula	Name	System	Class	Sign	2V	2E	Orientation	Lit.
	$C_{33}H_{32}O_4$	Benzyl santoate	R	Bi	+	85° 57' (red)		Ax pl a(100); Z∥c	(G)
	$C_{33}H_{36}ON_2I_2.2H_2O$	Cinchonidine ethiodide methiodide	R	Bi		73° 36'		Ax pl b(010); Z∥a	(G)
	$C_{23}H_{28}O_2N_2$	Quinidine ethyl alcoholate	R	Bi			78° 30'		(G)
	$C_{31}H_{32}O_3S_2$	Menthyl thioxanthic anhydride	R	Bi	-	85° 6'		Ax pl b(010); X∥a	(G)
	$C_{22}H_{18}ONBr$	Bromomethyltriphenyl pyrrolone	M	Bi	+	70° 15'	122° 55'	Ax pl b(010); Z apprx ⊥a(10 $\bar{1}$)	(G)
	$C_{21}H_{19}O_3N$	p-Toluyl anisohydroxamic benzoate (α-mod)	M	Bi	+	64° 32.5'	120° 48'	Ax pl ⊥b(010); Z∧a = about 60° in obtuse ∠β	(G)
	$C_{21}H_{19}O_3N$	Anisoyl benzohydroxamic p-toluate (α-mod.)	M	Bi	+	78° 39'		Ax pl ½c(001); Z∥a	(G)
	$C_{21}H_{19}O_3N$	Anisoyl p-toluhydroxamic benzoate	M	Bi		84° 55'		X⊥b	(G)
	$C_{21}H_{19}O_3N$	Benzoyl p-toluhydroxamic anisate	M	Bi		68° 32'	145°	Ax pl b(010); X∧c = 33° in obtuse ∠β	(G)
	$C_{21}H_{19}O_3N$	Benzoyl anisohydroxamic p-toluate	M	Bi	+	71° 12'		Ax pl b(010)	(G)
	$C_{21}H_{19}O_3N$	Benzoyl anisohydroxamic anisate	M	Bi			16° 42'	Ax pl ⊥b(010); Z∧c = 54° 50' in obtuse ∠β	(G)
	$C_{23}H_{24}O_2N_2 H_2O$	Methylene bisantipyrine	M	Bi		76° 30'		Ax pl b(010); Z∧c = 56° in obtuse ∠β	(G)
	$C_{23}H_{37}O_4NI H_2O$	Methyl trimethylcolchidimethanate methiodide	R	Bi		72° (apprx)		Ax pl a(100); Z∥b	(G)
5818	$C_{24}H_{18}$	1, 3, 5-Triphenylbenzene	R	Bi		0° 50'	18° 25'	Ax pl b(010); X∥c	(G)
	$C_{24}H_{21}ON$	Ethyltriphenylpyrrolone (β-mod)	M	Bi			17° 20'	Ax pl ⊥b(010); X∧c = 64° in obtuse ∠β	(G)
	$C_{25}H_{23}ON$	Propyltriphenylpyrrolone (α-mod)	R	Bi	+	65° 50'	135° 30'	Ax pl a(100); Z∥b	(G)
	$C_{24}H_{16}O_{10}$	Lepranthine	M	Bi				Ax pl b(010)	(G)
	$C_{26}H_{22}O$	Tetraphenyleneplnacoline	M	Bi		80° (apprx)		Ax pl b(010); X∧c = 50° (apprx) in obtuse ∠β	(G)
	$C_{26}H_{23}O_4N$	d-Benzoylbulbocapnine	R	Bi		78° 44'	108° 58'	Ax pl c(001), X∥b	(G)
	$C_{24}H_{26}O_4N_2$	Strychnine ethyl carbonate	?	Bi	+		30° (apprx)		(17)
	$C_{27}H_{26}O_6N_2$	Cinchonine phenylglycolate	R	Bi	+			Ax pl b(010); Z∥c	(G)
	$C_{27}H_{46}Br_2$	Cholestene dibromide (St mod.)	R	Bi	+		45°	Ax pl a(100); Z∥c	(G)
	$C_{28}H_{20}O_4$	Stilbeneglycol dibenzoate	M	Bi	+	85° 58'		Ax pl ⊥b(010); Z∥b	(G)
	$C_{23}H_{26}O_4N_2 3H_2O$	Brucine valerianate	M	Bi			86° (apprx)	Ax pl ⊥b(010)	(G)
	$C_{22}H_{36}O_2$	Gurjum resin	Tri	Bi		86° 6'			(G)
5961	$C_{28}H_{46}O_2$	Cholesteryl formate	M	Bi	+			Ax pl b(010), Z∧c = 21° 30'	(G)
	$C_{30}H_{26}O_8N_2S_2$	α-Napthylamine naphthalene-1, 5-disulfonate		Bi					(1)
	$C_{30}H_{26}O_8N_2S_2$	α-Napthylamine naphthalene-1, 6-disulfonate	M,	Bi	—		Large		(1)
	$C_{30}H_{26}O_8N_2S_2$	α-Napthylamide naphthalene-2, 6-disulfonate		Bi.	-		Large		(1)
	$C_{30}H_{26}O_8N_2S_2$	α-Napthylamine naphthalene-2, 7-disulfonate		Bi	+				(1)
	$C_{30}H_{26}O_8N_2S_2$	β-Napthylamine naphthalene-1, 5-disulfonate (normal salt)		Bi	+		75° 5' (obs) 77° 6' (calc)		(1)
	$C_{30}H_{26}O_8N_2S_2$	β-Napthylamine naphthalene-1, 5-disulfonate (acid salt)		Bi			Large		(1)
	$C_{30}H_{26}O_8N_2S_2$	β-Napthylamine naphthalene-1, 6-disulfonate		Bi.	--		Large		(1)
	$C_{30}H_{26}O_8N_2S_2$	β-Napthylamine naphthalene-2, 6-disulfonate		Bi	+		70° 5'		(1)
	$C_{30}H_{26}O_8N_2S_2$	β-Napthylamine naphthalene-2, 7-disulfonate		Bi	-		Large	Bxo ⊥plates	(1)
	$C_{30}H_{48}$	d-α-Amyrilene	R	Bi	+	72° 12'		Ax pl c(001), Z∥a	(G)
	$C_{30}H_{48}$	d-β-Amyrilene	R,	Bi	+	22° 21.5'	35° 26.5'	Ax pl c(001), Z∥b	(G)
	$C_{30}H_{50}O$	α-Isodypnopinacoline	R	Bi	+			Ax pl a(100), Z∥c	(G)
	$C_{32}H_{26}$	Tetraphenylethanebenzene	M	Bi		60° (apprx)		Ax. pl ⊥b(010)	(G)
	$C_{33}H_{32}O_2$	Dypnopinacone	M	Bi		26° (apprx)			(G)
	$C_{32}H_{37}O_{12}$	Tetrarin	Tri	Bi	--	33° (apprx)		Ax pl ⊥a(100)	(G)
6062 1	$C_{34}H_{40}O_{10}N_2S.7H_2O$	Morphine sulfate	R.	Bi	—	69° 37' (red)		Ax. pl. b(010); X∥a	(G)
6067	$C_{34}H_{47}O_{11}N$	Aconitine	R.	Bi	+	56° 10'		Ax. pl. b(010); Z∥a	(G)
6075	$C_{34}H_{50}O_2$	Cholesterol benzoate	Tet.	Un					(G)
	$C_{40}H_{52}O_2N_2Se$	Cinchonine selenate ethyl alcoholate	M.	Bi.			77° 40'		(G)
	$C_{42}H_{48}O_8N_4S 3.5H_2O$	Amarine sulfate	M	Bi.	+	60° 57'		Ax. pl. ⊥b(010); Z∧c = 80° in obtuse ∠β	(G)
	$C_{42}H_{46}O_8N_4Se 5H_2O$	Strychnine selenate	M.	Bi.	+		14°	Ax. pl. ⊥b(010); Z∧c = 34° in acute ∠β	(G)
	$C_{42}H_{46}O_8N_4S.5H_2O$	Strychnine sulfate	M.	Bi.	+		16° 30'	Ax. pl. ⊥b(010); Z∧c = 32° 43' in obtuse ∠β	(G)
	$C_{42}H_{62}O_4$	Zeorine	H	Un					(G)

LITERATURE

(For a key to the periodicals see end of volume)

(G) Groth, *Chemische Krystallographie*, 5 vols 1906-1919 **(L-B)** Landolt-Börnstein-Roth-Scheel, *Physikalisch-Chemische Tabellen*, 5th Ed., 1923

(1) Ambler, *46*, **12**: 1081, 20. **(2)** Artini, *72*, **46**: 475, 13 **(3)** Artini, *22*, **26 I**: 392; 17. **(4)** Beckenkamp, *94*, **40**: 597, 05. **(5)** Boger, *94*, **57**: 303, 22. **(6)** Bleicher, *94*, **51**: 504, 13 **(7)** Drugman, *94*, **50**: 579, 12. **(7,5)** Duffour, *6*, **30**: 169, 13. **(8)** Ehrlich and Pistschimuka, *25*, **45**: 2436, 12 **(8,5)** Gatewood, *1*, **47**: 411, 25 **(9)** Gilta, *27*, **31**: 250, 22

(10) Goldschmidt, *94*, **56**: 123, 15 **(11)** Gossner, *94*, **53**: 488, 14 **(12)** Hartley, Drugman *et al*, *4*, **108**: 751, 13 **(13)** Heilbron and Wilson, *4*, **101**: 1480, 12 **(14)** Hudson and Chernoff, *1*, **60**: 1007, 18 **(15)** Jaeger, *64P*, **18**: 49; 15 **(16)** Jamieson and Wherry, *1*, **43**: 136, 20 **(17)** Jungfleisch, *34*, **155**: 801, 12 **(18)** Keenan, *O* **(19)** Kraggs, *1*, **123**: 77, 23

(20) Mielcitner, *94*, **55**: 51; 19. **(21)** Moore and Gatewood, *1*, **45**: 144, 23 **(22)** Müller, *4*, **107**: 874; 15. **(23)** Nelson, *1*, **41**: 1115; 19. **(24)** Nelson, *1*, **41**: 2122; 19. **(25)** Nelson and Dawson, *1*, **45**: 2180; 23. **(26)** Ornstein and Pratt, *11*, **47**: 95; 12. **(27)** Robinson and Jones, *4*, **101**: 64; 12 Stefi, *94*, **54**: 343; 14. **(29)** Steinmetz, *94*, **54**: 467; 15.

(30) Steinmetz, *94*, **55**: 375; 16 **(31)** Stortenbeker, *70*, **32**: 226; 13. Thoms, *67*, **33**: 403; 12. **(34)** Wahl, *5*, **87A**: 371; 13. **(35)** Wherry, *128*, **8**: 321; 18. **(36)** Wherry, *1*, **43**: 126; 20. **(37)** Wherry, O. **(38)** Wherry and Hann, *128*, **12**: 291; 22. **(39)** Wherry and Yanovsky, *128*, **9**: 507, 19 **(40)** Wherry and Yanovsky, *1*, **40**: 1065; 18. **(41)** Widmer, *94*, **60**: 181, 21 **(42)** Hayman, Wagner and Holden, *284*, **14**: 388; 25.

Extensive data are given in Fedorow, *"Das Krystallreich. Tabellen zur Krystallochemischen Analyse,"* 1920, this is vol 36 of *Mémoires de l'Académie des Sciences de Russie* (VIII series)

X-RAY DIFFRACTION DATA FROM CRYSTALS AND LIQUIDS

R. W. G. WYCKOFF

Introduction.—To find a given substance, consult Table **A** for all elementary substances, **B** for all chemical compounds, **D** for all alloys which are not definite chemical compounds, **C** for all liquids, and **J** for solid solutions of salts

Except for the spacing observations given in Tables **C'** and **C**, there are recorded below only such observations as can be made to yield dimensions for at least a possible unit cell. The structure types of some of the simpler unit cells are shown in Figs. 1-11. The mode of designating these structures and other coordinate groups giving atomic positions is that described in Wyckoff, *"The Structure of Crystals,"* Chemical Catalog Co, New York, 1924.

ABBREVIATIONS

2a, 4b, 8f, (4b, 4c), (4b, 4d), (32b, 48c), etc. refer to the correspondingly numbered coordinate groups in Wyckoff, l c. and *Analytical Expression of the Results of the Theory of Space Groups* (Washington, 1922).

a_0, b_0, c_0	Edge length of unit cell along the *a-, b-,* and *c-*crystallographic axes, respectively.
α	The angle between the three equivalent axes of a rhombohedral unit, in a triclinic crystal, the angle between the *b-* and *c-*axes.
B.-c.	Body-centered type of structure. The cubic B-c. arrangement (*2a*) is shown in Fig. 1.
β	Angle between the *a-* and *c-*axes
C.-p.	The hexagonal close-packed type of atomic arrangement (*d*) (see Fig. 3).
γ	Angle between the *a-* and *b-*axes in a triclinic crystal
2Ci	Holohedral symmetry class, monoclinic system. 2Ci-*m* (C_{2h}^m) as under T.
3Ci	Second sort hexagonal tetartohedral symmetry class, rhombohedral division, hexagonal system 3Ci-*m* (C_{3i}^m) and 3Ci-*m* (*n*) as under T.
4C	Tetartohedral symmetry class, tetragonal system. 4C-*m* (C_4^m) as under T.
6Ci	Paramorphic hemihedral symmetry class, hexagonal division, hexagonal system. 6Ci-*m* (C_{6h}^m) as under T.
Dia.	Diamond type (8f.) of atomic arrangement (see Fig. 4).
2D	Enantiomorphic hemihedral symmetry class, orthorhombic(rhombic) system. 2D-*m* (V^m), as under T.
2Di	Holohedral symmetry class, orthorhombic system. 2Di-*m* (V_h^m) and 2Di-*m* (*n*) as under T.
3D	Enantiomorphic hemihedral symmetry class, rhombohedral division, hexagonal system. 3D-*m* (D_3^m) and 3D-*m* (*n*) as under T.

3Di	Holohedral symmetry class, rhombohedral division, hexagonal system. 3Di-*m* (D_{3d}^m) and 3Di-*m* (*n*) as under T.
4d	Second sort hemihedral symmetry class, tetragonal system. 4d-*m* (V_d^m) and 4d-*m* (*n*) as under T.
4D	Enantiomorphic hemihedral symmetry class, tetragonal system. 4D-*m* (D_4^m) as under T.
4Di	Holohedral symmetry class, tetragonal system. 4Di-*m* (D_{4h}^m) and 4Di-*m* (*n*) as under T.
6Di	Holohedral symmetry class, hexagonal division, hexagonal system. 6Di-*m* (D_{6h}^m) and 6Di-*m* (*n*) as under T.
2c	Hemimorphic hemihedral symmetry class, orthorhombic system. 2c-*m* (C_{2v}^m) as under T.
3c	Hemimorphic hemihedral symmetry class, rhombohedral division, hexagonal system. 3c-*m* (C_{3v}^m) and 3c-*m* (*n*) as under T.
6c	Hemimorphic hemihedral symmetry class, hexagonal division, hexagonal system. 6c-*m* (C_{6v}^m) and 6c-*m* (*n*) as under T.
F.-c.	Face-centered type of structure. Cubic F.-c. arrangement (4b) shown in Fig. 2.
Oi	Holohedral symmetry class, cubic system. Oi-*m* (O_h^m) and Oi-*m* (*n*) as under T.
P. S.	Possible structure. Used to designate those atomic arrangements which may be correct but for which additional results are needed or desirable.
P. U. C.	Possible unit cell. Used to designate those crystals for which the selected unit cells may be correct but which require additional experimental or theoretical treatment.
S. P.	Sample compressed.
T	Tetartohedral symmetry class, cubic system. T-*m* = n^{th} space group having this symmetry (= T^m). T-*m* (*n*) = n^{th} atomic arrangement under T-*m*. For instance T-3(c) is seen by reference to Wyckoff (*Analytical expression*, p. 122), to be arrangement 8a. Similarly 4Di-7 (c) is the coordinate pair 0½u; ½00 (*ibid.*, p. 93).
Te	Hemimorphic hemihedral (tetrahedral) symmetry class, cubic system. Te-*m* (T_d^m) and Te-*m* (*n*) as under T.
Ti	Paramorphic hemihedral (pyritohedral) symmetry class, cubic system. Ti-*m* (T_h^m) and Ti-*m* (*n*) have meanings analogous to those of similar symbols under T.
u, or v	Variable *x, y* or *z* parameter.

Fig. 1.—The unit cube of the body-centered cubic arrangement (2a). The coordinates of the atomic positions associated with this cell are 000; $\frac{1}{2}\frac{1}{2}\frac{1}{2}$.

Fig. 2.—The unit cube of the face-centered cubic arrangement (4b). The coordinates of the atomic positions associated with this cell are 000, $\frac{1}{2}\frac{1}{2}0$; $\frac{1}{2}0\frac{1}{2}$; $0\frac{1}{2}\frac{1}{2}$.

Fig. 3.— The unit cell of the hexagonal close-packed arrangement (d). The coordinates of the atomic positions associated with this cell are 000, $\frac{1}{3}$ $\frac{2}{3}$ $\frac{1}{2}$.

Fig. 4.—The unit cube of the diamond cubic arrangement (8f). The coordinates of the atomic positions associated with this cell are 000, $\frac{1}{2}\frac{1}{2}0$; $\frac{1}{2}0\frac{1}{2}$; $0\frac{1}{2}\frac{1}{2}$; $\frac{1}{4}\frac{1}{4}\frac{1}{4}$; $\frac{1}{4}\frac{3}{4}\frac{3}{4}$; $\frac{3}{4}\frac{1}{4}\frac{3}{4}$, $\frac{3}{4}\frac{3}{4}\frac{1}{4}$

Fig. 5.— The unit cube of the NaCl-arrangement (4b, 4c). The atoms in positions 4b are shown as annuli; those in 4c as black circles. The coordinates of 4c are $0\frac{1}{2}0$; $\frac{1}{2}00$; $00\frac{1}{2}$; $\frac{1}{2}\frac{1}{2}\frac{1}{2}$

Fig. 6.—The unit cube of the CsCl-arrangement (1a, 1b). Atoms of one sort, in 1a, are shown as annuli; the other kind of atom, in 1b, appears as a black circle.

Fig. 7. The unit cube of the ZnS-arrangement (4b, 4d). The atoms in position 4d appear as black circles; their coordinates are $\frac{1}{4}\frac{1}{4}\frac{1}{4}$; $\frac{1}{4}\frac{3}{4}\frac{3}{4}$, $\frac{3}{4}$ $\frac{1}{4}\frac{3}{4}$; $\frac{3}{4}$ $\frac{3}{4}\frac{1}{4}$.

Fig. 8.—The unit cell of the ZnO-arrangement (c'). The coordinates of equivalent atomic positions are 000, $\frac{2}{3}$ $\frac{1}{3}$ $\frac{1}{2}$ and 00w; $\frac{2}{3}$, $\frac{1}{3}$, $v + \frac{1}{2}$.

Fig. 9.— The unit cell of the CaF₂-arrangement (4b, 8c). The atoms in positions 8c, shown as black circles, have the coordinates $\frac{1}{4}\frac{1}{4}\frac{1}{4}$; $\frac{1}{4}\frac{3}{4}\frac{3}{4}$, $\frac{3}{4}\frac{1}{4}\frac{3}{4}$, $\frac{3}{4}\frac{3}{4}\frac{1}{4}$, $\frac{3}{4}\frac{3}{4}\frac{3}{4}$, $\frac{1}{4}\frac{1}{4}\frac{3}{4}$; $\frac{3}{4}\frac{1}{4}\frac{1}{4}$; $\frac{1}{4}\frac{3}{4}\frac{1}{4}$.

Fig. 10. The unit cube of the Cu₂O-arrangement (2a, 4d). The atoms in positions 4d are shown as annuli, those in 2a appear as black circles.

Fig. 11 —The unit cell of the hexagonal Mn(OH)₂ arrangement (h). The coordinates of the equivalent atomic positions in the unit are 000 and $\frac{1}{3}$ $\frac{2}{3}u$; $\frac{2}{3}$ $\frac{1}{3}u$.

A-Table.—Elements

Chemical symbol	Crystal system	Structure type	Space group	Unit cell Size, Å a_0	c_0	Molecules	Calculated density	Lit. and remarks
A	C.	F.-c (4b)		5 43		4	1 64$_5$	(227) (temp. ca. −253°)
Ag	C.	F.-c.(4b)		4 07$_9$		4	10.49	(82, 142, 165, 218, 235, 240, 241, 265, 329, 371)
Al	C.	F -c.(4b)		4 04$_3$		4	2.69$_2$	(84, 127, 128, 141, 197, 206, 216, 241, 329, 366, 361)
As	H.	3Di-5(c)	3Di-5	4 14$_2$; 54° 7'		2	5 7$_5$	(43, 366) u. = 0.22$_6$, probably correct
Au	C.	F.-c.(4b)		4 06$_4$		1	19 4	(82, 84, 142, 165, 218, 241, 329, 371)
Be	H.	C.-p (d)	6Di-4?	2 28$_3$	3 60$_7$	2	1 82$_8$	(163)
Bi*	H.	3Di-5(c)	3Di-5	4 72$_6$; 57° 16'		2	9.8$_6$	(82, 118, 139, 140, 142, 166, 193)
C-din.	C.	Dia.(8f)	Oi-7	3 56		8	3 51	(52, 59, 60, 128)
Graph.†	H.	6e-4(a, b)	6e-4?	2 46	6 79	4	2 22	(14, 88, 89, 105, 119, 128, 262, 310)
Ca	C.	F -c.(4b)		5 56		1	1 53$_8$	(134, 135)
Cd	H.	C.-p.(d)	6Di-4?	2 98	5 63	2	8 5$_6$	(134, 136, 229)
Ce	C.	F.-c.(4b)		5 12		4	6.9$_0$	(137)
	H.	C.-p (d)	6Di-4?	3 65	5 96	2	6 7$_3$	(137). Existence (?) (224)
Co	C.	F -c (4b)		3 55$_4$		4	8 67	(131, 136), cf. (224)
	H.	C -p (d)	6Di-4?	2 51$_4$	4 10$_5$	2	8 6$_6$	(131, 136), cf. (224)
Cr	C.	B -c (2a)		2 87$_5$		2	7 2$_2$	(131, 136, 201, 206)
Cu	C.	F.-c.(4b)		3 60$_3$		4	8 9$_5$	(46, 82, 84, 141, 145, 196, 197, 198, 199, 200, 329, 374, 371)
Fe-α	C.	B.-c (2a)		2 85$_5$		2	7.9$_2$	(82, 84, 122, 128, 131, 168, 196, 250, 253, 254, 255, 256, 362)
Fe-β	C.	B -c (2a)		2 90 at 800°		2	7 55	
Fe-γ	C.	F.-c (4b)		3 63 at 1100° / 3 68 at 1425°		4	7 70 at 1100° / 7 40 at 1425°	No structural inversion, α to β (250, 253, 254, 255, 256, 257)
Fe-δ	C.	B.-c (2a)		2 93 at 1425°		2	7.3$_3$	
Ga				Symmetry said to be not cubic				(285)
Ge	C.	Dia (8f)	Oi-7	5 62		8	5 3$_8$	(14, 138)
Hf	H.	C.-p (d)	6Di-4?	3 32	5 46	2	11 3	(324, 379)
Hg				Two different structures have been deduced				(2, 170)
In	Tet.?	?		4 58	4 86	4	7 4$_3$	(134, 136) P. U. C.
Ir	C.	F -c (4b)		3 82$_3$		4	22 8	(134, 136, 284)
K	C.	B -c.(2a)		5 20 at −150°		2	0 917 at −150°	(162). Approximate only
Li	C.	B.-c (2a)		3 50		2	0 534	(32, 33, 128)
Mg	H.	C.-p.(d)	6Di-4?	3 22	5 23	2	1.70$_9$	(36, 128, 129, 196)
Mn (α)	C.?			8 89		56?	7 2$_1$	(350) P. U. C.
Mn (β)	C.?			6 28$_9$		20?	7 2$_9$	(350) P. U. C.
Mn (γ)	Tet.?			3 77$_4$	3 53$_3$	4	7 2$_1$	(350, 368) P. U. C.
Mo	C.	B.-c.(2a)		3 14$_3$		2	10 2$_0$	(82, 136, 236, 329)
Na	C.	B.-c.(2a)		4 30		2	0 954	(128)
Nb	C.?			4 19		1		(366) P. U. C. Impure
Ni	C.	F.-c.(4b)		3 49$_9$		4	9 04	(36, 82, 84, 128, 131, 136, 168, 206, 260, 299, 329, 360, 361)
Os	H.	C.-p.(d)		2 71$_4$	4 32	2	22.8	(137)
P (black)	H.			5 96; 60° 16'		8		(392) P. S. like As
Pb	C.	F.-c.(4b)		4 92$_0$		4	11 4$_8$	(82, 84, 156, 196, 206, 241, 329, 340)
Pd	C.	F.-c.(4b)		3 85$_9$		4	12 25	(134, 136, 164, 167, 329, 393)
Pt	C.	F.-c.(4b)		3 91$_3$		4	21 5	(82, 134, 136, 142, 329, 393)
Rh	C.	F.-c.(4b)		3 82$_0$		4	12 2	(136, 393)
Ru	H.	C.-p.(d)	6Di-4?	2 68$_6$	4 27$_2$	2	12 6	(134, 136)
S	R.		2Di-24	10 61	24 56	128	2 02	(61, 314) b_0 = 12.87
Sb	H.	3Di-5(c)	3Di-5	4 50$_0$; 56° 37'		2	6 7$_3$	(140, 193) u. = 0.23$_1$
Se	H.	3D-4(a) (or 3D-6(a))	3D-4 or 3D-6	4 34	4 95	3	4 8$_6$	(42, 232, 308, 366) u. = 0 21$_6$. P. S.
Si	C.	Dia.(8f)	Oi-7	5 42		8	2 32	(88, 107, 108, 127, 128, 153, 154)
Sn (gray)	C.	Dia.(8f)	Oi-7	6 46		8	5.81	(29, 30, 31), cf. (206)
(white)	Tet.	4Di-19(a)	4Di-19?	5 82$_4$	3.16$_5$	4	7.3$_0$	(29, 30, 31, 172, 173, 174, 206, 238)
Ta	C.	B.-c.(2a)		3 27$_2$		2	17.1	(25, 134, 136)

Chemical symbol	Crystal system	Structure type	Space group	Unit cell Size Å a₀	Unit cell Size Å c₀	Molecules	Calculated density	Lit. and remarks
Te	H.	3D-4(a) or 3D-6(a)	3D-4 or 3D-6	4.44	5.90	3	6.26	(42, 232, 308, 366) u = 0.269. P. 8.
Th	C.	F.-c.(4b)		5.04		4	12.0	(36, 137)
Ti	H.	C.-p.(d)	6Di-4?	2.92	4.67	2	4.58	(36, 137, 201)
Tl	H.?	C.-p.(d)?	6Di-4(?)	3.47	5.52	2	11.7	(25, 156). Correct unit uncertain
	Tet.(?)			4.75	5.40			
U				Said to be not cubic				(25)
V	C.	B.-c.(2a)		3.04		2	5.98	(138)
W	C.	B.-c.(2a)		3.155		2	19.3	(67, 82, 84, 87, 136, 374)
Zn	H.	C.-p.(d)	6Di-4?	2.657	4.948	2	7.04	(134, 136, 206, 229, 346)
Zr	H.	C.-p.(d)	6Di-4?	3.23	5.14	2	6.47	(137, 379)

* u = 0.237. (142, 61 early editions) give incorrect structures

† u for 6e-4 (a) = 0. u for 6e-4 (b) = ¼

𝔅-Table.—Standard Arrangement p. p. 96

Chemical symbol	Crystal system	Structure type	Space group	Unit cell, size, Å a₀	Unit cell, size, Å c₀	M	Calculated density	Lit	Additional data and remarks
H₂O	H			4.52	7.32	4	0.91₅	(54, 90, 114, 210, 213)	P. U. C. Atomic arrangement not yet known with certainty.
HCl	C	F.-c ?		5.5₀, −168°C		4	1.45	(228)	
11 N₂O	C	(4f)	T-4	5.77		4	1.51	(233, 358)	u₀ = 0.22₈, distance O-N = 1.06Å P. 8.
NH₃	C	[4f, T-4(b)]	T-4	5.19(ca 80°)		4	0.81	(338)	u = 0.22
NH₄Cl (high)	C	NaCl-like		6.5₃(250°)		4	1.27	(30)	
NH₄Cl (low)	C	CsCl-like		3.86₈		1	1.52₈	(30, 120, 244, 250)	
N₂H₄Cl₂	C.	FeS₂-like (8h, 8h)	Th-6	7.89		4	1.41	(281)	u_N = ca. 0.04, u_Cl = 0.27
NH₄Br (high)	C	NaCl-like		6.90(250°)		4	1.97	(30)	
NH₄Br (low)	C	CsCl-like		4.04₇		1	2.43₈	(30, 120, 244)	
NH₄I	C.	NaCl-like		7.24₄		4	2.51₇	(20, 120, 243)	
(NH₄)₂SO₄	R.		2Di-16	5.95	7.73	4	1.80	(185)	b₀ = 10.5₈
12 PH₄I	Tet.	4D-7(a, c)	4Di-7	6.34	4.62	2	2.88	(94)	u_I = 0.40 ± 0.01
(NH₄)H₂PO₄	Tet.		4d-12	7.48	7.55	4	1.80	(342)	N atoms at 4d-12(a); P at 4d-12 (b)
As₂O₃	C.	(32b, 48c)	Oh-7	11.06		16	3.86	(41)	u_As = 0.89₈, v₀ = 0.21
Sb₂O₃	C.	(32b, 48c)	Oh-7	11.14		16	5.57	(41)	u_Sb = 0.88₆, v₀ = 0.23
16 CO₂	C	(4b, 8h)	Th-6	5.62		4	1.64	(317, 318, 358, 362)	u₀ uncertain. Liquid air-temperature

For other carbon compounds belonging here v. the ℭ-Table infra

Chemical symbol	Crystal system	Structure type	Space group	Unit cell a₀	Unit cell c₀	M	Calculated density	Lit	Additional data and remarks
SiO₂ (β-quartz)	H.	6D-4 6D-5 }(c, j)	6D-4 & 6D-5	5.01	5.47	3	2.50	(331, 332, 359)	u = 0.19₇
SiO₂ (low quartz)	H		3D-3 & 3D-5 or 3D-4 & 3D-6	4.903	5.39₈	3	2.64₈	(21, 42, 169, 227, 331)	P. U. C. a₀-spacing for quartz very accurately determined.
SiO₂ (β-cristobalite)	C.	(8f, 16b)	Oh-7 ?	7.12(290°)		8	2.20	(268, 377, 380)	
(NH₄)₂SiF₆	C	(4b, 8e, 24a)	Oh-5	8.38		4	2.00	(38)	u_F = 0.205
SiC, I	H.			3.09₈	37.9	15	3.15	(282)	Complex structure assigned
SiC, II	H.		6C-6 ?	3.09₈	15.17	6	3.15	(347, 348)	C at 6C-6(a) if u = 0 and 6C-6(b), if u = ¼ and ¾. Si at 6C-6(a) if u' = ¼ and 6C-6(b) if u' = 0.29 and 0.95 P. 8.
SiC, III	H.			3.09₈	10.10	4	3.16	(390)	C at 000; 00½; ½ ½ ⅓; ½ ½ ⅚. Si at 00u; 0, 0, u + ½; ½, ½, u + ⅓, ½, ½, u + ⅚, u = ca. ¹/₁₆. P. 8.
TiO₂ (rutile)	Tet.	4Di-14(a, f)	4Di-14	4.58	2.98	2	4.21	(83, 113, 241, 263)	
TiO₂ (anatase)	Tet.			5.27	9.37	8	4.05	(242)	P. U. C.
Ti₂O₃	H.	3Di-6(c, e)	3Di-6	5.37, 56° 48'		2	4.67	(351)	
TiN	C.	NaCl(4b, 4c)		4.23†		4	5.4₀?	(13, 306)	The later determination gives a₀ = 4.40
TiC	C.	NaCl(4b, 4c)		4.29?		4	5.01?	(13, 306)	The later determination gives a₀ = 4.60
21 ZrO₂	C.	CaF₂(4b, 8e)	Oh-5	5.08		4	6.1₉	(13)	P. 8. Other data (88) conflict. 2 modifications?
ZrB₂	H.	Mn(OH)₂(A)	3Di-3	3.68	5.85	1	3.73	(13)	P. 8. u = ca. 0.25
ZrBe₂	H.	Mn(OH)₂(h)	3Di-3	3.79	6.18	1	5.3₈	(13)	P. 8. u = ca. 0.25
ZrN	C	NaCl(4b, 4c)		4.61		4	7.1	(13, 306)	P. 8
(NH₄)₃ZrF₇	C.	(4d, 4e, 12a, 24u)	Oh-4	9.35		4	2.2₈	(13)	0.15 < u_N < 0.21; 0.42 < u_F < 0.48, 0.23 < v_F < 0.28
ZrC	C.	NaCl(4b, 4c)		4.7₈		4	6.4	(13, 306)	P. 8.
ZrSiO₄	Tet.			9.20	5.87	8	4.85	(341)	P. U. C.

Chemical symbol	Crystal system	Structure type	Space group	Unit cell, size, Å a₀	c₀	M	Calculated density	Lit.	Additional data and remarks
SnO	Tet.	4Dh·7(a, c)?		3 77	4 77	2	6 5₈	(300)	
SnO₂	Tet			4 72	3 16	2	7 07	(83, 341, 363)	P. U. C.
SnI₄	C.	Ti-6(c, d)	Ti-6	12 2₈		8	4 52	(96, 175)	$u_{Sn} = 0.129$, $u_I = 0.25₈$, $x =$
(NH₄)₂SnCl₆	C.	(4b, 8e, 24a)	Oₕ-5	10 0₈		4	2 39	(92)	0 009, y = 0.001, z = 0.25₈
23 PbO	Tet	4Dh-7(a, c)		3 99	5 01	2	9 2₈	(97, 300)	$u_{Cl} = 0.24₈$ and < 0.25
PbO₂	Tet.	4Dh-14(a, f)	4Dh-14	4 97	3 40	2	9 40	(345, 386)	$u_{Pb}[4Dh-7(c)] = 0.24$
PbF₂(β)	C.	CaF₂(4b, 8e)	Oₕ-5	5 93		4	7 76	(340)	
PbS	C	NaCl(4b, 4c)		5 97		4	7 4₂	(61, 76, 154, 340, 357)	
PbSe	C.	NaCl(4b, 4c)		6 14		4	8 17	(357, 366)	
PbTe	C	NaCl(4b, 4c)		6 34		4	8 67	(357)	
Pb(NO₃)₂	C.	(4b, 8A, Ti-6(24))	Ti-6	7 84		4	4 54	(191, 245)	
ThO₂	C.	CaF₂(4b, 8e)	Oₕ-5	5 59		4	9 9₈	(13, 83, 111)	Another determination of a_1
Ga₂O₃	H.	3Dh-6(c, e)	3Dₕ-6	5 28₁, 55° 35'		2	6 6₂	(381)	(288) varies widely from this.
In₂O₃	C		Oₕ-10	10 12		16	7 07	(381)	
(Ga, In)₂O₃	C.		Oₕ-10	9 76		16		(381)	39 mol. % In₂O₃
Tl₂O₃	C		Oₕ-10	10 57		16	10 2	(381)	
TlCl	C	CsCl(1a, 1b)	Oₕ-1	3 84		1	6 9₈	(85, 239, 369)	
TlBr	C	CsCl(1a, 1b)	Oₕ-1	3 97		1	7 44	(239, 369)	
ZnO	H	ZnO(e')	6e-4	3 25	5 23	2	5 6₁	(4, 7, 51, 61, 121, 249)	
Zn(BrO₃)₂·6H₂O	C	(4b, 8A, Ti-6(24))	Ti-6	10 3₁		4	2 59	(278)	
α-ZnS (wurtzite)	H.	ZnO(e')	6e-4	3 84	6 28	2	4 01	(9, 51, 381)	$u_s = ca.⅜$
β-ZnS (blende)	C.	ZnS(4b, 4d)	Te-2	5 43		4	4 02	(47, 103, 108, 154)	
ZnSe	C	ZnS(4b, 4d)	Te 2	5 65		4	5 29	(80)	
ZnCO₃	H.	3Dh-6(a, b, e)	3Dₕ-6	5 82, 48° 23'		2	4 54	(160)	
29 CdO	C	NaCl(4b, 4c)		4 72		4	8 0₈	(86, 217)	
CdF₂	C	CaF₂(4b, 8e)	Oₕ-5	5 40		4	6 30	(340)	
CdI₂	H.	Mn(OH)₂(A)	3Dₕ-3	4 ₂4	6 84	1	5 6₇	(29)	0 23 < u_I < 0.253
α-CdS	H	ZnO(e')	6e-4	4 14	6 72		4 78	(51, 381)	$u_s = ca ⅜$
β-CdS	C	ZnS(4b, 4d)	Te-2	5 82		4	4 84	(381)	
HgₐCl₂	Tet	4Dh-17(e)		4 47	10 89	2	7 16	(344)	$u_{Hg} = ⅙$, $u_{Cl} = ⅜$ P. S.
HgₐBr₂	Tet	4Dh-17(e)		4 65	11 10	2	7 71	(344)	$u_{Hg} = ⅙$, $u_{Br} = ⅜$ P. S.
HgI₂	Tet			4.35₆	12.34	2	6.4₀	(197)	
HgₐI₂	Tet.	4Dh-17(e)		4 92	11 61	2	7 6₈	(344)	$u_{Hg} = ⅙$, $u_I = ⅜$ P. S.
HgS (metacinnabarite)	C.	ZnS(4b, 4d)	Te-2	5 84		4	7 7₁	(150, 151, 154, 336, 337, 365, 366)	
HgS (cinnabar)	H		3D-4 & 3D-6	4 16	9 54	3	8 1₂	(150, 357, 365, 366)	P S suggested
CuO	Tri			3 74	4 67	4	6 4₈	(188)	P S This suggested structure resembles NaCl. $b_0 = r_0$ α = 85° 21', β = 86° 25'; γ = 93° 35'
Cu₂O	C	Cu₂O(2a, 4d)	Oₕ-4	4 2₈		2	6 0₂	(61, 113, 188)	
CuCl	C	ZnS(4b, 4d)	Te-2	5 4₀		4	4 1₈	(76, 293)	
CuBr	C	ZnS(4b, 4d)	Te-2	5 7₈		4	4 9₈	(76, 293)	
CuI	C	ZnS(4b, 4d)	Te-2	6 07		4	5 6₂	(8, 76, 293)	
CuₐSe	C.	CaF₂(4b, 8e)	Oₕ-5	5 75		4	7 1₆	(80)	
Cu₃Zn₈	C.			4 01				(34) cf (197)	Correctness in doubt
32 Ag₂O	C.	Cu₂O(2a, 4d)	Oₕ-4	4 72		2	7 27	(76, 88, 161, 277)	
AgCl	C.	NaCl(4b, 4c)		5 54		4	5 56	(76, 264, 265)	
AgBr	C.	NaCl(4b, 4c)		5 77		4	6 45	(76, 264, 265)	
AgI	H.	ZnO(e')	6e-4	4 59	7 5₀	2	5 66	(9, 8, 265)	
AgI	C.	ZnS (4b, 4d)	Te-2	6 49		4	5 67	(76, 264, 265)	
Ag₃PO₄	C.	(2a, 6f, 8a)	Te-4	6 00		2	6 37	(257)	
Ag₃AsO₄	C	(2a, 6f, 8a)	Te-4	6 12		2	6 66	(257)	
(4AgI:CuI) miersite	C.	ZnS(4b, 4d)	Te-2	6 3₈		4		(8)	A solid solution of AgI and CuI. Exact composition unknown
(NH₄)₂PtCl₆	C	(4b, 8e, 24a)	Oₕ-5	9 8₄		4	3 08	(292)	0 22 < u_{Cl} < 0.24
PtAs₂ (sperrylite)	C.	FeS₂(4b, 8A)	Ti-6	5 94	*	4		(357)	Composition unknown
(NH₄)₂PdCl₆	Tet	4Dh-1(a, e, j)	4Dh-1	7 21	4 26	1	2 12	(95)	$u_{Cl} = 0.23$
MnO	C.	NaCl(4b, 4c)		4 40		4	5 5₀	(187)	
MnO₂	Tet.			4 44	2 89	2	5 04	(214)	Pyrolusite gives the same pattern as polianite
Mn(OH)₂	H	Mn(OH)₂(A)	3Dₕ-3	3 34	4 6₈	1		(2)	Dimensions of this unit calculated from the density $\rho = 3.26$. $u_o = ca.0 22$
MnS	C.	NaCl(4b, 4c)		5 21		4	4 06	(272)	$u_o = 0 40₀$. Size of unit cell calculated from the best available density, [ρ = 3.38(162)]
MnS₂	C.	FeS₂(4b, 8A)	Ti-6	6 1₈		4		(104, 106)	
MnCO₃	H.	3Dh-6(a, b, e)	3Dₕ-6	5 84, 47° 45'		2	3 79	(47, 270)	C atoms at (a); $u_o = 0.27$
43 FeO	C	NaCl(4b, 4c)		4 29₄		4	5 9₉	(322)	
Fe₂O₃	H.	3Dh-6(c, e)	3Dₕ-6	5 42; 55° 17'		2	5 2₈	(61, 81, 191, 205, 351)	$u_{Fe} = 0.105 \pm 0.001$; $u_o = 0.292 \pm 0.007$ $u_o = ca.0.37$
Fe₃O₄	C	(8f, 16c, 32b)	Oₕ-7	8 37		8	5 21	(50, 121, 189, 394)	
FeS (troilite)	H.	6e-4(a, b)		3 43	5 79	2	4 90	(356, 391)	If $u_{Fe} = 0$, $u_s = ca.¼$. If $u = ¼$ exactly, the space group is 6Dₕ-4

Chemical symbol	Crystal system	Structure type	Space group	Unit cell, size, Å a_0	Unit cell, size, Å c_0	M	Calculated density	Lit.	Additional data and remarks
FeS₂ (pyrite)	C.	FeS₂(4b, 8h)	Tₕ-6	5 38		4	5 0₂	(47, 104, 106, 287)	$u_s = 0.386$
FeS + S₂	H.	6e-4(a, b)		3 43	5 68	2		(356, 291)	Artificial and natural pyrrhotites containing excess sulfur
FeSe	H.	6e-4(a, b)		3 61	5 87	2		(356)	39.4% Fe (weight)
FeSe + Se₂	H.	6e-4(a, b)		3 51	5 55	2		(356)	35 0% Fe (weight)
Fe(S, Se)	H.	6e-4(a, b)		3 54	5 91	2		(356)	49.8% (weight) Fe, 12 0% S, 38.2% Se
(NH₄)₂FeF₆	C.	(4b, 4c, 8e, 24a)	Oₕ-5	9 10		4	1 96	(203)	N atoms at (4c) and (8e). 0.187 < u_F < 0.217, best around 0.21
NH₄Fe(SO₄)₂.12H₂O	C	(4b, 4c, 8A, 8h, Tₕ-6 (24))	Tₕ-6	12 14		4	1 81	(246)	
FeₓC	R.			4 52	6 74	4	7 6₇	(5, 6, 7, 254, 261)	Cementite and cohenite are identical in structure. Atomic arrangement unknown. $b_0 = 5.07$
FeCO₃	H.	3D₁-6(a, b, e)	3D₁-6	5 82, 47° 45′		2	3 86	(47, 270)	C atoms at (a); $u_0 = 0.27$ probably
FeS₄	C.			4 48		4	6 1₈	(207)	Probably tetartohedral; atomic arrangement unknown
FeS₈	Tet.			2 69	5 0₈	1	5 0₂	(207)	P. U. C., structure unknown
FeCuS₂	Tet	4d-5(r, a, g)?	4d-5 ?	5 2₃	5 15	2		(66, 118)	Fe atoms at (e). $u_0 = ca. 0.21$. Probably correct structure.
CoO	C.	NaCl(4b, 4c)		4 2₄		4	6 4₉	(351)	
CoS	H.	6e-4(a, b)		3 37	5 11	2	5 9₄	(356)	
CoAsS	C.	FeS₂-like(4f)	T-4	5 65		4	6 07	(183, 387)	Reflection microscopic results (181) suggest that this structure may not be correct
(Fe, Co)S (synthetic)	H.	6e-4(a, b)		3 36	5 29	2		(356)	Composition = ca. 50 atomic % FeS
45 NiO	C.	NaCl(4b, 4c)		4 17₃		4	6 7₈	(74, 86, 299, 351, 353, 360)	
NiS (synthetic)	H.	6e-4(a, b)		3 42	5 30	2	5 5₈	(356)	$u_S = ca. ½$ taking $u_{Ni} = 0$
NiS (millerite)	H.	3e-5(b, b)	3e-5	5 64, 116° 36′		3		(356)	Possible atomic positions are suggested
NiS₂	C ?			4 0₈		1		(356)	P. U. C.
NiSe	H	6e-4(a, b)		3 66	5 3₃	2		(356)	
Ni(NO₃)₂·6NH₃	C.	(4b, 8A, Tₕ-6(24))	Tₕ-6	10 9₄		4	1 43	(273)	u_N in (8A) = $ca. ½$, y_N and z_N = ca. 0, x_0 and y_0 = ca. ½.
NiCl₂·6NH₃	C.	(4b, 8e, 24a)	Oₕ-5	10 0₉		4	1 49	(274)	$u_N = 0.24$
NiBr₂·6NH₃	C.	(4b, 8e, 24a)	Oₕ-5	10 4₈		4	1 84	(274)	
NiI₂·6NH₃	C.	(4b, 8e, 24a)	Oₕ-5	11 0₁		4	2 05	(274)	$u_N = 0.24$
NiAs	H.	6e-4(a, b)		3 61	5 03	2		(5, 356, 291)	Niccolite from Eisleben.
NiAsS (gersdorffite)	C.	FeS₂-like(4f)	T-4	5 68		4		(357, 366)	
NiSb	H.	6e-4(a, b)		3 9₂	5 11	2	8 7₂	(356, 291)	For the mineral breithauptite from Andreasberg $a_0 = 3 90$, $c_0 = 5.09$
NiSbS (ullmanite)	C.	FeS₂-like(4f)	T-4	5 91		4		(387)	Composition unknown
(Ni, Fe)S (synthetic)	H.	6e-4(a, b)		3 40₈	5 54₀	2		(356)	S = 37.8%, Fe = 33.9%, Ni = 28.3% (weight)
(Ni, Fe)S (synthetic)	H.	6e-4(a, b)		3 40₈	5 43₄	2		(356)	S = 38.4%, Fe = 28 7%, Ni = 32.8% (weight)
(Ni, Fe)S (pentlandite)	C.		Oₕ-5 ?	10 0₀		32		(356)	(8J, 24a, 32a) with u_{Fe} (24a) = ca. ½ and $u_S = ca. ½$ gives fair agreement. Various compositions
Cr₂O₃	H.	3D₁-6(c, e)	3D₁-6	5 3₈; 54° 58′		2	5 2₈	(351)	$u_0 = 0.62₁$
MoS₂	H	6D₁-4(c, f)	6D₁-4	3 15	12 30	2	5 00	(99, 211)	
(NH₄)₂MoO₂F₄	C.	(4b, 4c, 8e, 24a)	Oₕ-5 ?	9 10		4	2 23	(203)	N atoms at (4c) and (8e). F + O at (24a). 0.194 < $u_{F,0}$ < 0.220
PbMoO₄	Tet.			3 85	6 02	1		(91)	P. U. C.
Ag₂MoO₄	C.	(8f, 16c, 32b)	Oₕ-7	9 26		8	6 2₈	(276)	0.34 < u_0 < 0.40
49 UO₂	C.	CaF₂(4b, 8e)	Oₕ-5	5 47		4	10 8₉	(13, 111)	
UO₂(NO₃)₂·6H₂O	R.		2D₁-17	13 15	11 42	4	2 75	(68, 304)	U atoms probably at 2D₁-17 (c) with $u = 0.13$. $b_0 = 8.02$
V₂O₃	H.	3D₁-6(c, e)	3D₁-6	5 43, 53° 53′		2	5 0₉	(351)	
VN	C.	NaCl(4b, 4c)		4 28		4	5 47	(306)	
VC	C.	NaCl(4b, 4c)		4 30		4	5 2₈	(306)	
CbN	C.	NaCl(4b, 4c)		4 41		4	8 2₈	(306)	
CbC	C.	NaCl(4b, 4c)		4 40		4	8 1₄	(306)	
TaN	H.	ZnO(a')	6e-4	3 05	4 94	2	16 2	(12)	P. S. Cf. (307) which gives conflicting results
TaC	C.	NaCl(4b, 4c)		4 5₈		4	13 7	(12, 306)	
B₂H₆	H.			4 54	8 69	2	0 58₉	(349)	B atoms probably at 6D₁-4 (f) with $u = ca. 0.10$. Temperature not stated
55 Al₂O₃	H.	3D₁-6(c, e)	3D₁-6	5 12, 55° 17′		2	3 96	(61, 81, 181, 205, 351)	The α-form. $u_{Al} = 0 105 ± 0.001$; $u_0 = 0.303 ± 0 003$

Chemical symbol	Crystal system	Structure type	Space group	Unit cell, size, Å		M	Calculated density	Lit.	Additional data and remarks
				a_0	c_0				
AlN	H.	ZnO(e')	6e-4	3 11	4.98	2	3 24	(195)	$u = 0.38 \pm 0.01$
(NH₄)₃AlF₆	C.	(4b, 4c, 8c, 24a)	O₅-5	8 40		4	2 17	(203)	N atoms at (4c) and (8e). 0.154
									$< u_y < 0.200$
NH₄Al(SO₄)₂.12H₂O	C.	(4b, 4c, 8A, 8A, Ti-6 (24))	Ti-6	12 0₀		4	1 76	(245, 252)	
AlSb	C.	ZnS(4b, 4d)	Te-2	6 13		4	4 26	(293)	
Al₂F₂(SiO₄) topas	R		2D₁-16	4 64	8 37	4		(185)	Topas from San Luis Potosi, Mexico; $b_0 = 8.78$
CuAl	H	F.-c.?		3 89, 94° 36′		4		(141, 197, 258)	This structure may be incorrect
Cu₃Al	C.	F.-c.		3 47		4		(24) cf. (141)	Probably incorrect
CuAl₂	Tet.	B.-c.		6 05	4 88	4	4 35	(141, 197, 258)	Atomic arrangement unknown
(Fe″, Mn″)₃ Al₂(SiO₄)₃ (garnet)	C.		O₁-10	11 4₀		8		(190)	67 atomic % of ferrous iron
NiAl	C.	CsCl(1a, 1b)?		2 82		1	6.2₅	(24)	More work needed
56 Sc₂O₃	C.		O₁-10	9 79		16	3 89	(351)	
Sc N	C.	NaCl(4b, 4c)		4 44		4	4 4₆	(306)	
(Sc, In)₂O₃	C.		O₁-10	9 90		16		(351)	66.8 mol. % Sc₂O₃
(Al, Sc)₂O₃	C.		O₁-10	9 22		16		(351)	Composition unknown
Y₂O₃	C.		O₁-10	10 56		16	5 07	(351)	
Y₂PO₄	Tet.			9 60	5 94	8	4 44	(242)	P. U. C.
(Yt, Tl)₂O₃	C.		O₁-10	10 53		16		(351)	50 weight % Y₂O₃
(Yt, Bi)₂O₃	C.		O₁-10	10.72		16		(351)	37.4 mol % Bi₂O₃
La₂O₃	H.			3 94₅	6 15₁	1	6 4₈	(351)	
CeO₂	C.	CaF₂(4b, 8e)	O₁-5	5 41		4	7 1₈	(83, 111)	
Ce₂O₃	H.			3.88₀	6 05₇	1	6.8₆	(351)	
60 Pr₂O₃	H.			3 851	5 99₆	1	7.0₇	(351)	
Pr₆O₁₁	C.			10 98		?		(352)	P. U. C.
Nd₂O₃	H			3 84₁	6 00₉	1	7.2₈	(351)	
Sm₂O₃	C.		O₁ 10	10 85		16	7 2₁	(351)	
Eu₂O₃	C.		O₁-10	10 84		16	7 2₉	(351)	
Gd₂O₃	C.		O₁-10	10 79		16	7 6₂	(351)	
66 Tb₂O₃	C		O₁-10	10 70		16	7 9₀	(351)	
Tb₄O₇?	C			10 55		?		(352)	P.U.C. "Brown terbium oxide"
Dy₂O₃	C.		O₁-10	10 63		16	8 2₀	(351)	
Ho₂O₃	C.		O₁-10	10 58		16	8.3₅	(351)	
Er₂O₃	C.		O₁-10	10 54		16	8 6₄	(351)	
Tu₂O₃	C		O₁-10	10 52		16	8 7₇	(351)	
Yb₂O₃	C.		O₁-10	10 39		16	9 3₀	(351)	
Lu₂O₃	C.		O₁-10	10 37		16	9 4₃	(351)	
(NH₄)₃HfF₇	C	(4d, 4e, 12a, 24u)	O₁-4	9 40		4		(117)	Contains 15% (NH₄)₃ZrF₇
75 BeO	H	ZnO(e')	6e-4	2 70	4 39	2	2 98	(109, 163, 333, 364)	u_0 ca. ¼.
BeO(C₂H₃O₂)₆	C.			15 7₂		8	1 38	(56, 62)	A possible atomic arrangement suggested
BeO(C₃H₅O₂)₆	M.			16 0₀	9 1₅	2	1 26	(62)	P. U. C. $b_0 = 9.7_4, \beta = 116° 7′$
MgO	C.	NaCl(4b, 4c)		4 20₅		4	3 59	(86, 107, 109, 110, 121, 132, 322, 271, 287)	
Mg(OH)₂	H.	Mn(OH)₂(h)	3D₁-3	3 11	4 73	1	2.43	(3, 5, 159)	
MgF₂	Tet.	4D₁-14(a, f)	4D₁-14	4 66	3 08	2	3.11	(326, 345, 367)	
MgS	C	NaCl(4b, 4c)		5 08		4	2 84	(125)	$u_y = 0.30$
MgCO₃	H.	3D₁-6(a, b, e)	3D₁-6	5 61, 48° 12′		2	3.10	(160)	
Mg₂Si	C.	CaF₂(4b, 8e)	O₁-5	6 39		4	1 94	(298)	
Mg₂Sn	C.	CaF₂(4b, 8e)	O₁-5	6 78		4	3 54	(202, 370)	
Mg₂Pb	C.			6.75		4	5 47	(370)	Structure probably CaF₂(4b, 8e)
(Mg, Fe″)₂SiO₄ olivine	R.		2D₁-5	4 77	6 00	4		(28, 212)	14 atomic % of ferrous iron. $b_0 = 10.28$
Al₂Mg₃	C.			4 80			2 62	(24)	More work needed
MgAl₂O₄	C	(8f, 16c, 32b)	O₁ 7	8 07		8		(50, 189)	$u_0 = 0.37$. Value of a_0 calculated from the best available density ($\rho = 3.57$)
77 CaO	C.	NaCl(4b, 4c)		4 79		4	3 37	(79, 86, 107, 109)	
Ca(OH)₂	H.	Mn(OH)₂(h)	3D₁-3	3 52	4 93	1	2 31	(158)	
CaF₂	C.	CaF₂(4b, 8e)	O₁-5	5 46		4	3 17	(47, 76, 107, 108)	
CaS	C.	NaCl(4b, 4c)		5 68		4	2 60	(79, 125)	
CaSO₄	R.		2D₁-17	6 21	6 96	4		(326)	Anhydrite, not analysed. $b_0 = 6.95$
CaS₂O₃.6H₂O	Tri.							(15)	Some unreduced measurements have been recorded for this salt
CaSe	C	NaCl(4b, 4c)		5 91		4	3 81	(79)	
Ca(NO₃)₂	C.	(4b, 8A, Ti-6(24))	Ti-6	7 60		4	2 47	(245)	
Ca(F, Cl)Ca₄(PO₄)₃ apatite	H.		6C₁ 2	9 4₁	6 88	2		(122)	Composition unknown
CaCO₃ (calcite)	H.	3D₁-6(a, b, e)	3D₁-6	6 36; 46° 6′		2		(47, 49, 179, 221, 270)	C atoms at (a). $u_0 = 0.25$. A wave length standard
CaCO₃ (aragonite)	R.	2D₁-16(c, c, c, d)?	2D₁-16	4 94	5 72	4	2 94	(58, 286)	A possible atomic arrangement has been suggested. $b_0 = 7.94$
Ca(HCOO)₂	R.		2D₁-5 ?	10 16	6 20	8	2 03	(322)	P. U. C.
CaTiO₃	C.?			7.68		8		(343)	P. U. C. (?) More work necessary
CaWO₄	Tet.			3.64	5.54	1		(91)	P. U C.

Chemical symbol	Crystal system	Structure	Space group	Unit cell, size, Å a₀	c₀	M	Calculated density	Lit	Additional data and remarks
CaMg(CO₃)₂ (dolomite)	H.	3C₁-2(a, b, c, f)	3C₁-2	6 02, 47° 7'		1	2 84	(61, 249, 313)	
CaMg(SiO₃)₂ (diopside)	M.		2C₁-6	9 71	5 24	4	3 28	(291)	b₀ = 8.89, β = 74° 10'
Ca(Mg, Fe)(CO₃)₂	H.	3C₁-2(a, b, c, f)	3C₁-2	6 02, 47° 7'		1	3 15	(249)	30 atomic % of ferrous iron
78 SrO	C.	NaCl(4b, 4c)		5 10		4	5 1s	(107, 109)	
SrF₂	C.	CaF₂(4b, 8e)	O₁-5	5 86		4	4 12	(12)	
SrCl₂	C	CaF₂(4b, 8e)	O₁-5	7 00		4	3 05	(341)	
SrS	C.	NaCl(4b, 4c)		5 87		4	3 90	(128)	
SrSe	C.	NaCl(4b, 4c)		6 23		4	4 55	(230, 231, 306)	
Sr(NO₃)₂	C.	(4b, 8A, T₁-6(24))	T₁-6	7 81		4	2 93	(191, 245)	
BaO	C.	NaCl(4b, 4c)		5 50		4	6 0s	(107, 109)	
BaF₂	C	CaF₂(4b, 8e)	O₁-5	6 20		4	4 86	(76)	
BaS	C	NaCl(4b, 4c)		6 35		4	4 37	(128)	
BaSO₄	R.		2D₁-16	8 89s	7 17o	4	4 43₂	(1, 290, 326, 327, 334, 335)	b₀ = 5 44s
BaSe	C.	NaCl(4b, 4c)		6 62		4	4 93	(231, 306)	
Ba(NO₃)₂	C.	(4b, 8A, T₁-6(24))	T₁-6	8 11		4	3 23	(191, 245)	Approx atomic positions are said to be u_N, x₀ and y₀ = ca. ¼, z₀ = ca. 0
81 Li₂O	C.	CaF₂(4b, 8e)	O₁-5	4 61		4	2 01	(35)	
LiH	C.	NaCl(4b, 4c)		4 10		4	0 76	(34)	
LiF	C	NaCl(4b, 4c)		4 01		4	2 65	(76, 66, 132, 367)	
LiCl	C	NaCl(4b, 4c)		5 14		4	2 06	(75, 194, 219)	
LiBr	C.	NaCl(4b, 4c)		5 49		4	3 46	(76, 194, 219)	
LiI	C	NaCl(4b, 4c)		6 00		4	4 0₉	(76, 194, 219, 294)	
Li₂S	C	CaF₂(4b, 8e)	O₁-5	5 70		4	1 64	(239)	
Li₂CsO₄	R?			6 58	6 61	4	2 15	(25)	b₀ = 7 74. P U C.
LiCHO₂	M?			7 61	4 87	4	1 53	(25)	b₀ = 6.03, β = 95° 42'. P.U.C., S. P.
LiC₂H₃O₂	R ?			12 8o	7 43	12	1 17	(25)	b₀ = 11 6s P. U. C., S. P.
LiC₃H₅O₂	R ?			16 9s	9 45	16	1 0s	(25)	b₀ = 12 15. P. U. C., S. P.
LiC₃H₅O₂ crotonate	H ?			24 8	10 7	48	1 27	(25)	P. U. C., S P.
LiC₄H₇O₂ butyrate	H ?			27 7	10 1	48	1 07	(25)	P. U. C., S. P.
LiC₄H₇O₂ isobutyrate	Tet ?			19 7s	9 2s	24	1 01	(25)	P. U. C., S. P.
LiC₅H₉O₂ valerate	Tet ?			24 5	9 4	32	1 01	(25)	P. U. C., S. P.
LiC₅H₉O₂ isovalerate	R ?			11 7o	6 93	4	1 00	(25)	b₀ = 8 70. P. U. C., S. P.
LiC₅H₉O₂ trimethylacetate	C ?			18 5s		36	1 00	(25)	P. U. C., S. P.
LiC₇H₁₃O₂ heptylate	Tet ?			27 4	9 3	32	1 02	(25)	P. U. C., S. P.
LiC₈H₁₅O₂ caprylate	H ?			42 1	10 9	72	1 05	(25)	P. U. C., S. P.
LiC₉H₁₇O₂ nonylate	Tet ?			36 6	9 3	48	1 04	(25)	P U. C., S P.
LiC₁₁H₁₉O₂ undecylenate	H ?			52 6	9 5	72	0 99	(25)	P. U. C., S. P.
LiC₁₁H₂₁O₂ undecylate	Tet ?			41 8	9 2	48	0 04	(25)	P. U. C., S. P.
LiC₁₂H₂₃O₂ laurate	Tet ?			28 3	11 7	24	0 87	(25)	P. U. C., S. P.
LiC₁₈H₃₃O₂ oleate	H ?			64 6	9 5	72	0 99	(25)	P. U. C
LiC₁₈H₃₅O₂ stearate	H ?			62 5	9 8	72	1 04	(25)	P. U C
82 NaF	C.	NaCl(4b, 4c)		4 62		4	2 81	(75, 76, 209)	
NaHF₂	H.	3D₁-5(a, b, c)?	3D₁-5	5 17, 30° 44'		1	2 01	(211)	Na at (a), u_F = 0.42. P. S.
NaCl	C.	NaCl(4b, 4c)		5 628		4		(44, 45, 47)	One of the fundamental wave length standards
NaClO₃	C.	(4f, 4f, T-4(12))	T-4	6 56		4	2 49	(95, 143, 144, 147, 148, 149, 246, 247, 266)	u_Na = ca. 0 06, u_Cl = ca. 0.41. Different positions have been suggested for the O atoms
NaBr	C.	NaCl(4b, 4c)		5 94		4	3 24	(75, 76, 273)	
NaBrO₃	C	(4f, 4f, T-4(12))	T-4	6 71		4	3 30	(95, 143, 148, 149, 163, 246, 247)	u_Na = ca. 0.09, u_Br = ca. 0.41. Different positions have been suggested for the O atoms
NaI	C.	NaCl(4b, 4c)		6 46		4	3 67	(75, 76, 273)	
Na₂S	C	CaF₂(4b, 8e)	O₁-5	6 53		4	1 85	(239)	
NaN₃	H.	3D₁-5(a, b, c)	3D₁-5	5 48₁, 38° 43'		1	1.83s	(396)	u = 0.42s
NaNO₂	H	3D₁-6(a, b, c)	3D₁-6	6 32 48° 6'		2	2 19	(47, 367)	N atoms at (a). u₀ = 0.25
NaH(C₂H₃O₂)₂	C.		T₁-7?	15 9s		24	1 38	(279)	
NaC₂H₃O₂ v. Table C'.								(202)	Apparently very complicated
NaCd₂	C.			5 40	8 81	2			u_Al < 0 10; O positions not known
NaSb(AlO₃)₃	H.	6D₁-4(a or b, d, f, etc)	6D₁-4	5 33		4	2 53	(10)	
83 KF	C	NaCl(4b, 4c)		5 67	6 81	4	2 35	(75, 76, 132, 273)	
KHF₂	Tet.	4D₁-18(a, h)	4D₁-18	6 28o		4	1 98₇	(40)	u_F = 0 14 ± 0.01. The H atoms may have arrangement 4Di-18 (d)
KCl	C.	NaCl(4b, 4c)		6 57s		4	1 98₇	(44, 75, 76, 120)	
KBr	C	NaCl(4b, 4c)		7 05₂		4	2 76o	(44, 75, 120, 273)	
KI	C.	NaCl(4b, 4c)		9 36		4	3 12₄	(69, 70, 71, 75, 76, 120, 132, 273, 263, 366)	
KI₃	M.					4		(69, 70, 71)	P.U.C. b₀ and c₀ approx. = a₀, and β approx. = 90°.
K₂SO₄	R.		2D₁-16	5 73	7 42	4	2.70	(192, 276)	b₀ = 10.01
KN₃	Tet	4D₁-18(a, d, h)	4D₁-18	6 09₄	7.05s	4	2.04s	(396)	u = 013s
KH₂PO₄	Tet.	4d-12	4d-12	7 40	6 96	4	2.36	(342)	K atoms at 4d-12(a); P at 4d-12(b)
KCN	C.	NaCl-like		6 55		4	1 53	(27, 72, 73)	

Chemical symbol	Crystal system	Structure type	Space group	Unit cell, size, Å		M	Calculated density	Lit.	Additional data and remarks
				a_0	c_0				
KCNO	Tet.			6.07a	7.03o	4	2.06a	(396)	Structure similar to KN₃
KH₂C₄O₄Cl (H chloromaleate)	R.		2D-16(?)	7.62	10.95	8		(398)	b_0 = 15.74
KC₄H₅O₆ s. Table C'.									
K₂SnCl₆	C.	(4b, 8c, 24a)	Oₕ-5	9 96		4	2 74	(92)	u_{Cl} = 0.24₅ and < 0.25
K₂Zn(CN)₄	C.	(8f, 16c, 32b)	Oₕ-7	12 54		8	1 66	(93)	u_C = ca. 0.34, u_N = ca. 0.40, ½(u_C + u_N) = 0.37
K₂Cd(CN)₄	C.	(8f, 16c, 32b)	Oₕ-7	12 84		8	1 84	(93)	½(u_C + u_N) = 0.37
K₂Hg(CN)₄	C	(8f, 16c, 32b)	Oₕ-7	12 76		8	2 43	(93)	½(u_C + u_N) = 0.37
K₂PtCl₄	Tet.	4Dₕ-1(a, e, j)	4Dₕ-1	6 99	4 13	1	3 40	(96)	0.233 < u_{Cl} < 0.238
K₂PtCl₆	C.	(4b, 8c, 24a)	Oₕ-5	9 7		4	3 5	(219, 320)	Assigned value, u_{Cl} = 0 16, probably incorrect
K₂PdCl₄	Tet.	4Dₕ-1(a, e, j)	4Dₕ-1	7 04	4 10	1	2 65	(96)	u_{Cl} = 0.23
KCr(SO₄)₂.12H₂O	C	(4b, 4c, 8A, 8A, Tₕ-6 (24))	Tₕ-6	11 9a		4	1 97	(242)	
KAl(SO₄)₂.12H₂O	C	(4b, 4c, 8A, 8A, Tₕ-6 (24))	Tₕ-6	12 0a		4	1 81	(186, 237, 248, 282)	
KAlSi₃O₈ (adularia)	M		2Cₕ-3	8 57	7 23	4		(314)	b_0 = 13.01, β = 116° 7' Composition unknown
KLiSO₄	H.		6C 6?	5 13	8 60	2	2 39	(330)	P. U. C. An atomic arrangement is suggested
84 RbF	C ?	CsCl(1a, 1b)?		3 66?		1?		(78, 209, 294)	Structure probably incorrect
RbCl	C.	NaCl(4b, 4c)		6 571		4	2 81₂	(78, 102, 273, 366)	
RbBr	C	NaCl(4b, 4c)		6 86a		4	3 36₉	(74, 78, 120)	
RbI	C.	NaCl(4b, 4c)		7 32₅		4	3 56₆	(77, 78, 120, 273)	
Rb₂SO₄	R.		2Dₕ-16	5 95	7 78	4	3.66	(192)	b_0 = 10.3₉
CsF	C.	NaCl(4b, 4c)		6 01		4	4 62	(78, 209)	
CsCl	C.	CsCl(1a, 1b)	Oₕ-1	4 11o		1	3 99₉	(78, 85, 120)	
CsBr	C.	CsCl(1a, 1b)	Oₕ-1	4 29		1	4 45	(77, 78, 273)	
CsI	C.	CsCl(1a, 1b)	Oₕ-1	4.56₂		1	4 51₄	(69, 70, 71, 75, 78, 273)	
CsI₃	R.			6 82	11 01	4	4 51	(177, 178, 179, 325)	b_0 = 9.9₅
CsCl₂I	H.	3Dₕ-5(a, b, c)	3Dₕ-5	5 46; 70° 42'		1	3 88	(266)	I probably at (b); u_{Cl} = 0.31
CsBr₂I	R.		2Dₕ-16	6 57	10 66	4	4 29	(177, 178, 179, 325)	b_0 = 9.18
Cs₂SO₄	R.		2Dₕ-16	6 22	8 20	4	4 30	(192)	b_0 = 10.8₅
Tourmaline	H.		3e-1 3e-2	16 2a	7 2₆			(182)	P. U. C. Composition unknown
R'AlSi₂O₈ and R''Al₂Si₂O₈	Tri. and M							(116)	Unreduced powder- and Laue-photographs have been prepared from various feldspars

C-Table.—The C-Arrangement. See also Table C' infra

Chemical formula	Name	Crystal system	Unit cell, size, A			M	Calculated density	Lit.	Remarks
			a_0	b_0	c_0				
CH₄N₂O	Urea	Tet	5 63		4 7o	2	1 33	(25, 175)	Space group 4d-3
C₂H₂O₄	Oxalic acid	R	6.46	7 79	6 02	4	1 96	(315)	Space group 2Dₕ-15
C₂H₆	Ethane	H	4 46		8 19	2	0 70a	(349)	C atoms probably at 6Dₕ-4(f) with u = ca. 0 10 Temperature not stated.
C₂H₆N₂O	N-Methylurea .	R.	5 63	5 64	4 70	4?		(171)	Space group 2D-4?
C₂H₇NO	Acetaldehyde ammonia	H.	8 18, α = 84° 50'			6		(171, 316)	Space group 3Dₕ-5?
C₂H₆O₆	Oxalic acid dihydrate	M	6 05	3 57	11 9	2	1.68	(315)	Space group 2Cₕ-5. β = 106° 12'
C₃H₈N₂O	1, 2-Dimethylurea	R	4 53	10 9	5 14	2		(171)	Space group 2e-7?
C₄H₂O₃	Maleic anhydride	R	6 58	11 4a	5 90	4	1.44	(25)	P. U. C., 8 P
C₄H₂O₄	Acetylenedicarboxylic acid	M?	7.88	9 04	6 62	4	1.70	(25)	β = 111° 6'. P. U. C., 8 P
C₄H₄NIO₂	Iodosuccinimide	Tet	6 29		15 5a	4	2.41	(383)	P. U C. Space group 4C-2 and 4C-4?
C₄H₄O₃	Succinic anhydride	R	6 95	11 64	5.41	4	1 51	(296)	P. U C., cf. (25)
C₄H₄O₄	Maleic acid .	M	7 49	10 14	7.12	4	1 46	(25, 399)	β = 117° 7'. Space group 2Cₕ-5(?)
C₄H₅NO₂	Succinimide. .	R	7 50	9 60	12.75	8	1.42	(296)	P. U. C. Space group 2Dₕ-1?
C₄H₄O₄	Fumaric acid	T.	7 56	15 00	6.20	6		(399)	α = 90° 40', β = 88° 30', γ = 89° 48'
C₄H₆O₄	Succinic acid	M.	5 07	8 92	5 53	2		(296)	β = 91° 20'. P. U. C., cf (25)
C₄H₆O₆	dl-Tartaric acid.	Tri	14 8₂	9 74	4.90	4		(17)	α = 82° 20'; β = 122° 56'; γ = 111° 52'. P. U. C.
C₄H₆O₆	d-Tartaric acid.	M	7.70	6 04	6 20	2	1 76	(16)	β = 100° 17', cf (25)
C₅H₈N₄O₁₂	Pentaerythritol tetranitrate	Tet	13.2		6.66	4	1 80	(362)	Space group 4Dₕ-7
C₅H₁₂O₄	Pentaerythritol.	Tet.	6 16		8.76	2		(25, 176, 395)	Space group 4e-9
C₆H₄N₂O₄	o-Dinitrobenzene	M.	7 95	13 0	7.45	4		(25)	β = 112° 7'. P. U. C.
C₆H₄O₂	Quinone.....	M.	11 4o	6 43	6.85	4	1 40	(25)	β = 93° 20'. P. U. C., 8. P.
C₆H₆	Benzene......	R.	9 76	7 39	6.85	4	1 04	(44, 101, 372)	P. U. C., measurements at −20°C
C₆H₆O₂	Resorcinol .	R	9 56	10 2a	5 64	4		(53, 55)	P U C., cf. (25)
C₆H₆O₂	Hydroquinol. ...	M	13 5a	5 22	8 13	4		(53)	β = 107°. P. U. C.
		H	10 9a		7 55	6	1 39	(25)	P. U. C., Latter S. P.
(C₆H₁₀O₅)x	Cellulose and starch.........		Powder photographs have been obtained and possible units have been suggested					(124, 234)	

Chemical formula	Name	Crystal system	Unit cell, size, Å a_s	b_s	c_s	M	Calculated density	Lit.	Remarks
$C_6H_{12}N_4$	Hexamethylenetetramine	C.	7.02			2	1.33s	(100, 112)	u_N = ca. 0.12, u_C = ca. 0.23s. Structure type (8a, 12a); space group T6-4
$C_6H_{14}O_6$	d(l)-Mannitol.	R.	10 3s	8 1	4 5s	2	1.55	(27)	P U. C
$C_7H_6O_2$	Benzoic acid .	M.	5.44	5 18	21 6	4		(55)	β = 97° 5', P U C.
$C_4H_7NO_4$	Ammonium hydrogen fumarate...	T.	7 00	7 44	6 56	2		(396)	α = 107° 1', β = 117° 55', γ = 69° 16'
$C_4H_5ClN_2O_4$	Ammonium chlorofumarate.	M.	9 30	6 70	6 73s	2		(396)	β = 108° 25', Space group 2C-2(?).
$C_7H_6O_3$	Salicylic acid .	M.	11 5s	11 2s	4 93	4	1 58	(55)	β = 91° 22'. P. U. C
$C_7H_{14}O_6$	α-Methyl glycoside .	R.	10.8s	14 6s	5 61	4	1.46	(35)	P U C
$C_8H_4O_3$	o-Phthalic anhydride	R	7 74	13 6s	5 86	4	1.54	(25)	P. U. C., S. P.
$C_8H_6O_4$	o-Phthalic acid	M.	9.33	7.13	5 10	2	1.60	(25) cf (81)	β = 94° 36'. P. U. C, S. P.
$C_8H_{16}O_4$	Metaldehyde. .	Tet.	10 36	4.10		8		(171, 316)	Space group 4C-5?
$C_9H_8O_2$	trans-Cinnamic acid	M.	11 6s	14 1s	4 26	4	1 40	(55)	β = 98° 36'. P. U. C., S. P.
$C_9H_{10}O_2$	Hydrocinnamic acid	M	12 9s	9 20	6 98	4	1 23	(55)	β = 103° 36'. P. U C, S. P.
$C_{10}H_8$	Naphthalene .	M.	8 34	5 98	8 68	2		(52, 57)	β = 122° 44'. P U. C., cf. (25)
$C_{10}H_8O$	α-Naphthol	M	13 1	4 9	13 4	4	1 22	(53)	P. U. C. β = 117° 10'
$C_{10}H_8O$	β-Naphthol	M	11.70	4 28	17 4	4	1 22	(53)	P U. C. β = 119° 48'
$C_{12}H_{10}$	Acenaphthene	R.	8 32	14 1s	7 26	4	1 19	(52)	P. U. C.
$C_{12}H_{10}N_2$	Azobenzene	M	12 5s	5 2s	8 38	2	1 23	(28)	β = 116° P U. C.
$C_{12}H_{12}N_2$	Hydrazobenzene .	R.	11 1s	9.93	9 33	4	1 17	(28)	P U. C., S P.
$C_{12}H_{22}O_{11}$	Sacoharose. .	M	10 6s	8.7s	8 0s	2	1 57	(27)	β = 105° 44' P. U. C.
$C_{12}H_{24}O_2$	Lauric acid	Tet.?	28.3		11 4	24	0 86	(25)	P U. C., S. P. See Table C'.
$C_{14}H_8O_2$	Anthraquinone	R	12.0s	15 0s	2 69	2	1.40	(25)	P U C., S. P.
$C_{14}H_{10}$	Anthracene.	M	8 58	6 02	11.18	2	1 25	(52, 57)	β = 125°. P. U. C, cf. (25)
$C_{14}H_{10}$	Phenanthrene	M.	9.56	6 72	7 55	2	1 18	(25)	β = 92°. P. U. C., S. P.
$C_{14}H_{10}O_2$	Benzil	H	8.15		13 4s	3	1 41	(27)	P U. C.
$C_{14}H_{12}$	Stilbene	M	9 6	8 9	12.6	4	1 25	(27)	β = 118° 40'. P. U. C.
$C_{14}H_{14}$	Dibenzyl.	M	12.7	6 1	7.4	2	1 18	(27)	β = 119°. P. U. C.
$C_{14}H_{28}O_2$	Myristic acid	H?	57 4		11 4	72	0 83	(25)	P U C., see Table C'.
$C_{16}H_{10}N_2O_2$	Indigotin	H.	20 2		12 1s	12	1 20	(25)	P U C, Measurements also on S. P.
$C_{16}H_{32}O_2$	Palmitic acid.... .	H.?	60. 0		11 0	72	0 88	(25)	P. U. C, see Table C'.
$C_{18}H_{34}O_2$	Elaidic acid	Tet ?	26.5		10 s	16	0.98	(25)	P. U. C, S P., see Table C'.
$C_{18}H_{36}O_2$	Stearic acid	H.?	62.0		10.7	72	0.94	(25)	P. U. C., S. P, see Table C'.
$C_{19}H_{16}$	Triphenylmethane	R.	14.5s	25.6s	7.42	4		(23, 26) cf. (177, 178)	
$C_{19}H_{16}O$	Triphenylcarbinol	H.	16.5		8.8	6	1 23	(27)	P. U. C.
$C_{36}H_{70}O_4$	α, α'-Distearin.	H ?	81.5		10 8	48	0 82	(25)	P U C, S P

C'-Table.—Long Chain Compounds

Arrangement by Classes

1. Aliphatic Hydrocarbons (320, 401)

Formula	Maximum spacing, Å d_1	Spacings of broad lines, Å d_2	d_3	d_4	d_5	d_6	d_7
$C_{17}H_{36}$	24.3	4 25	3 93		2.54	2.32	
$C_{18}H_{38}\alpha$	25.9		4.0				
$C_{18}H_{38}\beta$	23.9	4 58	3.80	3.66	2.61		2 05
$C_{19}H_{40}$	26.9	4 22	3 84		2 52	2 25	
$C_{20}H_{42}\alpha$	28.0		3 9				
$C_{20}H_{42}\beta$	26.2	4.63	3.82	3.61	2 59	2 12	2 03
$C_{21}H_{44}$	29.4s	4.17	3.77	3 01	2 50	2 25	
$C_{22}H_{46}$	32.2						
$C_{24}H_{50}$	33.0s	4 18	3 80	3 02	2.50	2 25	
$C_{27}H_{56}$	37.1	4.17	3.77	3 01	2 51	2 25	
$C_{31}H_{64}$	43.0	4.14	3 74	2.99	2.49	2 21	
$C_{35}H_{72}$	47.7						

Formula	Max. spacing	Formula	Max. spacing
$C_{22}H_{46}$(?)	30.6	$C_{30}H_{62}$	40 4
$C_{24}H_{50}$	32.9	$C_{31}H_{64}$	41 6*
$C_{26}H_{52}$	34.3		42 9†
$C_{26}H_{54}$	35 6	$C_{32}H_{66}$	42.7
$C_{28}H_{58}$	37.7	$C_{34}H_{70}$	45.3
$C_{29}H_{60}$	39 4		

Specimens for (320) pressed, those for (401) melted on glass plates only.

* Melted.   † Pressed.

2. Aromatic Hydrocarbons

$C_{24}H_{42}$, Octadecylbenzene, d_1 = 49.2₀(225)

3. Aliphatic Acids
a. Monobasic

Formula	Name	Maximum spacing, Å d_1	Broad line spacing, Å d_2	d_3	d_4	d_5	Lit.
CH_2O_2	Formic	5 19					(309)
$C_2H_4O_2$	Acetic	6.66					(309)
$C_3H_6O_2$	Propionic	6.75	4.03			3.43	(309)
$C_4H_8O_2$	Butyric	9.65	4.09	3.65		3.45	(309)
$C_5H_{10}O_2$	Valeric	10.1(?)					(309)
$C_6H_{12}O_2$	Caproic	14.6	4.14	3.65		3.47	(309)
$C_7H_{14}O_2$	Heptoic	16.4	4.29	3.75	3.97	3.49	(309)
$C_8H_{16}O_2$	Caprylic	19.0	4.14	3.65		3.48	(309, 354)
$C_9H_{18}O_2$	Nonylic	22.9	4.22	3.71	3.97	3.48	(309)
$C_{10}H_{20}O_2$	Capric	23.3	4.14	3.73			(354, 309, 274)
$C_{11}H_{22}O_2$	Undecylic	25.8					(185)
$C_{12}H_{24}O_2$	Lauric	27.0	4.11	3.68			(184, 354)
$C_{14}H_{28}O_2$	Myristic	32.2	4.12	3 72			(184, 354)
$C_{15}H_{30}O_2$	Pentadecylic	36.2	4.00	3.76			(185)
$C_{16}H_{32}O_2$	Palmitic	34.7	4.08	3.65			(184, 354)
$C_{17}H_{34}O_2$	Margaric	39.2	4.05	3.77			(185)
$C_{18}H_{34}O_2$	Oleic	36.2(?)					(185)
$C_{18}H_{34}O_2$	Isoleic	35.9					(185)
$C_{18}H_{34}O_2$	Elaidic	48.3	4.03	3.65			(185)

3. Aliphatic Acids. a. Monobasic.—(Continued)

Formula	Name	Maximum spacing, Å d_1	Broad line spacing Å				Lit.
			d_2	d_3	d_4	d_5	
$C_{18}H_{36}O_2$	Stearic	38.7	4 05	3 62			(184, 354)
$C_{22}H_{42}O_2$	Erucic	46 3	4 22	3 72			(185)
$C_{22}H_{42}O_2$	Brassidic	59 9	4 25	3.72			(185)
$C_{22}H_{44}O_2$	Behenic	47 8	4 10	3 66			(184)

b. Dibasic

$C_4H_6O_4$	Succinic	4 5					(354)
$C_6H_{10}O_4$	Adipic	7 0					(354)
$C_7H_{12}O_4$	Pimelic	7 6					(354)
$C_8H_{14}O_4$	Suberic	9 3					(354)
$C_9H_{16}O_4$	Azelaic	9 6					(354)
$C_{10}H_{18}O_4$	Sebacic	11 4					(354)

4. Salts

Formula	Name	Maximum spacing Å d_1	Broad line spacing Å				Lit
			d_2	d_3	d_4	d_5	
$PbC_{12}H_{22}O_4$	Caproate	20 0					(355)
$PbC_{16}H_{30}O_4$	Caprylate	25 4					(355)
$PbC_{20}H_{38}O_4$	Caprate	30 6					(355)
$PbC_{24}H_{46}O_4$	Laurate	35.8					(355)
$PbC_{28}H_{54}O_4$	Myristate	41.2					(355)
$PbC_{32}H_{62}O_4$	Palmitate	46 3					(355)
$PbC_{36}H_{66}O_4$	Oleate	37.5; 29 8					(355)
$PbC_{36}H_{68}O_4$	Elaidate	50 0					(355)
$PbC_{36}H_{70}O_4$	Stearate	51 3					(355)
$NaC_{12}H_{23}O_2$	Laurate	33.5	4 22	4 88			(208)
$NaC_{14}H_{27}O_2$	Myristate	38 5	4 18	4.9			(208)
$NaC_{16}H_{31}O_2$	Palmitate	43 5	4 15	4.9			(208)
$NaC_{18}H_{33}O_2$	Oleate	43 5					(63)

Similar results obtained with K and NH_4 oleates

5. Esters

Formula	Name	d_1	Broad line spacing			Lit.
$C_{17}H_{34}O_2$	Methyl palmitate	22.0	4 07	3 72		(225)
$C_{18}H_{19}N_3O_4$	Ethyl p-azoxybenzoate	16 2	d_1 = 19.9 in the "smectic" state			(321)
$C_{18}H_{36}O_2$	Ethyl palmitate	23 2	4 07	3 67		(225)
$C_{19}H_{38}O_2$	Methyl stearate	24 0	4 07	3 74		(225)
$C_{20}H_{40}O_2$	Ethyl stearate	25 2	4 11	3 69		(225)
$C_{24}H_{48}O_2$	Octyl palmitate	30 4	4 10	3 72		(225)
$C_{22}H_{44}O_2$	Cetyl palmitate	40 4	4 05	3 69		(225)
$C_{44}H_{104}O_6$	Glycerol margarate	48 0				(355)

6. Ketones (319)

Formula	Name	Maximum spacing Å d_1
$C_{13}H_{26}O$	Di-n-hexyl	18 7
$C_{14}H_{30}O$	Methyl-n-tridecyl	42.4
$C_{17}H_{34}O$	Methyl n-pentadecyl	47.6
$C_{18}H_{36}O$	Methyl n-hexadecyl	50.0
$C_{18}H_{36}O$	Ethyl n-pentadecyl	25.2
$C_{18}H_{36}O$	Hexyl n-undecyl	25.2
$C_{19}H_{38}O$	Methyl n-heptadecyl	52.9
$C_{19}H_{38}O$	Propyl n-pentadecyl	26.3
$C_{20}H_{40}O$	Ethyl n-heptadecyl	27 3
$C_{21}H_{42}O$	Propyl n-heptadecyl	28.9
$C_{22}H_{44}O$	Hexyl n-pentadecyl	31.1
$C_{23}H_{46}O$	Di-n-undecyl	31.6
$C_{24}H_{48}O*$	Hexyl n-heptadecyl	33.6
$C_{27}H_{54}O$	Di-n-tridecyl	37.0
$C_{31}H_{62}O$	Di-n-pentadecyl	41.1
$C_{35}H_{70}O$	Di-n-heptadecyl	47.2

* A few orders of 30.8Å also present.

7. Phenols (225)

$C_{22}H_{38}O$	p-Hexadecyl	46.5
$C_{24}H_{42}O$	p-Octadecyl	51.3

TABLE D.—Alloys

(a) Non-ferrous. Standard Arrangement. All Compositions in Atomic %

Pb-Sn.—0 to 3.6% Sn alloys are F.-c. cubic (like Pb) with a_0 decreasing to 4.931Å, taking a_0 for Pb as 4.942Å. 10% – 95% Sn alloys are mixtures of the Pb-like and Sn structures. 95% – 100% Sn alloys show no measurable distortion in size or shape of the Sn unit cell (206).

Hg-Sn.—The structure varies, as follows, with the atomic % of Hg: 0 to ±2%, Tet.-Sn structure I; 2% I, with traces of "Hexagonal" amalgam, (composition unknown) structure II; 5%, I and II; 6%, trace of I with II; 6 to ±17%, II; ±17 to 33%, II and liquid alloy (229).

Hg-Pb.—A 20% Hg alloy had the F.-c. cubic structure (4b) of Pb, with a unit cell length 1.6% less than that of Pb (229).

Hg-Zn.—Two structures, the hexagonal Zn structure (d), and an "hexagonal" structure belonging to an amalgam of unknown composition. The relative intensities of the patterns of these two phases are as follows (229):

Atomic % Hg	0	10	20	35
Zn structure	strong	medium	weak	absent
"Amalgam" structure	absent	medium	strong	strong

Hg-Cd.—An 18% Hg amalgam gave a pattern substantially the same as that of Cd; 37 and 50% Hg amalgams yield a different pattern (229).

Cu-Si.—Though Si has the smaller atomic volume the unit cube of Cu which has dissolved Si is larger than that of pure Cu. No data available (84).

Cu-Sn.—Figure 12a. Black circles: metal melted in air; open circles: metal melted in vacuum (18, 372).

Cu-Zn.—Figure 13. Unless otherwise stated on the figure these data are from (198). Cf. (12, 199, 255, 375, 371) which gives a different structure for γ-brass.

Ag-Sn.—Solution of Sn increases the Ag unit though its atomic volume is less. No data available (84).

Ag-Zn.—The observed phases are the same as those for Cu-Zn alloys (371).

Phase	Composition wt. % Zn	Symmetry	Structure	a_0 Å	c_0 Å	No. atoms in unit cell
β	38 25	Cubic	(1a, 1b)	3.156		2
γ	50 3	Cubic		9.327		52 37
ε	{ 60 5	Hexagonal	Mg-like	2 818	4 456	2
	78 1	Hexagonal	Mg-like	2 815	4 382	2
η	Hexagonal close-packed with Zn-like structure					

Ag-Cu.—Broken series of solid solutions. Both components F -c cubic (4b) (**370**).

At. % Cu	0	4	9 2	16-80	96 4	100
a_0	4 06	4 05	4 03	Superimposed patterns of Ag and Cu	3 61	3 61

Au-Zn.—These alloys show all the phases of Cu-Zn alloys and two additional (**371**).

Phase	Composition wt. % Zn	Symmetry	Structure	a_0 Å	c_0 Å	No atoms in unit cell
β	30 2	Cubic	(1a, 1b)	3 146		2
γ	{ 36 9	Cubic		9 268		52 97
	41 1	Cubic		9 223		51.96
ε	{ 67 5	Hexagonal	Mg-like	2 809	4 377	2
	72 3	Hexagonal	Mg-like	2 809	4 369	2
η	95 0	Hexagonal	Zn-like	2 674	4 887	2
γ' (AuZn₃)?	50 2	Cubic	?	7 880		32
γ''			may be cubic			

Au-Cu.—Figure 12 (**18, 145, 361**).

Au-Ag.—Data conflicting. Probably an unbroken series of solid solutions, though marked variations from this relation have been reported. Figure 16 (**18, 165, 239, 372**).

Ir-Os.—A single alloy of unknown composition was found to be C.-p. Hex. (**11**).

Pd-H.—Data conflicting. One result (**295, 376**) shows that the Pd unit is swelled by an amount proportional to the quantity of occluded H (**79**). The other study (**164**) shows a discontinuous absorption of H in the sense that some crystals may be saturated though others in the same material have not begun to absorb gas. The length, a_0, of the edge of the unit cube of the saturated solution was found to vary between 4.000Å and 4.039Å with values usually not less than 4.023Å.

Pd-Cu and Pd-Au.—Figures 20 and 19 (**301**).

Pd-Ag.—(**15**) Figure 17 (**165**).

Mn-Cu.—67 % Cu is F.-c. cubic, like Cu, and has $a_0 = 3.615$Å, taking a_0 for Cu as 3.60Å (**18**). 70 % Cu is said to give $a_0 = 3.70$Å (**200, 384**).

Ni-Cu.—Figure 15 (**18, 197, 361, 370**).

Cr-Ni.—100 % to 40 % Ni alloys are F.-c. cubic (like Ni) with values of a_0 which change proportionately to the % of Cr added from 3.521Å (for Ni) to 3.576Å (**206**).

W-Mo.—(**67**) Said to show an unbroken series of solid solutions. No numerical data available (**18**). No lines (**86**) have been found from a 1:1 alloy to indicate the existence of a compound W-Mo (**239**).

Al-Zn.—0 to 20 % Zn alloys are F.-c. cubic (like Al), a_0 changing from 4.043Å (for Al) to 4.034Å. 20 % – 95 % Zn alloys show mixtures of cubic Al and hexagonal Zn structures. 95 % – 100 % Zn alloys are C.-p. hexagonal with no measurable distortion from size or shape of the Zn unit cell (**206**).

Al-Cu.—Figure 14. The data on this figure are from (**22, 141, 197, 258**).

Al-Ag.—The dissolving of Al in Ag increases the unit cube in the latter, though Al has a smaller atomic volume. No numerical data available (**84**).

Al-Mn-Cu.—Heussler Alloys. Alloy 15.9 % Al, 23.9 % Mn, 60.3 % Cu is said to be F.-c. cubic with $a_0 = 3.70$Å. Alloys 14.3 % Al, 28.6 % Mn, 57.1 % Cu is said to be a mixture of the preceding structure with a smaller amount of a B.-c. cubic phase having $a_0 = 2.98$Å (**12, 197**).

Mg-Sn.—0 to 67 % Mg give the superimposed patterns of Sn and Mg₂Sn; 67 – 100 % Mg yield the superimposed patterns of Mg₂Sn and Mg. No evidence of solid solution (**370**).

Mg-Pb.—0 to 67 % Mg give the superimposed patterns of Pb and PbMg₂; 67–100 % Mg yield the superimposed patterns of PbMg₂ and Mg. No evidence of solid solution (**370**).

Mg-Al.—91.2 % Al is F.-c cubic (1b) with $a_0 = 4.106$Å, taking a_0 for Al as 4.05Å. 7.3 % Al is C.-p. hexagonal (d) with $a_0 = 3.151$Å, $c = 5.23$Å, taking a_0 for Mg as 3.17Å and $c_0 = 5.17$Å (**197**).

(b) Ferrous Alloys

Fe-C Steels.—(1) Austenitic Steels. Structure that of γ-Fe, F.-c. cubic (4b) (**250 259**)

Composition, wt. %	a_0 in Å	Remarks
(1) 1.25 % C, quenched at 750°C	3 601	Contains also martensite.
(2) 1 98 % C, quenched at 1100°C	3 629	Contains also martensite.
(3)* 1.34 % C, 12.1 % Mn, 0 52 % Si, 0.1 % P	3 621	
(2) quenched at 750°C	3 606	A mixture of austenite and martensite.
(4) 1.18 % C, 24.3 % Ni, 6.05 % Mn quenched from 1000°C	3 64	
(5) 0.24 % C, 25.2 % Ni, quenched from 1000°C	3 56	

* Density calculations thought to indicate that C is present in interstitial solid solution in steel No. (3)

(2) Martensite Steels. Structure that of α-Fe, B.-c. cubic (2a) (**19, 122, 250 258**).

	a_0	
(5) Chilled subsequently in liquid air	2 81	Partly martensite and partly austenite.
(2)	2 90	Martensite lines very diffuse.
(1)	2 88	Martensite lines very diffuse.
(6) 0.80 % C quenched in oil from 750°C	2 89	Martensite lines very diffuse.
(7) 0.80 % C, 0.14 % Cr, 0.35 % Mn, 0.19 % Si	2 851	Broad lines, less intense than from Fe.
(8) 1.31 % C, 0.12 % Cr, 0.24 % Mn, 0.17 % Si	2 851	Density calculations from this steel thought to indicate that C isomorphously replaces Fe unless martensite is annealed when it is a mixture of α-Fe with cementite.

Fe-Si.—(**207, 252, 389**).

Weight % Si	0-15	17-30	33	40	50	75-100
Phases	Fe	Fe + FeSi	FeSi	FeSi + FeSi₂	FeSi₂	FeSi₂ + Si

Cu and Au are both F-c cubic They form an unbroken series of solid solutions

FIG. 12.—The diffraction data on Cu-Au alloys.

FIG. 12a.—The diffraction data on Cu-Sn alloys.

The α- brasses have the copper structure The value of a_o varies as shown below. (Face-centered cubic structure)

β-brass gives the pattern of a body-centered cube containing 2 molecules of Cu Zn and having a_o=2.95 Å. No difference was detected in samples quenched from the β and β'regions.

γ- brass gives a pattern attributed to a rhombo-hedral arrangement having a_o=4.14 and c/a=0.649. Another study (258) found Laue and powder photographs giving a body-centered cubic structure for 62, 65, and 67% Zn alloys.

ε-brass gives a pattern which has been in-terpreted as hexagonal with a_o=2.72 A° and c/a = 1.58.

Zinc structure; Hexagonal Close-packed

FIG. 13.—The diffraction data on brasses.

α-phase. Copper structure with values of a_o shown below. Face-centered cubic

A fine-grained mixture of α- and γ¹ phases. Preliminary attempts to study phases stable at high temperatures are yet inconclusive.

Pattern of γ¹ phase. Cubic with a_o values as shown above. Composition of crystalline phase uncertain.

It has been considered that an alloy of this composition is the compound CuAl and a structure has been assigned to it. Probably incorrect.

A mixture of γ¹ and θ- phases.

Pure θ- phase. Compound $CuAl_2$.

Mixtures of θ- phase $(CuAl_2)$ and Al patterns in varying proportions.

Al pattern. Face centered cubic

FIG. 14.—The diffraction data on Cu-Al alloys.

F-c cubic. Unbroken series of solid solutions.

FIG. 15.—The diffraction data on Cu-Ni alloys.

after Bain

FIG. 16.—The diffraction data on Ag-Au alloys.

FIG. 17.—The diffraction data on Ag-Pd alloys.

FIG. 18.— The diffraction data on Fe-Ni alloys.

FIG. 19.—The diffraction data on Au-Pd alloys.

FIG. 20.—The diffraction data on Cu-Pd alloys.

FIG. 21a.—The diffraction data on solid solutions of the alkali halides

Fe-Mn.—These alloys are said to have the following structures No numerical data available [18].

Atomic % Mn.	0-30	30 60	60 100
Structure.	B.-c. cubic (2a)	F.-c. cubic (4b)	Complex Mn

Fe-Co.—No numerical data available [12].

Weight % Co	0-80	85	90 98	98 100
Structure.	B.-c. cubic (2a)	B.-c.(2a) with F.-c. (4b)	F.-c. cubic (4b)	F.-c (4b) with C.-p. hex

Fe-Ni.—The best available data are shown in Fig. 18 The fused alloys were swaged, drawn and rolled into thin tapes. Spacings from photographs of these specimens without further treatment are shown as open circles, results after (1) annealing at 900-950°C followed by slow cooling, black circles; (2) after an additional heating to 600°C followed by rapid cooling in the air, crosses; and (3) after cooling for a time in liquid air following (1), triangles [12, 168].

Fe-Cr.—Interpretation of data uncertain [18].

Fe-W and **Fe-Mo.**—It is said that Fe dissolves a few atomic percents of each of these metals without apparent alteration in the size of the unit cell. In each case a 1:1 compound is formed. No numerical data available [18]. •

FIG. 21b.—The diffraction data on solid solutions of the alkali halides.

𝕏-TABLE.—THE POSITIONS OF X-RAY DIFFRACTION BANDS FROM LIQUIDS

Angle of Deviation and Wave Length, λ, of X-rays Used

Liquid	A		N₂		O₂	
Angle, deg	13 0; 18.9	27	11.3; 17 0	12 5; 19.5	27
λ, in Å	0 712	1 54	0 712	0 712	1.54
Lit .	(304)	(303)	(304)	(303)	(302)

Liquid	H₂O	CS₂	HCOOH	CH₃CHO Acetaldehyde
Angle, deg	13.4	29	13 2	24	22.7
λ, in Å	0 712	1 54	0 712	1 54	1 54
Lit .	(304)	(303)	(304)	(303)	(373)

Liquid	C_2H_6OH	$C_4H_8O_2$ Butyric acid	$C_4H_8O_2$ Ethyl acetate	$(C_2H_6)_2O$
Angle, deg	22	20 7; 36 5	20 7	19
λ, in Å	1.54	1 54	1 54	1 54
Lit	(303)	(373)	(373)	(303)

Liquid	C_6H_6		$(C_2H_4O)_3$ Paraldehyde	C_6H_6CHO Benzaldehyde
Angle, deg	8 5	18	23 3	19 3; 44 4
λ, in Å	0 712	1 54	1 54	1 54
Lit	(301)	(302, 303)	(373)	(373)

Liquid	C_5H_{12}	C_9H_{12} Mesitylene	$C_{14}H_{12}O_2$ Benzyl benzoate
Angle, deg	8 1	4 1; 6 2	18 3; 42 7; 65 8
λ, in Å	0 712	0 712	1 54
Lit	(301)	(301)	(373)

3-Table.—Data on Solid Solutions of Salts

Alkali Halides.—For data on the solutions NH_4I-NH_4Br, NH_4I-KI, $NH_4Br-KBr$, $RbCl-NH_4Cl$, $NH_4Cl-KCl$, $KCl-RbCl$, $KCl-KBr$, $CsCl-NH_4Cl$, NH_4Br-NH_4Cl, $RbCl-NH_4Cl$ see Fig. 21 (120). For additional data on KBr-KCl see (387, 388).

AgCl-NaCl (387).—Broken series of solid solutions. Quenched preparations: Both patterns present together.

	Composition mol % AgCl	a_0 Å
Annealed	100	5 53
	75	5 54
	50	5 57

AgCl-AgBr (402).—Both structures like NaCl (4b, 4c). Unbroken series of solid solutions

Composition mol % AgCl	a_0 Å
0	5 77
20	5 72
40	5 68
50	5 65
60	5 63
80	5 59
100	5 54

AgBr-AgI (402).—Broken series of solid solutions

Composition mol % AgI	a_0				
	Fused and slowly cooled		Fused and quenched		Precipitated
	Structure (4b, 4c)	Structure (4b, 4d)	Structure (4b, 4c)	Structure (4b, 4d)	Structure (4b, 4c)
0	5 768		5 768		5 768
10	5.814		5 816		5 806
20	5 842		5 854		5 84
30	5 86		5 876		5 878
40	5 896	(6 47)	5 908		
50	5 912	(6 47)	5 932		
60	5 918	6 47	5 96	(6 48)	
70	6 014	6 48	5 956	6 48	
	5 946				
	5 994				
80	5.916	6.47	(5 892)	(6 48)	
90		6.472	5.898	6 483	
95		6 481		6 487	
100		6 493		6 493	

LITERATURE

(For a key to the periodicals see end of volume)

(1) Allison, *12*, *8*: 261; 24. (2) Alsen and Aminoff, *207*, **44**: 124; 22 (3) Aminoff, *207*, **41**: 407; 19 (4) Aminoff, *94*, **56**: 495; 21 (5) Aminoff, *94*, **56**: 506, 21. (6) Aminoff, *94*, **57**: 180; 22. (7) Aminoff, *94*, **57**: 204, 22 (8) Aminoff, *207*, **44**: 444; 22. (9) Aminoff, *94*, **56**: 203; 23.

(10) Aminoff, *94*, **60**: 262; 24. (11) Aminoff, *94*, **56**: 510; 21. (12) Andrews, *2*, **18**: 245; 21. (13) van Arkel, *208*, **4**: 286; 24 (14) Asahara, *210*, 1, 23, 22 *209*, 1: 35; 22 (15) Astbury, *58*, **112**: 53; 23. (16) Astbury, *5*, **102A**: 506, 23 (17) Astbury, *5*, **104A**: 219; 23. (18) Bain, *33*, **28**: 21, 576, 23 (19) Bain, *80*, preprint; Feb 24.

(20) Bartlett and Langmuir, *1*, **43**: 84; 21. (21) Beckenkamp, *95*, **110**: 290, 20 (22) Becker, *95*, **15**: 303; 23. (23) Becker, *96*, **24**: 65; 24. (24) Becker and Ebert, *96*, **16**: 165; 23. (25) Becker and Jancke, 7, **99**: 242; 21 (26) Becker and Rose, *96*, **17**: 351; 23. (27) Becker and Rose, *96*, **14**: 369, 23 (28) Berndt, *211*, **36**: 1,21 (29) Bijl and Kolkmeijer, *176*, **15**: 1264, 18

(30) Bijl and Kolkmeijer, *64P*, **21**: 405, 19 (31) Bijl and Kolkmeijer, *64P*, **21**: 494, 501, 19 (32) Bijvoet, *70*, **42**: 859, 23. (33) Bijvoet and Karssen, *64P*, **23**: 1365; 21 (34) Bijvoet and Karssen, *64P*, **25**: 27; 22 (35) Bijvoet and Karssen, *70*, **45**: 680; 24. (36) Bohlin, *8*, **61**: 421; 20 (37) Bozorth, *1*, **44**: 317; 22. (38) Bozorth, *1*, **44**: 1066; 22. (39) Bozorth, *1*, **44**: 2232, 22.

(40) Bozorth, *1*, **45**: 2128, 23. (41) Bozorth, *1*, **45**: 1621; 23. (42) Bradley, *3*, **45**: 477; 24 (43) Bradley, *3*, **47**: 657; 24 (44) Bragg, *5*, **89**: 248, 13 (45) Bragg, *5*, **89**: 246; 13 (46) Bragg, *5*, **28**: 355; 14. (47) Bragg, *5*, **89A**: 468, 14 (48) Bragg, *5*, **89A**: 575; 14 (49) Bragg, *62*, **215**: 253, 15 (50) Bragg, *58*, **95**: 561; 15 3, **30**: 305, 15 (51) Bragg, *3*, **89**: 647; 20 (52) Bragg, *67*, **33**: 304, 21 (53) Bragg, *67*, **34**: 35; 21 (54) Bragg, *67*, **34**: 98; 22 (55) Bragg, *4*, **121**: 2766, 21 (56) Bragg, *58*, **111**: 532; 23 (57) Bragg, *67*, **36**: 167, 23 (58) Bragg, *5*, **105A**: 16; 24. (59) Bragg and Bragg, *58*, **91**: 557, 13.

(60) Bragg and Bragg, *5*, **89**: 277; 13. (61) Bragg and Bragg, *X-rays and Crystal Structure*, 4th ed , (London, 24) (62) Bragg and Morgan, *5*, **104A**: 437, 23 (63) de Broglie and Friedel, *34*, **176**: 738, 23. (64) Broomé, *63*, **24**: 124, 23 (65) Burdick and Ellis, *197*, **3**: 644; 17. 1, **39**: 2518, 17 (66) Burdick and Owen, *1*, **40**: 1749; 18. (67) Burger, *208*, **2**: 114; 22 (68) Clark, *1*, **46**: 372, 24. (69) Clark and Duane, *2*, **20**: 85; 22.

(70) Clark and Duane, *48*, **7**: 455; 23. (71) Clark and Duane, *197*, **9**: 126; 23 (72) Cooper, *58*, **107**: 745, 21. (73) Cooper, *58*, **110**: 544; 22 (74) Davey, *2*, **17**: 402; 21 (75) Davey, *2*, **18**: 102; 21 (76) Davey, *2*, **19**: 248; 22 (77) Davey, *2*, **19**: 538; 22 (78) Davey, *2*, **21**: 143; 23 (79) Davey, *2*, **21**: 213, 23

(80) Davey, *2*, **21**: 380; 23. (81) Davey, *2*, **21**: 716; 23 (82) Davey, *2*, **23**: 292; 24 (83) Davey, *2*, **23**: 763, 24 (84) Davey, *212*, **6**: 375, 24 (85) Davey and Wick, *2*, **17**: 403; 21 (86) Davey and Hoffman, *2*, **18**: 333, 20 (87) Debye, *63*, **18**: 483, 17. (88) Debye and Scherrer, *63*, **17**: 277; 16 (89) Debye and Scherrer, *63*, **18**: 291, 17

(90) Dennison, *2*, **17**: 20, 21 *135*, **122**: 54; 21 (91) Dickinson, *1*, **43**: 85; 20 (92) Dickinson, *1*, **44**: 276; 22 (93) Dickinson, *1*, **44**: 774; 22. (94) Dickinson, *1*, **44**: 1489; 22 (95) Dickinson, *1*, **44**: 2404, 22. (96) Dickinson, *1*, **45**: 958; 23 (97) Dickinson and Friauf, *1*, **46**: 2457; 24 (98) Dickinson and Goodhue, *1*, **43**: 2045; 21. (99) Dickinson and Pauling, *1*, **45**, 1466, 23.

(100) Dickinson and Raymond, *1*, **45**: 22, 23. (101) Eastman, *1*, **46**: 917; 24 (102) Espig, *211*, **36**: 53; 21 (103) Ewald, *8*, **44**: 257; 14. (104) Ewald, *63*, **15**: 399, 14 (105) Ewald, *213*, **1914**: 325. (106) Ewald and Friedrich, *8*, **44**: 1183, 14 (107) Gerlach, *63*, **22**: 557; 21 (108) Gerlach, *63*, **23**: 114. 22 (109) Gerlach, *96*, **9**: 184, 22

(110) Gerlach and Pauli, *96*, **7**: 116, 21 (111) Goldschmidt and Thomassen, *214*, No **2**: 5, 23 (112) Gonell and Mark, 7, **107**: 181, 23 (113) Greenwood, *3*, **43**: 654, 24 (114) Gross, *189*, **1919**: 201 (115) Gross and Gross, *190*, **43**: 113, 23 (116) Hadding, *215*, **17**: No. 6, 20 (117) Hassel and Mark, *96*, **27**: 89; 24 (118) Hassel and Mark, *96*, **23**: 269; 24. (119) Hassel and Mark, *96*, **25**: 317; 24.

(120) Havighurst, Mack and Blake, *1*, **46**: 2368; 24. (121) Hedvall, *19*, **8**: No. 11; 22. *95*, **130**: 327; 22 (122) Heindlhofer, *2*, **24**: 426; 24. (123) Hentschel, *189*, **1923**: 609 (124) Herzog, Jancke and Polanyi, *96*, **3**: 196, 343, 20 (125) Holgersson, *95*, **126**: 179; 23 (126) Hull, *2*, **9**: 84; 17 (127) Hull, *2*, **9**: 564; 17 (128) Hull, *2*, **10**: 661; 17. (129) Hull, *197*, **3**: 470; 17.

(130) Hull, *2*, **13**: 292; 19 (131) Hull, *2*, **14**: 540; 19 (132) Hull, *119*, **33**: 1171; 19 (133) Hull, *2*, **15**: 545; 20. (134) Hull, *166*, **53**: 227; 20. (135) Hull, *2*, **17**: 42; 21. (136) Hull, *2*, **17**: 571; 21. (137) Hull, *2*, **18**: 88; 21 (138) Hull, *2*, **20**: 113; 22. (139) James, *3*, **42**: 193; 21.

(140) James and Tunstall, *3*, **40**: 233; 20 (141) Jette, Phragmen and Westgren, *47*, **31**: 193; 24 (142) Kahler, *2*, **18**: 210; 21. (143) Karssen, *70*, **42**: 904; 23. (144) Kiby, *96*, **17**: 213; 23 (145) Kirchner, *8*, **69**: 59; 22. (146) Kolkmeijer, *64P*, **26**: 125 *64V*, **31**: 155; 22 (147) Kolkmeijer, Bijvoet and Karssen, *64P*, **23**: 644; 21 (148) Kolkmeijer, Bijvoet and Karssen, *96*, **14**: 291; 23. (149) Kolkmeijer, Bijvoet and Karssen, *96*, **20**: 82; 23.

(1.0) Kolkmeijer, Bijvoet and Karssen, 70, 42: 677; 24. (151) Kolkmeijer, Bijvoet and Karssen, 70, 42: 894; 24. (152) Kulasewski, 211, 38: 81, 21 (153) Kustner and Remy, 63, 24: 25; 23. (154) Lehmann, 94, 60: 379; 24 (155) Leonhardt, 189, 1923: 641. 94, 59: 216; 24. (156) Levi, 59, 1: 137, 24. (157) Levi, 78, 57: 619; 24. (158) Levi, 216, 6: 333, 24. (159) Levi and Ferrari, 28, 33: 397; 21.

(160) Levi and Ferrari, 28, 33: 516; 24. (161) Levi and Quilico, 56, 54: 154; 24 (162) McKeehan, 197, 8: 254; 22. (163) McKeehan, 197, 8: 270, 22. (164) McKeehan, 2, 20: 82; 22. (165) McKeehan, 2, 20: 424; 22 (166) McKeehan, 143, 196: 59; 23. (167) McKeehan, 2, 21: 334; 23 (168) McKeehan, 2, 21: 402; 23. (169) McKeehan, 2, 21: 503; 23.

(170) McKeehan, 2, 19: 444; 22. (171) Mark, 25, 57: 1820; 24 (172) Mark and Polyani, 96, 12: 75; 23. (173) Mark and Polyani, 96, 22: 200, 24 (174) Mark, Polyani and Schmid, 218, 11: 256; 23 (175) Mark and Weissenberg, 96, 16: 1; 23 (176) Mark and Weissenberg, 96, 17: 301; 23 (177) Mark and Weissenberg, 96, 17: 347; 23 (178) Mark and Weissenberg, 96, 24: 68; 24. (179) Mauguin, 54, 176: 1331; 23

(180) Mauguin, 54, 176: 1483; 23 (181) Mauguin, 54, 176: 785; 24. (182) Mauguin, 54, 176: 1913, 24 (183) Mechling, 211, 36: 37, 21 (184) Müller, 4, 123: 2043, 23 (185) Müller and Shearer, 4, 123: 3156; 23 (186) Niggli, 63, 19: 225, 18 (187) Niggli, 94, 56: 213, 22 (188) Niggli, 94, 57: 253; 22 (189) Nishikawa, 219, 8: 199; 15

(190) Nishikawa, 219, 9: 194; 17 (191) Nishikawa and Hudnuki, 219, 9 107; 17. (192) Ogg and Hopwood, 3, 32: 518, 16 (193) Ogg, 3, 42: 164. 21. (194) Ott, 63, 24: 209, 23 (195) Ott, 96, 22: 201, 24 (196) Owen and Preston, 67, 35: 101; 23. (197) Owen and Preston, 67, 36: 14; 23 (198) Owen and Preston, 67, 36: 49; 23 (199) Owen and Preston, 67, 36: 94; 23.

(200) Patterson, 45, 16: 689; 24 (201) Patterson, 2, 25: 581, 25 (202) Pauling, 1, 45: 2777; 23 (203) Pauling, 1, 46: 2738; 24. (204) Pauling and Dickinson, 1, 46: 1615; 24 (205) Pauling and Hendricks, 1, 47: 781, 25 (206) Phebus and Blake, 2, 25: 107; 25 (207) Phragmen, 220, 1923: 121 (208) Piper and Grindley, 67, 35: 269; 23 36: 31; 23. (209) Posnjak and Wyckoff, 128, 12: 248, 22.

(210) Rinne, 221, 69: 57, 17. (211) Rinne, Hentschel and Leonhardt, 94, 58: 629; 23. (212) Rinne, Leonhardt and Hentschel, 94, 59: 548; 24 (213) St John, 197, 4: 193; 18. (214) St John, 2, 21: 389, 23 (215) Scacchi, 222, 1: 187; 90. (216) Scacchi, 63, 19: 23, 18 (217) Scherrer, 94, 57: 186; 22. (218) Scherrer, B66: 387 (219) Scherrer and Stoll, 93, 121: 319, 22.

(220) Scherrer and Stoll, 149, 4: 232; 22. (221) Schiebold, 211, 36: 65; 19 (222) Schiebold, 94, 56: 430, 21. (223) Schneiderhohn, Mikroscopische Bestimmung von Erzen, p 196 (Berlin, 22) (224) Schumacher and Lucas, 1, 46: 1167; 24. (225) Shearer, 4, 123: 3152, 23. (226) Siegbahn and Dolejsek, 96, 10: 159; 22. (227) Simon and von Simson, 96, 25: 160, 24. (228) Simon and von Simson, 96, 21: 168, 24. (229) von Simson, 7, 109: 183; 24.

(230) Slattery, 2, 20: 84; 22. (231) Slattery, 2, 21: 213, 23 (232) Slattery, 2, 21: 378, 23. (233) de Smedt and Keesom, B60: 117, 24 (234) Sponsler, 225, 5: 757; 23. (235) Steinberg, 2, 21: 22, 23 (236) Stoll, 149, 3: 546, 21. (237) Terada, 219, 7: 292; 14 (238) van Arkel, 64V, 32: 197; 23 64P, 27: 97; 24 (239) van Arkel, 208, 4: 33 24

(240) Vegard, 5, 31: 83, 16 (241) Vegard, 5, 32: 65; 16 (242) Vegard, 5, 32: 505, 16. (243) Vegard, 5, 33: 395; 17. (244) Vegard, 96, 5: 17, 21 (245) Vegard, 96, 9: 395; 22 (246) Vegard, 96, 12: 289, 22 (247) Vegard, 96, 13: 379; 23. (248) Vegard, 8, 54: 146, 18. (249) Weber, 94, 57: 398; 22

(250) Westgren, 140, 103: 303; 21. (251) Westgren, 220, 105: 401, 21 (252) Westgren, 224, 1923: 223. (253) Westgren and Lindh, 7, 98: 181; 21 (254) Westgren and Phragmen, 140, 105: 241; 22 (255) Westgren and Phragmen, 7, 102: 1; 22. (256) Westgren and Phragmen, 220, 1923: 449 (257) Westgren and Phragmen, 140, 109: 159; 24. (258) Westgren and Phragmen, 58, 113: 122; 24. (259) Wever, 226, 3: 45, 21

(260) Wever, 226, 3: 17; 22. (261) Wever, 226, 4: 67; 22 (262) Wever, 226, 4: 81; 22. (263) Williams, 5, 93: 418; 17. (264) Wilsey, 3, 42: 262, 21. (265) Wilsey, 3, 46: 487; 23. (266) Wulff, 94, 57: 190; 22 (267) Wyckoff, O (268) Wyckoff, 1, 42: 1100; 20. (269) Wyckoff, 2, 16: 149, 20

(270) Wyckoff, 12, 50: 317; 20. (271) Wyckoff, 12, 1: 138; 21 (272) Wyckoff, 12, 2: 239; 21. (273) Wyckoff, 128, 11: 429; 21. (274) Wyckoff, 1, 44: 1239; 22. (275) Wyckoff, 1, 44: 1260; 22 (276) Wyckoff, 1, 44: 1994, 22 (277) Wyckoff, 12, 3: 184; 22. (278) Wyckoff, 12, 4: 188; 22 (279) Wyckoff, 12, 4: 193, 22.

(280) Wyckoff, 12, 4: 469; 22. (281) Wyckoff, 12, 5: 15; 23 (282) Wyckoff, 12, 5: 209; 23. 94, 57: 595; 23. (283) Wyckoff, 12, 6: 277; 23 (284)

Wyckoff, 94, 50: 55; 23 (285) Wyckoff, 128, 14: 121; 24. (286) Wyckoff, 12, 9: 145; 25 94, 61: 425; 25. (287) Wyckoff, 12, 10: 107; 25. (288) Wyckoff, 12, 9: 448, 25. (289) Wyckoff and Merwin, 12, 8: 447; 24.

(290) Wyckoff and Merwin, 12, 9: 286, 25 94, 61: 5; 25. (291) Wyckoff and Merwin, 12, 9: 379; 25 (292) Wyckoff and Posnjak, 1, 43: 2292; 21. (293) Wyckoff and Posnjak, 1, 44: 30; 22 (294) Wyckoff and Posnjak, 128, 13: 393, 23 (295) Yamada, 5, 45: 241, 23 (296) Yardley, 5, 105A: 451; 24 (297) Young, 5, 46: 291; 23 (298) Owen and Preston, 67, 36: 311, 24. (299) Levi and Tacchini, 56, 55: 28, 25.

(300) Levi, 59, 1: 335, 24 (301) Holgersson and Sedstrom, 8, 75: 143; 24. (302) Hewlett, 2, 20: 688, 22 (303) Debye and Scherrer, 188, 16: 16. (304) Keesom and DeSmedt, 64P, 25: 118, 22 (305) Keesom and DeSmedt, 64P, 26: 112, 23 (306) Becker and Ebert, 96, 31: 268; 25 (307) Havighurst, Mack and Blake, 1, 47: 29, 25 (308) Slattery, 2, 25: 333; 25. (309) Gibbs, 4, 125: 2022, 24

(310) Bernal, 5, 106A: 749, 24 (311) Hassel, 94, 61: 92; 25. (312) Wasastjerna, 138, 2, No 14 2, 25 (313) Schiebold, 94, 57: 579; 25. (314) Mark and Wigner, 7, 111: 398; 24 (315) Hoffman and Mark, 7, 111: 321; 24. (316) Hassel and Mark, 7, 111: 357, 24 (317) Keesom and DeSmedt, 64V, 33: 571; 24 (318) Mark and Pohland, 94, 61: 293; 25 (319) Saville and Shearer, 4, 127: 591; 25.

(320) Müller and Saville, 4, 127: 599; 25 (321) Friedel, 54, 180: 269; 25. (322) Wyckoff and Crittenden, 1, 47: 2876, 25. (323) Yardley, 269, 20: 296; 25 (324) Hevesy, 386, 2: 1, 25 (325) Bozorth and Pauling, 1, 47: 1561; 25. (326) Rinne, Hentschel and Schiebold, 94, 61: 164, 25 (327) Pauling and Emmett, 1, 47: 1026, 25 (328) Buckley and Vernon, 5, 49: 945; 25. (329) Davey 2, 26: 753, 25.

(330) Bradley, 3, 49: 1225, 25 (331) Wyckoff, 166, 62: 496; 25. (332) Gibbs, 5, 107A: 561, 25 (333) Zachariasen, Norsk geologisk tidsskrift, 8: 189: 25. (334) James and Wood, 315, 69: 24-25 (335) Wyckoff and Merwin, 94, 61: 152, 25 (336) Wyckoff, Bijvoet and Karssen, 64P, 27: 847; 24. (337) Kolkmeijer, Bijvoet and Karssen, 64V, 33: 327, 24 (338) Mark and Pohland, 94, 61: 532, 25 (339) Claassen, 70, 44: 790, 25.

(340) Kolderup, Bergens Museums Aarbok, 1924-25. (341) Mark and Tolksdorf, 96, 33: 681, 25 (342) Hassel, 156, 49: 465, 25 (343) Levi and Natta, 22, 2: 39, 25 (344) Havighurst, 12, 10: 15, 25 (345) van Arkel, 208, 5: 162; 25 (346) Peirce, Anderson, and Van Dyck, 145, 200: 363; 25. (347) Ott, 218, 13: 76, 25. (348) Ott, 94, 61: 515; 25 (349) Mark and Pohland, 94, 62: 103; 25

(350) Westgren and Phragmen, 96, 33: 77; 25 (351) Goldschmidt, Barth and Lunde, Skrifter Norske Videnskaps Akademi, No 7; 25. (352) Goldschmidt, Ulrich and Barth, Skrifter Norske Videnskaps Akademi, No 5, 25. (353) Brentano, 67, 37: 52, 25 (354) Trillat, 54, 180: 1320; 25. (355) Trillat 54, 180: 1838, 25 (356) Alsen, 207, 47: 19; 25 (357) Ramsdell, 328, 10: 281, 25. (358) de Smedt and Keesom, 64V, 33: 888; 24 (359) Bragg, 105, 9: 272, 25

(360) Clark, Asbury and Wick, 1, 47: 2661, 25 (361) Lange, 8, 76: 476; 25. (362) Blake, 2, 26: 60, 25 (363) Knaggs, 209, 30: 346; 25. (364) Aminoff, 94, 62: 113, 25 (365) Buckley and Vernon, 209, 30: 382; 25. (366) Olshausen, 94, 61: 463, 25 (367) Ferrari, 22, 1: 664; 25. (368) Bradley, 5, 50: 1018, 25 (369) Lunde, 7, 117: 51; 25

(370) Sacklowski, 8, 77: 211, 25 (371) Westgren and Phragmen, 5, 50: 311; 25 (372) Weiss, 5, 108A: 643, 25 (373) de Smedt, 186, 10: 366; 24. (374) Davey and Wilson, Proc Am Phys Soc, Nov. 27, 1925 (375) Nakamura, 210, 2: 287, 25 (376) Ösawa, 159, 14: 43, 25. (377) Wyckoff, 94, 62: 189; 25 (378) Broome, 94, 62: 325; 25. (379) Noethling and Tolksdorf, 94, 62: 255, 25

(380) Selyakov, Strutinskii and Krasnikov, 90, 23: 53; 25. (381) Ulrich and Zachariasen, 94, 62: 260; 25 (382) de Smedt and Keesom, 94, 62: 312; 25. (383) Ott, 94, 62: 201, 25 (384) Patterson, 2, 23: 552; 24. (385) Yardley, 5, 105A: 542; 25. (386) Ferrari, 22, 341: 186; 25. (387) Broome, 95, 142: 60; 25. (388) Sasahara, 520, 2: 277; 25 (389) Phragmen, 77, 45: 209; 25.

(390) Ott, 218, 13: 644; 25. (391) Jong, 208, 5: 194; 25. (392) Linck and Jong, 95, 147: 288; 25. (393) Goldschmidt, Barth and Lunde, Skrifter Norske Videnskaps. Akad. Oslo 1, Mat Nat. Kl: No. 7; 25. (394) Wyckoff and Crittenden, 1, 47: 2866; 25. (395) Huggins and Hendricks, 1, 48: 164; 26. (396) Hendricks and Pauling, 1, 47: 2904; 25. (397) Havighurst, 12, 10: 556, 25. (398) Yardley, 5, 50: 864; 25. (399) Yardley, 4, 127: 2207, 25

(400) Wyckoff, 12, 11: 101; 26 (401) Piper, Brown and Dyment, 4, 127: 2194; 20. (402) Wilsey, 143, 200: 739, 25.

SOME NUMERICAL DATA PERTAINING TO DISPERSOIDOLOGY

P. P. von Weimarn

From the large and heterogeneous mass of numerical data recorded in the literature of "Colloids," it seems desirable to present here only some selected illustrative examples of results of physical measurements which meet the following requirements: (1) The composition of the system is definite, reproducible, and exactly known; (2) all of the essential variables which affect the system are understood and are accurately controlled or measured; (3) the system, its behavior, and the resulting quantitative data are reproducible in the hands of any investigator working under *these same* controllable conditions; and (4) the examples selected shall be illustrative of some general law describing the behavior of dispersed systems.

As meeting the above conditions, the following examples have been selected and are presented in graphical form. Concise explanations are given in connection with the graphs. For a detailed description, explanation, discussion, and bibliography, the reader is referred to von Weimarn, Chem. Rev. **2**: 217; 25.

THE PRECIPITATION LAWS

Figures 1–9 illustrate the following precipitation laws: With increasing concentration of the reacting solutions, the average size of the precipitated crystalline individuals (*not their aggregates*) (1) passes through a maximum during, and (2) decreases continually after the completion of, the process of direct crystallization; (3) for the same absolute concentration of the reacting solutions (*other conditions being equal*), with decreasing solubility of a substance (Fig. 4; *cf.* Fig. 13), the average size of the precipitated crystals also decreases.

Figures 10–13 show that, if the aggregation of the individual ultramicrocrystals has not *proceeded too far*, the second law of precipitation remains valid; and besides they illustrate the law: (4) With increasing viscosity of the dispersion medium, the average size of the particles decreases (Fig. 12) [3, 4]; *cf.* [1].

The following general remarks apply to the figures: (1) The dispersion medium is indicated thus (60 vol. % C₂H₅OH); (2) mixing was brought about in all cases by pouring and shaking. The direction of pouring is indicated by the arrow. (3) In Figs. 1–9, the volumes mixed in each experiment satisfied the relation, concentration × volume = a constant (approx.), for a given dispersion medium; (4) the time, t_0, represents the period (ca. 10–15 min) required for the operations of sampling and photomicrographing; (5) all data shown are the averages of at least two independent experiments.

1. Precipitation of Ag₂SO₄.—*Reaction.*—2AgNO₃ + MnSO₄ = Ag₂SO₄ + Mn(NO₃)₂ (Figs. 1–7). In Figs. 4–5, per liter of final

solution, C = Ag₂SO₄ produced by the reaction and S = its solubility, both in g-equivalents [5].

2. Precipitation of AgC₂H₃O₂.—*Reaction.*—AgNO₃ + KC₂H₃O₂ = AgC₂H₃O₂ + KNO₃ (Figs. 8–9) [6]. These curves show the effect of time; the periods of time for the four curves are the same in both figures.

3. Precipitation of Se.—*Reaction.*—(a) 5 cc of aniline (an.) containing m mg of Se are poured into 100 cc of 93.5 wt. % C₂H₅OH (alc.) or (Fig. 13) mixtures thereof with an. or (Fig. 12) glycerol (gl.). t = 20° (Figs. 10–13 a curves) [7]. (b) As in (a) but with quinoline (q.) instead of aniline and using 90 wt. % C₂H₅OH (Figs. 10–13 b curves) [7].

4. Effects of Salts Dissolved in the Dispersion Medium on the Duration of Life of Dispersoidal Solutions.—(a) BaSO₄ *Reaction.*—50 cc (2a + 2x equiv.) BaR₂ + 50 cc (2a equiv.) MnSO₄ = 1 equiv. BaSO₄ + 1 equiv. MnR₂ + x equiv. BaR₂ Dispersion medium, 63 wt. % C₂H₅OH (Figs. 14–17) [8].

(b) *S.*—Dispersoidal solution of sulfur prepared by the method of grinding with grape-sugar. Ca. 25 mg S per liter of H₂O, particles ca. 85μμ (Figs. 18–23). C = millimols salt per liter. The dotted horizontal is for C = 0. To the right of the dotted vertical (Fig. 23) the disperse phase begins to dissolve by chemical action [10]; *cf.* [2].

(c) Al(OH)₃.—Prepared as in (b) *supra*. Ca. 55 mg Al₂O₃.3H₂O per liter of H₂O; particles ca. 90μμ (Fig. 24). The dotted horizontal is for C = 0. Dissolving begins at points marked with crosses [11]; *cf.* [2].

5. Adsorption and Solubility of Salts.—Adsorbent used—BaSO₄ extra pure; 20 g used per 100 cc of the salt solution. After shaking the solution with the adsorbent for 10 min, 24 hr. were allowed for the precipitate to settle. Fifty cc of the upper clear layer were used for analysis. Because partial dispergation occurred in the case of BaCl₂ in dilute C₂H₅OH solutions, these were centrifuged before analysis (Fig. 25) [9].

LITERATURE

(For a key to the periodicals see end of volume)

[1] Odén, *55*, **26**: 120; 20. [2] Ostwald, *55*, **26**: 28, 69; 20. [3] von Weimarn, *55*, **38**: 267, 624; 06. [4] von Weimarn, *55*, **38**: 933, 1400; 06. *55*, **2**: 76; 07. *287*, **18**: 44; 23. [5] von Weimarn, Aoki and Kataoka, O. *In part in* von Weimarn, *288*, **2**: 199; 24. [6] von Weimarn and Hori, O. [7] von Weimarn and Morishima, O. *In part in* von Weimarn *55*, **36**: 10; 25. [8] von Weimarn and Otsuka, O. *In part in* von Weimarn, *55*, **38**: 234; 23. [9] von Weimarn, Schochara and Takashige, O. *In part in* von Weimarn, *55*, **38**: 242, 23. [10] von Weimarn and Utsino, *55*, **36**: 265; 25. [11] von Weimarn and Utsino, O.

Fig. 1
Fig. 2
Fig. 3
Fig. 4
Fig. 5
Fig. 6
Fig. 7
Fig. 8
Fig. 9
Fig. 10
Fig. 12
Fig. 13

Fig. 11

Fig. 18

Fig. 14

Fig. 15

Fig. 19

Fig. 16

Fig. 17

Fig. 20

Fig. 21

Fig. 22

Fig. 23

Fig.24

Fig. 25

SWEETENING AGENTS. RELATIVE SWEETENING POWER

C. F. WALTON, JR.

The relative sweetness of various substances is usually cited in comparison with sucrose as unity. Since the concentration of the standard sucrose solution employed by different investigators has varied from 1 to 10%, and since the degree of sweetness does not decrease proportionately with dilution, the values reported in the literature vary accordingly, and are difficult to arrange accurately in numerical order. The following table, therefore, indicates only the approximate degree of sweetness, as reported by different investigators employing a variable procedure.

RELATIVE DEGREE OF SWEETNESS

(Sucrose = 1.0)

Name	Formula	Degree of sweetness	Lit.
Lactose	$C_{12}H_{22}O_{11}$	0.27–0.28	(26)
Dulcitol	$C_6H_{14}O_6$	0.41	(26)
Mannitol	$C_6H_{14}O_6$	0.45	(26)
Sorbitol	$C_6H_{14}O_6$	0.48	(26)
Glycerol	$C_3H_8O_3$	0.48	(26)
Glycol	$C_2H_6O_2$	0.49	(26)
Dextrose (d-glucose)	$C_6H_{12}O_6$	0.50–0.60	(10, 26, 29)
Maltose	$C_{12}H_{22}O_{11}$	0.60	(26, 29)

RELATIVE DEGREE OF SWEETNESS.—*(Continued)*

Name	Formula	Degree of sweetness	Lit.
Invert sugar (dextrose + levulose)	$C_6H_{12}O_6 + C_6H_{12}O_6$	0 78–0.95	(10, 26, 29)
Sucrose	$C_{12}H_{22}O_{11}$	1 00	(10, 26, 29)
Levulose (*d*-fructose)	$C_6H_{12}O_6$	1 03–1.50	(10, 26, 29)
p-Anisylurea	$CH_3OC_6H_4NHCONH_2$	18	(5)
Chloroform	$CHCl_3$	40	(31)
Glucin	Mixture	100	(11)
p-Methylsaccharin	$CH_3C_6H_3COSO_2NH$	200	(19)
Dulcin (*p*-phenetylurea)	$C_2H_5OC_6H_4NHCONH_2$	70–350	(11, 26)
6-Chlorosaccharin	$ClC_6H_3COSO_2NH$	100–350	(19)
n-Hexylchloromalonamid	n-$C_6H_{11}CCl(CONH_2)_2$	300	(11)
Saccharin (*o*-benzosulfonimid)	$C_6H_4COSO_2NH$	200–700	(11, 26)
Perillaldehyde α-anti-aldoxime (peryllartine)	$C_6H_5C(CH_2)CH_2CHNOH$	2000	(16)

LITERATURE

(For a key to the periodicals see end of volume)

The following list contains certain general references on methods of testing relative sweetening power, etc.

(1) Auerbach, *818*, *10*: 710; 22. (2) Barral and Rane, *871*, *56*: 712; 18 (3) Barral and Rane, *283*, *17*: 16; 20. (4) Becker and Herzog, *303*, *52*: 496, 07. (5) Boedecker and Rosenbuch, *273*, *30*: 251; 20. (6) Braun and Rawics, *25*, *49*: 799; 16. (7) Cohn, Die Organischen Geschmackstoffe (Siemanroth, Berlin, 1914). (8) Cohn, *196*, *22*: 1; 16. (9) Cohn, *274*, *55*: 735, 703; 14.

(10) Deerr, *275*, *24*: 481; 22. (11) Dox and Houston, *1*, *46*: 1278; 24. (12) Dyson, *276*, *11*: 572; 24. (13) Foerster, *282*, *25*: 400; 11. (14) Fränkel, Arzneimittelsynthese (Springer, Berlin, 1921): 134–53. (15) Furukawa, Japanese Patent 35332; 19. (16) Furukawa, *41*, *41*: 706, 979; 20 (17) Hermann, *13*, *429*: 163; 22. (18) Holleman, *70*, *40*: 446; 21. (19) Holleman, *70*, *43*: 839; 23.

(20) Holleman and Choufoer, *64V*, *33*: 307; 24. (21) Kionka and Sträts, *277*, *95*: 241; 22. (22) Kodama, *41*, *41*: 495; 20. (23) Lasarev, *278*, *194*: 293; 22. (24) Oertley and Myers, *1*, *41*: 855; 19. (25) Ogilvie, *275*, *24*: 288, 22. (26) Paul, *279*, *43*: 137; 22. (27) Paul, *280*, *26*: 610; 22. (28) Pauli, *205*, *125*: 97; 21. (29) Sale and Skinner, *45*, *14*: 522; 22.

(30) Speckam, *273*, *32*: 83; 22. (31) Sternberg, *281*, *36*: 272; 05. (32) Zuntz, *95*, *23*: 385; 10.

ODORIFEROUS MATERIALS

II. ZWAARDEMAKER

The unit used for expressing odor is the *olfacty*, the normal stimulus threshold for a given odor.

The characteristic grouping giving rise to odor is termed odoriphore (a), also called aromatophore (Klimout, 1897) and osmophore (Rupe, 1900). The principal odoriphores are: $-C(:O)O-$ Alkyl, esters; $-C(:O)H$, aldehydes; $-CO$, ketones; Alkyl O-Alkyl, ethers; $=C-OH$, alcohols; $-C(:O)OH$, acids; $-NO_2$, nitrites; $-CN$, nitriles; ⟨ ⟩$-$,terpenes; ⟨ ⟩, pinenes; $-S-S-$, sulfides; $-As-As-$, arsenides; $-As-O-As$, cacodyls; $-Hal.$, halogens; ⟨ ⟩N, pyridine; ⟩NH, pyrrole.

CLASSIFICATION

LINNÉ, MODIFIED BY ZWAARDEMAKER

Type	Key letter
Odores aetherei Lorry (Ethereal)	A
Odores aromatici Linné (Aromatic):	
1. Almond	B
2. Camphoric	C
3. Citric	D
Odores fragantes Linné (Balsam):	
1. Floral	E
2. Lilylike	F
3. Vanillin	G
Odores ambrosiae Linné (Musk)	H
Allyl	I
Cacodylic	J
Odores empyreumatic Haller (Empyreumatic)	K
Odores hircini Linné (Caprylic)	L
Odores tetri Linné (Narcotic)	M
Odores nauseois Linné (Nauseous)	N

Intensity.—The intensity of the odor of an odorivector (5) depends on (1) its volatility from dilute solution, (2) its rate of diffusion, (3) its absorption by a humid surface and (4) its solubility in liquids. (All odorous substances are soluble in oil (2).) The significance of an odor as a reflex stimulus depends on physiological, its pleasing or repulsive value on psychological conditions.

VOLATILITY OF ODOR FROM PARAFFINIC SOLUTIONS (4)

Substance	Concn. per cent	Volatility 10^{-6} g per min
Ethyl sulfide (I)	1	0.14
Scatole (N)	1	0.18
Valeric acid (L)	0.1	0.28
Guaiacol (K)	1	0.5
Pyridine (M)	10	0.93
Isoamyl acetate (A)	5	3.6
Terpineol (C)	25	7.5
Nitrobenzene (B)	50	9.2

DIFFUSION IN FREE AIR IN NEIGHBORHOOD OF SOURCE (10)

	cc per sec		cc per sec
Eugenol (C)	1.3	Ethyl ether (A)	4.4
Camphor (C)	2 1	Ethylacetone (A)	10

Extremes—ethyl acetate (A) and naphthalene (K). The anemodispersibility of odors depends on the size of the cloud and the velocity of the wind.

Spray Electricity.—All odorous substances lower the surface tension of water and therefore produce static electricity by spraying an aqueous solution of the odorivector against a disc well insulated with amber and paraffin. The value is expressed as 10^{-10} coulomb per cc of a saturated solution.

Substance	10^{-10} coulombs	Lit.
Cumidine (K)..	0.2	(12)
Aniline (K)	0.4	(6)
Toluidine (K) . .	0.4	(6)
Xylidine (K)	0.9	(6)
Scatole (N).	1 0	(12)
Trinitroisobutyltoluene (H) ..	1	(12)
Pseudocumene (K)....	3 4	(2)
Ethyl acetate (A). .	3 5	(2)
Xylene (K)	3 8	(6)
Aniline (K)	4 8	(2)
Toluene (K)	5 1	(2)
Thymol (C) . .	6 5	(2)
Benzene (K)	7 5	(2)
Toluidine (K)	7 9	(2)
Xylidine (K).	9 3	(2)
Nitrobenzene (B)	9 6	(2)
Vanillin (G).	10	(2)
Dimethylaniline (K) .	11 6	(6)
Benzaldehyde (B)	12 4	(2)
Anisaldehyde (G)	14 8	(2)
Phenol (K)..	15 2	(2)

Substance	10^{-10} coulombs	Lit.
Xylenol (K).	17	(2)
Ethyl alcohol (A)	17.2	(2)
Cresol (K)	19 1	(12)
Camphor (C)	20.3	(12)
Heliotropin (F)	44	(2)
Vanillin (G)	47	(12)
Heliotropin (F)	52	(12)
Acetone (A)	60	(12)
Guaiacol (K)	81.1	(2)
Carvacrol (C)	82.3	(2)
Terpineol (E)	89.1	(2)
Amyl acetate (A)	96.4	(2)
Ethyl acetate (A)	122	(12)
Guaiacol (K)	289	(12)
Terpineol (E)	296	(12)
Citral (D)	360	(12)
Methyl anthranilate (E)	602	(12)

RELATION BETWEEN SPRAY ELECTRICITY AND CONCENTRATION OF AQUEOUS SOLUTIONS (12)

	CHARGE IN 10^{-10} COULOMBS PER CC						
Degree of saturation	1	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{32}$	$\frac{1}{64}$
Coumarin	6 5	2	0 5	0			
Heliotropin	52	22	10	2	1.4	1.4	0
Vanillin	72	32	6	2	0.5	0	

ADSORPTION OF ODORS BY SURFACES EXPRESSED AS THE DURATION OF THE AFTER EFFECT FOLLOWING AN EXPOSURE TO A CONTINUOUS STREAM OF ODORIFEROUS AIR FOR 5 MINUTES (11). THE TERM sec DENOTES A FEW SECONDS, m = MINUTE, d = DAY, h = HOUR, min = SOME MINUTES

	Aluminium	Copper	Glass	Gold	Iron	Lead	Nickel	Porcelain	Silver	Steel	Tin	Zinc
Ethyl disulfide	1 m	sec	sec	sec	sec	1 m	sec	2 m	sec	sec	sec	sec
Guaiacol. .	15 m	3 m	1 m	12 m	8 m	sec	5 m	5 m	0	7 m	8 m	25 m
Ionone .	2 5 d	2 d	sec		4 d	1 d	2 d	sec	sec	4 d	min	
Isoamyl acetate	0	0	0	0	sec	0	sec	15 m	0	2 m	0	sec
Muscon	1 d	4 d	1 d	2 d	min	12 d	4 9 d	sec	2 d	sec	4 d	3 d
Nitrobenzene..	sec	sec	sec	sec	sec	sec	sec	8 m	sec	sec	sec	sec
Pyridine .	0	2 m	0	0	45 m	sec	sec	5 m	0	30 m	0.5 m	2.5 m
Scatole	9 d	3 d	1.5 h	1 5 d	10 d	10 d	3 5 d	0	1 d	20 d	7 d	14 d
Terpineol	0	sec	0	0	sec	0	0	5 m	sec	4 m	0	0
Valeric acid	3 m	0	30 m	sec	0	0	sec	0	5 m	0	2 m	0

Destruction of Odors by Ultraviolet Light.—The values are expressed as number of minutes required to reduce the odor in air from 2 to 1 olfacty by the radiation from a quartz mercury lamp (7).

Substance	Time	Substance	Time
Apiol (C)...	0 10	Methyl salicylate (C)	0 30
Valeric acid (L)..	0 10	Trimethylamine (J)	0 30
Menthol (C)..	0 15	Methyl nonyl ketone (C)	0 35
Ethyl sulfide (I)	0.25	Thymol (C)	0 40
Carvacrol (C)..... ...	0:25	Borneol (C)	0 45
Bornyl acetate (C).	0.30	Isoamyl acetate (A).	0 45
Caproic acid (L)	0 30	Pyridine (M)	0 45

Substance	Time	Substance	Time
Safrol (C) .	0 50	Methylheptenone (A)...	2.30
Salicylaldehyde (C) .	0 50	Eugenol (C)...	3
Scatole (N)... . .	0 50	Styrone (F)	3
Citral (D) . . .	0 55	Coumarin (G) . .	3.30
Indole (N)	1 0	Ethyl isovalerate (A) ..	4
Aniline (K) .	1 40	Cresol (K)....	5
Methyl anthranilate (E).	1 45	Ethyl butyrate (A)	5
Methyl butyrate (A)	2 0	Terpineol (E)	5·
Vanillin (G).	2 0	Chloroform (A).....	6
Citronellol (E)...... ..	2.30	Ethyl succinate. ..	6
Eucalyptol (C)...	2 30	Anethol (C) .	6.30
Isobutyl alcohol (K) .	2 30	Linalyl acetate (D) .	7

ODORIMETRY

The olfacty of an odor is the threshold or minimum perceptible concentration expressed in gms per cc which multiplied by $6.06 \times 10^{21}/M$, where M is the molecular weight, gives molecules per cc.

The authorities quoted are: Backman (1); Berthelot (2); Fischer and Peuzoldt (3); Henning (4); Hermanides (5); Huyer (6); Ohma (7); Passy (8); Tempelaar (9); van Wartenberg (10); Zwaardemaker (11).

Compound		Molecules per cc = A · 10^x		Authority
Name	Formula	A	x	
Ionone (F)	C₁₃H₂₀O	16 / 32	5 / 5	4 / 9
Ethyl bisulfide (I)	C₄H₁₀S	15	6	9
Scatole (N)	C₉H₉N	16 / 18	6 / 6	5 / 9
Vanillin (G)	C₈H₈O₃	20	6	8
Trinitroisobutyltoluene (H)	C₁₁H₁₃N₃O₆	21	6	9
Coumarin (G)	C₉H₆O₂	33	6	9
Citral (D)	C₁₀H₁₆O	40	6	8
Valeric acid (L)	C₅H₁₀O₂	47	6	4
Butyric acid (L)	C₄H₈O₂	69	6	8
Isoamyl alcohol (K)	C₅H₁₂O	69	6	8
Vanillin (G)	C₈H₈O₃	72	6	9
Valeric acid (D)	C₅H₁₀O₂	12	7	9
Heptylic acid (C)	C₇H₁₄O₂	16	7	8
Guaiacol (K)	C₇H₈O₂	18 / 20	7 / 7	5 / 9
Citral (D)	C₁₀H₁₆O	20	7	8
Methyl anthranilate (E)	C₈H₉NO₂	24	7	9
Nitrobenzene (B)	C₆H₅NO₂	32	7	4
Heliotropine (F)	C₈H₆O₃	40	7	4
Coumarin (G)	C₉H₆O₂	41	7	8
Iodoform	CHI₃	42	7	2
Bromoform	CHBr₃	48	7	8
Osmium tetroxide	OsO₄	48	7	10
Oenanthyl alcohol (C)	C₇H₁₆O	52	7	8
Valeric acid (D)	C₅H₁₀O₂	59	7	8
Cinnamaldehyde (C)	C₉H₈O	64	7	9
Nonylic acid (E)	C₉H₁₈O₂	77	7	8
Isobutyl alcohol	C₄H₁₀O	82	7	8
Thymol (C)	C₁₀H₁₄O	15	8	9
Capric acid (L)	C₁₀H₂₀O₂	18	8	8
Heliotropine (F)	C₈H₆O₃	20	8	8
Nitrobenzene (B)	C₆H₅NO₂	20 / 20	8 / 8	5 / 9
Borneol (C)	C₁₀H₁₈O	20	8	9
Coumarin (G)	C₉H₆O₂	21	8	8
Eucalyptol (C)	C₁₀H₁₈O	22	8	9
Citral (D)	C₁₀H₁₆O	25	8	9
Linalyl acetate (D)	C₁₂H₂₀O₂	29	8	9
Lauric acid (C)	C₁₂H₂₄O₂	30	8	8
Pyridine (M)	C₅H₅N	31	8	9
Pulegon (M)	C₁₀H₁₆O	33	8	9
Eucalyptol (C)	C₁₀H₁₈O	39	8	7
Heliotropine (F)	C₈H₆O₃	40	8	8
Carvacrol (C)	C₁₀H₁₄O	40	8	9
Propionic acid	C₃H₆O₂	41	8	8

Compound		Molecules per cc = A · 10^x		Authority
Name	Formula	A	x	
Durol (K)	C₁₀H₁₄	41	8	1
Isoamyl acetate (A)	C₇H₁₄O₂	42 / 42	8 / 8	5 / 9
Safrol (C)	C₁₀H₁₀O₂	48	8	7
Citral (D)	C₁₀H₁₆O	52	8	7
Anethol (C)	C₁₀H₁₂O	57	8	9
Methyl butyrate (A)	C₅H₁₀O₂	58	8	9
Terpineol (E)	C₁₀H₁₈O	79	8	9
Eugenol (C)	C₁₀H₁₂O₂	85	8	7
Pseudocumene (K)	C₉H₁₂	10	9	1
Bornyl acetate (C)	C₁₂H₂₀O₂	14	9	9
Methylheptenone (A)	C₈H₁₄O	15	9	9
Ethyl butyrate (A)	C₆H₁₂O₂	15	9	9
Methyl acetate (A)	C₃H₆O₂	16	9	11
Carvone (C)	C₁₀H₁₄O	22	9	9
Caproic acid (L)	C₆H₁₂O₂	27	9	8
Ethyl succinate (A)	C₈H₁₄O₄	28	9	9
Methyl salicylate (C)	C₈H₈O₃	39	9	9
Xylene (K)	C₈H₁₀	46	9	1
Cresol (K)	C₇H₈O	50	9	9
Methylnonyl ketone (C)	C₁₁H₂₂O	61	9	9
Ethyl ether (A)	C₄H₁₀O	61	9	4
Aniline (K)	C₆H₇N	63	9	9
Camphor (C)	C₁₀H₁₆O	64	9	8
Amyl alcohol (K)	C₅H₁₂O	69	9	8
Safrol (C)	C₁₀H₁₀O₂	75	9	9
Phenol (K)	C₆H₆O	77	9	4
Butyl alcohol (K)	C₄H₁₀O	82	9	8
Ethyl ether (A)	C₄H₁₀O	82	9	8
Fenchone (C)	C₁₀H₁₆O	92	9	9
Acetaldehyde (A)	C₂H₄O	96	9	9
Citronellol (E)	C₁₀H₂₀O	11	10	9
Valeric acid (L)	C₅H₁₀O₂	12	10	5
Toluene (K)	C₇H₈	13	10	1
Ethyl isovalerate (A)	C₇H₁₄O₂	21	10	9
Trimethylamine (J)	C₃H₉N	22	10	9
Phenol (K)	C₆H₆O	26	10	9
Benzene (K)	C₆H₆	41	10	1
Acetone (A)	C₃H₆O	42	10	11
Acetic acid (L)	C₂H₄O₂	50	10	8
Propyl alcohol (K)	C₃H₈O	51	10	8
Acetic acid (L)	C₂H₄O₂	71	10	9
Toluidine (K)	C₇H₉N	79	10	6
Xylidine (K)	C₈H₁₁N	10	11	6
Toluidine (K)	C₇H₉N	15 / 16	11 / 11	6 / 6
Menthol (C)	C₁₀H₂₀O	26	11	9
Aniline (K)	C₆H₇N	30	11	6
Formic acid	CH₂O₂	33	11	8
Terpineol (E)	C₁₀H₁₈O	73	11	5
Pyridine (M)	C₅H₅N	12	12	5
Ethyl alcohol (A)	C₂H₆O	24 / 33	12 / 12	4 / 4
Formic acid	CH₂O₂	84	12	9
Methyl alcohol	CH₄O	11	13	9
Methyl alcohol	CH₄O	19	13	8
Apiol (C)	C₁₂H₁₄O₄	17	15	9

VALUE OF AN OLFACTY EXPRESSED AS DEGREE OF SATURATION OF AIR WITH THE ODORIVECTOR

Substance	% Saturation	Substance	% Saturation
Eucalyptol	0.058	Methyl alcohol	1.388
Eugenol	0.144	Toluidine	1.515
Toluene	0.158	Ethyl alcohol	2 5
Benzene	0 169		

VALUE OF AN OLFACTY IN CM OF THE ZWAARDEMAKER OLFACTOMETER

The constants of Zwaardemaker olfactometer are: width of cylinder, 0.8 cm; length, 10 cm; contents, 50 cc; air contact per cc of cylinder, 2.5 cm²; velocity of air in the air tube, 100 cc per sec (exposure, 0.33 sec).

MINIMUM PERCEPTIBLE IN CM OF OLFACTOMETER SCALE
Saturated solutions (9)

Substance	cm	Substance	cm
Terpineol—H₂O	0 01	Caproic acid—H₂O	0.10
Ethyl propionate—H₂O	0.02	Trinitroisobutyltoluene -	
Ionone—H₂O	0 02	H₂O	0 10
Camphor—H₂O	0 07	Guaiacol—H₂O	0 20
		Trimethylamine—Paraffin	0 20

Aqueous solutions (10)

Substance	Concentration Wt. %	cm
Pyridine	0 05	0 1
Ethyl disulfide	0 02	0 5
Citral	0 01	0 2

Aqueous solutions (10).—(Continued)

Substance	Concentration Wt. %	cm
Scatole	0.01	0.4
Valeric acid	0.01	0.5
Isoamyl acetate	0.01	0.7
Guaiacol	0 0007	1 0

Paraffin solutions (11)

Substance	Concentration Wt. %	cm	Substance	Concentration Wt. %	cm
Borneol	1 0	0 001	Citral	1.0	0.09
Cadaverine	0 1	0 001	Isoamyl acetate	0.5	0.29
Scatole	0 1	0 002	Guaiacol	0.1	0.62
Ethyl sulfide	0 01	0 01	Ionone	0.0004	0.62
Pyridine	1 0	0 03	Safrol	3.0	1.12
Valeric acid	0 01	0 04	Terpineol	2.5	1.60
Nitrobenzene	5 0	0 06			

LITERATURE

(For a key to the periodicals see end of volume)

(1) Allison and Katz, 45, 11: 336; 19   (2) Backman, 278, 168: 351; 17.   (3) Henning, Der Geruch, 1924   (4) Hermanides, Thesis, Utrecht, 1907.   (5) Heynnx, Thesis, Brussels, 1919.   (6) Huyer, Thesis, Utrecht, 1917.   (7) Tempelaar, Thesis, Utrecht, 1913   (8) Zwaardemaker, Physiol. des Geruchs, 1895: 247   (9) Zwaardemaker, Arch. Anat. Physiol., Physiol. Abt., 1908: 48.
(10) Zwaardemaker, In Abderhalden, Handb biol. Arbeitsm. 5, pt. 7: 455; 23.   (11) Zwaardemaker, In Tigerstedt, Handb Physiol. 2, pt. 1: 49; 14.   (12) Zwaardemaker, Arch. Neerl. Physiol., 1 : 347; 17.

RADIOACTIVITY

S. C. LIND, SPECIAL EDITOR

1923 INTERNATIONAL TABLE
RADIOACTIVE ELEMENTS AND THEIR CONSTANTS

λ $(sec)^{-1}$ is the *radioactive constant of transformation.*

$$dQ = -\lambda Q dt, \qquad Q = Q_0 e^{-\lambda t}, \qquad \log_{10}\frac{Q_0}{Q} = 0.4343\lambda t,$$

in which Q_0 is the initial quantity and Q the quantity remaining after a time t (seconds).

$\lambda = -\frac{dQ}{Q}\frac{1}{dt}$ represents the fraction of the element transformed, reduced to the unit of time.

In the case of a double transformation, the values between brackets [ ] refer to the constants corresponding with the separate branches; the constant for both branches not being put between brackets.

The sign (?) indicates that the value has been indirectly deduced from the range of the α-rays expelled.

$\theta = \frac{1}{\lambda}$ is the *average life* of the radioactive atoms.

T is *the half period*, *i.e.*, the time in which the quantity of radio-element is diminished to one half:

$$\lambda T = -\log_e 0.5 = 0.69315 \text{ and } \theta = 1.443T$$

Radiation.—The brackets ( ) indicate that the radiation is relatively feeble.

REMARKS CONCERNING THE NOMENCLATURE

It is desirable that the nomenclature adopted by the international commission should be accepted universally but that now put forward for the present year is provisional, to serve as a basis of discussion with the view to the adoption ultimately of a standard nomenclature.

The most important points are:

1. The three radioactive emanations have been given the names **radon, actinon,** and **thoron,** with the symbols Rn, An, Tn, to **suggest** both their origin and their chemical character as members of the family of the rare gases of which the valency is zero;

2. In the branches which occur at the C members the sign (') **has** been used to indicate the products resulting from the emission of β-rays (isotopes of polonium) and the sign (") to indicate the products resulting from the emission of α-rays (isotopes of thallium);

3. The ultimate products have been indicated by the letter Ω.

EXPLANATION OF THE NOTES

NOTE 1.—*Uranium I.*—The value given for θ is that obtained from the equation:

$$\theta = \frac{1}{\lambda} = 2440 \times 0.97 \times 3 \times 10^6 \times \frac{226}{238} = 6.75 \times 10^9$$

in which the number 2440 represents the average life of radium in years, the number 0.97 the branching coefficient and $3 \times 10^6 \times \frac{226}{238}$ is the ratio between the numbers of atoms of uranium and radium in equilibrium in minerals.

If the actinium series is independent from that of uranium I, λ cannot be calculated by this method.

The value of λ obtained by the direct counting of the α-particles from a compound of uranium is 4.57×10^{-18} from which $\theta = 7 \times 10^9$ years and $T = 4.8 \times 10^9$ years.

NOTE 2.—*Uranium X_2* is also called brevium.

NOTE 3.—*Radon* replaces the names *radium emanation* and *niton* (the latter of which was proposed by Sir William Ramsay).

NOTE 4.—*Radium C* undergoes a double disintegration: 99.97% of the atoms emit β-rays and produce the substance Ra-C' which gives α-rays, and 0.03% of the atoms emit α-rays and produce the substance Ra-C'' which gives β-rays.

a_0 is the *range* in cm of the α-rays in air at 0°C and a pressure of 760 mm of mercury.

The range at $r°$ C. and under p mm of mercury is

$$a = \frac{a_0(273 + r)760}{273p}$$

V is the velocity of α or β-rays relatively to that of light. To convert to cm per sec multiply by 3×10^{10}.

For the α-rays:

$$V = 0.0342\, a^{\frac{1}{2}}$$

$\mu_{\beta Al}$ is the absorption coefficient of the β-rays in aluminium, the thickness being measured in cm.

$\mu_{\gamma Al}$ and $\mu_{\gamma Pb}$ are the absorption coefficients of the γ-rays in aluminium and lead respectively, the thickness being measured in cm; the latter is only given for the most penetrating type of γ-rays.

If I_0 is the initial intensity and I the intensity after the rays have traversed x cm of the absorbent:

$$I = I_0 e^{-\mu x} \qquad \log_{10}\frac{I_0}{I} = 0.4343\mu x$$

If D is the thickness corresponding with the absorption of one-half of the rays:

$$\mu D = 0.693$$

NOTE 5.—*Radium D* is also called radiolead.

NOTE 6.—*Radium C'''* is also called radium C_2.

NOTE 7.—*Uranium Y* is the first known member of the actinium series. It may be derived from Uranium I or Uranium II. In this case, 3% of the atoms of Uranium produce the actinium family, and 97% the radium family.

The hypothesis has also been put forward that the actinium series may be produced independently from a third (hypothetical) isotope of Uranium for which the name actinouranium has been proposed.

NOTE 8.—*Protoactinium* is also called eka-tantalum.

NOTE 9.—A new radioactive substance named uranium Z, and isotopic with protoactinium, accompanies uranium in minute quantity. (*25,* **54B**: 1131; 21). Its period is from 6 to 7 hours. It emits a β-radiation for which D_{Al} varies from: 0.0014 to 0.012. Its parent is an isotope of thorium, but it cannot yet be placed in the series.

NOTE 10.—*Actinon* is also called actinium emanation.

NOTE 11.—*Actinium C.* 0.2% of the α-rays emitted by this substance have a range $a_0 = 6.10$, instead of 5.12. From this it has been concluded that 0.2% of the atoms undergo a transformation by the emission of β-rays as is the case in the radium C and thorium C branches (*3,* **27**: 690; 14. **28**: 818; 14). Confirmatory evidence appears to be desirable.

NOTE 12.—*Actinium C''* is also called actinium D.

NOTE 13.—*Thorium.* The value given for λ is that obtained from the direct counting of the α-particles emitted by a compound of thorium. All the other values are less; the smallest being 0.55 of that given in the table and giving $\theta = 3.45 \times 10^{10}$ years and $T = 2.37 \times 10^{10}$ years (*63,* **19**: 259; 18).

NOTE 14.—*Thoron* is also called thorium emanation.

NOTE 15.—*Thorium C* undergoes a double disintegration: 65% of the atoms emit β-rays and produce the substance Th-C' which gives α-rays, and 35% emit α-rays and produce the substance Th-C'' which gives β-rays.

NOTE 16.—*Thorium C.* The value $a_0 = 4.69$ is that corresponding with $V = 0.0572$ which has been directly measured.

NOTE 17.—*Thorium C''* is also called thorium D.

NOTE 18.—*Potassium* and *rubidium* emit β-rays but show no other evidence of radioactivity.

T	$\theta = \frac{1}{\lambda}$	λ (sec)⁻¹	Name	Symbol	Atomic Wt.	Atomic No.	Iso-tope	Radiation	a_s	V	μ β Al	μ γ Al	μ γ Pb	Notes
SERIES OF URANIUM AND RADIUM															
4 67 × 10⁹ yrs	6 75 × 10⁹ yrs	4.7 × 10⁻¹⁸	Uranium I	U₁	238	92	U	α	2 37	0 0456				1
24 6 days	35.5 days	3 26 × 10⁻⁷	Uranium X₁	U-X₁	234	90	Th	β			463			
1 15 min	1.65 min	0.010	Uranium X₂	U-X₂	234	91	Pa	β (γ)			14 4	24; 0 7, 0 14	0 72	2
2 × 10⁶ yrs	3 × 10⁶ yrs	10⁻¹⁴ (?)	Uranium II	U₁₁	234	92	U	α	2 75	0 0479				
6 9 × 10⁴ yrs	10⁵ yrs	3.2 × 10⁻¹³	Ionium	Io	230	90	Th	α	2 85	0 0485				
1690 yrs	2440 yrs	1 30 × 10⁻¹¹	Radium	Ra	226	88	Ra	α (β + γ)	3 13	α 0 0500, β 0 52, 0 65	312	354, 16; 0 27		3
3 85 days	5 55 days	2 085 × 10⁻⁶	Radon	Rn	222	86	Rn	α	3 94	0 0540				
3 0 min	4.32 min	3 85 × 10⁻³	Radium A	Ra-A	218	84	Po	α	4 50	0 0565				
26 8 min	38.7 min	4.30 × 10⁻⁴	Radium B	Ra-B	214	82	Pb	β (γ)		0 36, 0 41; 0 63, 0 70; 0 74	13 1, 80	230, 40, 0 51		
19 5 min	28 1 min	5 92 × 10⁻⁴	Radium C	Ra-C	214	83	Bi	99 97%; β and γ		0 786, 0 862, 0 949, 0 957	13 2; 53	0 115	0 50	4
10⁻⁶ sec	10⁻⁶ sec	10⁶ (?)	Radium C'	Ra-C'	214	84	Po	α	6 57	0 0641				
16 5 min	23 8 yrs	1 33 × 10⁻⁹	Radium D	Ra-D	210	82	Pb	(β and γ)		0 33, 0 39	5500	45; 0 99		5
5 0 days	7 2 days	1 61 × 10⁻⁶	Radium E	Ra-E	210	83	Bi	β			43 3			
136 days	196 days	5 90 × 10⁻⁸	Radium F (Polonium)	Ra-F (Po)	210	84	Po	α (γ)	3 58	0 0523		585		
			Radium Ω (Lead)	Ra Ω' Pb²⁰⁶	206	82	Pb							
		[1 8 × 10⁻⁷]	Radium C	Ra-C	214	83	Bi	0 03% α	?					6
1 4 min	2 0 min	8 3 × 10⁻³	Radium C''	Ra-C''	210	81	Tl	β						
			Radium Ω'' (hypothetical)	Ra Ω''	210	82	Pb							
SERIES OF ACTINIUM															
			Uranium ?		?	92	U	α						7
1 04 days	1 5 days	7 8 × 10⁻⁶	Uranium Y	U-Y	?	90	Th	β			About 300			
1 2 × 10⁴ yrs	1 7 × 10⁴ yrs	1 9 × 10⁻¹²	Protoactinium	Pa	?	91	Pa	α	3 314	0 0510				8, 9
20 yrs	28 8 yrs	1 1 × 10⁻⁹	Actinium	Ac	?	89	Ac							
19 5 days	28.1 days	4 11 × 10⁻⁷	Radioactinium	Rd-Ac	?	90	Th	α (β)	4 36	α 0 0559; β 0 38, 0 43, 0 49, 0 53, 0 60, 0.67; 0 73	About 170	25, 0 19		
11 4 days	16 4 days	7 06 × 10⁻⁷	Actinium X	Ac-X	?	88	Ra	α	4 17	0 0550				
3 9 sec	5 6 sec	0 178	Actinon	An	?	86	Rn	α	5 40	0 0600				10
2 0 × 10⁻³ sec	2 9 × 10⁻³ sec	345	Actinium A	Ac-A	?	84	Po	α	6 16	0 0627				
36 1 min	,52 1 min	3 2 × 10⁻⁴	Actinium B	Ac-B	?	82	Pb	(β and γ)			Very large	120; 31; 0.45		11
2 15 min	3 10 min	5 37 × 10⁻³	Actinium C	Ac C	?	83	Bi	α	5 12	0 0589	28 5	0.198	1.2 to 1 8	12
4 71 min	6 83 min	2 44 × 10⁻³	Actinium C''	Ac-C''	?	81	Tl	β and γ						
			Actinium Ω'' (hypothetical)	Ac Ω''	?	82	Pb							
SERIES OF THORIUM															
1 31 × 10¹⁰ yrs	1.89 × 10¹⁰ yrs	1 68 × 10⁻¹⁸	Thorium	Th	232	90	Th	α	2 58	0 0469				13
6.7 yrs	9 67 yrs	3 28 × 10⁻⁹	Mesothorium 1	Ms-Th1	228	88	Ra							
6 2 hrs	8 9 hrs	3 12 × 10⁻⁵	Mesothorium 2	Ms-Th2	228	89	Ac	β and γ		0 37, 0 39, 0 43; 0 50; 0 57; 0 60; 0 66 and > 0 70	20 2 to 38 5	26; 0 116	0 62	
2 02 yrs	2 91 yrs	1 09 × 10⁻⁸	Radiothorium	Rd-Th	228	90	Th	α (β)	3 67	α 0 0527; β 0 47; 0 51				
3 64 days	5 25 days	2 20 × 10⁻⁶	Thorium X	Th-X	224	88	Ra	α	4 08	0.0546				14
54 sec	78 sec	0 0128	Thoron	Tn	220	86	Rn	α	4 74	0.574				
0 14 sec	0 20 sec	5.0	Thorium A	Th-A	216	84	Po	α	5 40	0 0600				
10 6 hrs	15 3 hrs	1 82 × 10⁻⁵	Thorium B	Th-B	212	82	Pb	β and γ		0 63; 0 72	110	160; 32; 0 36		15
60 min	87 min	1 92 × 10⁻⁴	Thorium C	Th-C	212	83	Bi	65% β		(C + C'') 0 29, 0 36; 0 93 to 0 95	14 4			
10⁻¹¹ sec	10⁻¹¹ sec	1.25 × 10 ⁴	Thorium C'	Th-C'	212	84	Po	α	8 16	0 0688				
		10¹¹ (?)	Thorium Ω' (Lead)	Th Ω' Pb²⁰⁸	208	82	Pb							
		[6.7 × 10 ⁴]	Thorium C	Th-C	212	83	Bi	35% α	{ 4 55 ¾4 69	} 0 0572				16
3 1 min	4 5 min	3 70 × 10 ³	Thorium C''	Th-C''	208	81	Tl	β and γ		(See Th-C)	21 6	0 096	0 46	17
			Thorium Ω'' (Lead)	Th Ω'' Pb²⁰⁸	208	82	Pb							
			Potassium	K	39 1	19	K	β			22 to 38			} 18
			Rubidium	Rb	85 5	37	Rb	β			308 to 347			

PHYSICAL PROPERTIES OF THE RADIOELEMENTS AND THEIR COMPOUNDS
(Except Ra, Th, U and Rn)

GEORG HEVESY

1. Atomic Weights.—Io (mixture of Io + Th), 231.51 [2]. RaΩ (=U-Pb), 206.04 [2]. ThΩ (=Th-Pb), 207.97.

2. Molecular Weights.—An (=Ac-Em), 220–232 [4]. Tn (=Th-Em), 201–210 [4]. Rate of effusion method.

3. Density [5].—RaΩ, 11.273 g cm^{-3} at 19.94°C.

4. Melting Point [26].—RaΩ', differs from Pb <0.05°.

5. Boiling Point [32]. —Ra-FH$_2$, 37°C.

6. Solubility.—S = solubility mol l^{-1}. $\alpha' = \dfrac{C_{Air}}{C_{H_2O}}$. An [14], $\alpha' = 2$ at 18°. Tn [15], $\alpha' = 1$ at 18°. Rn [16]. $S = 1.7989$ [15b] in H$_2$O at 25°. $S[$RaΩ'$(NO_3)_2] - S[$Pb$(NO_3)_2] < 10^{-4}$.

RELATIVE SOLUBILITY OF AN IN DIFFERENT SOLVENTS AT 18°

H$_2$O	Sat. KCl soln.	Conc. H$_2$SO$_4$	C$_2$H$_5$OH	C$_3$H$_7$OH	C$_2$H$_5$CHO	C$_6$H$_6$	C$_6$H$_5$CH$_3$	Kerosene	CS$_2$
1	0.9	0 95	1 11	1 6	1 7	1 7	1 8	1.9	2 1

7. Rate of Solution.

PERCENT DISSOLVED FROM SURFACE AT 18°

By H$_2$SO$_4$ in 15 sec [17]

H$_2$SO$_4$, equiv. per liter =	10^{-3}	10^{-2}	10^{-1}	1
Ra-B from glass	80	80	97	88
Ra-C from glass	28	60	88	99

By HNO$_3$ in 60 sec [18]

HNO$_3$, equiv. per liter =	0	10^{-5}	10^{-4}	10^{-3}	10^{-2}	10^{-1}	1
Th-B from quartz	60	61	60	80	81	83	84
Th-C from quartz	37	38	35	61	72	77	87

PERCENT RA-B AND RA-C DISSOLVED FROM GLASS SURFACE [17]

By H$_2$O in 5 min

t	Ra-B	Ra-C	t	Ra-B	Ra-C
0°	0 29	0 19	42°	0 78	0 67
17°	0 47	0 35	70°	0 97	0 91

By H$_2$SO$_4$ in 15 sec

t	Ra-B	Ra-C	t	Ra-B	Ra-C
0°	0 74	0 52	42°	0 895	0 71
17°	0 80	0 60	70°	0 96	0 81

8. Adsorption.—Ratio of molal conc. in gas at equilibrium to moles adsorbed per liter of charcoal at 18°, An [19] 0.05, Tn [20] 0.02. Percent of initial amount present (per 50 cc of solution) adsorbed by 1 g of adsorbent [21]. (a) By BaSO$_4$, from 0.1 N HCl, Th-B 81, Th-C 32; from 0.1 N KOH, Th-B 20, Th-C 64; from 0.1 N NH$_3$, Th-B 100, Th-C 86. (b) By Cr$_2$O$_3$, from 0.1 N HCl, Th-B 2.5, Th-C 69. (c) By AgBr, from 0.1 N HBr, Th-B 81, Th-C 34. (d) By BaSO$_4$, from 1 N HCl, Ra 80. (e) By Cr$_2$O$_3$, from 1 N HCl, Ra 0. (f) By AgCl, from 1 N HCl, Ra 0.

9. Vapor Pressure.—$p_{700°}$ for RaΩ' is 2% greater than for Pb [22].

10. Temperature of Volatilization.—Depends on nature of surface and chemical state of the radioactive element. v. [23, 24, 25].

11. Coefficient of Diffusion.

(a) In Gases at 76 cm and 15°

An, in	Air	H$_2$	CO$_2$	SO$_2$	A
Δ, cm^2 sec^{-1}. . .	0 098–0.123 [6, 7, 8, 9]	0 330 [7]	0 412 [8]	0 075 [7, 8]	0.062 [7]	0.107 [7]
Tn, in		Air		A	
Δ, cm^2 sec^{-1}. . .		0.085–0.103 [6, 7, 9]		0 084 [7]	

(b) The Cations in Water [10] at 18°

Ion	UX$_1^{++}$	Io^{++}	Ra-D^{++}	Ra-E^{++}	Ra-F^{++}	Ac^{+++}
Δ, cm^{-2} day^{-1}	0 4	0 33	0 65	0.45	0.76	0 46

Ion	AcX^{++}	Rd-Th^{++}	ThX^{++}	Th-B^{++}	Th-C^{+++}
Δ, cm^{-2} day^{-1}.	0 69	0 33	0 66	0.67	0 5

Th-CCl$_3$ in $\frac{1}{12}$ N NH$_3$, Δ = 0.37. Ra-FCl$_2$ in $\frac{1}{2}$ N NH$_3$, Δ = 0.19.

(c) In Metals. Δ in cm^{-2} Day^{-1}

	t	Δ
Th-B in Pb	343°	2.2 [11]
Ra-D in Pb	280°	<10^{-4} [12]
Ra-F in Pb	280°	<10^{-4} [12]
Ra-F in Au	470°	ca. 10^{-9} [13]
Ra-B + Ra-C in Ag	470°	3 8 × 10^{-7} [13]
Ra-B in Au	470°	8 2 × 10^{-7}
Ra-B in Pt	470°	3 4 × 10^{-7}

In re diffusion of Th-B in single crystals, in lead foils and in thallium foils v. [35].

12. Refractive Index [27].—$n_D^{25°}$ for cryst. RaΩ'(NO$_3$)$_2$ = 1.7814.

13. X-ray Spectra.—All lines of the L series and the Mα and Mβ lines of RaΩ' differ by less than 5 × 10^{-12} cm from the same lines for Pb [28].

14. Relative Ionic Mobilities [10].—In capillary tubes by comparison against Ra (Λ = 57.3 mhos).

Cation	Ra	Ra-C	Ra-D	Ra-E	Ra-F	AcX	ThX	Th-B	Th-C
Λ	57 3	54.5	61.9	61.9	68.8	56.1	58.0	55.4	54 0

15. Emf.—RaΩ' / N RaΩ'(NO$_3$)$_2$ // N Pb(NO$_3$)$_2$ / Pb. <0.1 millivolt [31].

16. Deposition Voltage.—From $\frac{1}{10}$ N HNO$_3$ containing 10^{-8} mole Ra-F, cathodic deposition occurs on Au electrodes at E_{Hg} = 0.35 volt, anodic at E_{Hg} = 1.05 volt [30].

LITERATURE AND REMARKS

(For the key to periodicals see end of volume)

[1] Hönigschmid, *9*, **22**: 21; 16. This mixture contained about 30 % Io and 70 % Th and was probably contaminated with some Th not present in the pure pitchblende (cf. Soddy and Hitchins, *3*, **47**: 1148; 24. Meyer and Ulrich, *76*, **182**: 279; 23). [2] Lowest value found. Higher values probably due to presence of lead. Richards and Lembert, *1*, **36**: 1329; 14. *95*, **88**: 429; 14. Hönigschmid and Horowitz, *75*, **183**: 2407; 14. *9*, **20**: 319; 14. Curie, *54*, **148**: 1676; 14. *198*, **34**: 586; 23. Richards, *Ann. Rep. Smithsonian Inst.* **1918**: 205. Richards and Putseys, *1*, **46**: 2954; 23. [3] Highest value found. Lower values probably due to presence of lead and RaΩ. Hönigschmid, *9*, **25**: 91; 19. Soddy, *4*, **105**: 1402; 14. *88*, **94**: 615; 15. **98**: 469; 17. **99**: 244; 17. [4] Leslie, *4*, **24**: 637; 12. *54*,

152: 328; 11. Marsden and Wood, 4, **29**: 945; 13. (⁸) Richards and Wadsworth, *1*, **38**: 221, 1658; 16. *Cf.* Soddy, *58*, **107**: 41; 21 Egerton and Lee, *5*, **103**: 487; 23. (⁶) Rutherford, "*Radioactivity*," Cambridge, 1913, p 387. (⁷) Russ, *4*, **17**: 540; 09. (⁸) B Bruhat, *199*, **6**: 67, 09 (⁹ Debierne, *199*, **4**: 213; 07. McLennan, *2*, **30**: 660; 10. Eckmann, *200*, **9**: 177; 12. Thomsen, *201*, **18**: 377; 09. Hevesy, *200*, **10**: 198; 13. (⁹) Leslie, *34*, **153**: 328; 11. Rutherford, *l c.*

(¹⁰) Hevesy, *63*, **14**: 49, 1202; 13. *4*, **26**: 586; 14. Paneth, *75*, **122**: 1636, 13. The radioelements probably present in colloidal state. (¹¹) Gröh and Hevesy, *8*, **63**: 85; 20. Diffusion rate of a mixture of Th-B and Pb in lead. Th-B used as indicator. (¹²) Gröh and Hevesy, *8*, **65**: 216, 21. Diffusion rate of a mixture of Ra-D and Pb in lead (¹³) Wertenstein and Dobrowolska, *51*, **4**: 324; 23. Diffusion rate of active deposit (probably of oxides). (¹⁴) Hevesy, *63*, **12**: 1214; 11. *50*, **16**: 429, 12 (¹⁵) Klaus, *63*, **6**: 820; 05. Boyle, *Macdonald Phys Build. Bull*, No **1**: 52, 10. α of short-lived An and Tn determined by making assumptions only partly justified. α of An and Tn probably practically identical with that of Rn. (¹⁶) Richards and Schumb, *1*, **40**: 1403; 18. The RaU' used contained some common lead, its atomic weight being 206 31 The solubility of common lead (at. wt. 207.19) was found by the same authors to be 1 7993 *Cf.* Fajans and Lembert, *93*, **95**: 297; 16. (¹⁷) Ramstedt, *147*, II· No. **31**; 13. *Cf.* Arrhenius, *199*, **7**: 228, 10 Godlewski, *199*, **10**: 250; 13. Schräder, *4*, **24**: 131; 12. Hevesy, *9*, **19**: 291; 13 (¹⁸) Hevesy and Rona, *7*, **89**: 294; 15. *In re* Ra-F, *cf* Paneth and Hevesy, *75*, **123**: 1050; 13. (¹⁹) Hevesy, *63*, **12**: 9; 12. *50*, **18**: 429; 12

(²⁰) Boyle, *4*, **17**: 389; 09. Ra-B and Th-B between Pb amalgam and Hg(NO₃)₂; *cf.* Z. Klemensievicz, *34*, **156**: 1889; 14. (²¹) Paneth, *63*, **18**: 924,

14 Horowitz and Paneth, *75*, **129**: 1819. 14 *In re* adsorption UX *cf.* Ebler and Rhyn, *25*: **84**: 2896, 21 A C Brown, *4*, **121**: 1738; 22. Freundlich and Wreschner, *7*, **106**: 366; 23. Adsorption of Ra-B, Ra-C, Th-B and Th-C Hevesy, *75*, **127**: 1787, 18 Cranston and Burnett, *4*, **119**: 2036; 21 **121**: 2880, 22 Paneth and Vorwerk, *?*, **101**: 445; 22. Fajans and Frankenberg, *7*, **106**: 255, 23 Absorption of Ra-F, Paneth, *55*, **13**: 1, 288; 13 Lachs and Werthenstein, *63*, **23**: 318; 22 Escher, *34*, **177**: 3, 172; 23 (²²) Egerton, *5*, **103**: 469; 23 (²³) Russell, *4*, **24**: 134; 12. *cf.* Schräder, *4*, **24**: 125, 12. (²⁴) St Lona, *63*, **17**: 6, 16 (²⁵) Wood, *5*, **91**: 543, 15 *Cf* Barrat and Wood, *67*, **26**: 218; 14 Wood, *4*, **28**: 808; 14. *In re* volatilization of Tn *cf* Fleck, *4*, **29**: 337; 15 and St Lona, *75*, **129**: 829; 15 Volatilization of RaFH₁ and of the hydrides of Ra-B, Th-B and Th-C, Paneth, *25*, **81**: 1704, 18 **83**: 1693, 20. *9*, **26**: 452; 20. (²⁶) Richards and Hall, *1*, **43**: 1550, 20 *cf* Lembert, *9*, **26**: 59, 20. (²⁷) Richards and Schumb, *1*, **40**: 1403; 18 For Pb(NO₃)₂, *n⁰₅* = 1.7815. (²⁸) Siegbahn and Stenstrom, *63*, **18**: 517, 17. *Cf* Duane and Shimizu, *197*, **5**: 198; 19 Cooksey and Cooksey, *2*, **16**: 327; 20. *In re* slight difference in the wave length of optical spectrum of ordinary Pb and mixtures of RaU and ordinary Pb, *cf* Aronberg, *197*, **3**: 710; 17. *21*, **47**: 96; 18. Harkins and Aronberg, *1*, **42**: 1328; 20. Merton, *5*, **99**: 87; 21. **100**: 81, 21 (²⁹) Hevesy, *4*, **26**: 410; 13 *63*, **14**: 49; 13.

(³⁰) Hevesy and Paneth, *75*, **123**: 161; 14. Meitner, *63*, **12**: 1094; 11. Hevesy, *4*, **23**: 628, 12 Wertensteinowa, *286*, **10**: No 6, 771; 17. On the deposition of Th-B and Ra-E, Paneth and Hevesy, *75*, **122**: 1037; 13. (³¹) Hevesy and Paneth, *75*, **124**: 381, 15 (³²) Paneth, *0* (³³) Fajans and Lembert, *93*, **95**: 297, 16 (³⁴) Richards and Schumb, *l c*. (³⁵) Hevesy and Obrutscheva, *58*, **115**: 674, 25.

ARTIFICIAL DISINTEGRATION OF THE ELEMENTS

G. RUDORF

Disintegration by the splitting off of positively charged hydrogen nuclei by the action of rapidly moving α-particles.

(a) Disintegration obtained with B, N, F, Ne, Na, Mg, Al, Si, P, S, Cl, A, K (¹, ², ³, ⁵).

(b) No disintegration obtained with H, He, Li, C, O, Ni, Cu, Zn, Se, Kr, Mo, Pd, Ag, Sn, X, Au, U (², ³, ⁵).

(c) Doubtful, Be (⁴, ⁵).

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Rutherford, *5*, **37**: 581; 19 *5*, **97**: 374; 20 (²) Rutherford and Chadwick, *5*, **42**: 809; 21 (³) Rutherford and Chadwick, *5*, **44**: 417, 22, also Rutherford, *4*, **121**: 400; 22. (⁴) Kirsch and Petterson, *75*, **132**: 299, 24 *5*, **47**: 500; 24. (⁵) Rutherford and Chadwick, *67*, **36**: 417; 24

RANGE OF EMITTED HYDROGEN NUCLEI (², ³, ⁵)

Element	Forward range in cms	Backward range in cms
B	58	38
N	40	18
F	65	48
Na	58	36
Al	90	67
P	65	49
Mg, Si, S, Cl, A, K	18-30	
Ne	16	

The values for B, F, Na, P are possibly somewhat in error (³) but are certainly greater than 40 (²).

ELECTRON EMISSION PRODUCED BY RADIATION FROM RADIOACTIVE SUBSTANCES

PIERRE AUGER

RELATIVE IONIZATION OF GASES BY Po α-RAYS HAVING A 3.8 CM RANGE(¹)

Gas	Air	O₂	N₂	CO₂	Illuminating gas
I	1	1 12	0 97	1 23	0 38

RELATIVE MOLECULAR IONIZATION OF GASES BY β AND γ RAYS (²)

Gas	Air	H₂	O₂	NH₃	N₂O	CO₂	C₂N₂	SO₂	CS₂	C₆H₁₂
Iβ	1	0.16	1.17	0 89	1.55	1.60	1 86	2 25	3 62	4 55
Iγ	1	.16	1 16	90	1 55	1.58	1 71	2 27	3 66	4 53

Gas	C₆H₆	CH₃OH	CH₃Br	CHCl₃	CH₃I	CCl₄	C₂H₄O
Iβ	3.95	1.69	3 73	4 94	5.11	6 28	2 12
Iγ	3.94	1 75	3 81	4 93	5.37	6 33	2 17

Gas	C₂H₅Cl	C₂H₅Br	C₂H₅I	(C₂H₅)₂O	Ni(CO)₄
Iβ	3.24	4.41	4.39	5.90	
Iγ	3 19	4 63	4.29	6.47	5 98

RESIDUAL IONIZATION AS DEPENDENT ON THE PRESSURE

Ionization from the walls (a secondary radiation) in air confined for 10 days. N_I = number of ions per cm³ per sec (³).

P, atm.	0	10	20	27	40	46	50	60
N_I	0	17	30	38	46	50	50	50

NUMBER OF ELECTRONS (δ-RAYS) LIBERATED BY α-RAYS

l = thickness of metal traversed. N_E = electrons emitted per incident particle (⁴).

$10^6\, l$ (g cm⁻²)	In Al						In Ag		In Au	
	81	162	243	324	410	492	570	28.5	591	12.3	1223
N_E	11 9	14 2	15 0	17 2	17 8	18 9	19 4	8.12	13.76	9.82	18.16

PAIRS OF IONS PRODUCED BY α-RAYS

If R_0 cms is the range of the α-particle in air, it will produce n pairs of ions. $n = n_0 R_0^{3/2}$, where $n_0 = 6.233 \times 10^4$. Direct measurement for Ra-C' gives $n = 2.20 \times 10^5$ (⁵).

ENERGY

Energy of electrons (Sec. β-rays) emitted by metals subjected to the action of γ-rays from Ra(C + E). Three groups of rays (⁶).

Metal	Pb	Pt	W	U	Ba
Atomic number	82	78	74	92	56
Energy of the secondary rays. Volts × 10⁻⁸.	1.49 2.03 2 60	1.58 2.12 2.69	1 66 2.20 2 76	1.22 1.74 2.31	2.53

SECONDARY β-RAY VELOCITIES

Pb subjected to the action of γ-rays from Ra-B has been found to emit the following secondary β-rays:

$$RH = \frac{mu^2}{e(1-\beta^2)} = 3610,\ 3250,\ 2990,\ 2735,\ 2225,\ 2130,\ 2000,$$

1935, 1825, 1750, 1620, 1560, 1400, 1240, 1150, 1010, 950, 820, 800 [8].

ABSORPTION

Absorption of the secondary β-rays emitted by metals when subjected to the radiation from Ra(B + C). μ_s for the hard rays, μ_w for the soft rays. Absorbing screen, Al [7].

Metal	Ag	Al	Au	Cu	Fe	Ni	Pb
μ_s, cm⁻¹	69	14	118	35	41	52	118
μ_w, cm⁻¹	207	52.5	345	105	165	165	345

LITERATURE

(For a key to the periodicals see end of volume)

(1) F. Hess and M. Horngate, 75, 129: 7; 20. (2) Klemann, 5, 79: 220, 07 (3) K. Melvina Downey, 8, 30: 186; 22. (4) H. Becker, 8, 75: 3, 217; 24. (5) H. Fonovits-Smereker, 75, 131: 355; 22. (6) Ellis, 5, 99: 261; 21 (7) A. Enderle, 75, 131: 9; 22. (8) Rutherford, Robinson and Rowlinson, 3, 28: 281; 16.

ENERGY OF RADIOACTIVE PROCESSES

STEFAN MEYER

HEAT PRODUCTION OF RADIOACTIVE SUBSTANCES

Joules per hour per gram of the radioactive element and the decay products in equilibrium therewith. (1 Joule = 0.2390 g-cal.)

Substance	Rays	Meyer & Hess(4)	Hess(2)	Rutherford & Robinson (7)
Ra	α and recoil		105 5	
Rn	α and recoil			105 0
Ra-A	α and recoil	573	467 7	119 7
Ra-B + Ra-C	α and recoil and β, γ			127 6
				211 3
Total		573	573	565

Substance	Heat	Lit.
Th	10.0 × 10⁻⁴	(5)
U	4 2 × 10⁻⁴	(6)
Pitchblende (ca. 64% U)	27 2 × 10⁻⁴	(6)

Ellis and Wooster (1) have determined the γ-heat effect of Ra-B to be 3.6; Ra-C, 32.2; total, 36 joules/h. Calculations of the heat effect of β-α and γ-rays have been made by Meitner (3) and Thibaud (8).

LITERATURE

(For a key to the periodicals see end of volume)

(1) Ellis and Wooster, 201, Feb. 2, 1925. (2) Hess, 75, 121: 1419; 12. (3) Meitner, 218, 12: 1146; 24. (4) Meyer and Hess, 75, 121: 603; 12. (5) Pegram and Webb, 8, 27: 18; 08. 199, 5: 271; 08. (6) Poole, 3, 19: 314; 10. 21: 58; 11. 22: 183; 12. (7) Rutherford and Robinson, 75, 121: 1491, 12 3, 25: 312; 13 (8) Thibaud, 34, 180: 1166, 25.

CHEMICAL EFFECTS OF α-PARTICLES

S. C. LIND AND D. C. BARDWELL

M is the total number of molecules reacting (on the left hand of the equation, first column); N is the total number of ion pairs produced in the reactants by α-particles.

$$\frac{M}{N} = \frac{\left(\frac{k\mu}{\lambda}\right)' \cdot V}{D \cdot F \cdot G \cdot H} \times 1.66 \times 10^3$$

V = volume in cm³ of, and D = diameter in cm of, the reaction sphere.

F = average intensity of ionization (1). G = specific molecular ionization (air = 1).

$H = (\alpha + R)/\alpha$ where α and R are α-ray and recoil atom effects resp. (2).

$$\left(\frac{k\mu}{\lambda}\right)' = \left(\ln\frac{P_1}{P_2}\right) + [E_o(e^{-\lambda t_1} - e^{-\lambda t_2})] \quad (3)$$

where E_o = initial radon (in curies), P = pressure (mm Hg), λ = decay constant of radon (in reciprocal days) and t = time (in days).

Where the quantity of gas in the reaction vessel at atmospheric pressure exceeds the air equivalent of a bulb 2.5 cm in diameter, the ionization is calculated by equations developed by W. Mund (17), slightly modified:¹

¹ The modified equation is derived by correcting the integration of Mund's function $\psi(r) = \int_0^{2R}(r-x)^{3/2}x^2dx$ (equation 5, p. 340). In the large bulbs used by Mund no error was introduced by employing his equation since $2R > r$.

$$I = N_0\,(1-e^{-\lambda t})k\bigg[r^{2/3} + \frac{1}{2}r'^{2/3} + \frac{1}{2}r''^{2/3} - \frac{3}{20R}\bigg\{3r^{5/3} + r'^{5/3} + r''^{5/3}$$
$$- 3(r - 2R)^{5/3} - (r' - 2R)^{5/3} - (r'' - 2R)^{5/3}\bigg\} + \frac{81r^{1/3}}{3520R^2} -$$
$$\frac{27}{160}(r-2R)^{2/3}\bigg\{\left(\frac{r-2R}{R}\right)^2 + \frac{3}{22}\left(\frac{r-2R}{R}\right)^3\bigg\}\bigg]$$

I = Number of ions produced by the three sets of α-particles in the time t.

N_2 = Number of atoms of radon present initially (t = 0) (1 curie = 1.772 × 10¹⁵ atoms Rn)

R = Radius of reaction bulb in cms.

λ = Decay constant of radon (as above)

$k = 6.67 \times 10^4 \dfrac{\text{ions}}{\text{cm}^{2/3}}$ = ionization constant per α-particle as a function of the range (5); $i = kr^{2/3}$ or $kr'^{2/3}$ or $kr''^{2/3}$ for Rn, Ra-A, and Ra-C, resp. (air at 760 mm and 0°C)

r, r', r'' = ranges of α-particles from Rn, Ra-A, and Ra-C, resp. Wourtzel's (13) M/N values are recalculated by the Mund equation

The values adopted for the number of α-particles per sec per g of radium, and the total ions from one α-particle of Ra-C in its completed path in air are respectively, for column (a) 3.72 × 10¹⁰ (4) and 2.37 × 10⁵ (5), and for (b) 3.40 × 10¹⁰ (6, 7) and 2.20 × 10⁵ (8). Other combinations of these numbers give intermediate values of M/N.

Reaction l = liquid, g = gas, s = solid	$\frac{M}{N}$		Lit.
	(a)	(b)	
$2H_2g + O_2g \rightarrow 2H_2Ol$ .... Dry or moist; at 25°C to −75°C	5.13	6.05	(9, 10)
$2H_2Ol \rightarrow 2H_2g + O_2g$	{ 0.86 1.05	1.01 1.24	(11) (11)
$2H_2Og \rightarrow 2H_2g + O_2g$	<0.01	<0.01	(11)
$2H_2O_2g \rightarrow 2H_2g + O_2g$	0.05	0.06	(11)
$CO_2 \rightarrow 1\%$ disappearance of gas, no decomposition products	5 × 10⁻³	6 × 10⁻³	(18)
$COg \rightarrow COg + C_nO_ms + Cs$	1.85	2.18	(18)
$2COg + O_2g \rightarrow 2CO_2g$ at room temperature	5.7	6 7	(18)
$2COg + O_2g \rightarrow 2CO_2s$ at liquid air temp.	>3.1	>3.7	(18)
$COg + H_2g \rightarrow$ carbohydrate s	3.13	3.7	(18)
$CO_2g + H_2g \rightarrow$ carbohydrate s + H_2Ol	1.44	1.70	(18)
$CO_2g + CH_4g \rightarrow$ carbohydrate s + H_2Ol	0.76	0.90	(10)
$CH_4g \rightarrow H_2g +$ hydrocarbons g,l and s..	2.0	2.4	(10)
$C_2H_6g \rightarrow H_2g +$ hydrocarbons g,l and s.	1.7	2.0	(10)
$C_3H_8g \rightarrow H_2g +$ hydrocarbons g,l and s.	1.5	1.8	(10)
$C_4H_{10}g + H_2g +$ hydrocarbons g,l and s	1.4	1.6	(10)
$CH_4g + 2O_2g \rightarrow CO_2g + H_2Ol....$	4.4	5.2	(10)
$CH_4g + 2O_2g + [1 mol\% (C_2H_5)_2Se] \rightarrow CO_2g + H_2Ol$	5.7	6.7	(10)
$2C_2H_6g + 7O_2g \rightarrow CO_2g + H_2Ol....$	6.8	8.0	(10)
$(CN)_2g \rightarrow$ { 5% to N_2g and Cs........ 95% to paracyanogen s....	} 7.8	9.2	(12)
	18°	1.01	1.19	(13)
	25°	1.0	1 2	(10)
$2NH_3g \rightarrow N_2g$ and $3H_2g$	108°	2.0	2 35	(13)
	220°	2.92	3.44	(13)
	315°	3.15	3.80	(13)
	18°	3.40	4.00	(13)
$H_2Sg \rightarrow H_2g + Ss$	95°	2.80	3.30	(13)
	220°	2.38	2.80	(13)
$H_2Ss \rightarrow H_2g + Ss$	−190°	3.?	4.?	(13)
	−78°	2.74	3 23	(13)
$N_2Og \rightarrow$ { $N_2g + O_2g$ .... $N_2g + NOg$	18°	2.21	2.61	(13)
	220°	2.95	3.48	(13)
$H_2g + Cl_2g \rightarrow 2HClg$	4000	4700	(14)
$2HClg \rightarrow H_2g + Cl_2g$	{ 0.76 1.24	0.90 1 46	(15) (10)
$H_2g + Br_2g \rightarrow 2HBrg$	0 54	0.64	(16)
$2HBrl \rightarrow H_2g + Br_2g$	2.6	3 1	(16)
KI in acid soln. → free I	0.76	0.90	(16)

Reaction l = liquid, g = gas, s = solid	$\frac{M}{N}$		Lit.
	(a)	(b)	
$xHCN \rightarrow (HCN)_xs + 5\% N_2g$	10.5	12.4	(18)
$C_2N_2g + O_2g \rightarrow$ { 63%→(CNO)_2s 37%→CO_2g + N_2g }	7.2	8.5	(10)
$C_2N_2g +$ { 67% C_2N_2→(HCN)_xs $H_2g \rightarrow$ 33% C_2N_2→(C_2N_2)_xs }	6.8	8.0	(10)
$C_2H_4g \rightarrow H_2g +$ hydrocarbons g, l, and s	5.0	5.9	(10)
$C_2H_4g \rightarrow (C_2H_4)_xs + 2\% H_2g$	19.5	23.0	(10)
$C_2H_4g \rightarrow (C_2H_4)_xs + 1 + \% H_2g$	20 5	24.2	(10)
$C_2H_4g + H_2g \rightarrow (C_2H_4)_xs$ (11% H_2 reacted)	19.6	23.1	(10)

Catalytic Effect of Inert Gases (10, 20, 21)

The $-M/N$ values in the table below give the total number of molecules of reactants disappearing for each ion pair of both catalyst and reactants. Example: $\frac{M_{C_2H_2}}{N(C_2H_2 + N_2)} = 18.7$, means that 18.7 molecules of C_2H_2 polymerize to $(C_2H_2)_xs$ for each ion pair whether formed in the reactant or in the catalyst. With the increasing ratio of catalyst to reactant, a decrease in the $-M/N$ is indicated—probably attributable to exhaustion effects. Values by the (a) method only are given.

Reactants	Catalysts							
	Pure gas	N_2	H	Ne	A	Xe	CO_2	H_2
C_2H_2	19 5	18 7 to 17.8	20 1 to 17.0	19.6 to 16.3	18.2 to 15 0	18.5	17.4	19.6
C_2N_2	7 2	7 2				7.2		reacts
HCN .	10 8	10 0				10.0		
$2H_2 + O_2$	5 13	5.0					reacts	
$2CO + O_2$	5 7				3 9		none	

LITERATURE

(For a key to the periodicals see end of volume)
(1) Lind and Bardwell, 1, 46: 2585; 23. (2) Lind and Bardwell, 1, 46: 2003; 24 (3) Lind, 50, 16: 592; 12. (4) Hess and Lawson, 75, 127: 405; 18. (5) Geiger, 5, 82A: 486; 09. (6) Rutherford and Geiger, 5, 81A: 141; 08. (7) Geiger and Werner, 96, 21: 187; 24. (8) Fonovits-Smereker, 75, 121: 355; 23. (9) Lind, 1, 41: 531; 19.
(10) Lind and Bardwell, O. (11) Duane and Scheuer, 199, 10: 33; 13. (12) Lind, Bardwell and Perry, O. (13) Wourtsel, 199, 269: 332; 19. (14) Bodenstein and Taylor, 1, 37: 24; 15. (15) Cameron and Ramsey, 4, 92: 965; 08. (16) Lind, 199, 8: 280; 11. (17) Mund, 527, 44: 336, 25. (18) Lind and Bardwell, 1, 47: 2675; 25. (19) Mund and Koch, 88, 34: 241; 25.
(20) Lind and Bardwell, 166, 62: 442, 25. (21) Ibid , 166, 62: 593; 25.

SATURATION CURRENT. ABSORPTION IN LIQUIDS AND SOLIDS
Stefan Meyer

SATURATION CURRENT AND NUMBER OF IONS FOR α-RADIATORS

The saturation current is $I_s = Zke$ where Z = number of α-particles per sec per unit mass, k = number of ion-pairs per α-particle and e = 4.774×10^{-10} es.

Number of Ions, k

Based on the values of Ra-C′ and the following alternative Z values for 1 g of Ra: (a) $Z_{Ra} = 3.72 \times 10^{10}$ (19, 25); (b) $Z_{Ra} = 3.45 \times 10^{10}$ (12).

$$k = A \times 10^9 \quad (9, 11, 13, 18, 45, 47)$$

Element	A		Element	A	
	(a)	(b)		(a)	(b)
U_I	1.16	1.25	An........	1.95	2.10
U_II........	1.27	1.37	Ac-A	2.12	2.28
Io.	1.31	1.41	Ac-C	1.88	2.03
Ra	1.36	1.47	Ac-C′.....	(2.09?)	(2.25?)
Rn	1.55	1.67	Th........	1.23	1.32
Ra-A	1.77	1.83	Rd-Th... .	1.53	1.64
Ra-C	(1.47?)	(1 58?)	Th-X	1.61	1.78 *
Ra-C′	2.20*	2.37*	Tn........	1.78	1.92
Po........ .	1.50	1.62	Th-A	1.92	2.07
Pa........ .	1.44	1.55	Th-C	1.71	1.85
Rd-Ac	1.69	1.82	Th-C′	2.54	2.73
AcX	1.61	1.74			

* Basic values.

The value of $Z_U = Z_{U_I} + U_{II}$ may be obtained from Z_{Ra} and the basic equilibrium ratio $Z_{Ra}/Z_U = 3.4 \times 10^{-7}$.

The value of Z_{Th} may be calculated from the decay constant of Th. For the following assumed values of the half-life, $T_{\frac{1}{2}}$, of Th we find for Z_{Th}: 1.25×10^{10} yrs, 4.5×10^3 α sec^{-1}; 1.65×10^{10}, 3.4 α sec^{-1}; and 2.2×10^{10}, 2.6 α sec^{-1}.

Saturation Current

1. (In Electrostatic Units) [2, 3, 4, 5, 6, 7, 8, 20, 26, 31, 32, 34, 43]

Element	U_I	U_{II}	Io	Ra	Rn	Ra-A	99 96% Ra-C'	Po
In equilibrium with 1 g Ra	$I_s =$	1 47	0 79	0 82	0 94	1 08	1 38	0 91
	$I_s \times 10^{-4} =$	4 32	2 38	2 43	2 75	3 02	3 91	2 64

2. On the basis of a branching ratio of 3% for the Ac family in equilibrium with 1 g Ra [1, 2, 10, 15, 16, 17, 23, 30, 33, 38, 41].

Element =	Pa	Rd-Ac	Ac-X	An	Ac-A	99 7% Ac-C
$I_s \times 10^{-4} =$	7 93	9 00	8 86	10 7	11 7	10 4

3. 1 g U in ores [i.e. U + 97% (Io→Ra-G) + 3% (Pa→Ac-D)] is equivalent to $I_s = 7.30$; 1 g (U₃O₈→Ra-G) to $I_s = 6.2$; and 1 g average ore with 50% U₃O₈ to $I_s = 3.1$.

4. 1 curie Rn is equivalent to $I_s = 2.75 \times 10^6$ and 1 curie Rn + ½(Ra-A + Ra-C') to $I_s = 6.2_2 \times 10^6$.

5. In equilibrium with 1 g Th and based on the following alternative Z values for 1 g Th: (a), $Z_{Th} = 4.5 \times 10^3$ α sec^{-1} and (b), $Z_{Th} = 3.4 \times 10^3$ α sec^{-1}.

Element		Th	Rd-Th	Th-X	Tn	Th-A	35% Th-C	65% Th-C'
$I_s =$	(a)	0 204	0 329	0 346	0 382	0 413	0 129	0 355
	(b)	0 200	0 248	0 261	0 289	0 312	0 097	0 268

RANGE OF α-PARTICLES IN LIQUIDS AND SOLIDS

All values in microns, $\mu = 10^{-4}$ cm

A. In Liquids

Liquid	From Po [36]								From Ra-C' [37, 48]		
	$C_2H_5OC_2H_5$	C_2H_5OH	CS_2	C_6H_6	$CHCl_3$	$C_6H_5NH_2$	H_2O	C_2H_5OH	C_2H_5OH	C_6H_6	C_6H_5N	H_2O
$R_{15°}$	43 0	37 1	36 7	36 3	34 3	33 0	32 0	27 9	7 05	70	63 9	60 0	50 5

B. In Solids

From Ra-C' [49, 50, 51]

Solid	Li	Mg	Al	Ca	Fe	Ni	Cu	Zn
$R_{15°}$	129 1	57 8	40 6	78 8	18 7	18 4	18 3	22.8

Solid	Ag	Cd	Sn	Pt	Au	Tl	Pb
$R_{15°}$	19 2	24.2	29 4	12 8	14 0	23 3	24 1

C. In Photographic Plates

Source	Ra-A	Rn-C'				Th-C'	Po
Type of plate	Ilford	Sigurd (Jahr)	Ilford				Sigurd
$R_{15°}$	34.8	50.0	50.7	54	48.2	27 7	23
Lit.	(21)	(36)	(21)	(21)	(22)	(36)	(35)

D. Pleochroitic Haloes v. (53)

STOPPING POWER EQUIVALENTS OF AIR AND METALS AT DIFFERENT PARTS OF THE PATH OF AN α-RAY

Milligrams per cm² of foil equivalent to 1 cm air lying between the distances given, measured from end of range. 15°C and 1 atm. (29).

Distances cms	0–1	1–2	2–3	3–4	4–5	5–6	6–7
Al	1 90	1 71	1 65	1 64	1.63	1.62	1 62
Ag	3 805	3 28	3 10	3.01	2.93	2.86	2 81
Au	6 10	4 84	4.44	4 25	4.06	3.96	3 91

INITIAL VELOCITIES OF RECOIL ATOMS

$u = A \times 10^7$ cm sec^{-1}

From	To	A =		From	To	A =
U_I	UX_1	2 39		An	Ac-A	3.36
U_{II}	Io	2.54		Ac-A	Ac-B	3.58
Io	Ra	2 62		Ac-C	Ac-C''	3.44
Ra	Rn	2 72		Ac-C'	Ac-D	3.61
Rn	Ra-A	2 96		Th	Ms-Th₁	2 40
Ra-A	Ra-B	3 16		Rd-Th	Th-X	2.86
Ra-C	Ra-C'''	2 99		Th-X	Tn	2 99
Ra-C'	Ra-D	3 66		Tn	Th-A	3 20
Po	Ra-G	3 08		Th-A	Th-B	3 39
Pa	Ac	2 74		Th-C	Th-C''	3 26
Rd-Ac	Ac-X	3 02		Th-C'	Th-D	3.97
Ac-X	An	3 01				

RANGES (PENETRATION) OF RECOIL ATOMS

Ra-A to Ra-B, 0.14 mm in air; 0.83 mm in H₂; ca. 20μμ in Ag (52).

Rn to Ra-A—Ra-C, ca. 10μμ in Cu and Ni [14, 40].

Th-C to Th-C'', at 15° and 1 atm., 0.55s mm in H₂; 0.129 mm in air (24).

Th-C to Th-D, 15° 1 atm., 0.96₃ mm in H₂; 0.224 mm in air (24).

THE McCOY NUMBER

The McCoy number is the ratio of the total α radiation to the uni-directional radiation per cm² from a U₃O₈ surface of α-saturated thickness. McCoy [27, 28] found 793 with $I_s = 1.74 \times 10^{-3}$ es per cm² U₃O₈ and St. Meyer and Paneth [34] found 790 with $I_s = 1 73 \times 10^{-3}$. These numbers are smaller than the theoretical.

LITERATURE

(For a key to the periodicals see end of volume)

(1) Antonoff, 3, 26: 1058; 13. 199, 10: 406; 13. (2) Boltwood, 12, 25: 269; 08. (3) Boltwood and Johnstone, 5, 40: 50; 20. (4) Brössler, 75, 129: 47; 20. (5) I Curie, 34, 176: 1462; 23 (6) Flamm and Mache, 75, 121: 227; 12 (7) Flamm and Mache, 75, 122: 535, 1539, 13. (8) Fonovits, 75, 126: 761; 19 (9) Fonovits-Smereker, 75, 131: 355; 22

(10) Fussler, 2, 9: 142, 17. (11) Geiger, 5, 83: 486; 09. (85: 505; 10. (12) Geiger and Werner, 96, 21: 187; 24. 88, 5: 12; 24. (13) Girard, 199, 10: 195; 13. (14) Godlewski, 75, 125: 137; 16. (15) Guy and Russell, 4, 122: 2618; 23. (16) Hahn and Meitner, 63, 20: 529; 19. 218, 7: 611. 25, 52: 1812, 19 (17) Hahn and Meitner, 25, 54: 69; 21. 96, 8: 202; 22. (18) Henderson, 3, 42: 538; 21. (19) Hess and Lawson, 75, 127: 405, 461, 535, 599; 18. 96, 24: 402; 24. 3, 45: 200; 24

(20) Hornyak, 75, 130: 135; 21. (21) Ikeuti, 5, 32: 129; 16. (22) Kinoshita and Ikeuti, 3, 29: 420; 15. (23) Kirsch, 75, 129: 309; 20. (24) Kolhörster, 96, 2: 257; 20. (25) Kovarik, 2, 23: 559; 24. (26) Maracineanu, 34, 177: 682; 23. (27) McCoy, 2, 20: 381; 05. 24: 124; 07. 5, 11: 177; 06. (28) McCoy and Ashman, 12, 26: 521; 08. 199, 5: 362; 08. (29) Marsden and Richardson, 3, 25: 184; 13.

(30) Meyer, 75, 129: 483, 20. (31) Meyer and Hess, 75, 120: 1187; 11. (32) Meyer and Hess, 75, 121: 603; 12. (33) Meyer and Hess, 75, 123: 909; 19. (34) Meyer and Paneth, 75, 121: 1403; 12. (35) Michl, 75, 122: 1955, 1965; 14. (36) Muhlestein, 149, 4: 38; 22. (37) Philipp, 96, 17: 23; 23. (38) Piccard and Kessler, 149, 5: 491; 23. (39) Richter, 75, 123: 539; 19.

(40) Rie, 75, 130: 283; 21. (41) Rona, 25, 55: 294; 22. (42) Russell and Widdowson, 5, 46: 915; 23. (43) Rutherford, "Radioactive Substances,"

p. 523; 13. (44) Rutherford, "*Radioactivity*," p. 156; 04. (45) Rutherford, *3*, 10: 193; 05. (46) Sahni, *3*, 29: 836; 15. (47) Taylor, *3*, 23: 670; 12. (48) Traubenberg and Philipp, *96*, 6: 404; 21. (49) Traubenberg, *96*, 6: 396; 21.

) Traubenberg, *65*, 21: 588; 20. (51) Traubenberg, *96*, 2: 268; 20 (52) Wertenstein, *Thesis Paris*, 13. (53) Duane, *241*, 61: 286; 22. Gudden, *218*, 12: 940; 24. *Diss Göttingen*, 19. *94*, 56: 422; 21. *96*, 26: 110; 24 Hirschi, *94*, 12: 939; 24. *242*, 64: 65; 19 68: 209; 20. Hovermann, *Diss. Göttingen*, 12. *190*, *Beil.*, 24: 321; 12. Joly, *Congr intern. Rad*

Bruxelles (1910) 1: 370; 11. "*Halley Lecture*," Oxford, Clarendon Press; 24. *58*, 99: 456, 476, 17. 109: 517, 578, 711; 22. 114: 160; 24. *94*, 12: 603; 24. *3*, 12: 381; 07. 19: 327; 10. *5*, 102: 682; 23. Joly and Fletcher, *5*, 19: 630; 10. Joly and Poole, *58*, 164: 92; 19. Joly and Rutherford, *5*, 23: 644; 13. Mennell, *58*, 82: 08; 09. Mügge, *188*, 11: 1; 23. 10: 78; 19 11: 110, 22. *189*, *71*; 65, 113, 142; 09. 69: 397; 07. Rudge, *58*, 88: 167; 11 Rutherford, *3*, 19: 192; 10. Rayleigh, *58*, 108: 279; 21. Schmidhuber, *Mitt Oberrhein geol. Ver. N.F.*, 6: 35; 15. Weber, *189*, 78: 388; 23

RADIOACTIVE RADIATIONS IN GASES

R. D. KLEEMAN

I. RANGE AND VELOCITY OF α-RAYS IN GASES AT 1 ATMOSPHERE

At $t°$ and 1 atm., $R_t = R_0 \dfrac{T}{273.1}$

RANGE IN AIR AT 0° AND 1 ATM. [13]

From	U$_I$	U$_{II}$	Io	Ra	Rn	Ra-A
R_0, cms	2 531	2 910	3 028	3 212	3 907	4 476

From	Ra-C'	Ra-C'$_1$*	Ra-C'$_2$*	Ra-F, Po	Pa	Rd-Ac
R_0, cms..	6 608	8 8	10.6	3 721	3 482	4 432

* Two new α-rays from Ra-C' by the scintillation method [24]

From	Ac-X	An	Ac-A	Ac-C'	Th	Rd-Th
R_0, cms	4 111	5 487	6 211	5 224	2 749	3 810

From	Th-X	Tn	Th-A	Th-C	Th-C'
R_0, cms	4 127	4 799	5 387	4 538	8 168

MEASURED RANGES IN OTHER GASES

Gas	From Ra-C' Air	O$_2$	H$_2$	He	From Po Air	O$_2$	H$_2$
R$_{18}$°	6 93 to 6 97	6.26	30.93	32.54	3 76 to 3 95	3 43	16 8
Lit.	(12, 15, 17, 27)	(27)	(27)	(27)	(9, 12, 14, 16, 18, 19, 20, 21, 22, 23, 27)	(21, 27)	(21, 27)

Gas	From Po He	N$_2$	CH$_4$	CO	CO$_2$	NO	SO$_2$	CH$_3$Br
R$_{18}$°	17 62	3 82	4 18	3 70	2 49	3 41	2 08	1 80
Lit	(27)	(21)	(21)	(21)	(21)	(21)	(21)	(21)

For range of recoil atoms, see p 368

Distribution of Ranges.—This follows a probability law. Thus the most probable range for a Ra-F ($=$Po) α-ray is 3.85 cm at 15° and 1 atm.; 90% lie between 3.75 and 3.95, and 60% between 3.8 and 3.9 (8). For long range particles from Th-C', Ac-C', and Ra-F, v. (2). I. Curie (8.5) found for a very narrow beam for Po, the range $R_{15}^{760} = 3.87$ cm, as against the much greater value of H. Geiger, $R_{15}^{760} = 3.925$ cm.

Velocity of α-particles.—The velocity, u, of any α-ray may be computed from the relation $u^3 = aR$ where a is a constant and R the length of the remaining path (11). Taking $u = 1.922 \times 10^9$ cm sec^{-1} (25) as the *initial* velocity of the α-particles from Ra-C', at 0° and 1 atmosphere in air, this becomes $u = 1.0246 \times 10^9 R^{1/3}$ where R is the range.

Example: R_0 for Th-C' in air is 8.168 cm (Table 1, *supra*). Hence $u = 1.0246 \times 10^9 \times \sqrt[3]{8.168} = 2.064$ cm sec^{-1}, the initial velocity.

The following values of $u \times 10^{-9}$ at 0° and 1 atm. have been directly measured: Ra-A, 1.690 (25); Ra-C', 1.922 (25); Po, 1.593 (7); Th-C, 1.714 (30); Th-C', 2.060 (30). S. Rosenblum (22.5) determined directly the ratio of the initial velocities of the α-particles from Th-C—Th-C' = 1.209.

For velocity of recoil atoms see p. 368.

II. NATURE OF PATH

The path of an α-particle may undergo sudden bends (4, 26, 29). The table gives the number of bends (whose angles lie between the limits $\theta_1 - \theta_2$) for path-lengths (between bends) within the limits $l_1 - l_2$, for 281 Ra-F α-rays in air containing 75% A. The unit of l is [1]$_{120}$ cm. 0° and 1 atm (3).

$\theta_1 - \theta_2 =$	20°-30°	30°-40°	40°-50°	50°-60°	60°-70°	70°-80°	80°-90°	90°-180°
3 7	11	20	22	8	13	7	6	8
7-15	21	17	16	5		7		5
15 30	12	8	7	2			5	
$\theta_1 - \theta_2 =$	10°-20°	20° 30°	30° 180°					
30-100	20	3	3					

The ionization along the path of a β particle varies inversely as the square of the velocity of the particle (28.5). The table gives the number, N_1, of ions produced by a ray per first cm of path (13.5). $e = 4.774 \times 10^{-10}$ es.

Source	Ac-C''	Th-C''	Ra-B	Ra-C	Ra-E	U
N_1	132	132	130	105	67	76

Coefficients of absorption, λ, of β rays in air and CO$_2$ at 1 atm. and 22° (18.5).

Substance	Ra-E	Ac-C''	Th-C''	U-X$_2$
Air, λ in cm^{-1}	0 0152	0.0091	0.0008	0.0065
Air, λ in (g/cm^2)$^{-1}$	12 70	7 60	5.68	5.43
CO$_2$, λ in cm^{-1}	0 0207	0 0175	0.0129	0.0114
CO$_2$, λ in (g/cm^2)$^{-1}$	16 31	9 62	7 08	6 26

Substance	U-X$_1$	Ra-D	Ra-D very soft	Th-B	Ac-B
Air, λ in cm^{-1}	0.12	0.097	0.64	0 090	0.31
Air, λ in (g/cm^2)$^{-1}$	100	81	535	75	260
CO$_2$, λ in cm^{-1}	0 23	0.183	1.69	0.142	
CO$_2$, λ in (g/cm^2)$^{-1}$	126	101	930	78	

Coefficient of absorption λ in cm $^{-1}$ of γ rays from Ra-C' in air at 1 atm. and 22° is 0.447 × 10^{-4} (17.5).

III. Ionization along the path of single α-ray of Ra-C in air at 12° and 1 atm Total number produced = 2 31 × 10⁵. e = 4.774 × 10⁵ es. Lit 27

IV. STOPPING POWER OF GASES

$S = \dfrac{R_{Gas}}{R_{Air}}$ for the same temperature and pressure [6].

1. Ionisation method [6].   2. Track-condensation method using Ra-F [31].   3. Scintillation method.   α-rays of $R_{11} = 6.15$ cm [1].

Gas	S	Method	Gas	S	Method
A	0 951 Ra-C'	1	CO	985 Ra-C'	1
	.934 Ra-A			976 Ra-A	
A	.930	3	CO	1 02 Ra-F	2
H₂	.24	1	CO₂	1.505 Ra-C'	1
H₂	.22 Ra-F	2		1.488 Ra-A	
He	.201	1	CO₂	1.52 Ra-F	2
He	.1757	3	CH₄	0.860 Ra-C'	1
Kr	1.330	3		.880 Ra-A	
N₂	.989 Ra-C'	1	CH₄	.91 Ra-F	2
	.982 Ra-A		CCl₄	4 00	1
N₂	.99 Ra-F	2	CS₂	2.18	1
Ne	.586	3	CHCl₃	3.16	1
O₂	1.064 Ra-C'	1	CH₃Br	2 03	1
	1.057 Ra-A		CH₃Br	2 04 Ra-F	2
O₂	1 08 Ra-F	2	CH₃I	2 58	1
Xe	1 804	3	C₂H₂	1 118 Ra-C'	1
Air	1.00	1		1 121 Ra-A	
H₂O	.77 Ra-F	2		1 122 Rn + Ra	
SO₂	1 82 Ra-F	2	C₂H₄	1 349 Ra-C'	1
N₂O	1.46	1		1 369 Ra-A	
N₂O	1.11 Ra-F	2		1 379 Rn	

Gas	S	Method	Gas	S	Method
	1 405 Ra		C₂H₆O	2.00	1
C₂H₄Cl	2 371 Ra-C'	1	C₄H₁₀O	3.437 Ra-C'	1
	2.385 Ra-A			3.471 Ra-A	
C₂H₅I	3 12	1	C₅H₁₂	3.544 Ra-C'	1
C₂H₆	1.514 Ra-C'	1		3.595 Ra-A	
	1 526 Ra-A		C₆H₆	3.33	1

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Bates, *5*, **106**: 622; 24.   (²) Bates and Rogers, *5*, **106**: 97; 24.   (³) Blackett, *5*, **102**: 294, 22.   (⁴) Blackett, *5*, **103**: 62; 23.   (⁵) Bragg, *8*, **10**: 617, 06.   *5*, **13**: 333; 07.   *Studies in Radioactivity*, p. 65, (Macmillan, London)   (⁶) Bragg and Kleeman, *5*, **10**: 318; 05.   (⁷) Curie, *34*, **176**: 220; 22   (⁸) Curie, *34*, **176**: 434; 23.   *51*, **4**: 170; 23.   (⁸·⁸) Curie, *350*, No. **212**; 25.   (⁹) Dawson, *75*, **124**: 509; 15.
(¹⁰) Geiger, *5*, **82**: 486; 09.   (¹¹) Geiger, *5*, **83**: 505; 10.   (¹²) Geiger, *96*, **8**: 45; 21.   (¹³) Geiger, *96*, **8**: 45; 21.   (¹³·⁵) Geiger and Kovarik, *3*, **22**: 604; 11.   (¹⁴) Geiger and Nuttall, *3*, **22**: 613; 11.   (¹⁵) Geiger and Nuttall, *3*, **24**: 647; 12.   (¹⁶) Hahn, *63*, **7**: 412, 456, 557; 06.   *5*, **11**. 793; **12**: 82, 244; 06.   (¹⁷) Henderson, *5*, **42**: 538; 21.   (¹⁷·⁵) Hess, *63*, **12**: 1000; 11.   (¹⁸) Kovarik, *199*, 11: 69; 14.   *2*, **3**: 148; 14.   (¹⁸·⁵) Kovarik, *2*, **6**: 419; 15.   (¹⁹) Kucera and Masek, *63*, **7**: 337, 630, 650; 06.
(²⁰) Levin, *63*, **7**: 519; 06.   (²¹) Van der Merwe, *3*, **45**: 379; 23.   (²²) Meyer, Hess and Paneth, *75*, **123**: 1459; 14.   (²²·⁵) Rosenblum, *34*, **180**: 1332; 25   (²³) Rothesteiner, *75*, **125**: 1257; 16.   (²⁴) Rutherford and Chadwick, *3*, **48**: 509; 24   (²⁵) Rutherford and Robinson, *3*, **28**: 552; 14.   (²⁶) Shimizu, *5*, **99**: 425, 432; 21.   (²⁷) Taylor, *3*, **26**: 402; 14.   (²⁸) Tunstall and Mackower, *3*, **29**: 259; 15.   (²⁸·⁵) Wilson, *5*, **88**: 240; 11.   (²⁹) Wilson, *5*, **87**: 277; 12.
(³⁰) Wood, *3*, **30**: 702; 15.

ABSORPTION AND DIFFUSION OF β-RAYS IN LIQUIDS AND SOLIDS

PIERRE AUGER

Absorption Coefficients.—If I_0 be the initial intensity, and I_x the intensity after screen thickness x is traversed, $I_x = I_0 e^{-\mu x}$ where μ, the absorption coefficient, varies slightly with the thickness traversed. d = density.

ABSORPTION BY AL

Source	Ra-D	Th-A	Ra-E	Ac-C	Th-D	Ra-C
μ, cm⁻¹	130	111.0	43 3	28 5	16 3	13 5
Lit			(12)			

Source	Ra-D very soft	Ra-B Soft	Ra-B Hard	Rb	Ra	U-X₁	U-X₂
μ, cm⁻¹	5500	91	13	347	312	500	15
Lit	(13)	(6)		(10)	(9)	(5)	(5)

ABSORPTION OF β-RAYS FROM U-X [11]

Screen material	Ag	Al	C	Ca	Cd	Fo	Ir	Mg	Ni	Pb
μ/d, cm² g⁻¹	7.31	4.1	3.75	6.3	7.4	6 61	9.5	4.0	6.35	9.75

Screen material	Rh	S	Sb	Sn	Ta	Zn	NH₄Cl	CaSO₄	SrSO₄
μ/d, cm² g⁻¹	7.0	4.52	7.74	7.6	8.9	6 4	5 2	4 95	6 50

Screen material	BaCl₂	BaSO₄	NaCl	KF	KCl	KBr	KI
μ/d, cm² g⁻¹	8.07	7.7	4.68	4.8	4 88	6.1	7.8

ABSORPTION OF β-RAYS OF RA-E [7]

Screen	C	Al	Cu	Mo	Ag	Sn
μ/d	15.8	16.9	19.2	21.0	21.7	22.1

If N is the atomic number of the screening element, $\mu/d = 15 + 0.142 \, N$.

RANGE IN ALUMINUM OF β-RAYS OF VARIOUS VELOCITIES (LINEAR EXTRAPOLATION) [15]

RH	1380	1930	2535	3170	3790	4400
Range in cm	0 018	0 064	0.124	0.189	0.279	0 360

RH	5026	6230	7490	8590	11 370
Range in cm	0.440	0 580	0 785	0.925	1.36

Velocity Decrease.—R = Radius of curvature of the β-ray in a magnetic field of N units and field force H gauss. ΔRH is the change in RH due to a screen of 0.01 g cm⁻² and is proportional to the velocity.   According to Bohr, $\dfrac{\Delta RH}{c^2} u^2 = $ a constant, K. u = the velocity of the particle, and c that of light [14].

DECREASE OF VELOCITY FOR β-RAYS FROM RA-B AND RA-C

RH	ΔRH	K	ΔRH	K	ΔRH	K
No screen	Mica screen		Sn screen		Au screen	
1392	138.1	34.8	89.2	22.8		
1660	101.4	34.7	67.4	23.4		
1925	78	33 1	56.8	24.1		
2235	72.6	36 2				
2960	66.7	43 5				
3260	59 2	41				
4840	47.3	39.9	37.6	31.7	32.2	27.3
5255	49 3	42.2	37.8	32.5		
5880	43.1	38	32.2	28.6	32.6	29
6160	41	36.7				
7060	38.4	35.4	30.2	27.8		

Dispersion of β-rays [2, 3, 6].

LITERATURE

(For a key to the periodicals see end of volume)

(¹) von Bœyer, *63*, *22*: 485; 12. (²) Bothe, *96*, *6*: 368; 23. (³) Crowther and Schonland, *5*, *100*: 526; 22. (⁴) Danysz, *51*, *2*: 949; 13. (⁵) Fajans and Göhring, *63*, *14*: 877; 13. (⁶) Fajans and Makower, *3*, *22*: 292; 12

(⁷) Fournier, *54*, *180*: 284; 25. (⁸) Geiger and Bothe, *96*, *6*: 205; 21. (⁹) Hahn and Meitner, *63*, *10*: 741; 09.
(¹⁰) Hahn and Rothenback, *63*, *20*: 197; 19. (¹¹) Jungenfeld, *63*, *14*: 807; 13. (¹²) Kovarik, *3*, *20*: 849; 10 (¹³) Meitner, *63*, *16*: 272; 15. (¹⁴) Rawlinson, *3*, *30*: 627; 15. (¹⁵) Varder, *8*, *29*: 725; 15. (¹⁶) Wilson, *8*, *96*: 141; 10.

WAVE LENGTHS OF γ-RAYS

E. von Schweidler

GENERAL RELATIONS

A wave length of λ milli-Ångstroms (10^{-3} Å $= 10^{-11}$ cm $= 1$ X-unit), corresponds to:

A Frequency $(\nu) = 2.9986 \times 10^{21}/\lambda$ sec^{-1}

An Energy $(E = h\nu) = 1.9653 \quad 10^{-5}/\lambda$ ergs

A Potential $\left(P = \dfrac{h\nu}{e}\right) = 1.2344 \quad 10^{7}/\lambda$ volts

The equivalent electron velocity as a fraction of the velocity of light,

$$(\beta) = \sqrt{1 - \frac{1}{\left(1 + \dfrac{24.288}{\lambda}\right)^2}}$$

$$h\nu = \frac{hc}{\lambda} = E = Pc = c^2 m_0 \left[\frac{1}{\sqrt{1 - \beta^2}} - 1\right].$$

See p. 17 for values of basic constants.

WAVE LENGTHS DETERMINED WITH CRYSTAL GRATINGS

φ = angle of reflexion, d = grating space = 2.814 Å for rock salt = 3.028 Å for calcite. $\lambda = 2d \sin \varphi$. Intensity indicated thus, s = small, m = moderate, g = great, vg = very great.

(a) Soft Radiations from Ra-B. Using rock salt (², ³). Corresponding to L-series of elements of atomic Nos. 82 and 83, according to Swinne (⁵) and Wagner (⁶).

λ, in 10^{-3} Å..	1365 m	1349 m	1315 s	1286 s	1266 s	1219 s	1196 m
φ, deg. min...	14° 00′	13° 52′	13° 31′	13° 14′	13° 00′	12° 31′	12° 16′
λ, in 10^{-3} Å..	1175 g	1141 m	1100 s	1074 s	1055 s	1029 m	1006 m
φ, deg. min.	12° 03′	11° 42′	11° 17′	11° 00′	10° 48′	10° 32′	10° 18′
λ, in 10^{-3} Å..	982 g	953 m	917 s	853 m	838 m	809 m	793 m
φ, deg. min...	10° 03′	9° 45′	9° 23′	8° 43′	8° 34′	8° 16′	8° 06′

(b) Hard Radiations from Ra-B + Ra-C, Sec. 1. Radiations from Ms-Th and its products, Sec. 2.

λ, in 10^{-3} Å ..		428	(393)	(324)	296	262	242	229	196
φ, deg. min	1. Using rock salt (⁴)	4° 22′	4° 00′	3° 18′	3° 00′	2° 40′	2° 28′	2° 20′	2° 00′
Remarks			Probably 2nd order spectrum to 196 and 159		K-series				

λ, in 10^{-3} Å		169 g	159 g	137	116	99 g	71	72	66
φ, in deg. min		1° 43′	1° 37′	1° 24′	1° 11′	1° 06′	43′	41′	37.5′
Remarks		K-line						Using calcite (¹⁵)	
		Ra-C?	Ra-B?						

λ, in 10^{-3} Å	58	48	37	28	2. Ms-Th (²⁰)	168 g	145 g	62 s	52 m
φ, deg. min	33′	27.5′	21′	16′					
Remarks	Using calcite (¹⁵)					to Rd-Th			to Th-B

WAVE LENGTHS CALCULATED FROM THE ENERGY OF β-RAYS

Primary γ-rays of energy E_γ produce in the disintegrating atom itself, or in other atoms, secondary β-rays of energy $E_\beta = E_\gamma - A$, where A is the work of removal and depends upon the level from

which the β-rays originate. Sometimes it is assumed that the β-rays are primary and produce secondary γ-rays of energy $E_\gamma = E_\beta$. The energy of the β-rays is obtained from their magnetic deflections.

λ, in 10^{-3} Å Lit		66 (¹⁴, ¹⁵)	Ra	Ra-B	230 (²⁶)	174 (²⁶)	155 (²⁶)	51.9 (²²)	51.3 m (¹⁶, ²⁰)
			Ra						

λ, in 10^{-3} Å Lit	48.0 s (²⁹)	42.6 (²²)	42.0 m (²⁶)	35.6 (²²)	35.2 g (²⁶)	Ra-C + (²⁶)	209? (²⁶)	52.1? (²⁶)

λ, in 10^{-3} Å Lit ..	49.8? (²⁶)	44.47 (²⁶)	28.0? (²⁶)	45.4 (¹⁶)	37.5 (¹⁶)	32.0 (¹⁶)	30.2 (²²)	29.0 (²⁹)

λ, in 10^{-3} Å Lit .	24.9 (¹⁶)	24.3 (²³)	21.2 (²²)	20.6 (²²)	20.4 (²⁹)	20.3 (²⁶)	16.2? (²⁹)	10.93 g (²⁹)

λ, in 10^{-3} Å Lit	10.0 s (²⁹)	9.03 g (²⁹)	7.00 s (²⁹)	6.94 g (²⁹)	5.56? g (²⁹)	Ra-D	209 (¹²)

λ, in 10^{-3} Å Lit...........	Ms-Th	171 (²²)	59.7 (²²)	53.0 (²²)	37.1 (²⁹)	37.0 (²²)	29.7 (²²)	26.9 g (²⁹)

λ, in 10^{-3} Å. Lit	Rd-Th	147 (¹³)	Th-B	52.9 g (²⁹)	52 (¹³)	41.6 (¹⁶)	41.3 s (²⁹)	Th-C	45.2 s (²⁹)

λ, in 10^{-3} Å . Lit	24.5 (¹⁶)	21.3 (²⁹)	13.6 g (²⁹)	13.5 g (²⁹)	12.8 m (²⁹)	Th-B + Th-C	4.84 (²⁴)	4.71 (²⁴)

EFFECTIVE WAVE LENGTHS CALCULATED FROM ABSORPTION AND SCATTERING

The ordinary or "apparent" absorption coefficient, $\mu' = \mu + \sigma$, where μ is the "true" or "fluorescent" absorption coefficient, and σ the coefficient of scattering. For dependence on wave length ν. Glocker (⁸); Compton (¹²); Wingårdh (²³); Warburton and Richtmyer (²⁴); Jauncy (²⁵); and Allen (³⁰).

γ-RAYS FROM RA-C

λ_{eff}, in 10^{-3} Å	<63	<60	120–60	80–30
Calc. from	Abs.	Abs.	Scat.	Abs.
Lit.....	(7)	(9)	(12a)	(10b)

λ_{eff}, in 10^{-3} Å	30–25	21	24	8	19	19.5
Calc. from	Scat.	Abs.	Abs.	Scat.	Scat.
Lit	(12b)	(31)	(23)	(32a, 32b)

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Shaw, *3*, *26*: 190; 13. (²) Rutherford and Andrade, *63*, *92*: 267; 13. (³) Rutherford and Andrade, *3*, *27*: 854; 14. (⁴) Rutherford and Andrade,

5, **28**: 263; 14.   (**5**) Swinne, *65*, **17**: 481; 16   (**6**) Wagner, *65*, **18**: 405, 432, 461, 468; 17.   (**7**) Rutherford, *5*, **84**: 153; 17.   (**8**) Glocker, *65*, **19**: 66; 18 *849*, **28**: 421; 19.   (**9**) Kohlrausch, *65*, **19**: 345; 18.
(**10a**) Treitel, *Diss Heidelberg*, 20.   (**10b**) Prehnger, *75*, **180**: 279; 21.   (**11**) Ellis, *5*, **99**: 261; 21.   (**12**) Compton, *5*, **18**: 296; 19.   *5*, **41**: 749; 770, 21.   (**13**) Meitner, *90*, **9**: 131, 145; 22.   (**14**) Meitner, *218*, **10**: 381; 22.   (**15**) Smekal, *96*, **10**: 275; 22.   (**16**) Ellis, *5*, **101**: 1, 22.   *201*, **21**: 121; 22.   *96*, **10**: 303; 22.   (**17**) Meitner, *96*, **11**: 35; 22.   (**18**) Kovarik, *5*, **19**: 433; 22   (**19**) Madgwick, *248*, **6**: 136, 21.
(**20**) Meitner, *90*, **17**: 54; 23   (**21**) Hahn and Meitner, *96*, **17**: 157, 23.   (**22**)

de Broglie and Cabrera, *54*, **174**: 939; 22.   *54*, **176**: 295; 23.   (**23**) Wingårdh, *96*, **20**: 315; 23.   (**24**) Warburton and Richtmyer, *5*, **22**: 539; 23. *5*, **22**: 291; 24.   (**25**) Jauncy, *5*, **22**: 233; 23.   (**26**) Ellis and Skinner, *5*, **105**: 60, 165, 185; 24.   (**27**) Smekal, *96*, **25**: 265; 24.   (**28a**) Hahn and Meitner, *96*, **26**: 161; 24.   (**28b**) Meitner, *96*, **26**: 169; 24.   (**29**) Thibaud, *54*, **173**: 1706; 24.   *54*, **179**: 165, 815, 1052, 1322; 24.   *54*, **180**: 138; 25. *£50*, **909**: 8; 24.
(**30**) Allen, *5*, **23**: 291; 24.   (**31**) Owen, Fleming and Fage, *67*, **34**: 355; 24. (**32a**) Ahmad, *5*, **105**: 507; 24.   (**32b**) Ahmad and Stoner, *5*, **106**: 8; 24. (**33**) Gray, *58*, **115**: 13, 86; 25.   (**34**) Black, *58*, **115**: 226; 25.

RADIOACTIVE RADIATIONS FROM ORDINARY METALS

R. B. MOORE

1. POTASSIUM AND RUBIDIUM

β-rays only are emitted spontaneously, the emission being an atomic property independent of the temperature

ACTIVITY OF K IN ARBITRARY UNITS [4]

Salt	K_2SO_4	KI	KBr	KCl	KF	KClO$_4$	KNO$_3$
%K...	44 91	23 58	32 87	52 18	67 32	28.91	28 69
Activity..	37 8	21	27 8	42 2	54 0	25 5	30 6
K/Act	118	112	118	124	123	110	126

ABSORPTION OF THE β-RADIATION [6]

λ = absorption coefficient cm^{-1}, d = density of absorbent

λ/d for β-rays from K		λ/d for β-rays from Rb	
By K$_2$SO$_4$	11 32	By Rb$_2$SO$_4$	96 7
By Sn (90% of the rays)	14	By paper (90% of the rays)	162
By Sn (10% of the rays).	90	By paper (10% of the rays)..	950

ABSORPTION OF β-RAYS FROM Rb BY PAPER [5]

W = wt. paper/cm^2. I$_0$, intensity of the initial radiation; I$_p$, that of the emergent radiation.

W	0	0 00153	0 00305	0 00458	0 00764	0 0107	0 0153	0 0198
I$_p$/I$_0$..	1	0 725	0 545	0 422	0 260	0 159	0 087	0 034

2. CAESIUM, SODIUM, LEAD, IRON AND ZINC

Cs and Na are not radioactive [8, 9, 10]. Ordinary Pb shows a slight, very old Pb only a trace of activity On account of their exceptionally small activity Fe and Zn are recommended for

construction of sensitive instruments for radioactive measurements. Ca, Ba, Sr, C, Cl, Br, Cu, Fe, Pb, Mg, Mn, Ni, Ag, Zn. W, Ta, La, Se, As, Sn, Au, Sb, Al and Hg are inactive [10].

3. NOTES

O) Hahn and M. Rothenbach [3] compared Rb salts of various ages but no difference in activity was detected. The Rb rays were found to be more penetrating than the β-rays of UX$_1$, but not so penetrating as those of Ra. The ratio of the intensity of the Rb rays to those of UX$_1$ is 1:15. The half-life of rubidium is calculated to be 10^{11} years and that of potassium 3 to 7 times greater. The absorption coefficient in Al of K is from 39.6 to 55.4 as foil thickness increases from 0.0135 to 0.0405 cm. Rb decreases from 593 to 522 as foil increases from 0.0017 to 0.0051 cm.

According to Bergwitz [1] the velocity of the Rb rays is 1.85 × 10^{-10} cm-sec^{-1}

Ringer [7] states that pure K and Rb give off homogeneous β-rays, the K rays having 10 times the penetrating power of the Rb rays. Harkins and Guy [10] give this figure as from 10 to 15 and state that the radiation from Rb is slightly heterogeneous.

Geiger [2] found that the saturation current from RbCl is the same at room temperature and at liquid-air temperatures.

LITERATURE

(For a key to the periodicals see end of volume)

(**1**) Bergwitz, *65*, **14**: 655; 13   (**2**) Geiger, *75*, **132**: 69; 23.   (**3**) Hahn and Rothenbach, *65*, **20**: 191; 19   (**4**) Henriot, *54*, **151**: 1751; 10.   (**5**) Henriot, *54*, **152**: 1384; 11.   (**6**) Henriot, Thesis, Paris, 1912.   (7) Ringer, *Onders Physiol Scheikunde* **1**: 24; 21.   (**8**) Ringer, *Arch. Neerland. physiol.* **7**: 141; 22.   (**9**) Zwaardemaker, *64P*, **26**: 575; 23.   (**10**) Harkins and Guy, *197*, **10**: 11, 25.

DISTRIBUTION OF RADIOACTIVE MATERIALS IN THE ATMOSPHERE, THE HYDROSPHERE AND THE LITHOSPHERE

HERMAN SCHLUNDT

TABLE OF CONTENTS

RADON IN THE ATMOSPHERE

Method A: Rn absorbed in charcoal.
Method B: Rn condensed with liquid air.
Method C: Rn directly determined in large ionization chamber.
Method D: Rn computed from active deposit on negatively charged wire.

Place	Micro-micro Curies (10^{-12} Curies) Rn per cubic meter	Method	Number of determinations	Lit.
Montreal, Can	24–127, Mean, 80	A		[21]
Montreal, Can	Mean, 60	A	50 during 1907–8	[22]
Cambridge, Eng......	35–350, Mean, 105	A	60 during 6 mos	[23]

Place	Micro-micro Curies (10⁻¹²) Curies) Rn per cubic meter	Method	Number of determinations	Lit.
Chicago, U. S. A.	45–200, Mean, 100	B	6	(1)
Manila, P. I.	71	A	30 during 1 year	(136)
Freiburg, Switzerland	54–305, Mean, 131	A or B		(78)
Innsbrück, Austria	40–1110, Mean, 433	C	49	(137)
Seeham, Austria	188	C		(116)
Tokyo, Japan	5	D		(49)
Pacific Ocean	1 3	D	Mean of 169, 1915–21	(66)
Atlantic Ocean	1 7	.	Mean of 79	(66)
Indian Ocean	1 3	..	Mean of 37	(66)
Southern Ocean S. of lat. 50°	0.3	...	Mean of 48	(66)
All accessible ocean areas.	1 2	..	Mean of 333	(66)
High seas	2 6	...	Mean of ca. 400*	(66)

* Includes some made relatively near large bodies of land

RADIOACTIVITY OF SPRING AND WELL WATERS AND SPRING GASES

$m\mu Cl^{-1}$ = Millimicrocuries (10^{-9} Curies) per liter

Ra, $\mu\mu gl^{-1}$ = Dissolved radium, micro-micro-grams (10^{-12} g) per liter

NORTH AMERICA

Source	t°C	mμCl⁻¹ Water	mμCl⁻¹ Gas	Ra, μμgl⁻¹	Lit.
CANADA					
Quebec					
Maskinonge	8	0 079	0 250	0 5	(99)
Radnor Forges	10	0 345		0 3	(99)
St. Benoit	11	0.028		0 0	(99)
St. Leon (Lupien)	8	0.148	0 46	0 8	(99)
St. Hyacinthe (Philudor)	8	0 106		46	(99)
St. Severe	8	0 087		2 8	(99)
Varennes	9	0 224	0 81	9 2	(99)
Ontario					
Borthwick, near Ottawa	11	0 140		8 4	(99)
Sulfur Spring, Caledonia Spr.	8	0 073		5 6	(99)
				15 0	(23)
Duncan Spring, Caledonia Spr.	9	0 053	0 204	5 6	(99)
Duncan Spring, Caledonia Spr.	9		0 42	18 0	(23)
Gas Spring, Caledonia Spr	8	0 090	0 306	8 4	(99)
Gas Spring, Caledonia Spr	8		0 62	15	(23)
White Sulfur Spring, Carsbad	9	0.09		0 8	(99)
Magic Spring	9	0.087		25	(99)
Soda Spring	9	0 081	0 23	1 1	(99)
Russell Lithia, Bourget	10	0.056		5 9	(99)
Alberta (Banff)					
Upper Hot Spring	46	0 221		8 6	(99)
Kidney Spring	39	0.392		8 5	(99)
Cave Spring	30	0 470	3 34	8.5	(99)
Basin Spring	35	0.232	2.37	8.5	(99)
Auto Road Spring	19	0.640	.	23.5	(99)

Source	t°C	mμCl⁻¹ Water	mμCl⁻¹ Gas	Ra, μμgl⁻¹	Lit.
British Columbia					
Fairmont Springs		3 5		100	(11)
Sinclair		4.0		tr.	(11)
UNITED STATES					
Arlington, R. I.					
Graphite Mine Spr		8.78			(79)
Williamstown, Mass.					
Wampanoag	22	0.22	7.3		(118)
Sherman Spring		0.04			(118)
Saratoga Spr., N. Y.					
Emperor	10	0 07	0 221	68	(71)
Hathorn No. 1	10	0 142	0 213	42	(71)
Geyser	10	0 039	0 034		(71)
Pump Well No. 4	12	0.231	0 078	21	(71)
Crystal Rock	10	0.88	0.847	9	(71)
Indiana					
Mean of 27 sprs	cold	0.75			(89)
French Lick					
Pluto Spring	13	0 54			(8)
Bowles Spring	10	1 78			(8)
Illinois					
Dixon Spr No. 2		2 93			(118)
Creal Spr. No. 3		0 84			(115)
Well, Joliet		0 39			(115)
Mt. Vernon Spring		0 18			(118)
Yellowstone Nat Pk					
Mammoth Hot Spr., Hot River	51	1 44		2 5*	(104)
Main Spring	71	none	none	3 8*	(104)
Apollinaris Spr	9	1 08			(104)
Nymph Spring, Tower Falls		0 23	6.5		(104)
Upper Geyser Basin, Bench Spring	86	0 22		124	(104)
Fish Cone, West Thumb				41.8	(104)
Lower Geyser Basin, Firehole Lake	85	0 28		204	(104)
Missouri					
Sweet Springs		0 81			(103)
Rollins Spring, Columbia		0 15			(103)
Hot Springs, Ark.					
Imperial Spring	61	9.03			(9)
Palace Spring	61	0 12			(9)
Avenue Spring	62	0.89			(9)
Twin Spring	62	2 22			(9)
Arsenic Spring	54	0 49			(9)
Horseshoe Spring	60	0.18			(9)
Liver Spring	8	0 59			(9)
Kidney Spring	13	3.63			(9)
Madison, Wisconsin					
Merrill Springs		0.49			(101)
Manitou, Colo.					
Shoshone Spring	15	3 38	12.7		(102)
Manitou Soda	15	1 25			(102)
Manitou Soda	15	0 268	1 62		(54)
Shoshone		1 66	15 52		(54)
Iron Soda Spring	15	0.24	1.15		(54)
Iron Soda Spring	15	1 53	1.07		(102)
Navajo Spring		1.37	3 4		(102)
Navajo Spring	22	1.21	3.3		(54)
Steamboat Springs, Colo.					
Soda	15	0.18	1.42		(102)
Soda	15	1.36	6 03		(54)

* Ra in 10⁻¹² g per g of residue.

Source	$t°C$	Water	Gas	Ra, $\mu\mu gl^{-1}$	Lit.
UNITED STATES.—(Cont'd)					
Steamboat Springs, Colo.— (Cont'd)					
Bath House...............	40	0 08		0.54	(102)
Bath House...............	40		0.79		(54)
Iron....................	24	0 99		3.71	(102)
Iron....................	24	0.91		3.50	(54)
Craddock, Glenwood Springs, Colo.........		2 21			(54)
Virginia					
Mean of 11 springs......		0 21			(120)
Ohio					
Mean of 9 springs.......	cold	0.34			(89)
Bloomington, Ind.					
Hottle Spring*..........		0.806			(90)

* Mean of 37 tests during 9 months.

EUROPE

Source	$t°C$	Gas	Water	Lit.
AUSTRIA				
Tauern Tunnel............		3 81*		(62)
Böckstein Valley..........		3.20†		(62)
Near Vienna				
Johannesbad.............	30	1 86	6 8	(63)
Haupt Quelle, Vöslau.....	23	0 29	1 07	(63)
Tyrol				
Magenquelle, Froy.......	6	17 6		(2)
Eisenquelle, Froy.......	8	4 5		(2)
Badequelle, Steinhof....	9	0 8		(2)
Herrenbadquelle, Fischau	19	0 23	0 80	(63)
Gastein				
Grabenbäckerquelle.....	36	55 5		(60, 61)
Elisabethstollen, Hauptquelle	47	53 3		(61)
Nordquelle..............	44	9 0		(61)
Rudolfstollen............	47	21.3		(61)
Franz Josephstollen......	41	34 6		(60, 61)
Reisacherstollen.........	36	84		(61)
Teichquelle, Tanbach.....		21 3		(61)
Melaniequelle, Radegund.		5 3		(132)
Annenquelle, Mariatrost...		0 36		(132)
Johannesbrunnen, Semmering...	5	1 27		(3)

* Mean of 101 springs; highest 23.7.
† Mean of 3 springs.

Source	Gas	Water	Lit.
BELGIUM			
Deloor Spa...............	1.45		(34)
Marie-Henriette Spa......	1.45		(34)
Prince de Conde I. Spa...	1.44	1 74	(34)
Tounelet, Spa...........	1 67	2 58	(34)
La Fraineuse Spa.........	2.43		(34)
Claire-Fagne Spa.........	2 1		(34)
Salmon E. superieure Spa.	3.31		(34)

Source	$t°C$	Water	Gas
CZECHO-SLOVAKIA (20, 51, 63, 139)			
Loimannsquelle, Fransenbad........	11	0.39	0 27
Salsquelle, Fransenbad............	11	0.05	
Mine water, St. Joachimsthal 60 m depth......................	6	13.5	
375 m depth....................	14	75.9	
500 m depth....................		163.8	448.0

Source	$t°C$	Water	Gas
Bernhardsbrunnen, Karlsbad.......	61	0.65	1.14
Mühlbrunnen, Karlsbad..........	39	12.9	38.6
Schlossbrunnen, Karlsbad.........	30	7.1	20.6
		· 3.61	
Hospitalquelle, Karlsbad.........	12	0.96	
Sprudel,* Karlsbad.............	71	0.16	0.36
Eisenquelle, Karlsbad...........	8	15.7	
		19.5	
Ferdinandsbrunnen, Marienbad.....	10	0.27	
Kreuzbrunnen, Marienbad.........	8	1.75	3.56
Marienquelle, Marienbad.........		0.71	
Waldquelle, Marienbad..........	7	1.87	4.47
Augenquelle, Teplitz Schönau......	22	1.28	
Riesenquelle, Dux.............		3.58	
Urquelle, Dux................	46	2.03	9.0

* 55×10^{-13} Ra per liter

Source	Water	Gas	Lit.
ENGLAND			
Nine Wells, Cambridge..........	0.130		(94)
Well, Dale's Brewery, Cambridge..	0.196		(94)
King's Well, Bath..............	1.73	33.65	(88)
Cross Spring, Bath............	1.19		(88)
Hetling Spring, Bath...........	1.70		(88)
Hospital Natural Baths, Buxton....	0.83	7.70	(64)
Gentlemen's Natural Baths, Buxton.	1.10		(64)

Source	$t°C$	Gas	Water	Lit.
FRANCE				
Choussy, La Bourboule..........		22.9	141.5	(52)
Choussy, La Bourboule..........		20.5	161.4	(53)
de la Grange, Beaucens.........		3.03	10 36	(52)
Chaude, Audinac............		0.14	0.59	(52)
Rivière, Chaudeau............		6.51	39.5	(12)
Dames, Plombières............		10.76		(12)
Lambinet, Plombières..........		15.96		(12)
Savonneuse, No. 2, Plombières....		7.47	35.1	(12)
Vauquelin, Plombières.........		4.83	86.4	(12)
Chaudes-Fontaines, Reherry......		4.1	19.8	(12)
Celestins, Vichy...............	44	0.653	4.1	(52)
Chomel, Vichy...............	44	0.653	4.1	(52)
Boussange, Vichy.............	42	0.103	0.60	(52)
Hôpital, Vichy..............	34	0.022	0.14	(52)
Condanny, Usson.............		0.563	34.5	(65)
Plaies, Usson................		0.663	1.9	(65)
d'Alun, Aix-les-Bains..........		4.1	25.8	(16)
Le Lymbe, Bourbon-Lancy......		1.5	14.6	(16)
Pavillon, Coutreville..........		0.51		(16)
Bordeu (Grande Source), Luchon...	43	16.1	134.8	(73)
Main Spring (Saline and H_2S), Uriage-les-Bains................		0.113		(8)
Gasseng, Columbières-sur Orb.....			6.69	(18)
Cabanel, Columbières-sur Orb......			2.22	(18)
Crémieu, Columbières-sur Orb.....			1.49	(12)
Viguerie, Ax.................			16.8	(73)
Savonneuse, Bains-les-Bains.......			25.6	(73)
Vielle, Eaux-Bonnes............			3.7	(73)
La Chaldette................			93.7	(73)
Romaine, Maizières...........			10.8	(73)
Souveraine, Vals-les-Bains........		1.047	5.08	(6)
Dominique, Vals-les-Bains........		8.80		(6)

Source	t°C	$m\mu Cl^{-1}$ Gas	$m\mu Cl^{-1}$ Water	Lit.
Caroline, Mont-Doré		0.34	2 49	[87]
Lepape, Bagnères-de-Luchon		41.5		[83]
Providence, Vernet-les-Bains	38	15.7	115.9	[83]
Santé, Vernet-les-Bains	37	2.7		[83]
Pastural, Les Escalades	27	3 5		[83]
Bassin Carré, Thuès-les-Bains	74	1.04	17.7	[83]
Saint-Victor, Royat	21	15.35	35 2	[83]
Hamel, Sail-les-Bains	34	11 5	50 2	[83]
Rouge, Saint-Nectair	21	0.54	2.2	[83]
Grande Source, Bagnoles-de-l'Orne			0 74	[56]
Chaude fontaine, Antoigny			3.86	[56]
Saint-Ursin, Lignières			1 57	[56]
Fontaine Minerale, St. Michel			0 44	[56]

Source	t°C	$m\mu Cl^{-1}$ Water	Lit.
GERMANY			
Schwarzwald Region			
Antoniusquelle, Antogast	cold	6.6	[20]
Büttquelle, Baden-Baden	24	51.3	[20]
Murquelle, Baden-Baden	59	9 8	[20]
Kirchenquelle, Baden-Baden	56	1 35	[20]
Hauptquelle, Badweiler	28	3 1	[20]
Gemeindequelle, Badweiler	23	4 2	[20]
Badquelle, Griesbach	cold	10 6	[20]
Sofienquelle, Petersthal	cold	1 76	[33]
Wenzelquelle, Rippoldsau	cold	0 86	[33]
Warme Quelle, Wildbad	36	1 35	[20]
Kalte Quelle, Wildbad	cold	0 08	[20]
Well, Heidelberg	27	2.15*	[7]
Wurttemberg			
Göppinger, Sauerbrunnen		1 27	[50]
Göppinger, Staufenbrunnen		0.57	[50]
Kursaal, Kanstatt		0 22	[50]
Karlsquelle, Mergentheim		0 08	[50]
Hirchquelle, Feinach		0 42	[50]
Wildbad		0.76	[50]
Hessen and Adjoining Regions			
Sprudel XII, Bad Nauheim	33	5 8†	[105]
Karlsbrunnen, Bad Nauheim	15	9 6†	[105]
Bad Homburg, Elizabethbrunnen	11	1 46†	[105]
Luisenbrunnen	11	0 84†	[105]
Wilhelmsbrunnen, Bad Soden	14	6 62†	[105]
Solbrunnen, Bad Soden	16	1 56†	[105]
Inselquelle, Kreuznach	13	7 42†	[105]
Theodorshalle, Kreuznach	7	3 06†	[105]
Hauptbrunnen, Münster am Stein	31	8 5†	[105]
Kochbrunnen, Wiesbaden	68	0 43‡	[39]
Adlerquelle, Wiesbaden	64	2 23‡	[39]
Schützenhofquelle, Wiesbaden	50	0 29‡	[39]
Racoczy, Kissingen		1.04†	[41]
Maxquelle, Kissingen		1 58†	[41]
Maxquelle, Dürkheim a.d. Haardt	20	0.69	[7]

* 1620×10^{-11} g Ra per liter of water.
† Values obtained by multiplying Mache units by $3 64 \times 10^{-10}$
‡ Values obtained by multiplying Mache units by 4.1×10^{-10}

Source	$m\mu Cl^{-1}$ Water	No. of samples	Lit.
Bavaria			
Alexanderbad	7.73	2 spr., 6 wells, 1 reservoir	[38]
Ebermanstadt and env	0.43	1½ spr., 2 w.	[38]

Source	$m\mu Cl^{-1}$ water	No. of samples	Lit.
Epprechstein and env	1.17	2 spr., 7 w., 2 reservoirs	[38]
Fichtelgebirge, Neubau	1.55	5 spr., 8 w.	[38]
Leinleiterthal	0.36	21 spr., 5 w.	[38]
Leupoldsdorf and env	25.0	6 spr., 2 w., 5 reservoirs	[38]
Schwarzenfeld and env	0.64	3 spr., 6 w.	[38]
Weisenthau	1.32	15 spr., 6 w.	[38]
Wolsenberg and env	4.87	17 springs	[38]
Wundsiedel and env	7 7	13 spr., 6 w., 1 reservoir	[38]
Saxony			
Wettingquelle, Brambach	826 2		[31]
	650 to 754		[89]
Trinkquelle, Oberschlema	688 to 920		[89]
Marx Semler Stollen, Oberschlema	288 to 330 at 10°C		[97]
Himmelfahrtstollen, Georgenthal	24 1		[97]
Olga Brunnen, Schneeberg	13 1		[97]
Rockelmann Quelle, Schwarzenberg	12 3		[97]

Source	t°C	$m\mu Cl^{-1}$ Water	$m\mu Cl^{-1}$ Gas	Lit.
HUNGARY				
Budapest				
Rakoczy, St. Lucasbad	42	7.40		[134]
Composite, 17 spr. Lucasbad		3.35	9.08	[135]
Trinkquelle, Kaiserbad	60	0.31		[134]
Grosse Quelle, Ritzenbad	43	3 16		[134]
Kerekmalom Quelle	20	0.11		[32]
Arpadquelle	23	0 046	0.624	[32]

Source	t°C	$m\mu Cl^{-1}$ Water	Lit.
ITALY			
Sorgente Montirone, Abano near Padua	87	2.05*	[20]
Upper Sulfur Therm, Aqui Piemont	72	0.28*	[20]
Fiuggi, Anticoli		8.02*	[20]
Surgonne Grotta, Battaglia near Padua	74	3.34*	[20]
Acidola, Castellamare	13	9.27*	[20]
Domenico Tricarico, Bagnoli near Naples	52	0.79*	[20]
Purgativo, Agnano near Naples	90	0.79*	[20]
Stabilimento, Porto d'Ischia	65	1.93*	[20]
Manzi I, Cassamicciola, Ischia	85	0.57	[20]
Old Roman Spring, Lacco Ameno, Ischia	57	152 5*	[20]
Fonte di Castello, Santa fiora	12	3.01	[77]
Fonte della Casella, Casteldelpiano	12	1.85	[77]
Acqua dei Bagnoli, Acidoso	14	3.29	[77]
Polla di Sotto, Bagnore	20	1.52	[77]
Sambuco, Montagna	8	2.08	[77]
Baleno Carcaiole, Uliveto		1.09	[75]
Pozzo delle Saline, Salsomaggiore		Gas = 8.6 / 4.41	[76]
Bagni di Casciana		0.0	[77]
Parlanti, Monsummano	31	Gas = 1.8 / 0.064	[92]

* Values obtained by multiplying Mache units by 4.1×10^{-10}.

Source	t°C	mμCl⁻¹ Water
NORWAY (86)		
Nasodden.		17 9
Sandsvar		12 9
Jellum, near Modum....		31.2
Tandberg estate, Simoa Valley		67.4
PORTUGAL (81)		
Sabroso, Sabroso (Vidago)		3.29
Fonte Romana, Fonte Romana		2.05
Da Bica, Ferez		8 20
Das Lamas, Cucos		10.4
RUMANIA (58)		
Orsova		
Hercules, Baile Herculane	46	0 19*
Regina Maria, Baile Herculane	60	0 22
RUSSIA (68)		
Essentuky No. 6, Caucasus		3.5
Batalinsky, Caucasus		0 6
SPAIN (15)		
Rivas, Gerona		0.33
Buitre, Seirra de Fuensante, Murcia		0 05
Garganton y Pisnolon, Sierra de Guadarrama		12 5
La Raja, Mazarron, Murcia		0 46
El Tubo, Mazarron, Murcia		0 48
Posa de Levante, Mazarron, Murcia		0 36
Medica Catalan, Mazarron, Murcia		0.68
SWEDEN (91, 119)		
Slottskallan, Upsala	7	1 8
Bourbrum, Upsala	6	1 55
Birjerjarlsg No. 120, Stockholm	6	14 6
Gamla (spring), Porla	7	1 77
Sofia (spring), Helsingborg	10	3 00
Villastaden (drilled well), Lidingon	8	17 06
Norrb, L. (well), Bodens fastning	5	70 6
Stockh l. (well), Vinterviken	10	67 2
Hermelinsgruf (well), Malmberget	3	2 75
Kalmar, l. (spring), Sodra Vi	6	14 1
Sanatorie parken (spring), Mosseberg	7	0 90

* Emanation content changes with season and even on same day.

Rock formation of source	No. samples	mμCl⁻¹ Water
SWEDEN.—(Continued)		
Boulders, morainal deposits	110	2 40
Diabase	10	0 70
Granite (Archean)	53	13 24
Granite (gneissic)	20	5 66
Granulite	11	10 2
Gray gneiss with granite intrusives.	6	6 11
Gneiss (granitic)	20	2 99
Iron-bearing gneiss	12	9 31
Limestone	42	0.78
Peat	16	1.18
Quartz porphyry	5	2 09
Sandstone	37	2 91
Slate	42	1 11
Syenite and granulitic syenite	15	15 46

Source	t°C	mμCl⁻¹ Water	Lit.
SWITZERLAND			
St. Placidus Spring, Disentis		4 66	(127)
Val Lunpegnia, Disentis	8	3.75	(117)

Source	t°C	mμCl⁻¹ Water	Lit.
Leuk	51	0.12	(127)
Waadt, Lavey		4.51	(117)
Paracelsusquelle, Engadine, St. Moritz.	5	0.57	(117)
Stollenquelle, Pfafers-Ragaz.	36	0.29	(117)
Sotsassquelle, Schuls		0.42	(117)
Carolaquelle, Tarast	7	0.46	(117)
Kurhaus, Acquarossa	25	1.24	(117)
Thomas, Val Sinestra	8	0.26	(117)
Les Trois Pigeons, Valangin		0.24	(80)
Come Girard, Locle		0.26	(80)
Vioulou, Paturage, Locle		0.37	(80)
Eplatures		0.15	(80)

ASIA

Source	t°C	mμCl⁻¹, Water
INDIA (122)		
Kaira District, Bombay		
Hot Spring	67	33.0 to 62.1
Cold Spring	28	33.9

Source	t°C	mμCl⁻¹	
		Water	Gas
JAPAN (42)			
Kami-no-yu, Tamatsukuri	64	1 08	10.18
Kami-no-yu, Misasa	71	51 69	
Kabu-yu, Misasa	45	3.72	22.82
Kaminoyu, Dogo	47	1.45	8.5
Tama-no-i, Dogo	cold	0.39	
Hirano, Tansan-sen	26	0.07	0.21
Gosho-no-yu, Kinosaki	60	3.06	
Ko-no-yu, Kinosaki	57	0.94	
Furosen, Beppu	58	0.07	
Kamigawara No. 1, Masutomi.	22	301.2	
Kuridaira No. 1, Masutomi	16	214 7	550 6
Yunosawa-Onsen, Innai-Yunosawa	41	0.43	
Takinoyu, Noboribetsu	72	0.074	
Yojo-Kwan-no-yu No. 1, Togo.	50	1.12	
Jizo-no-yu, Kusatsu.	57	0 057	0.065
Akakura-Onsen, Akakura.	62	0.43	
Ji-no-yu, Isobe	9	1 55	0.74
Arima-Onsen, Arima	52	0 92	
Maruyama-Kosen, Arima	19	3.01	
Zui-hoji-Onsen, Arima...	31	13 8	
Arifuku-Onsen, Arifuku	43	0.80	
Kizu-no-yu, Asama...	44	0.51	
O-yu, O-yu	57	1.13	trace
Kami-no-yu, Oyu	58	0.4	
Shimo-jyaya-no-yu, Sekigane	44	10.95	
Soto-no-yu, Katsura	29	0.31	
Yuntsumi-no-yu, Atsumi		0.40	
Awazu-Onsen, Awazu.	54	0.35	
Kami-no-moto-yu, Bobata	14	4.35	
Goshiki-Onsen No. 2, Goshiki	39	0.80	
Tsubataya-uchi-yu, Shibu.	48	0.11	
Hie-no-yu, Kaminoyana.	62	0 86	5.5
Shiotsu-no-Tsubo, Katayamazu	79	0.47	8.79
Gosho-no-yu A, Kinosaki.	63	2 67	
Koyabara-Onsen, Koyabara.	38	1 37	2.95
Murasugi-Kosen No. 1	26	18.04	
Osakaya-no-yu, Musashi.	45	1.17	11.8
Shirataki-no-yu, Nakabusa....	60	0.59	
Tsuru-no-yu, Mikko-Yumoto	62	0.85	
Shin-yu, Unzen...	38	0.85	

Source	t°C	mμCl⁻¹	
		Water	Gas
Ogawa-Onsen No. 2	49	1.01	
Omaki-Onsen, Omaka	49	0 48	
Taki-no-yu, Onogawa	70	2 37	
Umeka-no-yu, Owani	62	4 21	
Shigaku-Onsen, Shigaku	47	0 43	0 64
Ena-Kosen, Takayama	10	102 2	
Takarazuka-Tansan-sui, Takarazuka	19	1 20	0 72
Tochiomata-no-yu, Tochiomata	39	9 40	
Wakazaki-no-yu No. 1, Wakura	93	2 52	33 9
Yamanaka-Onsen, Yamanaka	45	0 62	
Yamashiro-Onsen	69	0 25	
Tottori-Onsen, Yoshikata	48	1 19	
Kasuga-Onsen, Teramadu	29	0 22	0 88
Kabu-yu, Yudani	32	1 54	8 65
Sento, Yukiku	67	0 23	3 34
Kabu-yu, Yummra	91	0 31	
Sagi-no-yu, Yunogo	38	0 31	1 95
Taki-no-yu, Yunokawa	50	0 74	8 23
Shinyu, Yunotsu	4	1 8	0 19

Source	t°C	mμCl⁻¹ Water	Lit.
PHILIPPINE ISLANDS			
Sibul Springs, Bulacan		1 28	(135)
Pansol Springs, Laguna		none	(135)
Bambangan Spr., Laguna		0 15	(135)
Adukpung Spr., Kiangan		1 33	(37)
Artesian Well, Batangas		2 11	(135)
Sinaba Spring, Laguna		1 3	(37)
Mairut Salt Spr., Bontoc	100	none	(37)
Salinas Salt Spring, Nueva Vizcaya	31	0 095	(37)

AFRICA			
Source	t°C	mμCl⁻¹ Water	
ALGERIA (85)			
Bains de la Reine, near Oran	50	13 1	
Louiye, A Hammam Bou Hadjar	44	22 4	
Hotel de Vichy, A Bou Hanifia	55	1 3	
d'Alma T'zoumoulal	17	5 3	

THE LITHOSPHERE

Uranium and Thorium Radioactive Minerals

The numbers following the name of the mineral represent weight percent of U, resp. Th. The qualitative chemical composition is indicated in parentheses ( ), the locality in brackets [ ], R = "rare earths;" aq. = "hydrous."

A. Aeschynite: U 0.3, Th 0–20 (RNbTiO$_x$). *Auerlite:* Th 61 (ThSiPO$_x$). *Autunite:* U 50 (UCaPO$_x$aq.).

B. Becquerelite: U 70 (UO$_x$aq.) [Belg. Congo] (111). *Blomstrandite:* U 22 (TaNbUO$_x$).

C. Calciothorite: Th 53 (RCaSiO$_x$aq.). *Carnotite:* U 53 (KUVO$_x$aq.). *Chalcolite:* (See Torbernite). *Clevite:* U 60; Th 4 (UThYO$_x$). *Curite:* U 73 (UPbO$_x$aq.) [Belg. Congo] (106).

D. Dewindtite: U 50 (PbUPO$_x$aq.) [Belg. Congo](108). *Dumontite:* U 56 (PbUPO$_x$aq.) [Belg. Congo] (114).

E. Ebigite: *Flutherite* (See Uranothallite). *Eliasite:* also Pittinite (See Gummite). *Erdmanite:* Th 9 (FeCaThBSiO$_x$). *Euxenite:* (Polycrase) U 5–15 (RNbTaO$_x$aq.).

F. Fergusonite: (Bragite, Tyrite, Yttrotantalite) U 1-7, Th 2–5 (RNbTaO$_x$). *Freyalite:* Th 24 (RThSiO$_x$aq.). *Fritzscheite:* (UMnVO$_x$aq.).

G. Gadolinite: Th < 1 (RO$_x$.SiO$_y$). *Gummite:* (Eliastite, Pittinite) U 60 (UPbCaSiO$_x$aq.).

H. Hatchettolite: U 13 (UCaNbTaO$_x$). *Hokutolite:* (PbBaSO$_4$) [Japan] (42).

J. Johannite: U 56 (CuUSO$_4$.aq.).

K. Kasolite: U 40 (PbU8iO$_x$aq.) [Belg. Congo] (107). *Kochelite* (See Fergusonite).

L. Liebigite: U 31 (UCaCO$_x$aq.).

M. Mackintoshite: U 20; Th 42 (RUThSiO$_x$aq.). *Medjidite:* (A variety of Uranopilite). *Mendeleeffite:* U 20 (UNbTiO$_x$) [Transbaikalia] (129). *Microlite:* U 1.6 (CaTaO$_x$). *Monazite:* Th 7 20 (RPO$_x$).

N. Naëgite: U 2 5; Th 45 (ZrRSiO$_x$) [Japan] (42). *Nivenite:* (See Uraninite). *Nohlite:* (See Samarskite).

O. Orangite: U 1–10; Th 65 (A variety of Thorite).

P. Parsonite: U 32 (PbUPO$_x$) [Belg. Congo] (112). *Phosphuranylite:* U 60 (UO$_x$PO$_x$aq.). *Pilbarite:* (PbUThSiO$_x$aq.). *Plumboniobate:* U 12 (PbUYNbO$_x$). *Pitchblende:* (See Uraninite). *Polycrase:* (See Euxenite). *Priorite:* (See Blomstrandite). *Pyrochlore:* Th 0 6 (RCaNbO$_x$).

R. Randite: (See Voglite). *Rowlandite:* U 0.4 (YSiO$_x$). *Rutherfordine:* U 65 (UO$_x$CO$_x$). *Rutherfordite:* (A variety of Fergusonite).

S. Samarskite: U 1-3 (RUNbTaO$_x$). *Schoepite:* (UO$_x$CO$_x$) [Belg. Congo]. *Schrockingerite:* (A variety of Voglite). *Sipylite:* U 3 (ErNbO$_x$). *Soddite:* U 71 (USiO$_x$aq.) [Belg. Congo] (110). *Stasite:* U 50 (PbOPO$_x$aq.) [Belg. Congo] (109). *Skaldowskite:* U 55 (MgUSiO$_x$aq.) [Belg. Congo] (113).

T. Thorogummite: U 18; Th 36 (UThPbSiO$_x$). *Thorianite:* U 12; Th 65 (RThUO$_x$). *Tritomite:* Th 5–8 (Th, Ce, Ca, Ta, B, F, SiO$_x$). *Torbernite:* U 50 (UCaPO$_x$aq.). *Trögerite:* U 53 (UAsO$_x$aq.). *Tscheffkinite:* Th 1–17 (RFeSiTiO$_x$). *Thysonite:* U 65 (U(OH)$_x$SO$_4$).

U. Uraninite: (Pitchblende) U 65–80; Th 1–8 (UO$_x$RUPbO$_x$). *Uranochalcite:* (A variety of Uranopolite). *Uraconite:* (A variety of Uranopolite). *Uranocircite:* U 47 (BaUPO$_x$aq.). *Uranophane:* U 55 (UCaSiO$_x$aq.). *Uranopolite:* U 64 (UO$_x$CaSO$_4$aq.). *Uranosphaerite:* U 42 (UO$_x$BiOUO$_x$aq.). *Uranospite:* U 49 (UCaAsO$_x$aq.). *Uranothallite:* U 32 (CaUCO$_x$aq.). *Uranothorite:* U 8; Th 52 (ThSiO$_x$).

V. Voglianite: (A variety of Uranopolite). *Voglite:* U 34 (CaCuUCO$_x$aq.).

W. Walpurgite: U 16 (BiUAsO$_x$aq.).

X. Xenotime: U 3; Th 0–2 (YPO$_x$).

Y. Yttrocrasite: U 2; Th 0-8 (YTiO$_x$). *Yttrotantalite:* U 0.5–2 (YNbTaO$_x$).

Z. Zuenerite: U 50 (CuUAsO$_x$aq.).

RADIOACTIVITY OF ROCKS

Ra unit = 10⁻¹² g Ra (element) per g. Th unit = 10⁻⁶ g Th (element) per g

IGNEOUS ROCKS

Name and locality	No. specimens	Ra mean	Lit.
Acidic Intrusives			
Charnockite			
Mysore State, India	3	0.09	(121)
Granite			
Mysore State, India	11	1.03	(121)
Dutch East Indies	5	4.9	(13)
Eisenach, Germany	1	3.5	(67)
Germany	7	9.8	(13)
France(1) Holland(2)	3	8.8	(13)
St. Francois Co., Mo., U. S. A	1	1.5	(100)
Ireland	10	2.0	(46)
Leinster, Ireland	28	1.7	(28)
Th mean =	28	7.0	

Name and locality	No. specimens	Ra mean	Lit.
Antartic region	2	0 4	(29)
Th mean =	2	2 6	
South Sea Islands	2	1 76	(26)
Sumatra(1) Bohemia(1)	2	26 1	(35)
Loetschberg Tunnel, Switz	7	2 3	(83)
Various localities	63	2 7	(48)
	1	1 63	(62)
	11	2 56	(123)
Th mean =	86	20 5	(82)
Monzonite			
Bella Monte, Tyrol, Austria	1	3 5	(13)
Pegmatite			
Mysore State, India	2	4 17	(121)
Porphyry			
Campbell Is., New Zealand	1	2 8	(26)
Various localities	10	2 8	(13)
Quartz			
Germany	3	16 0	(13)
Sumatra	1	1 3	(13)
Syenite			
Borneo and Molucca Island	13	1 58	(13)
Mount Royal, Canada	1	1 1	(25)
Vosges, France	1	13 2	(36)
Norway	3	2.46	(123)
Various localities	8	8.3	(13)
	23	3.9	(48)
Tinguaite			
Mount Royal, Canada	2	3 65	(25)
Tinguaite porphyry			
Germany	2	8 2	(13)

Basic Intrusives

Name and locality	No. specimens	Ra mean	Lit.
Diabase			
Borneo	2	0 85	(13)
Diabases and dolerites	8	1 0	(48)
New Zealand	1	0 43	(26)
Diabase and gabbro			
Germany	5	2 8	(13)
Diorite			
Borneo and Sumatra	4	0 78	(13)
Various localities	8	1 6	(48)
Dolerite			
Isle of Canna, Scotland	1	0 57	(123)
New Zealand	2	0 66	(26)
Dunite			
Loch Scavaig, Scotland	1	0 31	(123)
Essexite			
Mount Royal, Canada	1	0.26	(25)
Gabbro			
New Zealand	2	0 34	(26)
Gabbro and Norite	5	1.3	(48)
Greenstone			
Garrick Du, St. Ives, Eng	1	0 52	(123)
Hypersthenite	1	0 06	(121)
Peridotite			
Isle of Rum, Scotland	1	0.63	(123)
Porphyry			
New Zealand	1	0 99	(26)

Acid Extrusives

Name and locality	No. specimens	Ra mean	Lit.
Ash			
Krakatoa near Sumatra Th mean =	1	9.0	(82)
Kenyte			
Antartic region	4	2 29	(29)
Th mean =	4	12.0	
Lavas			
Various localities	18	3.4	(43)
Th mean =	15	24.0	
Liparite	2	4.7	(13)
Phonolite			
Kirchberg, Germany	1	0.9	(13)
Pitchstone			
Auckland Island, New Zealand	1	1.9	(26)
Dutch East Indies	2	0.6	(13)
Isle of Eigg, Scotland	1	1.53	(123)
Meissen, Germany	1	3.0	(13)
Rhyolite			
Yellowstone Park, U. S. A	6	2 21	(104)
Trachite			
Mt. Erebus, Antartic region	3	2.16	(29)
Th mean =	3	13.0	
Continental Europe	2	3.4	(13)
New Zealand	3	2.11	(26)
Transandine Tunnel	7	0 58	(27)
Th mean =	7	4.4	
Various localities	18	3.0	(48)
Tuff	2	2 9	(46)
Transandine Tunnel	12	0.92	(27)
Th mean =	10	5 87	

Basic Extrusives

Name and locality	No. specimens	Ra mean	Lit.
Anamesite			
Germany	2	1.8	(13)
Andesite			
Borneo and Molucca Is	13	1.58	(13)
Basalt			
Deccans and Antartic	14	2.0	(48)
Mt. Erebus, Antartic region	1	2 13	(29)
Th mean =	1	14 5	
Hebrides (mainly)	11	0 5	(48)
New Zealand	2	1.21	(26)
Various localities	6	0.47	(123, 125)
	6	2 2	(46)
	4	0.35	(126)
Lava			
Antartic region	7	0.58	(29)
Th mean =	7	4.7	
Vesuvius (1631-1906)	7	12.6	(43, 46)
Th mean =	6	53.4	(82)
Limburgite			
Germany	1	2.9	(67)
Melaphyre			
Oberstein, Germany	1	1.9	(13)
Tepharite	3	8.7	(67)
Trap			
Mysore State, India	43	0.21	(121)

METAMORPHIC ROCKS

Name and locality	Ra No. speci-mens	Ra Mean	Th No. speci-mens	Th Mean	Lit.
Amphibolite India					
Mysore State...........	1	0.82			(121)
Gneiss					
Freiburg, Ger...........	1	2.9			(67)
Various localities...	14	2 1	14	8 7	(48, 82)
Gneiss (granitic)					
Tauern Tunnel..........	11	3.41	7	17.7	(62)
Gneiss (porphyritic)					
Tauern Tunnel..........	9	4.34	9	41 0	(62)
Quartzite					
Various localities........			6	3 4	(45)
Villnos Gulch, Austria ..	1	54.7	1	5.79	(133)
Schist					
Lustre, Simplon Tunnel. .			1	10 4	(45)
St. Gothard Tunnel......	33	3.4	33	11.6	(47)
Schist (chlorite)					
Mysore St., India........	1	0 27			(121)
Schist (hornblende)					
Mysore St., India........	11	0.19			(131)
From mines, Mysore St., India................	17	0.25			(121)
Slate					
England...............	2	1.17			(124)
European..............			10	13.5	(45)
Germany...............	2	1.3			(13)
Tauern Tunnel..........	3	2.53	3	24 3	(62)
Slate (mica)					
From well boring, Beach-ville, Can........	1	1 6			(25)

SEDIMENTARY ROCKS

Name and locality	No. speci-mens	Ra mean	Th mean	Lit.
Clay				
Montreal, Canada........	2	1 17		(24)
England....................	3	0.79		(124)
England(1), Germany(1).	2		10 2	(45)
Coal				
Alabama, U. S. A..............	11	0 166		(55)
Lens, France..................	1	0 97	3 3	(74)
Frankenholz..................	1	0 04	0 3	(74)
Coal ash				
Alabama coals.................	11	2.15		(55)
Lens, France	1	8.8	30.	(74)
Frankenholz..................	1	2 0	15.	(74)
Flint				
Terling, Essex, Eng............	1	0 49		(124)
Grauwacke				
Wipperfurth, Germany..........	1		24.	(45)
Limestone				
Beachville, Ont., Can......... ..	6	1 02		(25)
Montreal, Canada..............	2	0 91		(25)
Deccan, India.................	1	0.25		(124)
England.....................	7	1.13		(124)
Germany(2), Ireland(1).........	3		2 3	(44)
New Zealand..................	2	0.37		(26)
Various localities..............	30		0.4	(44)

Name and locality	No. speci-mens	Ra mean	Th mean	Lit.
Limestone (oölithic)				
Yellowstone Park, U. S. A......	2	2.9		(104)
Marble and limestone				
Various localities..............	8	1.3		(13)
Sand (Saxicava)				
Montreal, Canada....	1	0.16		(24)
Sandstone	2	1.04		(124)
From 850 ft. borehole, Baarlo, Limburg, Holland.	8	1.66		(13)
Beachville, Canada. . .	1	0.50		(28)
Various localities	8		6.3	(45)

OCEANIC DEPOSITS

Name and locality	No. speci-mens	Ra mean	Lit.
Blue mud			
1210 fa. E. coast N. Amer	1	3.1	(138)
Calcareous mud			
2225 fa. E. of Society Islands	1	22.2	(138)
Globergina ooze			
1990 fa. Middle S. Atlantic. .. .	2	6.5	(138)
1825 fa. Pacific W. of South America .	1	7.4	(138)
570 fa. W. coast Ireland	2	6.3	(138)
2042 fa. Central Pacific	2	7.6	(138)
Radiolarian ooze			
Central Pacific....................	4	43.9	(138)
Red clay			
2740 fa. N. Atlantic, coast of Africa. ...	4	17.6	(138)
2350 fa. Central Pacific..	3	47.4	(138)
"Salt Lime" (gypsum from evap. sea water)	1	0.016	(130)
Sea Salt....	1	0.07	(124)
From evap. water of high seas	15	none	(40)

SOILS

	No. speci-mens	Ra mean	Lit.
Gravel—fine siftings			
Terling, Essex, Eng...	2	0.65	(124)
Surface loams			
7 localities in E. and S. parts of U. S. ...	7	1.97	(69)
Th mean =	5	4.5	(69)
Subsoils of above	7	1.52	(69)

Highest value for surface soils, 2.88; Lowest, 0.93	(69)
Highest value for subsoil, 3.8; Lowest 0.93	(69)
Loess, Heidelberg, 10.4 × 10⁻⁶ g Th per g	(45)
Mark, Ireland, 1.4 × 10⁻⁶ g Th per g	(45)

Highest value for surface soils, 2.88; Lowest, 0.93 (69)

Highest value for subsoil, 3.8; Lowest 0.93 (69)

Loess, Heidelberg, 10.4×10^{-6} g Th per g (45)

Mark, Ireland, 1.4×10^{-6} g Th per g (45)

ROCKS FROM TUNNELS

Rock and section of tunnel	No. of speci-mens	Units 10^{-12} g Ra per g	Units 10^{-6} g Th per g
The St. Gothard (47)			
Granites and gneiss			
Finsteraarhorn Massif..........	20	6.7	21.5
Altered sediments			
Unsernmulde.................	18	3.8	13.4
Tessinmulde..................	18	2.7	4.8
Schists, etc.			
St. Gothard Massif.	33	3.4	11.6
The Tauern, Austria (62)			
Granitic gneiss..	Ra 10, Th 7	3.41	17.7
Porphyritic granitic gneiss..........	Ra 13, Th 9	4.34	41.0

ROCKS FROM TUNNELS.—(Continued)

Rock and section of tunnel	No. of specimens	Units 10⁻¹² g Ra per g	Units 10⁻⁶ g Th per g
Slate	Ra 3, Th 3	2 53	24 3
The Loetschberg, Bernese Oberland, Switzerland[83]			
Anhydrite	2	3 4	
Aplete	2	2 5	
Granite	7	2 3	
Limestone	16	1 5	
Quartz porphyry	1	2 5	
Quartz sandstone	1	4 3	
Schists			
Feldspathic	3	2 7	
Hornblende	2	3 1	
Lustre	2	3 4	
Mica	2	2 1	
Quartz	12	2 4	
Talc	16	1 5	
(Unclassified)	16	2 5	
The Transandine, Argentine-Chile [27]			
Andesites	Ra 2, Th 1	0 71 / 0 79	4 1 / 5 6
Mean Ratio, Th-Ra = 7 × 10⁶			
Feldspathic Tuff	2	1 24	3 0
Trachytes	7	0 58	4 4
Tuff	Ra 8, Th 7	0 90	6 94

SPRING DEPOSITS

Country, name of spring, location	No. of specimens	Ra content*	Th content†	Remarks	Lit.
Austria					
Elizabethstollen, Gastein	1	2920	3970	Reissacherite	(62)
Rudolphstollen, Gastein	1	417, 300	1988		(62)
Vilnos Gulch	4	75	37 7	A sinter	(133)
England					
Hot Springs, Bath	1	381			(124)
France					
Chomel, Vichy	1	250		Ferruginous	(52)
Hôpital, Vichy	1	700		Black	(52)
Carnot, Santenay	1	1500			(52)
Neris	1	950	5100	Black	(52)
Luxeuil	1	660	1100	Manganous	(52)
Germany					
Badochquelle	1	4		Surface scum	(67)
Ems, Hessen-Nassau	4	0 63	35		(133)
Johanngeorgenstadt, Saxony	3	081	89	Mainly hydromorphite; Range of Ra content, 10–1300	(4)
Italy					
Fiuggi	1	5		Tufa	(84)
Russia	2	13 9	147		(14)
Borzhom Spring	2	13 9	147		(14)
United States					
Hatborn No. 1, Saratoga Springs, N. Y.	1	769			(71)

Country, name of spring, location	No. of specimens	Ra content*	Th content†	Remarks	Lit
Geyser, Saratoga Springs, N. Y.	1	17			(71)
Pump Well No. 4, Saratoga Springs, N. Y.	1	63			(71)
Palace Spring, Hot Springs, Arkansas	1	1724			(99)
Avenue Spring, Hot Springs, Arkansas	1	140			(99)
Horseshoe Spring, Hot Springs, Arkansas	1	2.3			(99)
Various springs, Hot Springs, Arkansas	11	175			(99)
Main Springs, Mammoth Hot Springs, Yellowstone	1	8.8		Travertine	(104)
Hot River, Mammoth Hot Springs, Yellowstone	1	8.1			(104)
Bench Springs, Upper Geyser Basin, Yellowstone	1	0.95			(104)
Fish Cone, West Thumb, Yellowstone	1	0 19			(104)
Fire Hole Lake, Lower Geyser Basin, Yellowstone	1	6 7			(104)
Doughty Springs, Delta Co., Colorado	2	1654			(100)

* Unit, 10⁻¹²g Ra per g.
† Unit, 10⁻⁶g Th per g.

METEORITES

Class and locality	Ra in 10⁻¹² g per g	Remarks	Lit.
Stony			
Dhurmsala, India	0 53		(123)
Coahuila, Coahuila, Mex.	7 69	Normal hexahydrite	(87)
Toluca, Xiquepeleo, Mex.	0 21	Medium octahedrite	(87)
Iron			
Augusta Co., Va., U. S. A.	0 0022	2 specimens	(125)
	none		(123)
Stone			
Various localities	0 75	Mean of 16 Range 2.17–0.073	(87)
Iron			
Various localities	0 69	Mean of 2	(87)
	none	Mean of 3	(87)

NATURAL GASES

Source and Locality	No. samples	Milli-micro-Curies (10⁻⁹ Curies) Ra per liter	Lit.
Canada			
Medicine Hat, Alberta	3	0.064	(97)
Suffield-Brooks Calgary	6	0.064	(97)

Source and Locality	No. samples	Millimicro-Curies (10⁻⁶ Curies) Ra per liter	Lit.
3 British Columbia wells		0 47	(97)
Brant, Anondoga, Ontario	4	0 42	(97)
Tilbury, Ontario		0 016	(97)
England			
Marsh gas, environs of Cambridge	10	0 3	(95)
France			
Alsace		7 1	(17)
Germany			
Nuengamme, Hamburg		0 24	(17)
Hungary			
Well No. 14, Bazna		0 043	(17)
Japan			
Well No. 22, Takiya		0 035	(42)
Rumania			
Well No. 103, Campina			(17)

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Ashman, *12*, **26**: 119; 08 (²) Bamberger, *75*, **116**: 1473; 07 (³) Bamberger, *75*, **117**: 1055; 08 (⁴) Bamberger and Weissenberger, *75*, **123**: 2065; 14. (⁵) Barnard (Schlundt), *266*, **16**: 228, 13 (⁶) de Beaujeu, *34*, **153**: 944; 11. (⁷) Becker, *93*, **131**: 209, 23. (⁸) Besson, *34*, **147**: 818, 08. (⁹) Boltwood, *12*, **20**: 128; 05

(¹⁰) Borne, *200*, **2**: 77, 142, 05 (¹¹) Boyle and McIntosh, *69*, **7**: 163; 13 (¹²) Brochet, *34*, **150**: 145, 291, 10. (¹³) Büchner, *200*, **10**: 516, 23 (¹⁴) Burkser, *53*, **47**: 21, 14 (¹⁵) del Castillo and de Rada, *132*, **11**: 236; 12. (¹⁶) Curie and Laborde, *34*, **142**: 1162, 06 (¹⁷) Czako, *93*, **82**: 249; 13. (¹⁸) Danne and Crémieu, *34*, **153**: 870; 11 (¹⁹) Dole, *U S Geol. Survey, Mineral Resources*, **2**: 393, 13

(²⁰) Engler and Sieveking, *93*, **53**: 1; 07 (²¹) Eve, *3*, **14**: 724; 07 (²²) Eve, *5*, **16**: 622; 08. (²³) Eve, *69*, **4**: 55; 10. (²⁴) Eve and McIntosh, *3*, **14**: 231; 07. (²⁵) Eve and McIntosh, *69*, **4**: 69; 10. (²⁶) Farr and Florance, *3*, **18**: 812; 09. (²⁷) Fletcher, *3*, **20**: 36; 10 (²⁸) Fletcher, *3*, **21**: 102, 11. (²⁹) Fletcher, *3*, **21**: 770; 11.

(³⁰) Fletcher, *5*, **23**: 279, 12. (³¹) Fresnius and Czapski, *136*, **35**: 722, 11 (³²) Frohner, *144*, **30**: 387; 12. (³³) Frommel, *93*, **53**: 1, 07 (³⁴) Gerard and Chauvin, *34*, **157**: 302; 13 (³⁵) Gockel, *"Die Radioaktivität von Boden und Quellen"* (1914). (³⁶) Gockel, *200*, **7**: 487; 10. (³⁷) Heise, *267*, **12**: 293; 17. (³⁸) Henrich, *63*, **8**: 112. 07. (³⁹) Henrich, *97*, **33**: 5, 13, 20; 20.

(⁴⁰) Hewlett, *268*, **22**: 173; 17. (⁴¹) Ishizu, *"The Mineral Springs of Japan"* (1915). *269*, **16**: 362; 13. (⁴²) Jentzsch, *63*, **8**: 887, 07 (⁴³) Joly, *3*,

(⁴³) *18*: 577; 09. (⁴⁴) Joly, *3*, **20**: 125; 10. (⁴⁵) Joly, *3*, **20**: 353; 10. (⁴⁶) Joly, *3*, **22**: 134; 11. (⁴⁷) Joly, *3*, **23**: 201; 12. (⁴⁸) Joly, *3*, **24**: 694; 12. (⁴⁹) Kinoshita, Nishikawa and Ono, *3*, **22**: 821; 11.

(⁵⁰) Koch, *63*, **7**: 806; 06. (⁵¹) Kulhöreter, *86*, **14**: 356; 12. (⁵²) Laborde and Lepape, *34*, **155**: 1902; 12 (⁵³) Lepape, *34*, **176**: 1702; 23. (⁵⁴) Lester, *18*, **44**: 621, 18. (⁵⁵) Lloyd and Cunningham, *11*, **50**: 47; 13. (⁵⁶) Loisel, *34*, **173**: 921; 21. (⁵⁷) Loisel and Castelnau, *34*, **173**: 1390; 21. (⁵⁸) Loisel and Michaelesco, *34*, **175**: 1054; 22. (⁵⁹) Ludwig, *63*, **26**: 180, 24.

(⁶⁰) Mache, *75*, **113**: 1329, 04 (⁶¹) Mache, *75*, **132**: 207; 24. (⁶²) Mache and Bamberger, *75*, **122**: 325; 14. (⁶³) Mache and St. Meyer, *75*, **114**: 355; 05 (⁶⁴) Makower, *135*, **106**: 135; 12 (⁶⁵) Massol, *34*, **156**: 373; 12 (⁶⁶) Mauchly, *268*, **29**: 187, 24. (⁶⁷) Meyer, *190*, **1**: 184; 16. *870*, **20**: 1, 14 (⁶⁸) Meaermtaku, *55*, **43**: 244, 11 (⁶⁹) Moore, *45*, **6**: 370; 14.

(⁷⁰) Moore and Kithl, *88*, No **70**: 13 (⁷¹) Moore and Whittemore, *45*, **6**: 552, 14. (⁷²) Moureu, *871*, **16**: 65; 11. (⁷³) Moureu and Lepape, *34*, **148**: 834, 09 (⁷⁴) Moureu and Lepape, *34*, **156**: 598; 14. (⁷⁵) Nasini, Marino, Ageno and Porlezza, *36*, **43**: 321, 13. (⁷⁶) Nasini and Porlezza, *24*, **76**: 725; 17 (⁷⁷) Nasini, Porlezza and Sborgi, *24*, **73**: 1572; 14. (⁷⁸) Oluje, *200*, **18**: 158; 18 (⁷⁹) Perkins, *166*, **43**: 806; 15.

(⁸⁰) Perret and Jacquerod, *149*, **45**: 277, 336, 418, 18. (⁸¹) Pinto, *C. R. Cong Int Radiol and Elect*, **1**: 79, 11. (⁸²) Poole, *3*, **29**: 483; 15. (⁸³) Poole, *3*, **40**: 466, 20 (⁸⁴) Porlezza and Norzi, *82*, **20**: 935; 11. (⁸⁵) Pouget and Chouchak, *34*, **177**: 1112, 1319, 23. (⁸⁶) Poulsson, *137*, No. **8**; 14 (⁸⁷) Quirke and Finkelstein, *18*, **44**: 237; 17. (⁸⁸) Ramsay, *135*, **106**: 134, 12 (⁸⁹) Ramsay, *18*, **40**: 309, 15.

(⁹⁰) Ramsay, *3*, **30**: 815, 25 (⁹¹) Sahlbom, *19*, **6**: 1; 16. (⁹²) Salvadori, *216*, **12**: 151; 23 (⁹³) Satterly, *3*, **20**: 1, 10. (⁹⁴) Satterly, *801*, **16**: 540, 10 (⁹⁵) Satterly, *201*, **16** 356, 11 (⁹⁶) Satterly and Elworthy, *Dept of Mines, Canada, Bull* **16** (1917). (⁹⁷) Satterly and McLennan, *69*, **12**: 153, 18 (⁹⁸) Schiffner, Weidig and Friedrich, *"Radioactiven Wässer in Sachsen"* (1912). (⁹⁹) Schlundt, *78*, **12**: 247; 07. *135*, **96**: 199, 07

(¹⁰⁰) Schlundt, *78*, **26**: 163, 14. (¹⁰¹) Schlundt, *278*, **16**: 1245; 09. (¹⁰²) Schlundt, *50*, **18**: 662, 14 (¹⁰³) Schlundt and Moore, *50*, **9**: 320; 05. (¹⁰⁴) Schlundt and Moore, *166*, No **395**; 09. (¹⁰⁵) Schmidt and Kurz, *63*, **7**: 209, 06 (¹⁰⁶) Schoep, *34*, **173**: 1186, 21. (¹⁰⁷) Schoep, *34*, **173**: 1476, 21. (¹⁰⁸) Schoep, *34*, **174**: 623, 22. (¹⁰⁹) Schoep, *34*, **174**: 875; 22. (¹¹⁰) Schoep, *34*, **174**: 1066, 22. (¹¹¹) Schoep, *34*, **174**: 1240; 22. (¹¹²) Schoep, *34*, **176**: 171, 23 (¹¹³) Schoep, *34*, **179**: 413; 24. (¹¹⁴) Schoep, *34*, **179**: 693, 24. (¹¹⁵) Scholl, *U of Ill Bul, Water Survey Series*, **16**: 111, 17 (¹¹⁶) Schwoidler, *75*, **129**: 919, 20. (¹¹⁷) Schwietzer, *149*, **27**: 256; 09 **30**: 46; 10 (¹¹⁸) Shrader, *2*, **3**: 339; 14. (¹¹⁹) Sjögren and Sahlbom, *19*, **3**: 1, 08

(¹²⁰) Skinner and Sale, *45*, **14**: 949; 22. (¹²¹) Smeeth and Watson, *3*, **35**: 206, 18 (¹²²) Steichen, *3*, **21**: 401, 16 (¹²³) Strutt, *5*, **77A**: 472; 06. (¹²⁴) Strutt, *5*, **78A**: 150, 06. (¹²⁵) Strutt, *5*, **80A**: 572; 08. (¹²⁶) Strutt, *5*, **84A**: 377; 10. (¹²⁷) de Sury, *149*, **24**: 101; 07. (¹²⁸) Ssilard, *34*, **154**: 982, 12. (¹²⁹) Versadsky, *34*, **176**: 993; 23.

(¹³⁰) Wade, *1*, **29**: 941; 07 (¹³¹) Watson and Pxl, *3*, **26**: 44; 14 (¹³²) Wellik, *77*, **117** 1191; 08. (¹³³) Weissenberger, *189*, **1914**: 481. (¹³⁴) Wenzelszky, *144*, **30**: 345; 12. (¹³⁵) Wright and Heise, *267*, **12**: 145; 17. (¹³⁶) Wright and Smith, *2*, **5**: 159, 15. (¹³⁷) Zlatarovic, *75*, **129**: 59; 20. (¹³⁸) Joly, *3*, **16**: 190, 18. (¹³⁹) Behounck, *51*, **6**: 48; 25.

AGES OF MINERALS AND ROCKS BASED ON RADIOACTIVE CHANGES

Roger C. Wells

There are a number of ways of estimating the ages of minerals by combining chemical and radioactive data, all based on the assumption that the law of each radioactive change is expressed by its constant, λ, over the periods and for the quantities of each element involved. The two principal methods employ the ratios of helium to uranium and thorium and of lead to uranium and thorium. The helium ratio is admitted to give minimum values on account of the loss of helium with lapse of time; and the lead ratio involves the assumption, or actual proof by means of an atomic weight determination, that the lead is wholly of radioactive origin. Associated rocks are generally assumed to be as old or older than the minerals found in them. Attempts have also been made to calculate the ages of rocks from determinations on bulk samples (Russell).

For the two methods mentioned the fundamental changes and data are:

$$U(238.17) \rightarrow Pb_U(206.06) + 8He(4.00)$$
$$Th(232.15) \rightarrow Pb_{Th}(208.00) + 6He(4.00)$$

One gram of uranium in equilibrium with its products gives 9.4×10^4 alpha particles per sec (15) or 1.96×10^{-11} gram He and 1.26×10^{-10} gram Pb$_U$ per year.

One gram of thorium in equilibrium with its products gives 2.7×10^4 alpha particles per sec, or 5.5×10^{-12} gram He and 4.8×10^{-11} gram Pb$_{Th}$ per year.

The ages of minerals may be calculated from the analytical data and the preceding information by simple proportion in the case of helium (equation 1) and also in the case of lead with sufficient accuracy for most purposes (equation 2), but if the percentage of lead is relatively large the theoretical relation is given by equation 3, where U, Th, Pb = percentage U, Th, Pb in the mineral.

$$(1) \quad Age = \frac{cm^3\ He/g}{U + 0.28Th} \times 910\ \text{million years}$$

(2) $Age = \dfrac{Pb}{U + 0.38Th} \times 7900$ million years

(3) $Age = \dfrac{\log (U + 0.38Th + 1.156Pb) - \log (U + 0.38Th)}{6.5 \times 10^{-5}}$

million years

Thorium minerals with Th/U greater than 3 are secondary

and younger than uranium minerals from the same geolo horizon[19]. Low lead ratios have little significance on accou of the ease with which certain minerals abstract lead from cir lating natural waters. The atomic weight of the lead should determined whenever possible in order to make certain that t lead is of radioactive origin. In general, only primary minerals suitable for age determinations.

Ages of Minerals from Helium Ratios by Equation (1)

(The values in parenthesis are calculated from the lead ratios for comparison)

Mineral	Geologic horizon	He cm^3/g	U Percent	Th Percent	Age million years	Lit
Phosphatic shark's teeth, Florida	Pliocene	1.7×10^{-6}	0.021	0	0.07	(23
Phosphatic shark's teeth, Felixtowe, Eng	Pliocene	1.6×10^{-6}	0.013	0	0.11	(23
Phosphatic nodules, Felixtowe, Eng	Pliocene	1.0×10^{-6}	0.0041	0	0.22	(23
Carnotite, Montrose Co., Colo.	Post Tertiary	0.01	2.53	0	3.6	(23
Zircon, Campbell 1, New Zealand	Tertiary	8.1×10^{-4}	0.029	0.07	1.5	(23
Pitchblende, Joachimsthal		0.107	62.4	0	1.6	(23
Sphaerosiderite, Germany	Oligocene	1.65×10^{-6}	0.00015	0.00017	7.6	(23
Zircon, Mayen, Eifel	Tertiary	1.14×10^{-4}	0.0108	0.00073	9.4	(23
Hematite, Co. Antrim, Ireland	Eocene	1.21×10^{-5}	0.00022	0.00073	26	(23
Zircon, Auvergne	Tertiary	2.12×10^{-4}	0.031	0	6.2	(23
Phosphatic nodules, Cambridge, Eng	Upper Cretaceous	3.0×10^{-5}	0.0091	0	3.0	(23
Phosphatic nodules, Bedfordshire	Lower Cretaceous	2.1×10^{-5}	0.0049	0	3.9	(23
Zircon, Cheyenne Canon, Colo.	Paleozoic	0.0193	0.109	0.10	128	(23
Hematite, Cumberland, Eng	Above Carboniferous	1.6×10^{-4}	0.0011	0	130	(26
Limonite, Forest of Dean	Carboniferous	1.5×10^{-4}	0.00087	0.00043	140	(23
Sipilite, Little Frier Mt., Va	Carboniferous (?)	0.59	2.42	4.33	147	(23
Euxenite, Arendal, Norway	Pre-Cambrian	0.73	2.41	2.39	210(1240)	(23
Samarskite, Mitchell Co., N. C	Carboniferous (?)	1.5	8.73	1.28	160	(23
Phosphatic nodules, Bala, England	Silurian	1.5×10^{-4}	0.0028	0	49	(23
Phosphatic limestone, Chirbury, Shropshire, Eng	Silurian	5.6×10^{-6}	0.0067	0	76	(23
Uraninite, Katanga	Pre-Silurian	8.88	77.76	0	104(665)	(4
Zircon, Brevig, Norway	Post-Devonian	0.0099	0.113	0.288	46	(23
Hematite, Caen	Devonian	9.8×10^{-5}	0.00037	0.0013	120	(23
Zircon, Green River, N. C	Paleozoic	0.0255	0.11	0.264	126	(23
Zircon, Ural Mts.	Paleozoic	0.030	0.0538	0.408	160	(23
Uraninite, Colo.	Tertiary	0.15	72.62		18(58)	(11
Uraninite, N. C.	Post-Cambrian	2.96	77.0	2.44	34(380)	(11
Thorianite, Sab. Province, Ceylon	Pegmatite in Charnokite Series	1.5	9.87	63.54	50(460)	(5
Thorianite, Galle Province, Ceylon	Pegmatite in Pre-Cambrian	9.3	20.6	57.55	230(400)	(23
Uraninite, Änneröd	Pre-Cambrian (?)	9.4	66.2	5.27	120(890)	(11
Uraninite, Portland, Conn.	Devonian (?)	19.2	72.0	8.79	230(290)	(11
Uraninite, Branchville, Conn.	Silurian (?)	21.0	74.3	5.72	250(400)	(11
Microlite, Amelia Court House, Va	Carboniferous (?)	0.05	1.60	0	280	(23
Cuprouranite, Cornwall	Devonian	0.10	50.9	0	1.8	(23
Orangite, Brevig, Norway	Middle Devonian	0.11	0.85	42.6	7.9(22)	(23
Zircon, Ural Mts.	Paleozoic	0.030	0.053	0.409	160	(23
Thorianite, Ceylon	Balangoda series	8.9	11.0	67.7	270(500)	(23
Zircon, Kimberly	Paleozoic	0.032	0.091	0.012	310	(23
Phosphatic nodules, Loch Broom	Pre-Cambrian	8.3×10^{-6}	0.084	0	9.0	(23
Gadolinite, Ytterby	Pre-Cambrian (?)	2.43	2.50	7.56	480	(23
Aeschynite, Ural Mts		0.98	2.12	7.19	210	(23
Cyrtolite, Llano Co., Texas	Pre-Cambrian (?)	1.15	3.11	4.44	240	(23
Uraninite, S. Dak	Pre-Cambrian (?)	4.35	66.90	1.89	59(540)	(4
Zircon, Ceylon	Ancient	0.0283	0.086	0.010	290	(23
Zircon (?), Renfrew Co., Ontario	Archaean	0.0114	0.0155	0.0008	660	(23
Aeschynite, Hitteroe, Norway		1.09	7.98	1.11	1200	(23

AGES OF MINERALS FROM LEAD RATIOS BY EQUATION (3)

Mineral	Geologic horizon	Pb Percent	U Percent	Th Percent	Th/U	Age million years	Lit.
Carnotite, Montrose Co., Colo	Tertiary	0.17	45.6			20	(11)
Johannite, Colo	Tertiary	0 76	47 2			123	(16)
Brannerite, Idaho	Tertiary	0 18	46 97	4.1	0.11	20	(9)
Uraninite, Gilpin Co., Colo	Tertiary	0 65	72 60			69	(11)
Thorite, Ceylon	Young mineral in pegmatite in Pre-Cambrian	2 86	72.00	8.70	0.12	280	(11)
Hatchettolite, Hybla, Ont	Pre-Cambrian (?)	0 50	13 72	0 46	0 03	270	(14)
Polycrase, Brasil	Pre-Devonian	0 59	5 49	4 59	0 84	600	(8)
Allanite, Blueberry Mtn., Mass	Young mineral in pegmatite	0 036	0 11	2 01	18 3	310	(17)
Freyalite, Brevig, Norway	Post-Devonian (Lawson)	0 0028	0 0526	6 330	120.3	8.8	(19)
Tritomite, Brevig, Norway	Post-Devonian (Lawson)	0 0026	0 0631	5 150	81.6	9.9	(19)
Thorite, Brevig, Norway	Post-Devonian (Lawson)	0 0196	0 4072	29 20	71 7	13.3	(19)
Thorite, Brevig, Norway	Post-Devonian (Lawson)	0 0810	0 7200	49 43	68.6	32.0	(19)
Thorite, Brevig, Norway	Post-Devonian (Lawson)	0 0760	0.7000	47 25	67.5	31 4	(19)
Orangite, Brevig, Norway	Post-Devonian (Lawson)	0 0570	1 2437	49 44	39 7	22.1	(19)
Orangite, Brevig, Norway	Post-Devonian (Lawson)	0 0542	1.1825	45 03	38 1	22.8	(19)
Homolite, Brevig, Norway	Post-Devonian (Lawson)	0 0121	0 2442	2 000	11.9	69.1	(19)
Mosandrite, Brevig, Norway	Post-Devonian (Lawson)	0 0024	0 0432	0 287	6 64	112	(19)
Eudidymite, Brevig, Norway	Middle Devonian	0 0007	0 0090	0 036	7 00	230	(19)
Eucolite, Brevig, Norway	Middle Devonian	0 0012	0 0170	0 040	2.35	280	(19)
Thorite, Brevig, Norway	Middle Devonian	0 4279	10 1040	14 20	1.41	210	(19)
Zircon, Brevig, Norway	Middle Devonian	0 0055	0 1460	0 114	0 78	220	(19)
Zircon, Brevig, Norway	Middle Devonian	0.0085	0 1941	0 082	0 42	280	(19)
Pyrochlore, Brevig, Norway	Middle Devonian	0 0093	0 1855	0 075	0.40	330	(19)
Aegerine, Brevig, Norway	Middle Devonian	0 0015	0 0253	0 007	0 28	400	(19)
Zircon, Brevig, Norway	Middle Devonian	0 0370	0 9310	0 141	0 15	280	(19)
Biotite, Brevig, Norway	Middle Devonian	0 0069	0 1602	0.017	0 11	310	(19)
Uraninite, Spruce Pine, N. C	Post-Cambrian (?)	3 90	77 01	2 44	0.03	380	(11)
Thorianite, Galle Province, Ceylon	Pegmatite in Pre-Cambrian	2 41	24.13	55 95	2 32	400	(19)
Betafite, Madagascar	Pegmatite, uncertain	0 35	22 58	0 98	0.04	120	(16)
Thorianite, Sa. Province, Ceylon	Pegmatite in Pre-Cambrian	2 09	9 87	63 54	6.45	460	(6, 19)
Uraninite, Branchville, Conn	Silurian (?)	4 03	73 00	6 09	0.81	400	(11)
Uraninite, Katanga	Pre-Silurian	6 51	77 76	0		620	(4)
Polycrase, Slättåkra, Sweden		0.85	8 45	3 08	0.36	650	(2)
Uraninite, Ånneröd, Norway	Pre-Cambrian (Moss district)	8.39	66.21	5 28	0.08	890	(11)
Uraninite, Elvestad	Pre-Cambrian (Moss district)	9.35	65.82	7.46	0.11	970	(11)
Ånnerödite	Pre-Cambrian (Moss district)	2 22	15 25	2.08	0.14	990	(2)
Mackintoshite, Llano Co., Tex	Pre-Cambrian (?)	3.47	19 75	39.83	2.02	730	(1)
Yttrocrasite, Llano Co., Tex	Pre-Cambrian (?)	0 45	2 28	7.69	3 38	640	(1)
Uraninite, Llano Co., Tex	Pre-Cambrian	9 43	56 45	6 65	1.18	1130	(1)
Uraninite, Llano Co., Tex	Pre-Cambrian	9 35	55.18	5 88	1.07	1150	(1)
Yttrialite, Llano Co., Tex	Pre-Cambrian	0 74	1 45	9 53	6.5	1040	(1)
Yttrialite, Llano Co., Tex	Pre-Cambrian	0 79	0 69	10.55	15.3	1190	(1)
Fergusonite, Ytterby, Sweden	Middle Pre-Cambrian	0 18	1 06			1200	(1)
Gadolinite, Ytterby, Sweden	Middle Pre-Cambrian	0 36	2 41			1100	(1)
Zircon, Ceylon	Pre-Cambrian	0 092	0.56	0 01	0.02	1150	(14)
Uraninite, Villeneuve, Quebec	Middle Pre-Cambrian	10.46	64.74	6 41	1 00	1110	(11)
Uraninite, Parry Sound, Ontario	Middle Pre-Cambrian	10 83	69.19	2 83	0.04	1090	(6)
Uraninite, Arendal, Norway	Pre-Cambrian (Arendal district)	10 16	61.27	3 65	0.06	1150	(11)
Uraninite, Black Hills, S. Dak	Pre-Cambrian	15.24	66.90	1 89	0 03	1540	(4)

LITERATURE

(For a key to the periodicals see end of volume)

(1) Barrell, *336*, **26**: 745, 17   (2) Blomstrand, in Dana's Mineralogy, p 741.   (3) Boltwood, *12*, **23**: 77; 07.   (4) C W Davis, U S Bureau of Mines, *0*.   (5) Dunstan and Blake, *5*, **76A**: 253, 05   (6) Ellsworth, *12*, **9**: 127, 25   (7) Geiger and Rutherford, *5*, **20**: 691, 10   (8) Hess and Henderson, *143*, **200**: 235, 25   (9) Hess and Wells, *143*, **189**: 225; 20.   (10) Hidden and Mackintosh, *12*, **36**: 481, 89

(11) Hillebrand, *12*, **43**: 390, 91   *156*, No **78**; 91.   (12) Hillebrand and Ransome, *12*, **10**: 120, 00   (13) Holmes, *5*, **85A**: 248; 11.   (14) Holmes

and Lawson, *3*, **26**: 823; 14.   (15) Kovarik and McKeehan, *357*, No. **81**; 25.   (16) Lacroix, *Mineralogie*, **V**: 93.   (17) Lane, *Trans. Lake Superior Mining Inst.*, Aug, 1925.   (18) Larsen and Brown, *323*, **2**: 75; 17   (19) Lawson, *75*, **126**: 721; 17.   (20) Russell, *Introduction to the Chemistry of Radioactive Substances*, Murray, London, 1922.

(21) Rutherford, *Radioactive Substances and Their Radiations*, Putnam's Sons, New York, 1913.   (22) Soddy and Hyman, *4*, **105**: 1402, 14   (23) Strutt, *5A*, **76**: 88, 05.   **50**: 572, 08   **51**: 272; 08.   **52**; 166, 09.   **53**; 96, 298, 09   **84**: 377; 10   (24) Walker, *Univ. Toronto Studies*, No **16**; 23.   (25) Wells, *Bibliography on the relation of radioactivity to geological problems*, National Research Council, 1924.

SELECTED PHYSICAL PROPERTIES OF STARS AND NEBULAE

Alfred H. Joy

Contents.—(A) Classification of stellar and nebular spectra; (B) Stellar temperatures, masses, and densities; (C) Stellar diameters. (Data pertaining to the solar spectra will be found with other spectroscopic data; consult index)

A. CLASSIFICATION OF STELLAR AND NEBULAR SPECTRA

The system[1] is that developed at Harvard College Observatory, as used by Miss Cannon in the Henry Draper Catalogue. Except where the exact nature of the spectral changes is not fully understood, decimal sub-classes, representing progressive steps toward the succeeding class, are used. In denoting objects by their catalogue numbers, the following abbreviations are used: B. D. = Bonn Durchmusterung; C. D. M. = Cordoba Durchmusterung; I. C. = Dreyer's Index Catalogue of nebulae and clusters; N. G. C. = New General Catalogue by Dreyer. The number, or numbers, following the abbreviation is the catalogue designation of the object.

Class P includes practically all the gaseous nebulae. Its unique characteristic is the appearance of lines from an unknown origin (nebulium). In addition there are many lines of H, He, C, He+, C+, and N+. All lines are bright and usually sharp. (The order of the Harvard (2) subdivisions should probably be reversed to indicate decreasing intensity of radiation)

Class	Typical object	Spectral criteria
Pa	I. C. 418	λ5007 and λ4959 faint, λ3869 not seen
Pb	Orion nebula	λ5007 and λ4959 stronger
Pc	I. C. 4997	λ4363 conspicuous
Pd	N. G. C. 6826	λ5007 and λ4959 strong
Pe	N. G. C. 7662	λ4686 present
Pf	N. G. C. 40	λ4686 strong

Wright (11) has divided these spectra into three classes: Class I, having λ4686 present, Class II, with λ4686 absent but λ3869 present, and Class III with both λ4686 and λ3869 absent.

Class O is distinguished by the presence of the Pickering series of ionized helium, upon a strong continuous spectrum with maximum intensity far in the violet. The elements present are H, He, He+, C+, N+, Mg+, O+, CIII, NIII, SiIII, OIII, SiIV. Broad emission bands occur in the earlier subdivisions. Few absorption lines are found in sub-classes Oa, Ob, Oc, which make up the group known as Wolf-Rayet stars. (The Harvard subclasses Od, Oe, and Oe5 which have absorption lines and in some cases narrow emission lines as well, are included in the subclasses O5 to O9 as suggested by H. H. Plaskett (7), the basis of classification being the absorption lines.)

[1] Adopted by International Astronomical Union. It defines a temperature scale which is linear within the present errors of measurement

Class	Typical object	Spectral criteria
Oa	B D. +35° 4013	Band λ4648 stronger than λ4686
Ob	B. D. +35° 4001	λ4686 stronger than λ4648
Oc	C. D. M. −41° 10972	Bands narrower. λ4686 twice λ4638
O5	B. D. +1° 1302	Pickering series very strong. H lines weak, λ4634 and λ4640 (NIII) present
O6	B D. +44° 3639	Neutral helium appears
O7	9 Sagittae	λ4471 (He), 1.4 × λ4541. λ4089 (SiIV), 0.8 × λ4097 (NIII)
O8	λOrionis	λ4481 (Mg+) appears
O9	10 Lacertae	H stronger, He weak. λ4471, 2.7 × λ4541. λ4089, 1.4 × λ4097

Class B is characterized by the presence of helium, which has its maximum intensity in B2. The principal elements are those of class O, with the addition, in the later sub-classes, of lines of the ionized atom of several of the metals, such as Sr, Ba, and Fe. The H and K lines of calcium are found in increasing strength in this class. The hydrogen lines increase through the sub-classes, reaching a strong maximum at Ao of the following class.

Class	Typical object	Spectral criteria
B0	ζ Orionis	Pickering series weak, λ4649 (OII), λ4116 (SiIV), and λ4089 (SiIV) maximum intensity
B1	β Canis Majoris	He more prominent than O and Si.
B2	γ Orionis	λ4116 not seen. λ4089 and λ4649 faint
B3	η Aurigae	Strongest lines are helium
B5	q Tauri	λ4128 and λ4131 (SiII) stronger than λ4121 (He). λ4481, 0.7 × λ4471
B8	β Orionis	λ4481 equal to λ4471
B9	λ Aquilae	H strong. He weak. Several prominent enhanced metallic lines

Classes A, F, G, K and M, which contain the largest numbers of the stars, show a gradual increase in the number and intensity of the lines of neutral metallic elements of the lower atomic weights, and a decrease in the intensity of lines due to ionized elements. Compounds produce bands in the later classes. The sun's spectrum is Go, and is intermediate between that of the white and the red stars.

Class	Typical object	Spectral criteria
Ao	α Lyrae	H maximum strength. Very few other lines except λ4481 (Mg+)
A5	ρ Sagittarii	K (Ca+) stronger than Hδ. λ4290 well marked. λ4481 weaker
Fo	σ Bootis,	K 3.0 × Hδ and equal to H + Hε

Class	Typical object	Spectral criteria
F5	α Canis Minoris	Fraunhofer band G first seen. Numerous solar lines
Go	α Aurigae	Solar type. H not conspicuous. G band well defined, Hδ = λ4226.
G5	η Piscium	Hγ fainter than λ4325
Ko	α Bootis	G band conspicuous, λ4226 strong. Hydrogen weaker
K5	α Tauri	λ4226 very wide. λ4254 and λ4274 (Cr) strong. Titanium bands very faint
Mo	β Andromedae	Titanium bands well marked
M5	α Herculis	Titanium bands very stong. Metallic lines fewer

Class R and N stars show the carbon bands in increasing strength. The more advanced stars of class N have very little light in the violet or blue portions of the spectrum. They are the reddest stars known. Typical stars: Class R, B. D. −10° 5057; Class N, 19 Piscium.

Class S spectra resemble those of class K5 except for the presence of bands of zirconium, and other peculiarities in the region near λ4650. The line λ4554 of Ba + is conspicuous

Class Q stars are the novae. Near maximum of outburst their spectra are characterized by numerous wide emission bands of hydrogen and helium, and by absorption lines of ionized elements, especially titanium and iron. As the star decreases in light, both absorption and emission lines of N and O become more prominent In the later stages, bright nebular bands appear; these are ultimately superseded by the bright bands of the Wolf-Rayet spectrum.

B. STELLAR TEMPERATURES, MASSES, AND DENSITIES

Giant stars are characterized by large mass, low density, and great total luminosity. Dwarf stars have smaller mass, higher density, and less total luminosity. Both are found in all classes, but the greatest contrasts between the two are found in the cooler stars of classes K and M. The continuous spectrum of dwarfs has its maximum shifted towards the violet, as compared with that of giants of the same spectral class, indicating that their absolute temperature is about 15% higher than that of the giants. Even with small dispersion, pronounced differences between giants and dwarfs may be noticed in the distribution of intensity in their line spectra. These differences probably arise from differences in the density gradients; they show a correlation with the absolute magnitude and mass of the stars. The low densities of giants favor the enhancement of those lines (absorption) which are produced under conditions of high excitation, such as the spark lines of the metals; the high density of dwarfs favor those produced by low excitation, such as the resonance lines of neutral atoms. The lines λ4077, λ4215 (ionized Sr) are much strengthened in giants, and weakened in dwarfs; the reverse is true of λ4226 (Ca), λ4454 (Ca), λ4607 (Sr).

STELLAR TEMPERATURES, MASSES AND DENSITIES

Units: Temperature, 1000°C abs.; Mass, Mass of Sun; Density, g/cm³.

Class	Effective temperature (giants*)					Mean mass (*)		Mean density (*)	
	A†	P‡	C§	S‖	F¶	Giants	Dwarfs	Giants	Dwarfs
Oa		23		23					
O5					30	50 (6)			
Bo		20	13	18	19	10			
B3					16	9		0.22	
B8	16					7	3	0.24	
Ao	14	11	8	12	10	7 0	6 0	0.16	0 36
A5		9				5 6	4 0	0.071	0.40
Fo		7 5		9	7 5	4 3	2 5	0 025	0 40
F5	6	7 2	6			3 2	1.5	0 0078	0.39
Go	5.8	6 5	6	7	6	2 6	1 0	0 0025	0 68
G5		4 5				2.8	0 76	0 000 87	1 2
Ko		3 7	4		4 5	3 0	0 68	0.000 18	1 3
K5	3	3 5	3 5		3 9	2 6	0 62	0.000 026	1 4
Mo		3	3	5	3	2 0	0 59	0.000 0096	5.4
M5	2 5	2 95		4					
N		2 3							

* Temperatures of dwarfs are 10 % to 20 % higher than giants of same class (indirect methods)

† Abbot ([1]). By radiometer

‡ Potsdam observations Wilsing *et al* ([10])

§ Coblentz ([3]) By thermocouple

‖ Saha ([8]) Calculated from initial appearance of certain spectral lines under pressure of 0 1 atmosphere (See note ¶)

¶ Fowler and Milne ([4]) Calculated from maximum intensity of certain spectral lines under pressure of 1 31 × 10⁻⁴ atmospheres, assuming 10 000° corresponds to maximum of Balmer lines of H These temperatures, and those of Saha, are for the reversing layer, true effective temperature is somewhat higher

STELLAR DIAMETERS

Unit: Linear Diameter, 10⁶ km.

Star	Class	Parallax	Diameter	
			Angular*	Linear
α Tauri	K5	0.055″	0 022″	60
α Orionis	M2	0 019	0 044	347
α Bootis	Ko	0 088	0 022	37
α Scorpii	M1	0 017	0.040	353

* Measured by means of interferometer ([5])

LITERATURE

(For a key to the periodicals see end of volume)

([1]) Abbot, *21*, **60**. 105, 24. ([2]) Cannon, *Harvard College Obs. Annals*, **76**: 19, 16. ([3]) Coblentz, *37A*, **17**: 725; 22. ([4]) Fowler and Milne, *Monthly Notices, R A S*, **83**: 403, 23. ([5]) Michelson and Pease, *21*, **53**: 249; 21. Pease, *Publ Ast Soc Pacific*, **33**: 171, 204, 21. **34**: 346; 22. ([6]) J. S. Plaskett, *Publ Domin Astrop Obs*, **2**: 298; 24 ([7]) H. H Plaskett, *Ibid.*, **1**: 366, 22 ([8]) Saha, *5*, **99**: 151, 21 ([9]) Seares, *21*, **55**: 202, 22. ([10]) Wilsing, Scheiner and Münch, *Publ Astrop Obs. Potsdam*, **24**: 21; 19. ([11]) Wright, *Publ Lick Obs*, **13**: 262; 18.

DISTRIBUTION OF STARS

FREDERICK H. SEARES

Restriction.—No account is here taken of globular star clusters nor of stars included in spiral nebulae, many of which contain objects whose essentially stellar character can no longer be doubted.

Apparent Distribution and Number.—Statistically considered, the stars are distributed over the face of the sky with a high degree of regularity, their numbers gradually increasing as the Milky

Way is approached from either side. The Milky Way defines what is very nearly a plane of symmetry, and for a first approximation, systematic difference between the two hemispheres, progressive changes in galactic longitude, and all local irregularities can be ignored. The resulting mean distribution, as found by Seares and van Rhijn, is shown in Table 1.

To apparent magnitude (see p. 39) $m = 13.5$ the results depend on data covering a large portion of the sky. From $m = 13.5$ to 18.5 they are derived from counts of stars on photographs of the 139 Selected Areas of Kapteyn between the North Pole and declination $-15°$. For still higher values of m, the values of log N_m are extrapolated, but the uncertainty consequent to the extrapolation itself is probably small. Excepting in low galactic latitudes, there is little or no systematic uncertainty arising from the particular choice of fields used for the counts. To $m = 16$ the magnitude scale is the mean of several closely accordant determinations made at different observatories, and is probably accurate within a few hundredths of a magnitude. Below this limit the scale depends wholly upon observations made at the Mount Wilson Observatory. Although this part of the scale has not been confirmed by independent measures made elsewhere, it

has been established by methods successfully used for the brighter stars.

The indicated total, to the twenty-first photographic magnitude, of all stars in the sky is 890 000 000, and to the twentieth visual magnitude, 1 000 000 000. Barring losses of light by absorption, scattering etc., the increase in log N_m for a uniform distribution of stars throughout space would be 0.6 per unit of magnitude. The observed increase nowhere attains this value; the stars thin out with increasing distance from the sun, and 'at great distances they thin out more rapidly than near the sun; these changes are most pronounced in the direction of the poles of the Milky Way. If the law of decreasing space density indicated by the stars accessible to observation holds for those beyond present telescopic reach, the total number of luminous stars in the galactic system must be of the order of 3×10^{10}.

TABLE 1.—LOGARITHMS OF NUMBERS (N_m) OF STARS, OF MAGNITUDES LESS THAN m, PER SQUARE DEGREE IN DIFFERENT GALACTIC LATITUDES [1]

Units: Last column; m = visual magnitude; average N_m = 1, if m = 8. Other columns; m = international photographic magnitude [2]; N_m = 1, if m = 8, Lat. = 0. Galactic pole: R. A. 12ʰ41ᵐ20ˢ, Dec. +27° 21' (1875) (Gould).

m	Log₁₀ N_m at latitude														Log₁₀ (average N_m) between latitudes				
	0°	5°	10°	15°	20°	25°	30°	35°	40°	50°	60°	70°	80°	90°	0°-20°	20°-40°	40°-90°	0°-90°	0°-90°(v)
4.0	2.19	2.17	2.12	2.05	3.99	3.93	3.87	3.82	3.78	3.74	3.71	3.69	3.67	3.66	2.12	3.73	3.94	3.94	2.11
4.5	2.42	2.40	2.35	2.28	2.22	2.16	2.10	2.05	2.01	3.97	3.94	3.92	3.90	3.88	2.35	2.11	3.96	2.17	2.35
5.0	2.65	2.63	2.58	2.51	2.45	2.39	2.33	2.28	2.24	2.20	2.17	2.15	2.13	2.12	2.58	2.34	2.19	2.40	2.60
5.5	2.88	2.86	2.80	2.74	2.68	2.62	2.56	2.51	2.47	2.43	2.40	2.38	2.36	2.34	2.80	2.57	2.41	2.63	2.83
6.0	1.11	1.08	1.03	2.97	2.90	2.84	2.79	2.74	2.70	2.65	2.62	2.60	2.58	2.57	1.03	2.80	2.64	2.85	1.07
6.5	1.33	1.31	1.26	1.19	1.13	1.07	1.01	2.97	2.92	2.88	2.85	2.83	2.80	2.79	1.26	1.03	2.86	1.08	1.31
7.0	1.56	1.53	1.48	1.42	1.35	1.29	1.21	1.19	1.15	1.10	1.07	1.05	1.02	1.01	1.48	1.25	1.09	1.30	1.54
7.5	1.78	1.76	1.70	1.64	1.57	1.52	1.46	1.41	1.37	1.32	1.29	1.27	1.24	1.23	1.70	1.47	1.31	1.52	1.77
8.0	0.00	1.98	1.92	1.86	1.79	1.74	1.68	1.64	1.59	1.54	1.51	1.48	1.46	1.44	1.92	1.69	1.53	1.74	0.00
8.5	0.23	0.20	0.14	0.08	0.01	1.95	1.90	1.85	1.81	1.76	1.73	1.69	1.67	1.65	0.14	1.91	1.74	1.96	0.23
9.0	0.45	0.42	0.36	0.29	0.22	0.17	0.12	0.07	0.03	1.98	1.94	1.90	1.88	1.86	0.36	0.13	1.96	0.18	0.45
9.5	0.67	0.64	0.57	0.50	0.44	0.38	0.33	0.28	0.24	0.19	0.15	0.11	0.08	0.06	0.58	0.34	0.16	0.39	0.68
10.0	0.89	0.85	0.79	0.72	0.65	0.59	0.54	0.50	0.45	0.40	0.35	0.30	0.28	0.26	0.79	0.55	0.37	0.60	0.90
10.5	1.10	1.07	1.00	0.93	0.86	0.80	0.75	0.70	0.66	0.60	0.55	0.50	0.47	0.45	1.00	0.76	0.57	0.81	1.11
11.0	1.32	1.28	1.21	1.14	1.06	1.01	0.96	0.91	0.86	0.80	0.74	0.69	0.65	0.64	1.22	0.96	0.76	1.02	1.32
11.5	1.53	1.49	1.42	1.34	1.27	1.21	1.16	1.11	1.06	0.99	0.92	0.87	0.84	0.82	1.43	1.17	0.95	1.22	1.53
12.0	1.74	1.70	1.63	1.54	1.47	1.41	1.36	1.30	1.25	1.18	1.11	1.05	1.01	1.00	1.63	1.36	1.14	1.42	1.74
12.5	1.96	1.91	1.83	1.75	1.67	1.61	1.55	1.49	1.44	1.36	1.28	1.23	1.18	1.17	1.84	1.56	1.32	1.62	1.94
13.0	2.16	2.12	2.04	1.95	1.87	1.80	1.74	1.68	1.62	1.54	1.46	1.39	1.35	1.33	2.04	1.75	1.50	1.82	2.14
13.5	2.37	2.32	2.24	2.14	2.06	1.99	1.92	1.86	1.80	1.71	1.62	1.56	1.51	1.49	2.24	1.93	1.67	2.01	2.34
14.0	2.57	2.52	2.43	2.34	2.24	2.17	2.10	2.03	1.97	1.88	1.78	1.72	1.67	1.65	2.44	2.11	1.83	2.20	2.52
14.5	2.77	2.72	2.63	2.52	2.43	2.34	2.27	2.20	2.14	2.04	1.94	1.87	1.82	1.80	2.63	2.29	1.99	2.38	2.71
15.0	2.96	2.91	2.82	2.71	2.60	2.51	2.44	2.36	2.30	2.19	2.09	2.01	1.96	1.94	2.82	2.45	2.14	2.56	2.89
15.5	3.15	3.10	3.01	2.89	2.77	2.68	2.60	2.52	2.45	2.34	2.24	2.15	2.10	2.08	3.01	2.62	2.29	2.73	3.07
16.0	3.33	3.28	3.19	3.07	2.94	2.84	2.75	2.67	2.60	2.48	2.37	2.29	2.23	2.21	3.19	2.77	2.43	2.90	3.24
16.5	3.51	3.46	3.37	3.24	3.10	2.99	2.90	2.81	2.74	2.61	2.50	2.42	2.36	2.34	3.37	2.92	2.56	3.07	3.40
17.0	3.68	3.64	3.54	3.41	3.26	3.14	3.04	2.95	2.87	2.74	2.63	2.54	2.48	2.46	3.54	3.07	2.69	3.23	3.56
17.5	3.85	3.81	3.71	3.57	3.41	3.28	3.17	3.08	3.00	2.86	2.75	2.66	2.60	2.57	3.70	3.20	2.81	3.39	3.71
18.0	4.01	3.97	3.87	3.73	3.56	3.42	3.30	3.20	3.12	2.98	2.86	2.77	2.71	2.68	3.86	3.34	2.93	3.54	3.86
18.5	4.16	4.12	4.03	3.88	3.70	3.55	3.42	3.32	3.23	3.08	2.97	2.88	2.82	2.79	4.02	3.46	3.04	3.68	4.00
19.0	4.32	4.28	4.18	4.02	3.84	3.67	3.54	3.43	3.34	3.19	3.08	2.98	2.92	2.89	4.17	3.59	3.14	3.82	4.13
19.5	4.46	4.42	4.32	4.16	3.97	3.79	3.65	3.53	3.44	3.29	3.17	3.07	3.01	2.98	4.31	3.70	3.24	3.96	4.26
20.0	4.60	4.56	4.46	4.29	4.09	3.90	3.75	3.63	3.53	3.38	3.26	3.16	3.10	3.07	4.45	3.81	3.33	4.09	4.38
20.5	4.74	4.69	4.59	4.42	4.21	4.01	3.85	3.72	3.62	3.46	3.34	3.25	3.18	3.15	4.58	3.91	3.42	4.21	
21.0	4.87	4.82	4.72	4.54	4.33	4.11	3.94	3.81	3.70	3.54	3.42	3.33	3.26	3.22	4.71	4.01	3.50	4.33	

Distribution of Intrinsic Brightness.—The range in intrinsic brightness among stars is enormous—at least twenty magnitudes, corresponding to an intensity ratio of 100 000 000 to 1. A knowledge of the frequencies of different luminosities among the stars in a given volume of space is essential (unless questionable assumptions are to be introduced) for the calculation of the space distribution of the stars. It is, however, difficult to obtain, and,

at present, the frequencies are but imperfectly known. By assuming that the mean parallaxes of stars of apparent magnitude m and proper motion μ can be represented by a linear function of m and log μ supposed to be valid for all magnitudes and proper motions, Kapteyn and van Rhijn derived for the distribution of the absolute magnitudes a Gaussian error curve whose ordinates are given in the second column of Table 2. Seares [4] has shown

that their adopted mean parallax formula does not represent the distances of the stars of large motion and faint apparent magnitude, all of which are of low luminosity. A revision of the parallax formula, still only provisionally determined, and a recalculation of the luminosity function from about 500 stars of large proper motion leads to the frequencies in the third column of Table 2.

TABLE 2.—APPROXIMATE LUMINOSITY FUNCTION

$\phi(M)$ = number of stars, absolute magnitude M, per cubic parsec in the neighborhood of the sun. Unit of distance for M is 10 parsecs. 1 parsec = 3.26 light years = 30.8 × 10^{12} km.

M	10 + Log₁₀ φ(M)		Diff.
	Kapteyn v. Rhijn (3)	Seares (4)	
−4.64	2.61		
−3.64	3.42		
−2.64	4.17		
−1.64	4.85		
−0.64	5.46	5 58	0 12
+0.36	6.00	6.16	0.16
1.36	6 47	6.66	0.19
2.36	6.88	7.05	0 17
3.36	7.21	7.34	0.13
4.36	7.47	7.58	0 11
5.36	7.67	7.74	0 07
6.36	7.80	7.84	0 04
7.36	7.85	7.87	0 02
8.36	7.84	7.86	0 02
9.36	7.76	7.88	0.12
10.36	7.61	7.92	0 31
11.36	7.39	8.06	0 67
12 36	7.10	8.11	1 01
13 36	6.75	8.11	1 36
14.36	6 3	8 13	1 8

For the stars of low luminosity, the departure of Seares' curve from the error curve, shown by the differences in the fourth column, is important and must be accepted as real, although quantitatively the results are still very uncertain. The possibility of a maximum within the range of absolute magnitude considered is not excluded, but any such maximum must be well below the Kapteyn-van Rhijn limit, $M = 7.7$. Since the frequencies of stars of very low luminosity are still unknown, it is impossible at present to express the luminosity function as a true frequency function.

Space Distribution of Stars.—The space distribution is defined by a density function, preferably in a form expressing the total number of stars per unit volume at different distances from the sun. At present, however, we must be content with so expressing the number of stars which are brighter than some limit of absolute magnitude.

Analytically, the problem is to determine the density function, $\Delta(\rho)$, from the integral equation

$$\frac{dN_m}{dm} = \omega \int_0^\infty \phi(M)\Delta(\rho)\rho^2 d\rho$$

where the left hand member can be found from the data in Table 1; ω is a constant, ρ = distance from sun. Since $\phi(M)$, for $M > 8$, is still very uncertain, the general solution cannot be found at present. Values of the density for the neighborhood of the sun (Table 3) can, however, be calculated incidentally in deriving the data in Table 2. Results in the second column of Table 3 ($M = 7.86$) are in good agreement with similar results by Kapteyn and van Rhijn; the other tabular values indicate what is to be expected for lower limiting values of M. The uncertainty of the luminosity function for $M > 8$ scarcely justifies the effort required to complete the table.

TABLE 3.—AVERAGE NUMBER OF STARS, BRIGHTER THAN ABSOLUTE MAGNITUDE M, PER CUBIC PARSEC AT DISTANCE ρ FROM SUN (4)

Unit of ρ is 1 parsec; of distance for M, 10 parsecs. 1 parsec = 3.26 light years = 30.8 × 10^{11} km.

Log₁₀ ρ \ M	7.86	8 86	9 86	10.86	11.86	12 86	13.86	14.86
0.9	0 028	0 035	0.042	0.050	0.060	0 073	0.087	0.098
1 1	.026	033	.040	.048	.058	.069	.078	
1 3	.024	030	035	.041				
1 5	.023	.028	033					
1 7	.022							
1.9	.020							
2 1	.017							
2 3	014							
2.5	.011							
2 7	.008							
2 9	004							

(Values based upon $\phi(M)$ for stars near the sun, and on the assumption that the relative frequencies of M are the same at all distances.)

Average densities for the whole sky give a very imperfect picture of the real distribution in space, as the latter varies greatly with galactic latitude. Broadly speaking, the surfaces of equal space density are concentric, and approximately similar, ellipsoids of revolution, similarly situated, with axes in the ratio of about 5 to 1. See Table 4.

TABLE 4.—RADII OF EQUIDENSITY ELLIPSOIDS(5)

$\Delta(\rho)$ = number of stars per cubic parsec at distance ρ from sun. (Values require revision for recent star counts (Table 1) and for error in luminosity function (cf. Table 2)).

Unit of radius = 1 parsec. 1 parsec = 3.26 light years = 30.8 × 10^{11} km. Latitude is galactic.

Δ(ρ)	Latitude	
	90°	0°
1.00	0	0
0.63	118	602
0.40	198	1010
0 25	296	1510
0 16	413	2106
0 100	553	2820
0 063	717	3656
0 040	902	4600

Size of the Galactic System.—At present we have no certain indication as to the distance of the most remote stars belonging to the galactic system; but if ordinary blue stars of absolute magnitude zero occur among the faintest objects listed in Table 1, the diameter of the system cannot be less than a million light years. Such objects are not to be expected in high galactic latitudes, where the stars of very faint apparent magnitude are almost certainly all dwarfs; but their occurrence in the Milky Way is by no means excluded. We have, indeed, strong, though not conclusive, evidence of the existence in the Milky Way of stars of zero absolute magnitude among those of the sixteenth apparent magnitude. The corresponding diameter of the system is a hundred thousand light years. This value may be accepted with some assurance as a lower limit for the size of the system in the plane of the Milky Way, exclusive of such objects as globular star clusters and spiral nebulae, whose relation to the general stellar system about us is not yet clearly defined.

Position of the Sun.—The symmetrical distribution of stars adopted in Table 1 tacitly assumes the sun to be at the center of the system. This is not actually the case, as is shown by systematic deviations from the adopted mean distribution. Shapley's (6)

value for the distance of the sun from the galactic plane is about 60 parsecs, to the north, which is certainly of the right order of magnitude. The sun's distance from the center is much less certain, and different estimates range from a few hundred to many thousand parsecs, according to the underlying assumptions and the method of attack. The question is much complicated by the fact that the sun lies within a local cluster whose members form a considerable fraction of the stars of the brighter apparent

magnitudes, and a final answer must await the detailed discussion of the distribution of faint stars in galactic longitude.

LITERATURE

(For a key to the periodicals see end of volume)

(¹) Seares and van Rhijn, *197*, **11**: 358; 25; a more detailed account appears in *21*, **42**: 320; 25. (²) *Trans. Internat. Astronomical Union*, **1**: 69; 28. (Stand-ard magnitudes of stars.) (³) Kapteyn and van Rhijn, *21*, **52**: 23; 20 (⁴) Seares, *21*, **59**: 310; 24. (⁵) Shapley, *21*, **49**: 333, 19. (⁶) Kapteyn, *21*, **55**: 302, 22.

DISTRIBUTION OF NEBULAE

Frederick H. Seares

The term nebula is applied to objects of such diversity of form, size, distance, and physical characteristics that any study of their distribution presupposes a consideration of the question of classification. The following general classification by Hubble provides for two mutually exclusive divisions, characterized by position in the sky as well as by physical peculiarities, and five sub-classes representing physical differences.

A GENERAL CLASSIFICATION OF NEBULAE

I. Galactic nebulae, characterized by (1) tendency to concentrate about the Milky Way, (2) conspicuous association with individual stars from which they probably derive their luminosity, (3) early-type spectra, either emission or absorption, depending upon the spectral type of the associated stars, and (4) smooth and cloudy or wispy texture. They include

(a) *Planetaries,* distinguished by symmetrical distribution of nebulosity about central stars, sharply defined edges, and emission spectra.

(b) *Diffuse nebulae,* clouds in low galactic latitudes, usually associated with early-type stars. This type ranges from luminous to dark and from semi-transparent to opaque. Subdivided into predominantly luminous, predominantly obscure, and conspicuously mixed

II. Non-galactic nebulae, characterized by (1) tendency to avoid the Milky Way, (2) no conspicuous association with stars, (3) late-type absorption spectra, and (4) usually a rotational symmetry about dominating non-stellar nuclei. They include

(a) *Elliptical nebulae,* amorphous objects whose forms can be represented as successive stages of an original globular mass flattening under the influence of increasing rotation

(b) *Spirals of two kinds, logarithmic and barred,* which, once formed, appear to develop along parallel lines, the arms unwinding and the granulation of the material becoming more and more conspicuous.

(c) *Irregular nebulae,* including a few non-galactic objects having no dominating nuclei and, significantly, showing no rotational symmetry.

Physically, the planetaries and diffuse nebulae, Ia and Ib, are distinct and apparently without genetic relationship, except that the planetaries, which, in some cases at least, seem to be late stages in the development of novae, may represent the catastrophic consequences of the penetration of a star within a nebulous cloud of the diffuse sub-class. The spirals IIb, on the other hand, are apparently an evolutionary development from elliptical nebulae, IIa, although it does not follow that all elliptical nebulae will necessarily become spirals. The few irregular nebulae, IIc, present features that might be expected in the case of spirals in the absence of or through the neutralization of dominating dynamical characteristics.

The distribution of the various classes of nebulae is not in general easily shown in tabular form. The following summary for each of the important sub-classes includes, however, references to diagrams which exhibit the main features of the distribution.

Ia. Planetary Nebulae.—In the whole sky only about 150 of these objects are known, many of which are so small as to be recognizable only from their gaseous emission spectra. The smallest objects are closely associated with the Milky Way, and show a marked concentration in the Aquila-Sagittarius region. With increasing size the mean galactic latitude increases, and the largest known objects, to the extent of a dozen or so, are scattered over the sky with some approach to uniformity (³, ⁶, ¹¹). This suggests that the linear distances of planetaries from the galactic plane are relatively small and that their angular diameters are correlated with their distances from the sun. Very small nebulae thus appear in low galactic latitudes because their distances from the sun are many times their distances from the galactic plane.

The actual distances of planetary nebulae are still very uncertain. Van Maanen (¹⁵) has measured the parallaxes of about 20 of these objects and finds distances ranging from 50 to a few hundred parsecs; but, as he points out, these values are in conflict with the fact that the radial velocities average about 30 km/sec, while the proper motions are apparently small, of the order of the parallaxes themselves.

Ib. Diffuse Nebulae.—The distant star clouds of the Milky Way define the galactic circle. A secondary galaxy, inclined some 12° to the galactic circle proper, is outlined by the bright helium stars of the much-flattened local cluster immediately surrounding the sun, most of whose members are within 500 parsecs (¹⁴). The diffuse nebulae outside the Magellanic Clouds, some hundreds in all,¹ are closely associated with the primary and secondary galactic circles (⁷). Since the mean galactic latitude of those following the primary galaxy is only about 2°, and since the space within the two circles is not well filled, the inference is that these nebulae are directly connected either with the Milky Way star clouds or with the local cluster, and that few are to be found in the intervening regions. We thus have a group of diffuse nebulae whose members are within a few hundred parsecs of the sun; the others, forming a widely scattered group associated with the Milky Way, are at distances probably to be counted in thousands of parsecs (¹⁰). Both groups include both luminous and dark nebulae; the luminous members of the two groups present somewhat different physical characteristics, most marked in their spectra, which may be either emission, or predominantly continuous or absorption in type. The continuous and absorption spectra occur mostly among the nearer objects connected with the local cluster. The luminous diffuse nebulae are conspicuously associated with stars of high temperature from which they derive their luminosity, either by excitation or reflection.

II. Non-galactic Nebulae.—The members of this class, consisting chiefly of the related sub-classes, elliptical nebulae (IIa) and spirals (IIb), are far more numerous than the galactic nebulae. On the whole, the elliptical nebulae out number the spirals many times; but if only bright objects are considered, the spirals are the more numerous. The distribution in galactic latitude is shown in

¹ Less than 200 luminous ones known; no complete list published (*p.* ⁷, ⁸); Most complete list of dark nebulae (182 small objects) is given by Barnard (⁹).

Table 1, which gives to limiting magnitude 18.6 on the international photographic scale the average number per square degree at various latitudes in each hemisphere. The data are compiled from Fath's list ([4]), based on Mount Wilson photographs (exposure time 1 hour with 60-inch reflector) of the 139 Selected Areas between the North Pole and declination −15°. That part of the northern galactic hemisphere within which nebulae are frequent is wholly covered. About one-half the southern hemisphere is included, but not the south pole itself. Fath's counts have been corrected for losses caused by poor definition in the corners of the negatives ([13]).

TABLE 1.—NON-GALACTIC NEBULAE: NUMBER PER SQUARE DEGREE([4])

Average number; international photographic magnitude ≤ 18.6; cf. Table 2 .

Galactic latitude	Hemisphere	
	N	S
5°	0 2	0 0
15	0 8	0 4
25	2 5	5 4
35	13 2	8 2
45	10 3	5 8
55	12 2	7 0
65	22 2	11 9
74	31	
83	(68)	

Fath's list includes all classes of nebulae, but the galactic nebulae are relatively so infrequent that it is practically one of non-galactic nebulae alone. These objects begin to appear at about 20° latitude and increase rapidly in the interval 20° to 35°. From 40° to 70° the numbers increase slowly. The concentration near the north galactic pole is very pronounced. Below latitude 70° the numbers in the southern hemisphere average about three-fourths those of the northern. The assumption of a similar ratio for the regions 70° to 90° leads to integrated totals of 170 000 and 128 000 for the northern and southern hemispheres, a round total of 300 000 for the whole sky (limiting phot. mag. for stars 18.6).

The summary in Table 2 emphasizes the dependence of the distribution on galactic latitude. The uncertainty in the average number per square degree in the region 70°–90° is considerable, and since the number of nebulae in this region is large (29% or 50 000 in the northern hemisphere), the total given for the whole sky is in doubt by many thousand. Curtis ([2]) has estimated the total (to an undetermined limiting magnitude) to be over 700 000. The difference in the estimates may arise from a difference in magnitude limits or from the fact that the fields counted by Curtis are not certainly representative of the sky as a whole

TABLE 2.—DISTRIBUTION OF NON-GALACTIC NEBULAE

Lat. = interval in galactic latitude. Sky = % area of sky. Neb. = % number of nebulae. N = northern, S = southern hemisphere.

Lat.	Sky	Neb.	
		N	S
0°–30°	50	7	15
30 70	44	64	56
70 90	6	29	29

The distribution of non-galactic nebulae is not, however, simply one of galactic latitude. Data collected by Hardcastle and Hinks ([5]) and by Reynolds ([12]) show marked irregularities in longitude, which seem to depend on the angular diameters of the nebulae. Thus objects with diameters > 10' are almost all in the hemisphere including galactic longitudes 50° to 230°. For diameters 5' to 10' the northern galactic hemisphere shows high frequencies in longitude 110° and 260° 270°, which become even more marked for diameters 2' to 5'. For still smaller nebulae, the distribution is again different. Fath's counts, including mostly very small and faint nebulae, show a band of high frequency crossing the northern galactic hemisphere approximately in longitudes 50° and 220°, with other irregularities suggesting a very complicated distribution

Nothing is known directly of the distances of elliptical nebulae, but their relationship with the spirals is so intimate that the distances of the two sub-classes must be regarded as of the same order Van Maanen's measures ([14]) of internal motion in spirals suggest distances of the order of 3000 to 30 000 light years. The application of Shapley's period-luminosity relation by Hubble ([9]) to numerous typical Cepheid variables discovered by him in the spirals Messier 31 (the Andromeda nebula) and Messier 33 leads to distances of about a million light years for these two objects. The applicability of the period-luminosity relation is assumed, but several lines of corroborative evidence strongly support the larger value of the distance. It is probable, however, that the zero point of the period-luminosity relation requires revision by an amount which would reduce these distances by about 40%.

LITERATURE

(For a key to the periodicals see end of volume)

(1) Barnard, 21, 49: 1, 19 (also consult index of other volumes). (2) Curtis, Publ Lick Obs 13: 15; 18 (3) Curtis, Ibid , 13: 60, 18. (4) Fath, Astronom Jour 28: 75, 14. (5) Hardcastle and Hinks, Monthly Notices, R. A. S. 74: 699, 14. (6) Hinks, Ibid , 71. 694; 11. (7) Hubble, 21, 56: 162; 22. (8) Hubble, 21, 56: 400, 22. (9) Hubble, Pop Astronomy 33: 252; 25. Observatory 48: 139, 25
(10) Lundmark, Publ Astron Soc Pacific, 34: 40; 22. (11) Perrine, 21, 46: 177, 17 (12) Reynolds, Monthly Notices, R A S. 81: 129, 20. 83: 147; 23 84: 76, 23 (13) Searcs, 21, 62: 108, 25. (14) Shapley, 21, 49: 311; 19 (15) van Maanen, Mt Wilson Contribs Nos. 237 (1922), 270 (1923), 280 (1925) (16) van Maanen, 21, 57: 274; 23.

MOTIONS OF THE STARS AND NEBULAE

GUSTAF STRÖMBERG

The *proper motion* of a star is defined as the angular motion, per year, referred to a certain fundamental system of apparently bright stars distributed uniformly over the sky The *radial motion* is determined by the Doppler shift for spectral lines of known wave-length. If the distance to a star is known, the three velocity-components of its *space-velocity* can be determined Proper motions and radial velocities are in general referred to the sun as origin, by correction for the periodic changes due to the earth's motion. The proper motions are in general very small; for the majority of the stars they are below 0 1″ per year. The largest proper motion is that of Barnard's star R. A. 17[h]

53 0[m], Dec. +4° 28′, (1900 0), which moves 10.27″ per year. The radial velocities are mostly below 40 km/sec, the largest being that of the variable star V X Herculis, which approaches the sun with a velocity of 390 km/sec. The spiral nebulae have even higher velocities, the highest being 1800 km/sec, recession, (N. G. C. 584).

SOLAR MOTION

The sun's motion relative to the stars can be determined either from proper motions, from radial velocities, or from space-velocities. The point in the sky towards which the sun is moving is called the sun's *apex*.

TABLE 1.—SOLAR APEX AND THE SUN'S VELOCITY

(Referred to apparently bright stars. Unit· velocity, km/sec)

R. A. 1900	Dec. 1900	Velo-city	Method	No. of stars	Lit.
18ʰ 03ᵐ	+34.3°		Proper Motions P. G. C.*	5413	(²)
18 11	+31.6		Proper Motions m <6.0†	4041	(⁵)
17 56	+32.3		Proper Motions P. G. C.	5943	(⁸)
17 54	+25.3	19.5	Rad. Vel. Lick Obs.	1193	(³)
18 2	+28 6	19.8	Rad. Vel. B to M	1596	(⁶)
18 4	+29.2	21.5	Rad. Vel. F to M	1405	(⁹)
18 11	+36.9	18.8	Space Vel. Giants	800	(¹⁰)
18 43	+20.5	31 7	Space Vel. Dwarfs	415	(¹⁰)
18 40	+32	29	Space Vel. of nearby stars	83	(⁷)

* Preliminary General Catalogue by L. Boss, Washington, 1910
† Stars brighter than the 6th magnitude (apparent).

Although the agreement between the different determinations is fairly good, a detailed study shows that the sun's motion can not be regarded as a constant vector. The A stars and giant stars in general give a small velocity for the sun; and dwarf stars, a much higher velocity.

AVERAGE PECULIAR MOTIONS OF THE STARS

After the effect of the sun's motion has been removed, the residual or "peculiar" velocities show certain regularities. The average peculiar velocities are different for stars of different spectral types, and vary also with the intrinsic brightness of the stars.

TABLE 2.—AVERAGE RESIDUAL RADIAL VELOCITIES (θ) OF STARS OF DIFFERENT SPECTRAL CLASSES (Sp) AND ABSOLUTE MAGNITUDES (M)

Unit of θ = 1 km/sec

Sp	M*	θ	Lit.	Sp	M*	θ	Lit.
O5 to O9	−3	20 7	(¹¹)	K	+1	18 4	(¹)
B	−1	6 5	(³)	K	+6	27 0	(¹)
A	+1	11 0	(¹¹)	M	+1	21.6	(¹)
F	+2	15.8	(¹)	M	+9	29.6	(¹¹)
G	+1	18 0	(¹)	Me†	0	40 1	(¹¹)
G	+5	26 3	(¹)	P‡	−	28 6	(¹¹)

* The apparent magnitude as observed from a distance of 10 parsecs
† Contains M stars with bright hydrogen-lines, all are variable stars of long period.
‡ Bright-line nebulae.

PREFERENTIAL MOTION

The peculiar velocities of the stars are not distributed at random. In general the stars show a tendency to move parallel to the galactic plane. To describe the distribution of the peculiar velocities, a distribution-function is adopted, which gives the relative numbers of stars moving in different directions and with different velocities. The simplest distribution-function is the spherical distribution-law,

$$F(xyz) = \frac{N}{(2\pi)^{\frac{3}{2}}\sigma^3}\, e^{-\frac{x^2+y^2+z^2}{2\sigma^2}}$$

where x, y, and z are the velocity-components referred to the "centroid" of the group. N is the number of stars in the group, and σ is the dispersion or the square-root of the mean of the squares of the velocity-components. The number of stars of velocity-components between $x \pm \frac{1}{2}dx, y \pm \frac{1}{2}dy, z \pm \frac{1}{2}dz$ is then given by $F(xyz)\,dxdydz$. In a spherical distribution, the frequency of a velocity is independent of its direction and only dependent upon its size. Spherical velocity-distributions occur for several classes of stars, but in general the distribution in

velocity-space is either flattened (B stars) or elongated (A, F, and dwarf stars). Two functions have been used to describe the elongated distribution. Kapteyn and Eddington have used a sum of two spherical functions and have regarded the stars as belonging to two intermingled systems, "two stream hypothesis." Schwarzschild has introduced the ellipsoidal distribution defined by the distribution-function

$$F(xyz) = \frac{N}{(2\pi)^{\frac{3}{2}}abc}\, e^{-\left(\frac{x^2}{2a^2}+\frac{y^2}{2b^2}+\frac{z^2}{2c^2}\right)}$$

with three principal dispersions a, b, and c, which define the three axes of the "velocity-ellipsoid." The velocity-components x, y, and z are here projected on the principal axes of this ellipsoid. The major axis of the velocity-ellipsoid corresponds to the line joining the two centers in the two stream theory. The direction of this fundamental axis, which is common in the two theories, is about R. A. 6ʰ 6ᵐ, Dec. + 9°, (true vertex). The dwarf stars give a somewhat higher declination for the true vertex.

In the analysis of proper motions, the two stream theory gives two vertices, which correspond to the directions of motion of the two streams relative to the sun. The coordinates of these vertices are R. A. 6ʰ 14ᵐ, Dec. −13° (first stream) and R. A. 19ʰ 16ᵐ, Dec. −60° (second stream).

Analyzing stellar motions on the basis of the two stream theory, we find a number of stars which cannot be regarded as belonging to either of the two streams. The B stars and stars of spectral class M, for instance, have a group-motion intermediate between the two streams. For this reason Halm has introduced a third stream (0 stream). But these streams taken together can be fairly well represented by an ellipsoidal distribution using a smaller number of parameters.

Charlier (⁴) has introduced a generalization of the ellipsoidal theory which makes it possible to take into account deviations from a strictly ellipsoidal distribution, but it is only when these deviations are small that this generalization is practicable.

MOVING CLUSTERS OR GROUPS

Several stars move nearly parallel to one another, the best known example being 5 of the 7 bright stars in the constellation Ursa Major. Another moving group or cluster is the Hyades in the constellation Taurus (Taurus Group). The proper motions of the stars belonging to such a group converge towards a point in the sky, the "convergent point," whose position in the sky gives the direction of motion of the group relative to the sun. The convergent point for 17 stars belonging to the Ursa Major Group is R. A. 20ʰ 30ᵐ, Dec. −40°; for the Taurus Group (39 stars) R. A. 6ʰ 7ᵐ, Dec. + 7°. A number of other moving groups are known.

THE GENERAL DISTRIBUTION OF COSMIC VELOCITIES

When the sun's motion is referred to different classes of objects it has been found that this motion is not a constant vector but varies greatly, from about 12 km/sec for the A stars and the Cepheids of long period up to 300 km/sec for the fast moving objects, the globular clusters and the spiral nebulae. A general relationship between group-motion and dispersion exists, which, according to Strömberg (¹¹), holds for all classes of objects, but with a small deviation for the B star system. This variation in group-motion produces an asymmetry in the velocity distribution, in such a way that all fast moving objects move, relative to the sun, towards the same hemisphere. This asymmetry defines an axis along which the group-motion increases with increasing internal velocity-dispersion. The direction of this axis is R. A. 8ʰ 39ᵐ, Dec. −57°, and the motion of objects with small velocity-dispersion relative to those of high velocity-dispersion is about 300 km/sec in the opposite direction. The group-motion of objects

with high velocity-dispersion is approximately the same as that of the globular clusters and spiral nebulae.

The general distribution of cosmic velocities can be approximately represented by a product of two symmetrical distributions S_1 and S_2. The first of these is a sum of concentric and co-axial ellipsoidal distributions, the velocity of the sun relative to the center of the distribution S_1 being 14.8 km/sec in the direction R. A. 17^h 43^m, Dec. $+22°$. The sun's motion relative to the second distribution, S_2, is 300 km/sec in the direction R. A. 20^h 28^m, Dec. $+56°$. The first distribution can be regarded as the velocity-distribution in our local system of stars, the second as a velocity-restriction in a universal world-frame of enormous dimensions. Other interpretations, however, may be possible.

LITERATURE

(For a key to the periodicals see end of volume)

[1] Adams, Strömberg and Joy, *1, 54: 9; 21. [2] Boss, *36, 26: 111; 10. [3] Campbell, *Lick Obs. Bull.* No. 196; 11. [4] Charlier, *Lund Observatorium, Meddelanden,* II: No 13; 15. [5] Charlier and Wicksell, *Ibid.,* II: No 12· 45, 15 [6] Gyllenberg, *Ibid.,* II: No. 13; 15. [7] Luyten, *Annals Harvard College Obs.* 85: No 5; 23 [8] Raymond, *36, 26: 191; 17. [9] Strömberg, *1, 47: 7, 18. [10] Strömberg, *1, 56: 265; 22 [11] Strömberg, *1, 61: 363; 25.

TIME

CHRONOLOGICAL ERAS
Gregorian Calendar

Era	Year	Begins, 1925 A. D.
Byzantine¶	7434	September 14
Diocletian¶	1642	September 11
Grecian*¶	2237	{ September 14 / October 14
Hegira	1344‡	July 21
Japanese	2585†	January 1
Jewish	5686‡	September 18
Julian calendar	1925	January 14
Julian period	6638§	January 14
Mohammedan	1344‡	July 21
Nabonassar¶	2674	May 12
Rome¶	2678‖	January 14
Seleucidae¶	2237	(See Grecian)

* In present-day usage of Syrians, begins in September or October depending upon the sect. In ancient usage of Damascus and Arabia Petraea, began with vernal equinox.
† The 14th year of period Taisho.
‡ Begins at sunset.
§ Julian day number of January 1, 1925 (Gregorian) is 2 424 152
‖ Since foundation of Rome, according to Varro
¶ Based upon Julian calendar.

TIME

Interval	Days*
Year:	
Tropical†	365 2422
Sidereal	365 2564
Anomalistic	365 2596
Month:	
Synodical†	29 530 59
Tropical	27 321 58
Sidereal	27 321 66
Day:	
Sidereal	0 997 2696

* Mean solar days.   † Ordinary

EQUATION OF TIME*
(Δ = mean − apparent)
Unit of Δ is minute. Time is Greenwich mean noon

Date	Δ	Date	Δ	Date	Δ
I 1	+ 3 4	V 11	−3 8	IX 18	− 5.6
6	5.8	16	−3 8	23	− 7.3
11	7 8	21	−3 7	28	− 9.0
16	9 7	26	−3.3	X 3	−10.7
21	11 3	31	−2 6	8	−12.2
26	12 6	VI 5	−1.8	13	−13.5
31	13 6	10	−1 0	18	−14.6
II 5	14 1	15	0 0	23	−15.5
10	14 4	20	+1.1	28	−16.1
15	14 3	25	2.2	XI 2	−16.3
20	14 0	30	3.2	7	−16.3
25	13 3	VII 5	4.2	12	−15.9
III 2	12 4	10	5.0	17	−15.1
7	11.4	15	5.6	22	−14.0
12	10 0	20	6.1	27	−12.5
17	8 7	25	6.3	XII 2	−10.7
22	7 2	30	6.3	7	− 8.8
27	5.7	VIII 4	6.0	12	− 6.5
IV 1	4 2	9	5.4	17	− 4.1
6	2.7	14	4.7	22	− 1.6
11	1 2	19	3.7	27	+ 0.9
16	+ 0 0	24	2.5	31	+ 2.8
21	− 1 2	29	+1 1		
26	− 2 2	IX 3	− 0 4		
V 1	− 2 9	8	−2.1		
6	− 3 4	13	−3.8		

* Δ is the amount by which mean time exceeds apparent time when it is noon at Greenwich, it is the excess of the right ascension of the actual sun over that of the mean sun at that instant. It varies continuously with the time, and does not exactly repeat its values in successive years; those given are average values for Greenwich mean noon of an ordinary year, and will seldom differ from the actual values for that time by as much as 0 2 min., except in January and December, when the difference may amount to 0.3 min. In leap years, all dates in the table after February must be reduced by one day.

SOLAR SYSTEM

ORBITAL DATA; SOLAR SYSTEM (1925)

Units: Distance, 10^6 km; period, tropical year

Planet	Distance*	Eccentricity	Inclination †	Mean longitude		Sidereal period
				Node‡	Perihelion	
☿ Mercury. .	57 9	0 2056	7° 0' 12.0"	47° 26' 32 1"	76° 17' 18.9"	0.24085
♀ Venus .	108 1	0 0068	3 23 38 0	76 0 16 7	130 30 56.8	0.61521
⊕ Earth	149 5	0 01674			101 39 2 3	1.00004
♂ Mars	227 8	0 0933	1 51 0 6	48 58 45.0	334 40 42.2	1 88089
♃ Jupiter	778	0 0484	1 18 26 4	99 41 26 3	13 6 51.4	11 862
♄ Saturn	1426	0 0558	2 29 28 7	113 0 5 7	91 34 42.0	29.458
♅ Uranus.	2869	0 0471	0 46 22 1	73 36 57 7	169 26 56.8	84.015
♆ Neptune	4196	0 00855	1 46 36 7	130 57 13 3	43 58 27.9	164 788

* Mean distance
† Angle between plane of orbit and plane of ecliptic
‡ Ascending node

CHARACTERISTICS OF MEMBERS OF SOLAR SYSTEM

Units: Linear diameter, 1000 km; density, g/cm^3; time, mean solar

Name	Diameter		Mass† × 10^6	Density	Sidereal rotation	Number satellites
	Linear	Angular*	Mass sun			
Mercury..	4 84	10 90"	0 1670	5 6		0
Venus.	12 19	1' 0 80	2.451	5 1		0
Earth.. .	12 76§		3 036‡	5 52	23 hr 56.07 min	1
Mars.	6 78	17 88	0 3233	3 9	24 37 4	0
Jupiter	142 7§	46 86§	954 8	1 4	9 8 hr	7
Saturn	120 8§	19 52§	285 6	0 7	10 2 hr	9
Uranus..	49 7	3 76	43 7	1 3+		4
Neptune	53 0	2 52	50 8	1 3		1
Sun‖	1391	31 59 26	1 001 341	1 4	25 3 da	
Moon	3 48	31 5 16¶	0 037**	3.3	27.32 da	

* At distance = difference mean distance sun to object and mean distance sun to Earth, nearly at distance of nearest approach to Earth.
† Includes satellite (or planetary) system, if any
‡ Mass of Earth alone = 2.999 × 10^{-6} mass of sun
§ Equatorial diameter Polar diameter Earth = 12 71, Jupiter = 133 2, 13 74", Saturn = 108 1, 17 46" Diameter of sphere of volume = Earth, is 12 74
‖ At mean distance of Earth, gravitational acceleration due to Sun is $k^2 = 2.9592 \times 10^{-4}$ (mean distance) per day² = 0 5926 cm per sec². For solar spectrum etc., see index
¶ At mean distance from Earth Apparent diameter varies, with distance, from 29 5' to 33 5'
** Moon alone. Mass Moon = 0 01227 mass Earth

SOLAR DATA

Inclination of equator to ecliptic, about	7°
Longitude of ascending node of equator	74 5'
Period of rotation, about	28 da*
Sun spot period, about	11 yr

TERRESTRIAL AND LUNAR DATA†

General precession (retro-
grade) . . . 50 2564" + 0 000222"(t - 1900) per yr
Obliquity of the ecliptic . 23° 27' 8 26" - 0 4684 "(t - 1900)

* From observations of sun spots near latitude 45°, spots near equator rotate in about 24 da; those near lat 80°, in 30 da
† For geodetic and geophysical data, see p 393.

Constant of notation.	9 21"	
Constant of aberration	20 47"	Paris conference values
Solar parallax	8 80"	

From parallax measurements	8.806"
From velocity of light	8 781
From mass of Earth.	8.762
From motion of Moon	. . . . . . 8 773
Equatorial horizontal parallax of Moon*	57' 2 70" (Brown)
Mean distance Earth to Moon.	384 403 km
Inclination of Moon's equator to ecliptic	1° 32 1"
Inclination of Moon's orbit to ecliptic, about	5°
Eccentricity of Moon's orbit (average). . .	0 055
Revolution of Moon's nodes (retrograde) .	18.6 yr

* Mean of greatest and least values; actual values vary from 53' to 61' ca.

COMPOSITION OF THE ATMOSPHERE

W. J. Humphreys

TABLE 1.—Composition of Dry Air at Sea-level (4, 5)

v = volume of the gas in volume V of dry air

Gas	N₂	O₂	A	CO₂	H₂*	Ne	He	Kr	Xe
$10^4 v/V$	7803	2099	94	3	1	0 123	0 04	0 005	0 0006

* Values found by analysis vary; the one here given is that accepted by Hann and the Recueil de Constantes Physiques.

TABLE 2.—Composition of Atmosphere at Various Levels

Computed from data of Table 1 on the assumptions: (1) at surface, H₂O vapor supplies 1.2% of the total number of gas molecules, (2) absolute humidity decreases rapidly to a negligible amount at about 10 km, (3) temperature = 11°C at sea-level, decreases normally (6°C per km) to −55°C at 11 km, remains constant above 11 km, (4) relative proportions of the gases, water vapor excepted, remains constant up to 11 km, (5) above 11 km, distribution is in accordance with their molecular weights (3). The amount of H₂ is in doubt (see note Table 1), especially above 11 km; it may become oxidized to H₂O before reaching the upper atmosphere.

v = volume of the gas contained in volume V of atmosphere. Unit of height = 1 km = 0 621 mi.; of pressure = 1 mm of Hg

Height	\multicolumn{7}{c	}{100 v/V}	Total pres-sure
	N₂	O₂	H₂O	A	CO₂	H₂	He	
140	0 01					99 15	0 84	0 0040
130	0 04					99 00	0 96	0 0046
120	0 19					98 74	1 07	0 0052
110	0 67	0 02	0 02			98 10	1 19	0 0059
100	2 95	0 11	0 05			95 58	1 31	0 0067
90	9 78	0.49	0.10			88 28	1 35	0 0081

Height	\multicolumn{6}{c	}{100 v/V}		Total pres-sure
	N₂	O₂	H₂O	A	CO₂	H₂	He	
80	32 18	1 85	0 17			64 70	1 10	0 0123
70	61 83	4 72	0 20	0 03		32 61	0 61	0 0274
60	81 22	7 69	0 15	0 03		10 68	0 23	0 0635
50	86 78	10 17	0 10	0 12		2 76	0 07	0 408
40	86 12	12 61	0 06	0 22		0 67	0.02	1 84
30	81 26	15 18	0 03	0 35	0 01	0 16	0 01	8 63
20	81 24	18 10	0 02	0 59	0 01	0 04		40 99
15	79 52	19 66	0 01	0 77	0 02	0 02		89 66
11	78 02	20 99	0 01	0 94	0 03	0 01		168 00
5	77 89	20 95	0 18	0 94	0 03	0 01		405.
0	77 08	20 75	1 20	0 93	0 03	0 01		760.

TABLE 3. Masses of the Atmosphere and Its Constituents

Based upon Table 1, the assumptions of Table 2, and the assumption that the average atmospheric pressure at the surface of the earth = 73 7 cm and at base of stratosphere = 14.5 cm (1, 2). Area of earth is taken as 51 × 10¹⁷ cm².

Total mass $M = m \times 10^8$ kg; 1000 kg = 1.102 tons (of 2000 lb.)

Gas	All	N₂	O₂	A	H₂O	CO₂	H₂	Ne	Kr	He	Xe
m	511	387	116	624	133	217	129	471	64	63	116
n	16	16	16	14	14	13	12	11	11	11	10

LITERATURE

(For a key to the periodicals see end of volume)

(1) Hann, *Lehrbuch der Meteorologie* (3rd ed.) (2) Humphreys, *Monthly Weather Review*, **49**: 341, 21 (3) Humphreys, *Physics of the Air*, p. 69; 20. (4) Ramsay, **5**, **80**: 599; 08 (5) Various authorities

MISCELLANEOUS GEODETIC DATA

W. D. Lambert

With certain exceptions which are especially noted, those of the following data which depend upon the dimensions of the earth have been calculated strictly in accordance with the INTERNATIONAL ELLIPSOID OF REFERENCE, adopted by the Section of Geodesy of the International Geodetic and Geophysical Union, meeting at Madrid, October 6 and 7, 1924. This ellipsoid is based upon the results obtained by J. F. Hayford (Supplementary Investigation in 1909 of the Figure of the Earth and Isostasy, Washington, 1910), but is not absolutely identical with Hayford's ellipsoid. (For some of the other spheroids that are used for geographical purposes, see Special Publication #100, U. S. Coast and Geodetic Survey. Recent attempts have been made to show that the actual figure of the earth can be represented more closely by an ellipsoid of three unequal axes, than by one of revolution, systematic departures from the latter being of the order of 100 to 200 meters in elevation and depression.)

If the positions of the two ends of a line are determined geodetically for any assumed spheroid of reference, the uncertainty in the length of the line as measured along the earth depends almost entirely upon the errors in the survey; for geodetic surveys of the highest class, the uncertainty is a little less than one in 100 000 and for an ordinary fair survey it is about four times as great. The proportional error in the straight-line distance is greater, mainly because the geoid does not coincide with the ellipsoid; these additional errors are not serious for a short line, but for two points almost diametrically opposite may amount to 100 or 200 meters.

If the end points are determined astronomically, the principal error in the computed length is due to the difference in the deflection of the plumb-line at the two points; unless the measured line is short, the average uncertainty so introduced is of the order of 200 meters, but may be much more, especially in rugged country.

Latitude.—The latitude of a place is defined as the angle which some line of reference makes with the equatorial plane. Four lines of reference, defining four distinct kinds of latitude, are used. Three of these lines pass through the place considered; viz., (1) The plumb-line, defining the *astronomical* latitude, (2) the normal to the spheroid of reference, defining the *geographical* latitude, and (3) the line to the center of the earth, defining the *geocentric* latitude. The fourth line of reference passes through the center of the earth and that point which is upon the circumscribed sphere (radius = equatorial radius of the spheroid) and at the same distance from the axis of rotation as is the point on the spheroid representing the place considered; this defines the *parametric*, or *reduced*, latitude.

Gravity.[1]—If the earth's sea-level surface were accurately represented by the International Ellipsoid of Reference, and if no attracting matter projected above this surface, then the variation of gravity at sea-level (γ_0) would be represented by the equations

$$\gamma_0 = \gamma_e(1 + 0.005\ 288 \sin^2\varphi - 0.000\ 006 \sin^2 2\varphi)$$
$$= \gamma_{45}(1 - 0.002\ 637 \cos 2\varphi + 0.000\ 006 \cos^2 2\varphi)$$

[1] The resultant acceleration arising from the gravitational attraction and the rotation of the earth.

where φ is the geographic latitude, and γ_e, γ_{45} are the values of γ_e at the equator and at latitude 45°, respectively. These equations differ slightly from that used in computing the table on p. 396; the latter corresponds to an ellipticity of 1/297.4.

Table 1.—Form and Size of the Earth

Based upon International Ellipsoid of Reference; accepted constants, from which the others are computed, are $a = 6\ 378\ 388$ meters, ellipticity $[= (a - b)/a] = 1/297$. The indicated uncertainties are estimates, by Lambert, based upon a consideration of systematic errors as well as of internal discordances.

a = semi-major axis	= 6 378 388(± 60)m
b = semi-minor axis	= 6 356 911 946 m
Radius of sphere of same area	= 6 371 227 7 m
Radius of sphere of same volume	= 6 371 221 3 m
Length of equatorial quadrant	= 10 019 148 4 m
Length of meridional quadrant	= 10 002 288 3 m
f = ellipticity = $\left(\dfrac{a-b}{a}\right)$	= 0 003 367 0034
$\dfrac{1}{f}$ = reciprocal of ellipticity	= 297 0(±0 4)
e^2 = (eccentricity)² = $f\left(\dfrac{2}{f}-1\right)$ = $\dfrac{a^2-b^2}{a^2}$	= 0 006 722 6700
Area of the ellipsoid	= 510 100 934 km²
Land area	= 148 847 000 km²
Ocean area	= 361 254 000 km²
Volume of the ellipsoid	= 1 083 319 78 × 10⁶ km³
Mass of the ellipsoid* ($d = 5 527$ g/cm³, p 395)	= 5 988 × 10²⁴ kg
Principal moments of inertia ($A = B < C$)†	
$A\ddagger = B\ddagger$	= 0 332 35 Ea^2
$C\ddagger$	= 0 333 44 Ea^2
$C - A$	= 0 001 0921 Ea^2
$\left(\dfrac{C-A}{C}\right) = \left(\dfrac{1}{305\ 12}\right)$	= 0 003 2774

* For discussion of variation of density with depth below surface, see Adams and Williamson, Smithsonian Annual Report, 1923, p 241

† E = mass of earth.

‡ Computed values vary but little with any admissible assumption regarding the constitution of the interior of the earth Values are based upon computations of De Sitter (84V, 27: 233, 24); ellipticity taken as 1/296 92

§ Deduced from precession of equinoxes, involves no hypothesis regarding constitution of interior of earth.

Table 2.—Distances upon Surface of the International Ellipsoid of Reference

M = length of meridian from equator to geographic latitude φ; S_m = length of meridian from latitude $(\varphi - \frac{1}{2}\Delta\varphi)$ to $(\varphi + \frac{1}{2}\Delta\varphi)$; S_p = length of arc of parallel for 1° of longitude at latitude φ. These may be computed by means of the equations: $M = a\varphi - b \sin 2\varphi + c \sin 4\varphi - d \sin 6\varphi$; $S_m = a\Delta\varphi - b \sin \Delta\varphi \cos 2\varphi + c \sin 2\Delta\varphi \cos 4\varphi - d \sin 3\Delta\varphi \cos 6\varphi$; S_m (for $\Delta\varphi = 1°$) = $a - b \cos 2\varphi + c \cos 4\varphi - d \cos 6\varphi$; $S_p = a \cos \varphi - b \cos 3\varphi + c \cos 5\varphi$; where the coefficients and their logarithms have the following values:

Unit of length = 1 meter; of angle = 1°

	M*		S_m*	
	Value	log₁₀	Value	log₁₀
a	111 136.537	5.045 856 86	111 136 537	5 045 856 86
b	16 107 035	4 207 015 6	32 214 069	4 508 045 6
c	16.976	1.229 84	33 952	1 530 87
d	0.022	2̄ 348	0 045	2̄.649

	S_m* for $\Delta\varphi = 1°$		S_p*	
	Value	log₁₀	Value	log₁₀
a	111 136.537	5.045 856 86	111 417.657	5.046 954 02
b	562.213	2.749 901	93.904	1.972 686
c	1.185	0 073 7	0.119	1̄.074 6
d	0 002	3̄ 37		

* Owing to uncertainty regarding the actual size of the earth, actual distances upon the earth at sea-level may differ from these computed distances by about 2 in 100 000 near the equator or the poles, by somewhat less in middle latitudes

Table 3.—Excess of Geographic Latitude (φ) over Geocentric (φ') and Parametric (θ) Latitudes

$$\varphi - \varphi' = a \sin 2\varphi - b \sin 4\varphi + c \sin 6\varphi$$
$$= a \sin 2\varphi' + b \sin 4\varphi' + c \sin 6\varphi'$$
$$\varphi - \theta = a' \sin 2\varphi - b' \sin 4\varphi + c' \sin 6\varphi$$
$$= a' \sin 2\theta + b' \sin 4\theta + c' \sin 6\theta$$

where the coefficients and their logarithms have the following values:

Unit of coefficients = 1″

	Value	log₁₀		Value	log₁₀
a	695 6635	2 842 3992	a'	347 8327	2 541 3704
b	1.1731	0 069 34	b'	0 2933	1̄ 467 29
c	0 0026	3̄.421	c'	0 0003	4̄ 52

Table 4.—Miscellaneous Terrestrial Data

Angular velocity of rotation	72 921 × 10⁻⁸ radians/sec*
Rotational energy	2 160 × 10³⁶ ergs
Rotational energy lost by tidal friction	1.1 × 10¹⁹ ergs/sec†
Work required to dissipate the material of the earth to infinity	2.46 × 10³⁹ ergs
Mean elevation of land above sea-level	825 m
Mean depth of the oceans	3681 m
Mean effective viscosity is not known, but perhaps between	10²⁰ and 10²⁵ poises‡

* Mean solar second.

† Jeffreys, 68, 221A: 239, 20; The Earth, Its Origin, History and Physical Constitution, 205–237,; 24 Heiskanen, 175, 18A: 1; 21

‡ Schweydar, Veröffentl des Preuss Geodät Inst, No 79; 19; Jeffreys, Monthly Notices, Roy Ast Soc., 78: 648; 15. 76: 84. 16 77: 449, 17; also The Earth, its Origin, History, and Physical Constitution, 222, 1924

Rigidity (μ). From the yielding of the solid portions (revealed by observations with horizontal pendulums), and on assumption of incompressibility, Schweydar (Zentralbureau Int. Erdmes., Neue Folge No. 38, 1921) deduces $\mu = 30.8 (1 - 0.90r^2/a^2) \times 10^{11}$ dynes/cm², and mean effective rigidity = 17.6 × 10¹¹ dynes/cm² (r = distance from center, a = mean radius). To allow for compressibility, these values must be increased by about 20% (Lambert, preliminary, unpublished computations); even then the value computed for the outer shell of half-radius thickness is much less than that deduced from earthquake data. (See Adams and Williamson, Smithsonian Annual Report, 1923.) The discrepancy may arise from Schweydar's assumption of high rigidity in the central portions, which may possibly behave as a fluid. (See Knott, 68, 39: 157; 19; Sieberg, Geologische, physikalische und angewandte Erdbebenkunde, 364; 23.)

GRAVITY DATA

Clarence H. Swick

This section includes: (A) The value of the gravitation constant; (B) the absolute determination upon which the tabulated values of the acceleration of gravity[1] rest; (C) values of the acceleration of gravity (g) at numerous stations well distributed over the surface of the earth, together with a table giving the values of g at sea-level and at various latitudes; and (D) means for computing the variation in g with the distance of the station above, or below, either the surface of the earth or sea-level. In preparing the data, valuable assistance was received from several colleagues. In particular should be mentioned Mr. W. D. Lambert's assistance with section D, and Miss Sarah Beall's and Mr. H. S. Rappleye's assistance with section C.

A. GRAVITATION CONSTANT

The best determinations of the gravitation constant (G)[2] are considered to be those by C. V. Boys [7] and by K. Braun [8]. Each used an improved form of the Cavendish apparatus; and they obtained almost identical results, the final values of the two determinations being the same to the fourth significant figure. They found

$$G = 6.658 \times 10^{-8} \text{ cm}^3 g^{-1} \text{sec}^{-2}$$

which requires that the mean density of earth $= 5.527$ g/cm^3.

B. BASIS OF REFERENCE

The observed values of gravity in Tables 1 and 2 are relative determinations in the Potsdam system, that is, they are based on

the value of 981.274 cm/sec^2 for the pendulum room of the Geodetic Institut in Potsdam, Germany. This value for Potsdam is the result of a large number of careful absolute determinations extending over a series of years. The degree of uncertainty in such absolute determinations is well illustrated by the fact that a similar series of absolute determinations at Vienna, Austria, gave a value 0.016 cm/sec^2 greater than the one above when referred to Potsdam by relative determinations.

All determinations of gravity should be based on the Potsdam system by means of relative determinations with some station already accurately based on that system. A table of 20 base stations on the Potsdam system is given in *Comptes Rendus l'Association Geodesique Internationale* for 1909, III:25. Most of these stations are included in Table 1.

C. ACCELERATION OF GRAVITY AT SELECTED STATIONS

The stations included in Table 1 are grouped (1) in the order America, Europe, Asia, Africa, Australia, and Oceanic; (2) generally, alphabetically according to countries (United States of America, first); (3) in each subdivision, the stations are arranged alphabetically. Numerals in parentheses, following the name of a subdivision or station refer to the bibliography, and indicate the source from which the data were obtained. If the effect of topography and of isostatic compensation has been computed on the uniform basis of compensation extending to a depth of 113.7 km, the amount of this computed effect is given in the column TC. This effect is the amount by which the actual value of the acceleration would exceed that obtained from Table 2, after correction for elevation by means of equation (1), if there were complete isostatic compensation and if the local distribution of matter were not anomalous.

TABLE 1.—ACCELERATION (*g*) OF GRAVITY, POTSDAM SYSTEM

(The effect of topography and of isostatic compensation = TC)

Units: Elevation (h), meters: g, cm/sec^2; TC, cm/sec^2

Station	Latitude	Longitude	h	g	TC	Station	Latitude	Longitude	h	g	TC
AMERICA						Madison, Wis (University of Wisconsin)	43° 4 6'	89° 24 0'	270	980.365	+0 003
United States [5,6]						Minneapolis, Minn (University of Minnesota)	44 58.7	93 13.9	256	980.597	-0.005
Albany, N Y. (Public School No. 24)	42° 39 1'	73° 46 1'	61	980 311	-0 006	Mount Hamilton, Calif (Lick Observatory)	37 20 4	121 38 6	1282	979 660	+0 120
Apalachicola, F l a (Weather Bureau)	29 43 5	84 58 8	4	979 322	+0 015	New Orleans, La (City Hall)	29 57 0	90 4 2	2	979 324	+0 013
Asheville, N C (Post-office)	35 35 9	82 33 3	670	979 603	+0 026	New York, N. Y. (Columbia University)	40 48 5	73 57 7	38	980 267	+0.011
Atlanta, Ga. (State Capitol)	33 45 0	84 23 3	324	979 521	+0 014	Norris Geyser Basin, Wyo (Yellowstone Park)	44 44 2	110 42 0	2276	979 950	+0 031
Austin, Tex (University)	30 17 2	97 44 2	189	979 283	-0 001	Pembina, N Dak (Public School)	48 58 1	97 14 9	243	980 917	-0 009
Baltimore, Md (Johns Hopkins University)	39 17 8	76 37 3	30	980 097	+0 006	Philadelphia, Pa (University of Pennsylvania)	39 57 1	75 11 7	16	980 196	+0 009
Bismarck, N Dak (Will School)	46 48 5	100 47 0	516	980 625	-0 005	Pierre, S Dak (High School)	44 21 9	100 20 8	454	980 427	-0 013
Boise, Idaho (High School)	43 37 2	116 12 3	821	980 212	-0 042	Pittsburgh, Pa (Second Ward School)	40 27 4	80 0 6	235	980 118	0 000
Calais, Me (High School)	45 11 2	67 16 9	38	980 641	+0 010	Point Isabel, Tex	26 4 7	97 12 4	8	979 076	+0.015
Cambridge, Mass (Harvard College Observatory)	42 22 8	71 7 8	14	980 398	+0 010	Portland, Oreg (Custom House)	45 31 4	122 40 7	8	980 646	-0 016
Charleston, W Va (High School)	38 20 9	81 37 7	184	979 936	-0 010	Potsdam, N Y. (Clarkson School of Technology)	44 40 1	74 58 8	130	980 571	-0 004
Charleston, S C. (S C Military Academy)	32 47 2	79 56 0	6	979 546	+0 016	Princeton, N J (Princeton University)	40 21 0	74 39 5	64	980 178	+0.013
Charlottesville, Va. (University of Virginia)	38 2 0	78 30 3	166	979 938	+0 002	Richmond, Va (Post-office)	37 32 2	77 26 1	30	979 960	+0 010
Chicago, Ill. (Univ of Chicago)	41 47 4	87 36 1	182	980 278	+0 007	St Louis, Mo (Washington University)	38 38 0	90 12 2	154	980 001	+0 001
Cincinnati, Ohio (Cincinnati Observatory)	39 8 3	84 25 3	245	980 001	+0 002	Salt Lake City, Utah (Temple Block)	40 46 1	111 53 8	1322	979 803	-0 041
Cleveland, Ohio (Adelbert College)	41 30 4	81 36 6	210	980 211	0 000	San Francisco, Calif. (Davidson Observatory)	37 47 5	122 25 7	114	979 965	+0.045
Colorado Springs, Colo. (Colorado College)	38 50 7	104 49 0	1841	979 190	-0 007	Sandpoint, Idaho (Farmington Central School)	48 16 4	116 33 3	637	980 680	-0 044
Denver, Colo (University of Denver)	39 40 6	104 56 9	1638	979 609	-0 015	Seattle, Wash (Washington State University)	47 39 6	122 18 3	58	980 733	-0.020
Dover, Del (Wilmington Conference Academy)	39 9 7	75 32 0	12	980 099	+0 013	Springfield, Ill. (Edwards Public School)	39 47 7	89 39 5	183	980.089	+0.005
El Paso, Tex (High School)	31 46 3	106 29 0	1146	979 124	+0 001	State College, Pa. (Chemistry Physics Building)	40 47 9	77 51 8	358	980.124	+0.010
Galveston, Tex. (Ball High School)	29 18 2	94 47 5	4	979 272	+0 007	Terre Haute, Ind (Rose Polytechnic Institute)	39 28 7	87 23 8	151	980 072	+0 001
Georgetown, Tex (South western University)	30 38 0	97 40 1	241	979 298	+0.002	Washington, D C (U S C. and G S, base station)	38 53 2	77 0 5	11	980 112	+0.004
Goldfield, Nev (High School)	37 42 2	117 14 5	1716	979 456	+0 027	Washington, D C (Bureau of Standards)	38 56.3	77 4 0	103	980 095	+0.012
Hartford, Conn (Jarvis Laboratory of Trinity College)	41 44 8	72 41 8	37	980 336	+0 008	Wilmington, N. C (Court House)	34 14 2	77 56 6	9	979 663	+0.023
Hinsdale, Mont (Public School)	48 23 8	107 5 3	661	980 739	-0 017	Worcester, Mass. (Worcester Polytechnic Institute)	42 16 5	71 48 5	170	980 324	+0.018
Hoboken, N J (Stevens Institute of Technology)	40 44	74 2	11	980 269	+0 008	Yavapai, Ariz (Yavapai Point)	36 3 9	112 7.1	2179	979 192	+0.034
Indianapolis, Ind (Postoffice)	39 45 9	86 8 8	217	980 090	+0 003	**Alaska** [4]					
Ithaca, N Y (Cornell University)	42 27 1	76 29 0	247	980 300	+0 005	Fort Egbert, Eagle City	64 47.4	141 12.4	269	982.183	-0.042
Kansas City, Mo (Franklin School)	39 5 8	94 35 4	278	979 990	-0 001	Percy Islands, Southeast Alaska	54 55.8	131 35.3	4	981.524	-0.013
Key West, Fla. (Post-office)	24 33 6	81 48 4	1	978 970	+0 035						
Lancaster, N H (High School)	44 29.5	71 34 3	261	980 486	+0 007						
Las Vegas, N Mex (Normal School).	35 35.8	105 12 1	1960	979 204	+0 017						
Little Rock, Ark. (Postoffice)	34 45.0	92 16.4	89	979 721	+0.001						

Station	Latitude	Longitude	h	g	TC
Point Young, Southeast Alaska	58° 11.5'	134° 33 4'	7	981 757	-0 054
Quiet Harbor, Southeast Alaska	56 14 1	132 39 6	4	981 624	-0 034
St. Michael	63 28 5	162 2 4	1	982 192	-0 004
St. Paul Island	57 7 3	170 16 6	10	981 726	+0 011
Canada (6, 10, 21, 22)					
Arctic Red River, N W. Ter	67 26 6	133 44 2	41	982 434	-0 026
Banff, Alta	51 10 9	115 34 5	1376	980 753	-0 012
Calgary, Alta	51 2.7	114 3 8	1044	980 823	-0 022
Charlottetown, P E. I	46 13 9	63 7 5	8	980 733	+0 013
Chipewyan, Alta	58 42 7	111 8 8	229	981 723	-0 012
Good Hope, N. W. Ter	66 15.3	128 38 2	59	982 340	-0 029
Halifax, N. S	44 40 8	63 33 8	9	980 574	+0 008
Kenora, Ont .	49 46 0	94 30 0	330	980 974	+0 018
Kingston, Ont. (City Hall)	44 14 6	76 28 8	79	980 530	+0 008
Liard River, B. C	59 58 7	123 17 5	160	981 790	-0 059
Moose Jaw, Sask	50 23 4	105 31 8	511	980 943	+0 003
Norman, N. W. Ter.	64 54.0	125 31 2	87	982 214	-0 036
Ottawa, Ont. (Dominion Observatory, base station)	45 23 6	75 43 0	83	980 618	0 000
Peace River, Alta	56 14.1	117 17 2	324	981 482	-0 038
Port Arthur, Ont. (Masonic Building)	48 26 0	89 13 0	189	980 820	-0 014
Providence, N W Ter .	61 21 2	117 39 2	156	981 955	-0 018
Resolution, N W Ter	61 10 1	113 40 5	152	981 942	-0 009
Revelstoke, B. C	50 59 8	118 11 8	453	980 903	-0 080
St Jérôme (Chateau Larose)	45 46 6	74 0 0	107	980 681	+0 006
St. John, N. B. (Meteorological Observatory)	45 16 0	66 5 0	33	980 663	+0 016
Sault Ste. Marie, Ont (City Hall)	46 30 4	84 19 2	186	980 680	-0 005
Simpson, N. W. Ter	61 51 6	121 20 8	132	982 004	-0 023
Sydney, N S	46 8 4	60 11 8	12	980 731	+0 014
Vancouver, B C	49 16 8	123 6 8	6	980 949	-0 016
Winnipeg, Man	49 54 4	97 8 0	231	980 990	+0 002
Woodstock, N B (Armoury)	46 9 0	67 34 5	56	980 699	+0 008
Woodstock, O n t . (Market)	43 8 6	80 47 0	299	980 352	-0 002
Central and South America (2)					
Bahía Blanca, Argentina .	38 47 1 8	62 15 9	2	980 061	
Buenos Aires, Argentina	34 36 5 8	58 22 2	2	979 669	
Bahia, Brazil	12 58 5 8	38 31 0	4	978 331	
Panama, Canal Zone	8 54 9	79 31 9	6	978 243	
Valdivia, Chile	39 53.4 8	73 28 3	10	979 920	
Valparaíso, Chile	33 1 8 8	71 38 5	60	979 609	
Callao, Peru	12 4 1 8	77 15 8	1	978 375	
Acajutla, Salvador.	13 34 7	89 50 4	12	978 303	
Montevideo, Uruguay	34 54 5 8	56 12 9	4	979 772	
Kanada *see* Canada.					
EUROPE					
Allemagne, *see* Germany					
Angleterre, *see* Great Britain.					
Austria (2, 6)					
Brenner	47 0 3	11 30.5 E	1372	980 353	
Dalaas	47 8	9 59 E	838	980 454	
Grafenstein	46 37	14 28 E	417	980 614	
Mixnitz..	47 19.8	15 22 E	445	980 657	
Ober-Drauburg	46 45	12 58 E	617	980 555	
Stilfserjoch (Stelvio Pass) .	46 31 8	10 27 4 E	2760	980 045	0 152
Vienna (base station)	48 12 7	16 21.5 E	183	980 860	
Waidhofen	47 57.7	14 46 7 E	352	980 750	
Wien (base station)	48 12.7	16 21 5 E	183	980 860	
Wolfsthal .	48 8.3	17 0 5 E	146	980 904	
Belgium (2)					
Brussels.	50 51 0	4 22 E.	102	981 112	
Czechoslovakia (2)					
Böhmerwald .	49 40 1	12 59 3 E.	537	980 921	
Cebon........	50 0.9	13 0.4 E.	822	980 906	

Station	Latitude	Longitude	h	g	TC
Karlowitz	49° 21 9'	18° 18 7'E.	510	980 890	
Mount Hora	49 10 3	15 42 4 E	710	980 845	
Rosenau	48 39 1	20 32 E.	281	980 871	
Denmark (2)					
Copenhagen (Sternwarte, base station)	55 41 2	12 34 7 E	14	981 559	
Frederikshavn	57 27 1	10 32 2 E	15	981 740	
Magleby	54 47 3	10 43 0 E	14	981 502	
Peders Kirke	55 1 6	14 58 8 E	12	981 533	
Trige	56 15 2	10 9 5 E	91	981 618	
Vinding .	55 40 3	9 34 5 E	78	981 575	
Deutschland, *see* Germany					
England, *see* Great Britain					
Espagna, *see* Spain					
Finland (2)					
Helsingfors (Observatory)	60 9 7	24 57 3 E	29	981 912	
Uleåborg	65 1 2	25 29 1 E.	9	982 202	
Viborg (Vipurin)	60 42 9	28 43 7 E	12	981 928	
Fiume (2)	45 20 0	14 25 8 E	10	980 630	
France (2, 3)					
Arcachon	44 39 6	1 10 4	24	980 586	
Aurillac, Lyceum	44 56 8	2 26 6 E	640	980 483	
Bayonne	43 29 7	1 28 0	3	980 475	
Bordeaux (Observatoire)	44 50 1	0 31 4	72	980 572	
Coutras	45 2 5	0 7 9	13	980 591	
Jonzac	45 26 7	0 26 0	35	980 647	
Langon	44 32 7	0 15 3	25	980 561	
Libons	49 50 0	2 45 E	106	981 038	
Lyon	45 41 0	4 47 E	286	980 629	
Marseille (Observatoire)	43 17 9	5 23 E	61	980 482	
Metz	49 7 0	6 10 7 L.	175	980 957	
Meudon (Observatoire)	48 48 3	2 13 9 E	130(?)	980 919	
Mont Blanc (Observatoire)	45 50	6 52 E	4807	979 401	
Mont-Louis	42 31 0	2 7 E	1620	979 906	
Nice (Observatoire)	43 42 8	7 18 E	367	980 471	
Paris (Observatoire, base station)	48 50 2	2 20 3 E	61	980 943	
Port-Vendres	42 50 9	3 6 E	25	980 456	
Rosendaël-les-Dunk	51 2 9	2 24 E	20	981 170	
Soulac	45 31 0	1 7.4	8	980 655	
Strasbourg (base station)	48 35 0	7 46 1 E	137	980 904	
Valence	44 56	4 53 E	125	980 562	
Germany (2, 6)					
Alter Bruch	50 45 7	15 44 6 E	917	980 930	+0 060
Bremen	53 5 0	8 49 2 E	0	981 341	
Brocken	51 48 0	10 37 E	1140	981 015	+0 088
Coburg	50 16 0	10 58 E	290	981 015	
Gottingen (Sternwarte)	51 32 0	9 57 E	162	981 176	
Grimmen	54 6 9	13 2 7 E	11	981 434	
Hamburg (Seewarte)	53 32 8	9 58 3 E	24	981 375	
Helgoland	54 10 8	7 53 1 E	51	981 410	
Immenstaad	47 40 0	9 22 1 E	403	980 709	
Jena	50 55 6	11 35 2 E	154	981 123	
Karlsruhe	49 0 7	8 24 7 E	114	980 967	
Kiel (Sternwarte)	54 20 5	10 9 E	41	981 464	
Kirchhain	51 38 3	13 33 5 E	98	981 235	
Kolberg	54 11 3	15 35 8 E	8	981 453	
Königsberg (Sternwarte)	54 42 8	20 29 8 E	22	981 477	
Leipzig	51 20.1	12 23.5 E	115	981 180	
Lüdenhausen	52 4 3	9 0.0 E	205	981 242	
Munich	48 8 7	11 36 6 E	525	980 733	
Münster	51 57.9	7 37.9 E	62	981 233	
Neumünster	54 4.4	10 0 E.	25	981.427	
Potsdam (Geodetic Institute, base station)	52 22 9	13 4.1 E	87	981.274	
Scharfenstein	51 50.0	10 36.0 E	623	981.130	+0 041
Schneekoppe	50 44 2	15 44.6 E	1605	980.776	+0 110
Sehlagrund	52 52 8	15 48.0 E	109	981 278	
Stuttgart	48 46 9	9 10.5 E	217	980 901	
Waldsee	47 55	9 45.3 E	590	980.706	

Station	Latitude	Longitude	h	g	TC
Great Britain (2)					
Edinburgh, Scotland (Observatory)	55° 57.4'	3° 9 4'	104	981 584	
Glasgow, Scotland (University)	55 31 5	4 14 0	61	981.605	
Greenwich, England (Observatory)	51 28 6	0 0 0	48	981 184	
Kew, England (Observatory)	51 28 1	0 19	5	981 144	
Plymouth, England	50 22.2	4 8 4	43	981 148	
Holland, see Netherlands					
Hungary (2)					
Budapest	47 29 5	19 3 6 E	108	980.852	
Kis-Komárom	46 32 9	17 10 7 E	115	980.745	
Italy (2, 6)					
Alba	44 42 0	8 2 3 E	169	980 444	
Arona	45 45 8	8 34 1 E	210	980 629	
Bologna (Università)	44 29 8	11 21 3 E	51	980 450	
Brenner (see Austria)					
Catania, Sicily	37 30.2	15 4 7 E	43	980 065	
Castellammare di Stabia	40 41 6	14 28 7 E	4	980 321	
Domo d'Ossola	46 7 0	8 18 4 E	276	980 598	
Florence	43 46 8	11 15 2 E	48	980 510	
Genoa (Instituto Idrografico)	44 25 1	8 55 3 E	93	980.573	
Livorno (Leghorn)	43 32 0	10 18 5 E	6	980 531	-0 018
Milan (Osservatorio)	45 28 0	9 11 5 E	141	980 569	
Padua (Osservatorio, base station)	45 24 0	11 52 3 E	19	980 658	
Palermo, Sicily	38 6 9	13 22 0 E	20	980 069	
Pola	44 51 8	13 50 7 E	28	980 626	
Praecchia	44 3 0	10 54 3 E	627	980 378	
Romagnano	45 38 1	8 23 8 E	266	980 620	
Rome	41 53 5	12 29.7 E	49	980 367	-0 012
San Remo	43 49 1	7 16.5 E	23	980 505	
Stilfserjoch, see Austria					
Stromboli, Lipari Is	38 48.2	15 14.1 E	48	980 212	
Turin	45 4 1	7 41 8 E	233	980 549	
Jugoslavia, see Yugoslavia					
Netherlands (24)					
Amsterdam (Université)	52 21 9	4 54 7 E	0	981 288	
Bergen op Zoom (Cathédrale)	51 29 7	4 17 3 E	10	981 212	
Breda (Académie Militaire)	51 35 5	4 46 5 E	1	981 213	
De Bilt (Institut Météorologique, base station)	52 6 2	5 10 7 E	2	981.207	
Delft (Institut Géodésique)	52 0 6	4 22.1 E	2	981 264	
Gronigen (Université)	53 13.2	6 34.0 E	5	981 348	
Hollander (Sanatorium Hellendoorn)	52 24.2	6 25 0 E	11	981 296	
Leeuwarden (Friesche Levensverzekering)	53 12 3	5 48 3 E	1	981 348	
Leiden (Observatoire)	52 0 4	4 29 1 E	2	981 273	
Maastricht (Hôtel de Ville)	50 51.2	5 41 6 E	49	981 110	
Middelburg (États Prov.)	51 30 0	3 36 8 E	8	981 215	
Oldenzaal (Église Plechelmi)	52 18 8	6 55.8 E	47	981 282	
Schoorl (École primaire)	52 42.1	4 41 6 E	9	981 312	
Sittard (Ambachtsschool)	50 59.8	5 51 6 E	48	981 148	
Sleen	52 46.5	6 48.1 E	16	981 318	
Terschelling (École Navale)	53 21 6	5 12.9 E	6	981 376	
Ubagsberg	50 51 0	5 57 2 E	191	981 108	
Utrecht (Observatoire)	52 5.2	5 7 8 E	5	981 203	
Weert (Église catholique)	51 15.3	5 42.5 E	33	981 161	
Winschoten	53 8 7	7 2.4 E	0	981.346	

Station	Latitude	Longitude	h	g	TC
Norway (3, 6)					
Bergen (Sternwarte)	60° 23.9'	5° 18.3' E	38	981.922	
Christiansund	63 6.6	7 44.2 E	20	982.175	
Dambaas	62 4.6	9 8.3 E	643	981.892	
Florö	61 33.8	5 2.4 E	10	982.071	
Langenaes	69 1 2	15 8.7 E	8	982.640	
Laredal	61 6.3	7 27.9 E	7	981.942	
Mehavn	71 1.3	27 47 E	10	982.688	
Oslo (Christiania) (Sternwarte, base station)	59 54 7	10 43.5 E	28	981.927	
Oxö	58 4 3	8 3.5 E	10	981.763	
Rörvik	64 51 9	11 14 3 E	10	982.313	
Sand	59 29 1	6 15.7 E	14	981 853	
Hannesjöen	66 1 3	12 38.8 E	12	982.351	
Sörvaagen	67 53 6	13 2 E	19	982.622	+0.0
Stavanger	58 58	5 44.3 E	11	981.845	
Triset	59 25.8	8 10.8 E	115	981.795	
Österreich, see Austria					
Olanda, see Netherlands					
Paesi Bassi, see Netherlands.					
Pays-Bas, see Netherlands					
Poland (2)					
Bedsin	50 19.3	19 8.7 E	256	981.058	
Kraków (Sternwarte)	50 3 9	19 57 6 E	205	981.054	
Lwów (Lemberg)	49 50 2	24 0.0 E	314	980.911	
Tuchla	48 55.2	23 29 E	540	980.789	
Portugal (15)					
Camposancos	41 53 2	8 49 0	9	980.383	
Lisbon	38 42 5	9 11 3	75	980.088	
Oporto	41 8 2	8 36.1	94	980 290	
Praia da Rocha	37 7 0	8 32 7	17	980.005	
Rumania (2)					
Bocsa	46 56.9	22 42 E	379	980 711	
Bucharest (Bucuresti)	44 24 6	26 6.8 E	83	980.553	
Elead	47 2 5	22 22 E	225	980.794	
Maros-Ludas (Ludos)	46 28 1	24 6 E	281	980.715	
Russia and Siberia (2, 11)					
Alexandropol	40 47 0	43 49 7 E	1519	979.785	
Archangel	64 34	40 31 0 E	5	982.278	
Astrakhan	46 21.0	48 2 7 E	-21	980.774	
Byelgorod	50 36.1	36 35 9 E	203	981.038	
Dagarskoje (Lake Baikal), Siberia	55 42 2	109 54 E	465	981 32	
Erivan	40 10 7	44 32 8 E	990	979.880	
Gorjátschinskoi, Siberia	52 59.4	108 18 0 E	470	981.178	
Irkutsk, Siberia (Meteorological Observatory)	52 16 5	104 16 5 E	470	981.096	
Kazan (Observatory)	55 47.4	49 7 3 E	70	981.572	
Kingisepp	59 22.5	28 35.7 E	16	981.858	
Leningrad, see St Petersburg.					
Lenkoran	38 45 6	48 51 5 E	-20	980.092	
Listvinichnoe, Siberia	51 51 0	104 52.5 E	465	981.051	
Moscow (Observatory)	55 45 3	37 34.3 E	139	981.562	
Novgorod	58 31 4	31 17.3 E	48	981.780	
Odessa	46 26 4	30 46 4 E	43	980.769	
Pulkova (base station)	59 46 3	30 19.7 E	71	981.899	
St. Petersburg (Leningrad)	59 56 5	30 17 7 E	3	981 929	
Schaitanskij	56 54 8	59 57.0 E	310	981 641	
Simbirsk	54 19.0	48 24.2 E	181	981.469	
Staraya Russa	57 59 4	31 22 E	23	981.747	
Tartu (Dorpat, Yuriev), (Observatory)	58 22 8	26 43.2 E	50	981.793	
Tiflis (Physical Observatory)	41 43.1	44 47.8 E	412	980.176	
Tver	56 51.2	35 50.9 E	136	981 607	
Verevye	58 40.8	32 42 0 E	113	981.794	
Volkhovo	59 4.2	31 46.2 E	21	981.826	
Vyshnij Volochok	57 35.1	34 33.1 E	164	981.695	
Vologda	59 13	39 53.0 E	118	981.837	
Schweden, see Sweden					
Schweiz, see Switzerland					
Scotland, see Great Britain					

Station	Latitude	Longitude	h	g	TC	Station	Latitude	Longitude	h	g	TC
Spain (18)						**Ungarn, see Hungary**					
Alcázar de San Juan..	39° 24.0′	3° 12.0′	648	979 933		**Ungheria, see Hungary.**					
Andújar............	38 3.0	4 3 0	207	979 943		**Yugoslavia (2)**					
Aranda de Duero ...	41 40.0	3 40.0	801	980 086		Marburg (Maribor)	46° 34′	15° 39′ E.	270	980.706	
Arbas...........	43 0.9	5 45 0	1329	980 132		Ragusa (Dubrovnik)	42 38 6	18 6 E	47	980.394	
Badajoz	38 53.0	6 58 0	188	980 050		Serajevo	43 48.2	18 19.7 E.	311	980.383	
Barcelona	41 25.0	2 7.0 E	407	980 240		**ASIA**					
Basa............	37 30.0	2 45.0	858	979 669		**Giappóne, see Japan**					
Cortegana........	37 54.0	6 47.0	765	979 895		**China (2)**					
Daroca	41 7.0	1 25 0	770	980 038		Hankow	30 35.5	114 17.5 E	73(?)	979.369	
Lérida.	41 37.0	0 38 0 E.	165	980 260		Hongkong	22 18 2	114 10 5 E.	33	978.771	
Llansá....	42 22.0	3 9 0 E.	6	980 431		Port Arthur	38 47 9	121 22.3 E.	1	980.128	
Málaga.	36 43.0	4 25 2	61	979 918		Shasi	30 18 1	112 14.8 E.	132(?)	979 303	
Plasencia	40 2.0	6 3 0	369	980 073		Weihaiwei	37 30 0	122 11 0 E.	1	979.093	
Puigcerdá	42 25.0	1 54 7 E	1190	980 055		Zikawei, Observatory	31 11 6	121 25.8 E.	4	979.437	
Roncal....	42 49.0	0 59 6	675	980 228		**India (4, 9)**					
Salamanca	40 58.0	5 39 0	805	980 057		Agra	27 10.3	78 1 1 E.	163	979.058	−0.018
Salou .	41 4 0	1 0 0 E.	2	980 268		Allahabad	25 25 9	81 55 E.	88	978.945	−0.021
San Fernando.	36 28.0	6 12 3	44	979 843		Badnur	21 54 2	77 54.2 E.	641	978.609	+0.018
Santander	43 29 1	3 49 0	10	980 503		Chatra.	24 12 7	88 23.4 E.	20	978.880	−0.019
Seville .	37 23 0	5 59 0	11	979 965		Colaba	18 53 8	72 48.8 E	10	978.643	0.000
Tarifa	36 0 0	5 37 0	29	979 748		Cuttack	20 29 1	85 52 0 E.	28	978.661	0.000
Toledo	39 51 0	4 1 0	520	980 015		Dehra Dun	30 19 5	78 3 2 E.	682	979.065	−0.060
Torrejón	38 0 1	0 39 1	2	980 032		Dollpur	26 42 0	77 54.8 E.	176	979.001	−0.015
Valencia .	39 29 0	0 23 0	6	980 127		Geaupur	28 33 0	77 42 0 E.	211	979.127	−0.035
Valladolid	41 39 0	4 43 0	695	980 111		Jacobabad	28 16 6	68 27.1 E.	56	979.188	−0.034
Vivero	43 39 0	7 35 0	12	980 553		Jalpaiguri	26 31 3	88 44.2 E.	82	978.924	−0.093
						Jubbulpore	23 8.9	79 59 E.	447	978.731	−0.003
Suede, see Sweden						Kalianpur	24 7.2	77 39.3 E.	537	978.779	+0.011
Suisse, see Switzerland.						Madras	13 4.1	80 14 9 E.	6	978.281	+0.040
Svézia, see Sweden.						Majhauli	26 17.8	83 58 E.	67	978.930	−0.037
Svizzera, see Switzerland.						Mian Mir	31 31.6	74 22.5 E.	216	979.355	−0.033
						Moghal Sarai	25 17 0	83 6 E.	78	978.921	−0.024
Sweden (2)						Montgomery	30 39 8	73 6.3 E	170	979.323	−0.019
Haparanda. . .	65 49 7	24 9.6 E.	4	982 327		Mussoorie (Camel's					
Hernösand ...	62 37 8	17 57 0 E	25	982 082		Back)	30 27 6	78 4.5 E.	2110	978.795	+0.082
Lund (Sternwarte)	55 41 9	13 11 3 E	32	981 564		Musaffarpur	26 7 1	85 25 E.	55	978.936	−0.038
Stockholm (Stern-						Quetta	30 12 2	67 0.7 E.	1682	978.853	+0.024
warte, base station)	59 20 6	18 3.5 E	45	981 843		Raipur	21 13 9	81 41 E.	304	978.614	+0.001
Upsala (Sternwarte) .	59 51 5	17 37 6 E.	20	981 910		Rajpur	30 21 2	78 5 8 E.	1012	979.004	−0.066
Switzerland (6, 23)						Sandakphu Peak	27 6 1	88 0.2 E.	3586	978.192	+0.141
Basel (base station)	47 33 6	7 34 8 E	277	980 788		Yercaud	11 46 9	78 12 5 E	1369	977.910	+0.116
Bern (Landestopo-						**Japan (3, 6)**					
graphie) ...	46 56 5	7 26 8 E	522	980 622		Aomori .	40 49	140 45 E.	1	980.325	
Bironico	46 7 4	8 55 7 E.	473	980 580		Chofu	34 0	131 0 E.	6	979 691	
Brusio	46 15.3	10 7.7 E.	721	980 429		Fukushima	37 45	140 27 E.	67	980.022	
Burgdorf (Techni-						Fukuyama	34 30	133 22 5 E.	3	979.711	
kums) .	47 3 5	7 37.2 E	558	980 633		Hachinohe	40 31	141 30 E.	21	980.359	+ .049
Chanrion (Klubhütte)	45 56.3	7 22.9 E	2435	980 107	+0 113	Hamada .	34 54	132 6 E.	3	979.768	
Eggishorn (Hotel						Hamamatsu	34 42 9	137 43 E.	31	979.750	
Jungfrau)	46 25 2	8 6.8 E	2187	980 169	+0 086	Himeji	34 50 1	134 42 E.	16(?)	979.754	
Frauenfeld (Kantons-						Kamakura	35 19.2	139 34 E.	13	979.779	
schule)	47 33 3	8 54 2 E	431	980 703		Kofu	35 39	138 35 E.	270	979 719	
Fribourg (Universität)	46 47 6	7 9.4 E.	633	980 584		Kurume	33 19.3	130 31 6 E.	11	979 618	
Gornergrat	45 59.0	7 46 8 E	3016	979 992	+0 165	Kyoto	35 1 6	135 47.1 E.	55	979.727	
Grand St. Bernard	45 52 1	7 10 4 E	2473	980 072	+0 131	Matsue	35 30	133 3 E.	23	979.812	
Geneva (Sternwarte)	46 12 0	6 9 2 E	402	980 592		Matsuyama	33 50	132 45 E.	19	979.607	
Gsteig (Hotel						Mizusawa	39 8.1	141 8 E.	61	980.159	
Sanetsch)	46 23.2	7 56 2 E	1185	980 396	−0 001	Nagasaki	32 44 7	129 52.3 E	30	979.594	
Landquart (Schul-						Nagoya	35 10.4	136 53 E	14	979.756	
haus)	46 57.8	9 32.6 E	520	980 523		Nikko .	36 44	139 38 E.	649	979.780	
Lausanne (Ecole de						Okazaki .	34 57.4	137 10 E.	25	979.764	
Chimie et de Physi-						Shizuoka .	34 58.4	138 23 E.	23	979.753	
que) ..	46 31.5	6 38 2 E	531	980 599		Tokyo (base station)	35 42 6	139 46 0 E.	18	979 803	
Les Verrières	46 54 3	6 28 8 E	928	980 573		Tsukuba	36 13 4	140 5 8 E.	870	979.781	
Lungern (Schulhaus)	46 47.1	8 9.6 E.	714	980 515		Uwajima	33 13	132 34 5 E.	2	979.597	
Lusern (Kantons-						Wakayama	34 14 2	135 11 0 E.	3	979.704	
schule)	47 3.0	8 18.2 E.	434	980 626		Yamada	34 29 6	136 42.8 E.	4	979.727	
Neuchatel (Stern-						Yamagata	38 15	140 16 E.	153	980.027	
warte)	47 0.1	6 57.3 E.	487	980 653	−0 026	**Siam (2, 3, 6)**					
Rivera	46 7.4	8 55.7 E	473	980 580		Bankok .	13 43.9	100 29.4 E.	7	978.278	
St. Maurice (Hotel du						**Siberia, (see Russia, p.**					
Simplon)	46 13.0	7 0.2 E	422	980 512	−0 130	**398).**					
Simplonhospis.	46 14.9	8 1 9 E.	1998	980 202	+0 076	**Turkestan (2, 6)**					
Sion (Collège)..	46 14.1	7 21.5 E.	514	980 480	−0 082	Derbent, Bokhara. ..	38 12 0	67 3.2 E.	1012	979.672	
Stilfserjoch, see Aus-						Kala Khum, Bokhara	38 27 3	70 46 5 E.	1345	979.462	−0.086
tria.						Samarkand .	39 39 1	66 58 7 E.	719	979 883	
Truns (Schulhaus) ..	46 44.6	8 59 4 E.	859	980.432		Sultan-Bend .. .	37 7 5	62 28 0 E.	272	979 798	
Zermatt	46 1.5	7 45.0 E.	1603	980 250	−0.007	Tashkent.	41 19.5	69 17 7 E.	478	980.088	
Zernes (Schloss). ...	46 42.0	10 5.8 E.	1473	980.308		Chardzhui (Interna-					
Zürich.	47 22.7	8 33.1 E.	463	980 676		tional Latitude Sta-					
Tscheco-Slovaquie, see Czechoslovakia.						tion).	39 6.2	63 36 1 E.	192	980.014	

Station	Latitude	Longitude	h	g	TC	Station	Latitude	Longitude	h	g	TC
AFRICA						Perth..	31° 57.1'S.	115° 50.5'E.	58	979 378	
Egypt and Anglo-Egyptian Sudan (10)						Sydney	33 51.7 S.	151 12.7 E	43	979 680	
Abu Hamed	19° 32 0'	33° 19 9'E	339	978 538		**OCEANIC**					
Aswan	24 5 1	32 53 1 E	97	978 879		**Atlantic Ocean a n d Mediterranean Sea** (2, 3, 6, 18)					
Atbara	17 41 9	33 58 9 E	354	978 421							
Helwan	29 51 5	31 20 4 E	104	979 295		Bastia, Corsica ..	42 41 2	9 27 E	20	980 519	
Khartum	15 36 6	32 32 9 E	383	978 308		Bridgetown, Barbados	13 4 3	59 36.5	2	978 340	
Luxor	25 43 1	32 39 3 E	82	978 982		Catania, Sicily	37 30 2	15 4.7 E	43	980 065	
Minia	28 5 8	30 45 5 E	42	979 155		Fornella, Balearic Islands	40 3 4	4 7.9 E	7	980.283	
Wadi Halfa	21 55 8	31 19 9 E	126	978 728		Ibiza, Balearic Islands	38 54 3	1 26.1 E.	3	980 146	
Red Sea (2)						Jamestown, St Helena	15 55 8	5 43.7	10	978.712	+0.177
Aden	12 47 3	44 59 3 E	5	978 327		Karajak G l a c i e r, Greenland	70 26 9	50 19 8	20	982 534	
Harmil Island, Dahlak Archipelago Eritrea	16 28 8	40 8 7 E	4	978 465		Kingston, Jamaica	17 57 7	76 47 3	2	978 591	
St. John Island (Zebirget)	23 35 8	36 12 0 E	6	979 026		Las Palmas, Canary Islands	28 7 0	15 26.0	8	979 385	
Mersa Dhiba	25 20 2	34 44 3 E	2	979 007		Palermo, Sicily	38 6 9	13 22 0 E	20	980.069	
Sherm Sheikh (Sinai)	27 51 1	34 16 9 E	2	979 174		Palma de Mallorca, Balearic Islands .	39 34 5	2 39.1 E	23	980 179	
Suez	29 56 0	32 33 4 L	3	979 307		Ponta Delgada, Azores	37 43 8	25 40 8	4	980 113	
Sudan, see Egypt						Reykjavik, Iceland	64 8 5	22 0.3	39	982 273	
Miscellaneous (2, 3)						St George, Bermuda	32 21	64 40	2	979.806	+0 218
Algiers (Observatory)	36 44 8	3 3 E	213	979 905		Santa Cruz de la Palma, Canary Islands	28 41 0	17 46 0	12	979.459	
Biserta, Tunisia	37 16 4	9 52 5 E	7	979 975		Stromboli, Lipari Islands	38 48 2	15 14 1 E	48	980 212	
Biskra, Algeria	34 50 9	5 43 E	137	979 617		Whales Point, Spitsbergen	77 30 4	20 58 8 E	458(?)	982 899	
Cape Town, U S Af (Observatory)	33 56 1 S	18 28 7 E	11	979 657		Valetta, Malta	35 53 8	14 31 3 E	62	979 887	
Dar-es-Salaam, Tanganyika Ter	6 49 0 S	39 18 0 E	7	978 117		**Indian Ocean,** see Pacific Ocean.					
Domjo Ndorobbo	3 08 8 S	35 13 2 E	1715	977 549		**Mediterranean Sea,** see Atlantic Ocean					
Freetown, Sierre Leone	8 29 4	13 14 3	65	978 200		**Pacific and I n d i a n Oceans** (2, 3, 6)					
E Uasso Nyiro, Kenya	1 53 1 S	36 8 2 E	676	977 737		Auckland, New Zealand	36 50 9 S	174 46 2 E	3	979.962	
Johannesburg, U S Af (Observatory)	26 10 9 S	28 4 5 E	1805	978 553		Batavia, Java (Observatory)	6 11 0 S	106 49 8 E	7	978.178	
Kampo, Cameroons, Fr Equat. Af	2 21 2	9 49 6 E	3	978 040		Hobart, Tasmania (Observatory)	42 53 6 S	147 22 0 E	58	980 441	
Laghwat, Algeria	33 47 7	2 53 E	755	979 356		Honolulu, Territory of Hawaii (Observatory)	21 18 1	157 51 S	6	978 946	+0 162
Langenburg, U S Af	9 35 8 S	34 8 6 E	477	977 907							
Libreville, Gabon, Fr Equat Af	0 22 3	9 27.2 E	2	977 999		Kudat, British North Borneo	6 53 0	116 50 7 E	2	978.149	
Loanda, Angola, Portuguese W Af .	8 48 6 S	13 14 1 E	4	978 212		Makassar, Celebes	5 7 3 S	119 24 5 E	2	978 138	
Lourenço M a r q u e s, Mozambique, Portuguese E Af (Observatory)	26 2 5 S	32 19 8 E	55	979 068		Manila, Philippines	14 34 7	120 38 6 E	3	978 360	
						Marau-Sound, Solomon Islands	9 49 1 S	160 48 5 E	3	978.349	
Lüderitz Bay, Southwest Af	26 38 8 S	15 9 7 E	2	979 103		Mauna Kea, Hawaiian Islands	19 49 2	155 28 8	3981	978.069	+0.469
Monrovia, Liberia	6 19 0	10 48 8	41	978 165		Numea, New Caledonia	22 16 6 S	166 27 8 E	2	978.877	
Mozambique, Portuguese E. Af	15 2 1 S	38 25 E	3	978 451		Singapore, S t r a i t s Settlements	1 16 5	103 50 3 E	21	978.082	
Ouled Rhamoun, Algeria	36 10 S	6 41 E	687	979 709		Port Vila, Sandwich Island, New Hebrides	17 45 0 S	168 19 0 E	3	978.637	
Pangani, Tanganyika Ter	5 25 8 S	38 58 8 E	7	978 039		Winter Q u a r t e r s, Kaiser Wilhelm II Land	66 2 2 S	89 38 1 E	1	982 388	
Rio del Rey, Nigeria	4 43 5	8 38 3 E	2	978 087							
Tangier, Morocco	35 46 5	5 48 6	63	979 737							
AUSTRALIA (2, 3, 19)											
Brisbane (Observatory) .	27 28 0 S	153 1 6 E	40	979 148							
Hobart, T a s m a n i a (Observatory)	42 53 6 S	147 22 0 E	58	980 441							
Melbourne (Observatory)	37 49 9 S	144 58 5 E	26	979 987							

TABLE 2.—ACCELERATION OF GRAVITY AT SEA-LEVEL (g_0)

$g_0 = 978.039 \ (1 + 0.005294 \sin^2 \varphi - 0.000\ 007 \sin^2 2\varphi)^*$; Bowie (6). φ = latitude. Unit of g_0 is cm/sec². Basis: Potsdam system

φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²	φ	g_0 cm/sec²
0° 00'	978.039	10° 00'	978 104	20° 00'	978 642	30° 00'	979.328	40° 00'	980 172	50° 00'	981 071	60° 00'	981 917	70° 00'	982 608	80° 00'	983.060
10	.039	10	199	10	.652	10	341	10	186	10	086	10	930	10	.618	10	.065
20	.039	20	205	20	661	20	354	20	201	20	100	20	943	20	.628	20	.070
30	.039	30	.210	30	.671	30	368	30	216	30	.115	30	956	30	637	30	.075
40	.040	40	215	40	681	40	381	40	231	40	130	40	969	40	.647	40	.080
50	.040	50	.221	50	691	50	.394	50	246	50	.145	50	982	50	.656	50	.085
1 00	978.041	11 00	978.227	21 00	978 701	31 00	979 407	41 00	980 261	51 00	981 160	61 00	981 995	71 00	982.665	81 00	983.080
10	.041	10	232	10	711	10	.420	10	276	10	.174	10	982 008	10	.675	10	.094
20	.042	20	.238	20	721	20	434	20	291	20	.189	20	020	20	.684	20	.099
30	.043	30	.244	30	.731	30	.447	30	306	30	.204	30	033	30	.693	30	.103
40	.043	40	.250	40	742	40	.460	40	321	40	.218	40	046	40	.702	40	.107
50	.044	50	256	50	.752	50	.474	50	336	50	233	50	058	50	.711	50	.112
2 00	978.045	12 00	978 262	22 00	978 762	32 00	979 487	42 00	980 350	52 00	981.248	62 00	982.071	72 00	982.720	82 00	983.116
10	.046	10	.268	10	773	10	.501	10	365	10	262	10	083	10	.729	10	.120
20	.048	20	.274	20	.783	20	515	20	380	20	277	20	096	20	.738	20	.124
30	.049	30	.280	30	794	30	528	30	395	30	292	30	108	30	.746	30	.128
40	.050	40	287	40	804	40	542	40	410	40	306	40	121	40	.755	40	.132
50	.052	50	.293	50	815	50	555	50	425	50	321	50	133	50	.764	50	.136
3 00	978 053	13 00	978 300	23 00	978 826	33 00	979 569	43 00	980 440	53 00	981 335	63 00	982 145	73 00	982.772	83 00	983.139
10	055	10	306	10	837	10	583	10	455	10	350	10	157	10	.780	10	.143
20	.056	20	313	20	848	20	597	20	471	20	364	20	169	20	.789	20	.147
30	.058	30	320	30	859	30	611	30	486	30	379	30	182	30	.797	30	.150
40	.060	40	327	40	870	40	624	40	501	40	393	40	194	40	.805	40	.153
50	.062	50	334	50	881	50	638	50	516	50	407	50	206	50	.813	50	.157
4 00	978.064	14 00	978 341	24 00	978 892	34 00	979 652	44 00	980 531	54 00	981 422	64 00	982 218	74 00	982.821	84 00	983.160
10	.066	10	.348	10	.903	10	666	10	546	10	436	10	230	10	.829	10	.163
20	.068	20	.355	20	914	20	680	20	561	20	450	20	.241	20	.837	20	.166
30	071	30	362	30	926	30	694	30	576	30	465	30	253	30	.845	30	.169
40	073	40	369	40	937	40	708	40	591	40	479	40	265	40	.853	40	.172
50	076	50	377	50	948	50	722	50	606	50	493	50	276	50	.861	50	.175
5 00	978 078	15 00	978 384	25 00	978 960	35 00	979 736	45 00	980 621	55 00	981.507	65 00	982 288	75 00	982.868	85 00	983 177
10	081	10	.392	10	971	10	751	10	636	10	521	10	300	10	.876	10	.180
20	083	20	399	20	983	20	765	20	651	20	536	20	311	20	883	20	.182
30	086	30	407	30	994	30	779	30	666	30	550	30	322	30	891	30	.185
40	089	40	.415	40	979 006	40	793	40	681	40	564	40	334	40	.898	40	.187
50	.092	50	423	50	018	50	807	50	696	50	578	50	345	50	.905	50	.189
6 00	978 095	16 00	978 430	26 00	979 030	36 00	979 822	46 00	980 711	56 00	981.592	66 00	982 356	76 00	982.912	86 00	983.191
10	098	10	438	10	042	10	836	10	726	10	606	10	368	10	.919	10	.193
20	102	20	.446	20	054	20	850	20	741	20	620	20	379	20	926	20	.195
30	.105	30	.455	30	065	30	865	30	757	30	634	30	390	30	.933	30	.197
40	108	40	.463	40	077	40	879	40	772	40	648	40	401	40	.940	40	.199
50	.112	50	.471	50	090	50	894	50	787	50	661	50	412	50	.947	50	.201
7 00	978.115	17 00	978 479	27 00	979 102	37 00	979 908	47 00	980 802	57 00	981 675	67 00	982 423	77 00	982.953	87 00	983.202
10	.119	10	.488	10	.114	10	922	10	817	10	689	10	434	10	.960	10	.204
20	.123	20	.496	20	126	20	937	20	832	20	703	20	.444	20	.967	20	.205
30	.127	30	.505	30	.138	30	951	30	847	30	716	30	455	30	973	30	.207
40	131	40	.514	40	151	40	966	40	862	40	730	40	466	40	979	40	.208
50	.135	50	.522	50	.163	50	981	50	877	50	744	50	.476	50	986	50	.209
8 00	978 139	18 00	978 531	28 00	979 175	38 00	979 995	48 00	980 892	58 00	981 757	68 00	982.487	78 00	982.992	88 00	983.210
10	143	10	.540	10	188	10	980 010	10	907	10	771	10	.497	10	.998	10	.211
20	147	20	549	20	200	20	024	20	922	20	784	20	.508	20	983 004	20	.212
30	152	30	.558	30	213	30	039	30	937	30	798	30	.518	30	.010	30	.213
40	.156	40	.567	40	.226	40	054	40	952	40	811	40	.528	40	016	40	.214
50	.160	50	.576	50	.238	50	068	50	967	50	825	50	.539	50	022	50	.215
9 00	978.165	19 00	978.585	29 00	979.251	39 00	980 083	49 00	980 981	59 00	981 838	69 00	982 549	79 00	983 027	89 00	983.215
10	.170	10	.594	10	264	10	098	10	996	10	851	10	559	10	.033	10	.216
20	.174	20	.604	20	277	20	113	20	981 011	20	865	20	569	20	038	20	.216
30	.179	30	613	30	.290	30	127	30	026	30	878	30	579	30	044	30	.216
40	.184	40	.623	40	.302	40	142	40	041	40	891	40	589	40	.049	40	.217
50	.189	50	632	50	.315	50	157	50	056	50	904	50	598	50	.055	50	.217
																90 00	983 217

* This formula differs slightly (not over one in 100 000) from that proposed by Helmert (14) and quite extensively used. A table similar to this, but based on Helmert's formula is given by Albrecht (1)

D. VARIATION OF GRAVITY WITH ELEVATION AND DEPTH

Elevation; Free Air Method.—If there were no matter projecting above the geoid and the geoid were a smooth ellipsoid of revolution, then the value (g_H) of the acceleration of gravity (cm/sec²) at a height H meters above the surface would be related [15, 16] to that (g_0) at the surface, as indicated by equation (1), in which φ is the latitude.

$$g_H = g_0 - (0.000\ 308\ 55 + 0.000\ 000\ 22\cos 2\varphi)H + 0.000\ 072 \left(\frac{H}{1000}\right)^2 \quad (1)$$

This is known as the free air correction. For most purposes it is sufficient to use the approximate formula (2).

$$g_H = g_0 - 0.000\ 3086\ H \quad (2)$$

If g_0 is taken from Table 2, the value of g_H obtained for any station by the use of equation (1) will agree fairly well with the true acceleration, if the surrounding topography is not too rugged. In a fairly flat country, the difference will be considerably less than 0.1 cm/sec², except in very rare cases; and even in a mountainous country, the difference will ordinarily be less than 0.2 cm/sec². For stations below sea-level, but not below the surface of the earth, the same formulae apply; but for such stations, H is negative.

More Exact Methods.—In mountainous country, the computed value will be practically as close to the true value as in flat country if an additional term is added to the right hand side of equation (1), to take account of the elevation of the place above or below the general level of the topography within a radius of, say, approximately 160 km. For every 10 m the place in question is above the general level, this term amounts to 0.001 cm/sec², and for every 10 m below the general level, it amounts to —0.001 cm/sec². In computing the height of a coast station above the general level, the water must be considered replaced by an equal mass of rock, of average surface density, resting on the bottom of the ocean.

If it is desired to obtain a somewhat better value for the computed gravity at a place, the correction term just mentioned must be replaced by a correction for topography and isostatic compensation, computed by the method of John F. Hayford [12].

A somewhat larger error should be expected in the computed values of gravity on oceanic islands than on the continents. The rocks forming these islands are evidently somewhat heavier than normal in many cases, or the ocean is over-compensated, and the observed values of gravity are therefore usually larger than the computed values. In such cases, an error of 0.3 cm/sec², or possibly even 0.4 cm/sec² in computed values may be expected.

Depth.—As the density of the crust is less than two-thirds the mean density of the earth, the acceleration of gravity increases as we advance into the crust. The mean rate of increase is 0.000 0851 cm/sec² per meter of depth. The actual rate at any place depends upon the density of the crustal material in that locality, and is approximately given by the formula [13, 17]

$$g_d = g_0 + (0.000\ 3086 - 0.000\ 0837\rho)d \quad (3)$$

where g_d = acceleration of gravity (cm/sec²) at the depth of d m, and ρ = density (g/cm³).

LITERATURE

(For a key to the periodicals see end of volume)

(1) Albrecht, *Formeln u. Hilfstafeln f geog. Ortsbestimmungen*, 296; 08. (2) Borrass, *C. R. 16th Conf Gen. Assoc. Geodesique Internat.* **3:** 09. (3) Borrass, *C. R. 17th Conf Gen Assoc. Geodesique Internat.* **2:** 261; 12. (4) Bowie, *217*, No **12:** 12. (5) Bowie, *217*, No. **99:** 15. (6) Bowie, *217*, No **40:** 17. (7) Boys, *69*, **186:** 1; 95. (8) Braun, *228*, **64:** 187; 96. (9) Couchman, *Survey of India*, Prof. Paper, No. **15:** 15.
(10) Curry, *Survey Dept. of Egypt*, Paper No. **15:** 13. (11) Gigitsky and Yachontow, *Bull. Astronomic Inst*, No **5:** 28; 24. (12) Hayford and Bowie, *217*, No **10:** 12. (13) Helmert, *Höhere Geodasie*, **2: 46**, 492; 84. (14) Helmert, *76*, 336; 01. (15) Helmert, *76*, 651; 03. (16) Helmert, *Encyklop math. Wiss*, **6** (1B): 97; 10. (17) *Ibid.*, p. 160. (18) Huclin, *Trav. d l'Entens Pesanteur en Espagne*, 24. (19) Love, *251*, **35:** 90; 22.
(20) McDiarmid, *Publ. Dominion Observatory*, **2:** 201; 15. (21) *Ibid.*, **3:** 357; 18. (22) Miller, *Jour. Roy. Astron. Soc. Canada*, **17:** 10, 197; 23. (23) Niethammer, *Astronomischgeodatische Arbeit. in d Schweis; Schweizerischen geodat Komm* **12, 13, 15, 16;** 1910, 1911, 1916, 1921 (24) Vening Meinesz, *Obs. Pendule dans Pays-Bas; Publ. Com. Geod Neerlandaisse*, 23.

AERODYNAMICS

L. J. BRIGGS AND H. L. DRYDEN

Problems in aerodynamics cannot be idealized with the same readiness as problems in mechanics. The side of a building may not be regarded as a thin, flat plate for the purpose of computing the force of the wind, and data for a cylinder of a particular length cannot be directly applied for computing the wind force on a cylinder of some other length. Nearby objects exert an influence which cannot be neglected.

Results obtained for a particular object can be applied strictly only to geometrically similar (definition 6) objects in similar surroundings. Many of the apparent discrepancies among the results of different experimenters are to be attributed to departures from geometrical similarity of the models, to the effects of the supports or other nearby objects, and to differences in the fine structure (turbulence) of the approximately steady air streams, rather than to errors in measuring the force or wind speed. It is not possible to discuss these matters in detail here, and there is no complete discussion available for reference.

SYMBOLS

A	Some specified area	C_M	Moment coefficient (see paragraph on air foils)	$N. A.$ $C. A.$	National Advisory Committee for Aeronautics, U. S. A.
A_r	Aspect ratio				
C	A coefficient	C_N	Coefficient of force normal to the plane of reference	n	Number of revolutions per second
C_{cp}	Coefficient of center of pressure			P_o	Power developed (output)
C_d	Coefficient of drag	C_P	Coefficient of power (input)	P_i	Power input to propeller
C_l	Coefficient of lift			$P R.$	Pitch ratio
				p	Pressure at a point on a surface
				p_s	Static pressure of the air
C_{Po}	Coefficient of power out-put				
C_Q	Coefficient of torque			Q	Torque
C_{Qo}	Coefficient of torque load (output)			Q_o	Torque load (output)
C_T	Coefficient of force parallel to the plane of reference			q	Dynamic pressure, as indicated by Pitot tube (Fig. 1)
C_t	Coefficient of thrust				
$C P.$	Center of pressure			q_o	$\rho V^2/2$ (= q if there is no compression of the air)
c	Length of chord of air foil			R	Reynold's number
D	Diameter			S	That dimension of the plane of reference which is at right angles to the wind = Span
F	Resultant wind force				
F_d	Drag = Component of F parallel to wind				
F_f	Frictional force				
F_l	Lift = Component of F normal to wind and to W			T	Temperature
F_N	Component of F normal to the plane of reference			t	Thickness
				V	Air speed relative to point considered
F_T	Component of F parallel to the plane of reference			V_i	Indicated air speed
F_t	Thrust of propeller			W	Width = That dimension of plane of ref-
F_s	Any component of F				
L	Some linear dimension				
M	Moment of F about forward (leading) edge				

ance which is normal	μ	Viscosity
to S; i.e., makes least	ρ	Density of air when un-
angle with wind		disturbed by bodies
s_o Distance in the plane of		moving relatively to it.
reference, from the	ρ_o	Conventionally chosen
leading edge, or its		"standard" value of ρ
projection to $C. P.$	ϕ	A definite but un-
η Efficiency		specified mathematical
θ_A Angle of attack		function

DEFINITIONS

1. Angle of Attack (θ_A) is the angle which the direction of the wind makes with the plane of reference; it is positive if the wind strikes what is the under side of this plane when the body is in its usual position.

2. Aspect ratio $(A_r) = S/W$.

3. Center of pressure $(C. P.)$ of a body is that point, in the plane of reference, about which the resultant moment of the pressures is zero.

4. Chord (c). See paragraph on airfoils.

5. Coefficient of center of pressure (C_{cp}).

$$C_{cp} = x_c/W; \text{ for airfoil, } C_{cp} = x_c/c.$$

FIG. 1.—Standard Pitot-static tubes.

A = British   B = Bureau of Standards   C = N A C A.

6. Geometrically similar systems. If two bodies together with their surroundings, are so related geometrically that one system corresponds exactly with a uniformly magnified image of the other, the two systems are said to be geometrically similar.

7. Indicated air speed (V_i) is defined by the relation $q = \rho V^2/2 = \rho_0 V_i^2/2$, where ρ_0 is the "standard" air density.

8. Mean temperature (T_m) of atmospheric air column below Z is that temperature for which the pressure at height Z in an isothermal column of air, pressure at bottom $= 760$ mm of mercury, would be that actually observed in the atmosphere at Z.

9. Pitch ratio $(P. R.)_x$ at any point of the blade of a propeller or of a wind-mill distant x from the axis of revolution is $(P. R.)_x = 2\pi x/D \tan \theta_x$, where D is the diameter of propeller or mill wheel, $\theta_x =$ angle which face of blade makes with plane of revolution. If $(P. R.)_x$ is independent of x, propeller has a constant pitch ratio; if θ_x is independent of x, it has a constant blade angle.

10. Reynold's number $(R) = VL\rho/\mu$, where L is some specified linear dimension. The choice of L depends upon the form of the object, and the problem. R is dimensionless.

CONSTANTS ASSUMED

Standard air density is $\rho_0 = 1.2255 \text{ kg/m}^3 (= 0.002377 \text{ slug/ft.}^3)$, which is essentially that of dry air, with normal CO_2 content, at 15°C and one atmosphere.

$\mu/\rho = 1.427 \times 10^{-5} \text{ m}^2/\text{sec} (= 1.535 \times 10^{-4} \text{ ft.}^2/\text{sec})$.

For geometrically similar systems $F_x = qL^2\phi(R) = CAq$ [43], where ϕ is independent of the actual size of the system, and q is the value of the dynamic pressure at some specified point. C is a function only of R and of the geometrical form of the system; its value is the same in every self-consistent system of units, and is independent of the actual size of the system. The data in the following tables and graphs apply when all surrounding bodies

are so far removed from the one considered that they produce no effect upon F_x.

Reduction of Observations.—To obtain true air speed from speed recorded by cup anemometer, use Table 1. Aerodynamic data are usually reduced to a standard air density (ρ_0). For q, this reduction can be effected by replacing the true air speed (V) by the indicated air speed (V_i) (definition 7), and in most cases the same procedure is amply sufficient for C. Example: If $V = 100$ ft./sec in air at 30°C and 754 mm of mercury, $V/V_i = 1.030$ (Fig. 2); hence $V_i = 97.1$ ft./sec and $q_0 = 11.20 \text{ lb./ft.}^2$ (Table 2). Owing to isentropic compression of air at this speed, the actual dynamic pressure (q) is $11.20/0.998$ (Table 3) $= 11.22 \text{ lb./ft.}^2 = 54.78 \text{ kg/m}^2$.

As a basis for the calibration of altimeters, and for use in the comparison of the performances of aircraft, it is assumed that (1) below a certain altitude (Z_i), the rate of decrease (a) of the temperature (T) with the altitude is a constant; (2) above Z_i, $a = 0$; (3) at $Z = 0$, pressure $= p_0$, temperature $= T_0$. The temperature at $Z_i = T_i$; the mean temperature below Z is T_m. All temperatures are reckoned from absolute zero. Then, if $Z < Z_i$,

$$T_m = aZ/\log_e(T_0/T); \text{ if } Z > Z_i, \quad T_m = Z/\left(\frac{1}{a}\log_e\frac{T_0}{T_i} + \frac{Z - Z_i}{T_i}\right),$$

and for any value of Z, $Z = K\frac{T_m}{T_0}\log_{10}\left(\frac{p_0}{p}\right)$.

The values of these constants define what is called the "standard" atmosphere. There is not entire agreement regarding the values which best represent the average atmospheric condition [28]. Those adopted by the governmental aeronautic organizations of the U. S. A. and by many of those of Europe are $T_0 = 288°C$, $T_i = 218°C$, $p_0 = 760$ mm of mercury, $a = 6.500 \times 10^{-3} °C/m$ $(= 1.9812 \times 10^{-3} °C/ft.)$, $Z_i = 10769$ m $(= 35332$ ft.$)$, $K = 19413.3$ m $(= 63691.8$ ft.$)$. These differ slightly from those adopted by the International Commission for Aerial Navigation (see p. 72).

TABLE 1.—ROBINSON CUP ANEMOMETER[*]

True air speed $= V$; recorded speed $= V_r$. If unit is 1 mi./hr, $\log_{10} V = 0.079 + 0.9012 \log_{10} V_r$.

Unit is 1 mi./hr $= 1.467$ ft./sec $= 0.4470$ m/sec

V_r	V	V_r	V	V_r	V	V_r	V
1	1 20	26	22 6	51	41 5	76	59.4
2	2.24	27	23 4	52	42 2	77	60.1
3	3.23	28	24.2	53	42.9	78	60.8
4	4 18	29	24 9	54	43.7	79	61.5
5	5 12	30	25 7	55	44.4	80	62.2
6	6 03	31	26 5	56	45.1	81	62.9
7	6 93	32	27 3	57	45.9	82	63.6
8	7.81	33	28 0	58	46.6	83	64.3
9	8 69	34	28 8	59	47.3	84	65.0
10	9 55	35	29.5	60	48 0	85	65.7
11	10 4	36	30.3	61	48 7	86	66.4
12	11.3	37	31 1	62	49.5	87	67.1
13	12 1	38	31 8	63	50.2	88	67.8
14	12.9	39	32.6	64	50.9	89	68.5
15	13 8	40	33.3	65	51.6	90	69.2
16	14.6	41	34.1	66	52 3	91	69.9
17	15 4	42	34 8	67	53 0	92	70.6
18	16 2	43	35.6	68	53.8	93	71.3
19	17.0	44	36.3	69	54.5	94	72.0
20	17.8	45	37.1	70	55.2	95	72.7
21	18.6	46	37.8	71	55.9	96	73.4
22	19.4	47	38.5	72	56.6	97	74.0
23	20 2	48	39.3	73	57.3	98	74.7
24	21.0	49	40 0	74	58.0	99	75.4
25	21 8	50	40.7	75	58.7	100	76.1

[*] U. S. Weather Bureau type; diameter of cups = 4 in.; centers of cups are 6 72 in. from axis; $V_r = 3$ times linear speed of centers of cups [3, 42, 43].

TABLE 2.—DYNAMIC PRESSURE ($q = q_0$) FOR INDICATED AIR SPEED V_i

Air compression is negligible, and $q = q_0 = \rho_0 V_i^2/2$ if $V_i < 30$ m/sec ($= 100$ ft./sec); for greater speeds, q exceeds q_0, see Table 3. Metric units are m, kg, sec. English units are ft., lb., sec. 1 lb./ft.$^2 = 4.882$ kg/m^2; 1 ft./sec $= 0.3048$ m/sec.

Metric		English	Metric		English	English							
q_0	V_i	q_0	q_0	V_i	q_0	V_i	q_0	V_i	q_0	V_i	q_0	V_i	q_0
0.063	1	0.00119	42 25	26	0 8038	51	3.093	76	6.868	101	12.13	126	18 88
0 250	2	0 00476	45 56	27	0 8668	52	3 215	77	7.050	102	12.37	127	19 18
0.562	3	0 01070	49 00	28	0.9322	53	3 340	78	7.234	103	12.61	128	19 48
1.00	4	0 0190	52 56	29	0 9999	54	3 467	79	7.421	104	12.86	129	19 79
1.56	5	0 0297	56 25	30	1 070	55	3 597	80	7.610	105	13.11	130	20 09
2.25	6	0 0428	60 06	31	1 143	56	3 729	81	7.801	106	13.36	131	20.40
3.06	7	0 0583	64 00	32	1 218	57	3 863	82	7.995	107	13.61	132	20 72
4.00	8	0 0761	68 06	33	1 295	58	4 000	83	8.191	108	13.87	133	21 03
5 06	9	0 0963	72 25	34	1 374	59	4 139	84	8.390	109	14.13	134	21 35
6 25	10	0 1189	76 56	35	1 457	60	4.280	85	8.591	110	14.39	135	21 67
7 56	11	0 1438	81 00	36	1 511	61	4 421	86	8 794	111	14.65	136	21.99
9 00	12	0 1712	85 56	37	1 628	62	4 571	87	9.000	112	14 91	137	22 32
10.56	13	0 2009	90 25	38	1 717	63	4 719	88	9 208	113	15 18	138	22 64
12 25	14	0 2330	95 06	39	1 808	64	4 870	89	9.418	114	15 45	139	22 97
14 06	15	0 2675	100 0	40	1 902	65	5 024	90	9.631	115	15.72	140	23 30
16 00	16	0 3044	105 1	41	1 999	66	5 179	91	9 846	116	16.00	141	23 64
18 06	17	0 3436	110 3	42	2 097	67	5 337	92	10 06	117	16.28	142	23.97
20 25	18	0 3852	115 6	43	2 198	68	5 498	93	10 28	118	16 56	143	24.31
22 56	19	0 4292	121 0	44	2 302	69	5.661	94	10 51	119	16 84	144	24 66
25 00	20	0 4756	126 6	45	2 408	70	5 826	95	10 73	120	17 12	145	25.00
27 56	21	0 5243	132 2	46	2 516	71	5 994	96	10 96	121	17.41	146	25.34
30.25	22	0 5755	138.1	47	2 627	72	6 164	97	11 18	122	17.70	147	25 69
33.06	23	0 6290	144 0	48	2 739	73	6.336	98	11 42	123	17.99	148	26.04
36 00	24	0 6849	150 1	49	2 855	74	6 511	99	11 65	124	18.28	149	26.40
39 06	25	0 7431	156 3	50	2 973	75	6 688	100	11 89	125	18 58	150	26 75

FIG. 2. Ratio of true air speed (V) to indicated air speed (V_i).

FIG. 3.—Air force: flat plates normal to wind.

TABLE 3. CORRECTION FOR ISENTROPIC COMPRESSION (63)

Metric (M) unit of V = 1 m/sec; English (E) = 100 ft sec

V		$\rho v^2/2q$	V		$\rho v^2/2q$
E	M	$= q_0/q$	E	M	$= q_0/q$
1	30	0 998	6	183	0 931
2	61	0 992	7	213	0 907
3	91	0.982	8	244	0 881
4	122	0 969	9	274	0 852
5	152	0 951	10	305	0 822

TABLE 4. WIND PRESSURE ON STRUCTURES

Reference plane (see below) is normal to wind. $F_N = C_N A q$; A = area of projection of object upon reference plane. Unit of $F_N/A = 1$ lb./ft.$^2 = 4.88$ kg/m^2

Object	C_N	F_N/A*
1. Long flat plate	2	30
2. Square flat plate	1 1	16
3. Rectangular prism (1:1:5) (75)	1.6	24
4. Long cylinder	0 8	12
5. Short cylinder	0.7	10

* For $V = 76$ mi per hr ($= 34$ m/per sec) true speed $= 100$ mi. per hr recorded by Robinson anemometer.

Contour intervals = 1mm water
Air speed = 10m/sec; q = 6.24 mm water

LOWER SURFACE   L = Leading edge, T = Trailing edge   UPPER SURFACE

Fig. 4.—Pressure distribution oblong, rectangular plate, inclined (12, 13).

Wind Pressure on Structures.—One must consider (1) maximum wind speed to which the structure will be subjected, (2) the value of the coefficient C_N, and (3) the effective exposed area. The first and the third depend upon local conditions; in the third, shielding effects are very important. The value of C_N should be determined from observations upon a model of the actual structure, as experiments upon flat plates are of little value for this purpose. Opinions differ regarding whether, in gusty winds, the maximum value of F_N is determined by the average or by the maximum value of V (20, 52). Approximate values of C_N for certain typical cases are given in Table 4, where reference plane for flat plate is surface of plate; for prism, its largest face; for cylinder, the plane through axis and normal to that which contains axis and direction of wind. Object (1) is comparable to such structures as wireless masts and long narrow bridge girders; (2) to thin square signboards; (3) to tall buildings; (4) to chimneys; (5) to cylindrical water tanks.

TABLE 5.—SURFACE FRICTION (F_f) ON THIN FLAT PLATES
(Standard density and viscosity)

$F_f \ (= \int \int f dA) = 0.0375 \ AqR^{-0.16} = F_0 A K_w K_v$ (5, 61) where A = total area (both sides) exposed to air stream, F_0 is a factor depending upon the density and viscosity of the air and upon the units employed, and K_w and K_v are numerical factors determined, respectively, by the width (W) of the plate in the direction of the stream, and by the speed (V). F_0 is independent of the ratio S/W, provided $0.5 < (S/W) < 2$; if $S/W = 30$, F_0 is 10% less than the value given in the table. For effect of roughness (it is great), and for variation of f from point to point see (22, 24, 32, 53, 54, 55, 62).

English units $F_0 = 0.0420$ lb /ft.^{2} Unit of $F_f = 1$ lb.; of $A = 1$ ft.^{2}; of $V = 1$ ft./sec				Metric units $F_v = 0.0311$ kg/m^2 Unit of $F_f = 1$ kg; of $A = 1$ m^2; of $V = 1$ m/sec			
W	K_w	V	K_v	W	K_w	V	K_v
1	1.413	10	0 014	1	1 000	10	1.000
2	1 273	20	0 051	2	0.901	20	3.605
3	1 198	30	0 108	3	0 848	30	7.633
4	1 147	40	0 184	4	0 812	40	13.00
5	1 110	50	0 277	5	0 786	50	19.64
6	1 080	60	0 389	6	0 764	60	27.52
7	1 055	70	0 517	7	0 747	70	36 60
8	1 034	80	0 662	8	0 732	80	46.85
9	1 016	90	0 823	9	0 719	90	58.26
10	1 000	100	1.000	10	0 708	100	70.80
11	0 986	110	1 193	11	0 698	110	84 45
12	0 973	120	1 401	12	0.689	120	99.19
13	0 961	130	1 625	13	0 681	130	115.0
14	0 951	140	1 864	14	0.673	140	131.9
15	0 941	150	2 117	15	0 666	150	149.9
20	0 901	160	2 386	20	0 638	160	168.9
30	0 848	170	2 669	30	0.600	170	188.9
40	0 812	180	2 967	40	0 575	180	210.0
50	0 786	190	3 279	50	0 556	190	232.1
100	0 708	200	3 605	100	0.501	200	255.2

FIG. 5.—Pressure distribution: thin circular disc normal to wind.

FIG. 7.—Coefficients: inclined, rectangular plates, A_r = 3. (See Table 6.)

FIG. 6.—Coefficients: square, inclined plates. (See Table 6; for notation, *v.* Fig. 7.)

FIG. 8.—Coefficients: inclined rectangular plates, A_r = 6. (See Table 6; for notation, *v.* Fig. 7.)

Fig. 10.—Pressure distribution: spheres

$F = CAq$; $A = \pi D^2/4$; $R = VD\rho/\mu$

A   Allen (Lit.1)
B₁ Bacon, free air (Lit.2)
B₂ Bacon, compressed air (Lit.2)
E   Eiffel (Lit.15)
    Pannell (Lit.38)
    abouchinski (Lit.49,50)
S   Bureau of Standards (Lit.84)
W   Wieselsberger (Lit.58,60)

$\text{Log}_{10} R \longrightarrow$

Fig. 9.—Air force: spheres.

TABLE 6.—EXPERIMENTAL DATA; FIGURES 6, 7, 8

Unit of S and W = 1 cm; of t = 1 mm; of TD = 1 m; of $R^\dagger$ = 1000

	Fig. 6				Fig. 7				Fig. 8			
	.	×	+	0	.	×	+	0	.	×	+	0
S	25	30.5	12	12	45	7.6	36		90	30 5	72	30.5
W	25	30.5	12	12	15	2 5	12		15	5 08	12	5.08
t	3	3.18	1.7		3	0.25	1.7		3	1 17	1 7	1.29
TD*	1.5	∞	2.0	1.2	1.5	0.6	2.0	1 5	1.37	2 0	1.37
R	210	382	55	42	126	10	55	126	64	55	64

* TD = tunnel diameter.
† R is dimensionless.

The flow about a sphere is extremely sensitive to slight changes in the method of support, and to the condition of turbulence of the air stream. Changes in C are associated with changes in the locus of the points at which the smooth flow leaves the surface, forming a highly turbulent region to the rear. The location of this locus is determined solely by the irregularities in the air stream, as there are no sharp edges or other geometrical feature which might serve to fix it.

Airfoils.—Aerodynamical characteristics are specified in the same manner as are those of plates. An airfoil's area and angle of attack are conventionally defined with reference to some specified plane. The area of the airfoil is defined as that of its normal projection upon this plane of reference. The length (c) of the projection upon this plane of any fore-and-aft section of the airfoil is called the chord of that section; it is the unit in terms of which all dimensions of that section are expressed. The form of the section is specified by the rectangular coordinates of points upon its boundary; the choice of axes is immaterial, although usually one axis is in the plane of reference. The aspect ratio (A_r) of the airfoil is defined as the ratio of length of span (S) to length of the chord. In addition to the coefficients considered for plates, the moment coefficient $C_M = M/(qAc)$, and the lift-drag ratio (F_l/F_d) are also of importance.

Data are usually given for A_r = 6. If $A_r > 3$, then for a given C_l, $\theta_A = \theta'_A + C_l/\pi A_r$ radians, and $C_d = C'_d + C_l^2/\pi A_r$; θ'_A and C'_d are values of θ_A and C_d when $A_r = \infty$; $C_l/\pi A_r$ and $C_l^2/\pi A_r$ are called the induced angle of attack and the induced coefficient of drag, respectively (25, 26, 42, 72).

For airfoils, C_l increases slightly, and C_d decreases very appreciably, as R is increased; C_{c_p} remains unchanged. The difference between the values of the coefficients for airfoils of the size used on aircraft and those for models of the size generally employed in laboratory tests, depends upon the form of the airfoil; for a thin, low cambered section (RAF 15), it is small; for a highly cambered section, it is large.

For the effects produced by placing one airfoil near another, as in a biplane combination see (26, 27, 36, 42, 74).

For a complete airplane, the drag introduced by the body, and the moment of tail lift, both vary appreciably with the size of the airplane (6, 67, 73).

$F = CAq$; $R = VD\rho/\mu$

A = Area of axial section of cylinder = LD; the plane of reference contains the axis and is perpendicular to the plane defined by axis and wind direction.

1. Length infinite and normal to wind; see curves.

Wieselsberger (60)

Reif (45)

2. Length infinite, axis at angle θ with wind

| | $R = 8100^*$ | | $32,500^{**}$ | |
θ	C_N	C_T	C_N	C_T
90°	1.13	0.	1.22	0
80	1.11	0.01		
70	1.02	0.02	1.03	0.04
60	0.88	0.03	0.91	0.04
50	0.70	0.03	0.72	0.04
40	0.50	0.04	0.53	0.04
30	0.32	0.04	0.32	0.04
20	0.151	0.04	$^*Lit.(46)$	
10	0.042	0.03	$^{**}Lit.(84)$	

3. Length (L) finite; normal to wind.
   D = diameter; $R = 80,000^*$ (Lit. 60)

L/D =	1	2	3	5	10	20	40	∞
C_N =	0.63	0.69	0.75	0.74	0.83	0.92	1.00	1.20

* Eiffel (Lit.16) finds that near this value of R, C for short cylinders decreases rapidly as R increases

LOG$_{10}$ R

Fig. 11.—Air force: non-rotating circular cylinders.

Fig. 12.—Pressure distribution: non-rotating circular cylinders.

Axis normal to wind. Dryden's cylinder was 18 diameters long and had end guards each 6 diameters long. (See also Lit. 55)

$Q = C_Q \rho n^2 D^4 L$

$A_r = 3.5$ Lafay (31)

$F_N = C_N Aq$; $A = LD$, L (= length of cylinder) is normal to wind (For direction of F_N, see Fig 14)

Circulation theory

Reid (44)

Lafay (31) $A_r = 3.5$

Rough, Smooth

$\dfrac{\pi n D}{V}$

Fig. 13.—Air force: rotating circular cylinders (Magnus effect).

Reid (44), Lafay (31), Rough, Smooth, $A_r = 3.5$

On circulation theory, θ = 90°

WIND

$\dfrac{\pi n D}{V}$

Fig. 14.—Direction of air force: rotating circular cylinders (Magnus effect).

Contours of $(p - p_s)$; unit = 0.1q

LOWER SURFACE   UPPER SURFACE

Section of airfoil

L is leading edge

Fig. 15.— Pressure distribution: airfoil (30).

Form of half section of strut (See Table 8)

$F_d = C_d A q$; $R = VB\rho/\mu$; $A = BL$
B = maximum value of $2y$
L = length of strut
Axis of section is parallel to wind

Effect of stagger (Lit 41)

θ	$C_d/(C_d)90$	F_l/F_d
10°	0.194	0.593
20	0.312	0.630
30	0.451	0.547
40	0.573	0.516
50	0.688	0.478
60	0.813	0.384
70	0.918	0.251
80	0.995	0.149
90	1.000	0.000

θ·angle length of strut makes with wind
$(C_d)90$ =value of C_d when $\theta = 90°$

U.S. Navy 3, and 4Z almost identical (See Table 8) U S NAVY 3
Variation of C_d with A_f is not definitely known (Lit 4)
Air resistance of struts is very small, and very sensitive
to slight changes in shape, in method of support, and
in turbulence of the air stream

Fig. 16.—Air force on long struts (40, 64, 76, 79).

$F_x = C_x A q$, $A = BL$ (see Fig 16)
for F_d, use curve C_d, for F_l, use C_l
$R = VB\rho/\mu = 21,000$

Strut with end (1) up, $\theta = \theta$
Strut with end (2) up, $\theta = -\theta$

Fig. 17.—Air force on strut 4Z: inclined (85), see
also (4).

TABLE 7.—CHARACTERISTICS OF AIRFOIL SECTIONS

$A_r = 6$; model 36 in. by 6 in.; $V = 40$ mi./hr; $R(= \rho Vc/\mu) = 181\,000$; tunnel diameter $= 7.5$ ft. [57]. θ_A is measured from reference plane AB (see Figs. 22, 23, 24); x and y are rectangular coordinates of points on surface of airfoil (y_u, y_l refer to upper and lower surface, respectively); x is measured in plane AB. Unit of x and of y is 1% of chord. For additional data for these and other sections see [12, 13, 14, 34, 37, 68, 69, 70, 73, 80, 81].

Form			Aerodynamical characteristics					
x	y_u	y_l	θ_A	C_l	C_d	F_l/F_d	x_r/c	C_M
0.00	0.30	+0.30						
1.25	1.90	−0.35						
2.50	2.85	−0.70	−4°	−0 18	0.025	−7.3	—	—
5.00	3.95	−1.05	−2°	−0 04	0.014	−2.8	—	—
7.50	4.65	−1.15	−1°	+0.03	0.013	+2.6	0.966	0.029
10.00	5.05	−1.20	0°	0.14	0.013	10.7	0.479	0.067
15.00	5.55	−0.85	1°	0 24	0.013	18.8	0.407	0.098
20.00	5.78	−0.55	2°	0.32	0.016	20.0	0.367	0.117
30.00	5.80	−0 10	4°	0.46	0.023	20.0	0.321	0.148
40.00	5.60	−0.03	6°	. 0.61	0.033	18.4	0.302	0.185
50.00	5.23	−0.24	8°	0.76	0.047	16 2	0.297	0 228
60.00	4.63	−0.50	10°	0.89	0 061	14.7	0.288	0.260
70.00	4.05	−0.65	12°	1 00	0 083	12.1	0.281	0.286
80.00	3.30	−0.65	14°	1.02	0.124	8.2	0.298	0.313
90.00	2.30	−0.30						
95.00	1.68	0.00						
100.00	0.65	+0.34						

FIG. 22.

0.00	0.00	0.00						
1.25	2.02	−1.65						
2.50	2.71	−2.45						
5.00	3.67	−3.46						
7.50	4.47	−4.10	−4°	−0.26	0.011	—	—	—
10.00	4.95	−4.57	−2°	−0.10	0 012	−8 8	—	—
15.00	5.37	−5.27	0°	+0.04	0.013	+3.1	0.197	0.008
20.00	5.60	−5 58	2°	0 18	0 015	12.4	0 224	0.040
30.00	5.60	−5.69	4°	0 33	0 020	17.2	0.229	0.076
40.00	5.32	−5.27	6°	0.50	0 028	17.5	0.241	0.121
50.00	4.68	−4.52	8°	0.65	0.040	16 2	0.242	0.159
60.00	3.72	−3.56	10°	0.78	0 054	14 6	0.244	0.193
70.00	2.61	−2.39	12°	0.88	0.076	11.6	0.246	0 220
80.00	1.60	−1.44	14°	0.73	0.170	4.3	0 234	0.181
90.00	0.69	−0.74	16°	0.70	0.239	2.9	0.382	0.293
95.00	0.37	−0.43						
100.00	0.16	−0.16						

FIG. 23.

| 0.00	3.61	3.61						
1.25	6.74	1.35						
2.50	7.98	0 80	−8°	−0.07	0.071	−0.9	—	—
5.00	9.86	0.35	−6°	+0.08	0.031	+2 6	1.410	0.109
7.50	11.32	0.18	−4°	0.22	0 024	9.4	0.684	0.150
10.00	12.40	0.09	−2°	0.37	0.026	14 3	0 507	0.188
15.00	13.83	0.00	0°	0.51	0.031	16.4	0.436	0.222
20.00	14.77	0.07	2°	0.66	0.039	16.9	0 396	0.261
30.00	15.36	0.21	4°	0.81	0.051	15.9	0.369	0.300
40.00	14.88	0.37	6°	0.96	0.067	14.3	0.348	0.336
50.00	13.47	0.54	8°	1.10	0.084	13.0	0.337	0.374
60.00	11.50	0.54	10°	1.23	0.104	11.8	0 323	0.403
70.00	9.27	0.54	12°	1.33	0.125	10.6	0.307	0.416
80.00	6.57	0.49	14°	1.42	0.148	9.6	0.312	0.454
90.00	3.61	0.27	16°	1.43	0.182	7.9	0 315	0.466
95.00	1.99	0.16	18°	1.42	0.213	6.7	0.327	0.486
100.00	0.36	0.00	20°	1.41	—	—		

FIG. 24.

TABLE 8.—FORM OF STRUTS; U. S. NAVY 3, BRITISH 4Z

(See Fig. 16) (These struts give as small a C_d as any)

Unit = axial length of section

x	2y		x	2y		x	2y	
	U.S.N. 3	4Z		U.S.N. 3	4Z		U.S.N. 3	4Z
0	0	0	0.250	0.240		0.700	0.184	0.182
0.025	0.092		0.300	0.247	0.250	0.750	0.164	
0 050	0.132	0.122	0.350	0.250		0.800	0.142	0.142
0 075	0 159		0.400	0 250	0.246	0.850	0.116	
0 100	0.180	0 182	0.450	0 250		0.900	0.085	0.094
0 125	0.197		0.500	0.240	0.234	0.950	0.049	
0 150	0 210		0.550	0.230		1.000	0.000	0.000
0.175	0 220		0.600	0.215	0.212			
0 200	0 229	0 240	0 650	0 201				

FIG. 18.—Durand's $F_2A_1S_1P_1$ propeller family. Pitch ratio constant. (Members differ only in pitch ratio.)

FIG. 19.—Characteristics of Durand propellers at a fixed point [8, 19].

Elongated stream-line solids of revolution have a small resultant drag, which varies greatly with turbulence of air stream, position of neighboring bodies, and slight changes in form. The area entering into the expression $F = CAq$, is generally taken either as the area of maximum section normal to the length, or as (volume)$^{2/3}$. C varies with the Reynold's number. When $A = $ (volume)$^{2/3}$, the minimum value of C for large values of R, and for bodies which are 4 to 5 diameters long, is of the order of 0.014. When $A = $ sectional area, the minimum value of C is of the order 0.03, and is obtained with bodies shorter than 4 diameters. Their equilibrium when parallel to the air stream is unstable; adding fins gives stability and greatly increases their drag [23, 35, 39].

Propellers.—Propellers are usually divided into families in which pitch-ratio and diameter are the only variables. Blade thickness and outline are usually determined largely by structural considerations; if the average thickness and width of blade are fixed, other variations have small effect upon attainable efficiency [6, 9, 15, 19, 65, 66, 71, 76, 77].

The characteristics of a propeller working at a fixed point may be expressed by two dimensionless coefficients, C_t and C_P, defined by the equations $F_t = C_t \rho n^2 D^4$ and $P_t = C_P \rho n^3 D^5$. For most propellers, there is, between C_t and C_P, a functional relation which is nearly independent of the design, provided large blade angles are not used [33]. In Fig. 19, the curve indicates the most favorable results; marked departures from the curve occur mainly with propellers of high pitch ratio, or of constant blade angle.

Fig. 20.—Characteristics of advancing Durand $F_1A_1S_1P_1$ propeller family [9].

The characteristics of propellers at various forward speeds (V) and speeds of rotation may be expressed by curves showing the relationships between three parameters. In Fig. 20, the parameters used are C_t, C_P, and $s$ or η, defined by the equation $F_t = C_t \rho V^2 D^2$, $P_t = C_P \rho n^3 D^4$, $s = V/Dn$; $\eta = C_t s^3 / C_P$, and D = diameter of the propeller. Useful range of C_t is 0.05 to 0.25; of C_P is 0.04 to 0.16. Data given are for propellers of two blades; increasing the number of blades, displaces the curves upwards and to the right.

Wind mills.—Quite different principles control the designing of wind mills which derive power from natural winds, and of those (such as the small wind mills used on airplanes for driving fuel pumps, etc.) which derive their power from the motion of a power driven craft. In the former, the controlling factor is the cost per unit of power developed; in the latter, it is the power consumed per unit of power, or torque load, developed.

LITERATURE

(For a key to the periodicals see end of volume)

[1] Allen, 5, 50: 323, 310; 00. [2] Bacon and Reid, 197, No. 126. [3] Bradfield, 199, No. 712. [4] Cowley, et. al., 300, No. 266. [5] Diehl, 196, No. 102. [6] Diehl, 197, No. 111. [7] Dines, 5, 48; 233; 90. [8] Durand and Lesley, 197, No. 30. [9] Ibid., No. 141.
[10] Durand, et al, 198, No. 4, appx. [11] Dryden, 31, No. 394; 20. [12] Eiffel, Resistance de l'air et l'aviation (Paris, Dunod et Pinat), 2nd ed., p. 42. [13] Ibid., p 150. [14] Ibid., p 231. [15] Eiffel, Nouvelle recherches sur la resistance de l'air et l'aviation (Paris, Duond et Pinat), p. 87. [16] Eiffel, Travaux Laboratoire aerodynamique Eiffel, 1915-18, p. 60. [17] Ibid., p 85. [18] Fage, 300, No. 106. [19] Fage, Airscrews in Theory and Experiment (London, Constable and Co), 1920.
[20] Fleming, Wind Pressure on Structures, 1915. [21] Föppl, 301, 4: 51;10. [22] Froude, 155, 1872: 118. 1874: 249. [23] Fuhrmann, 301, 5: 65; 11. [24] Gibbons, 197, No. 8, pt. 1. [25] Glauert, 199, No. 722. [26] Ibid., No. 849. [27] Ibid., No. 901. [28] Gregg, 197, No. 147. [29] Hunsaker, 308, 62: 4: 77; 16.
[30] Jones and Paterson, 300, No. 72. [31] Lafay, Rev. mecanique, 30: 417; 12 [32] Lanchester, 300, No. 149. [33] Margoulis, Les helicopteres. [34] Morse, U S. Air Service, Inf. Circ. No. 478. [35] Munk, 197, No. 184. [36] Ibid., No. 181. [37] Norton and Bacon, 197, No. 168. [38] Pannell, 300, No. 190. [39] Pannell and Jones, 300, No. 190.

Fig. 21.—Characteristics of two blade windmill [17].

[40] Powell, 300, No. 416. [41] Ibid., No. 899. [42] Prandtl, 197, No. 116. [43] Rayleigh, 300, No. 29. [44] Reid, 198, No. 209. [45] Relf, 300, No. 102. [46] Relf and Powell, 300, No. 307. [47] Riabouchinski, 308, 4: 43, 56; 12. [48] Ibid., 4: 113; 12. [49] Ibid., 5: 73; 14.
[50] Riabouchinski, 298, No. 44. [51] Stanton, 155, 156: 78; 03. [52] Ibid., 216: 34; 22. [53] Stanton, 115, 117: 718; 24. [54] Stanton and Marshall, 300, No. 621. [55] Taylor, 300, No. 191. [56] Ibid., No. 604. [57] Warner, E P, O. [58] Wieselsberger, Z. Flugtechnik Motorluftschiffahrt, 5: 140; 14. [59] Ibid, 6: 127; 15.
[60] Wieselsberger, 65, 22: 219; 22. [61] Wieselsberger, 304, 1: 120; 21. [62] Zahm, 5, 5: 58; 04. [63] Zahm and Smith, 197, No. 61. [64] Zahm, et. al., 197, No. 127. [65] Zahm, et. al, 300, Nos. 61, 62, 63, 64, 82, 128, 155, 259, 264, 265, 305, 316, 328, 331, 371, 382, 385, 390, 393, 401, 402, 408, 431, 437, 439, 433, 443, 444, 458, 460, 475, 540, 548, 571, 577, 596, 591, 594, 639. [66] Zahm, et. al., 299, Nos. 669, 765, 829, 830, 869, 870, 871, 881, 882, 884, 885, 887, 892. [67] Ibid, No. 900. [68] Zahm, et. al., 197, No. 93. [69] Ibid, No. 124.
[70] Zahm, et. al, 197, No. 182. [71] Ibid., Nos. 14, 64, 83, 109, 155,*175, 177, 182, 186, 196, 207. [72] Zahm, 304, 1: 37; 21. [73] Ibid., 1:71; 21. [74] Ibid., 2: 9, 10, 11; 23. [75] Ibid., 2: 33; 23. [76] Zahm, 303, 2: 3; 09. [77] Ibid., 4: 80; 12. [78] Zahm, Tech. Ber. Flugzeugmeisterei, 1, No. 4: 119; 17. [79] Ibid., 2, No. 1: 15; 18.
[80] Zahm, Tech. Ber. Flugzeugmeisterei, 1, No 5: 148; 17. [81] Ibid., 1, No. 6: 204; 17. [82] Zahm, Anemometry (3rd ed.) U. S. Weather Bureau, Inst. Div., Circ. D. [83] Zahm, 305, 37: 288; 97. [84] U. S. Bureau of Standards, O. [85] The National Physical Laboratory, O.

LITERATURE REFERENCES

LITERATURE REFERENCES

In all literature references cited in International Critical Tables the name of the journal or publication is indicated by means of a *Key Number* corresponding to the list given below. The numbers which follow this key number in a literature citation are, in the order named: (1) the volume, (2) the page, and (3) the last two figures of the year. Thus, 64*V*, **31**: 253; 22, indicates Verslag Koninklijke Akademie van Wettenschappen te Amsterdam, Vol 31, page 253, 1922. Series numbers are not given. Key Numbers referring to books and other non-serial publications are preceded by the letter *B*, and the volume number is given in Roman numerals. Thus, *B*10, **IV**: 191; 18 indicates Doelter, Handbuch der Mineralchemie, page 191 of Vol. 4 of the 1918 edition. The Key Number *O* is used to indicate "private communication from."

RÉFÉRENCES BIBLIOGRAPHIQUES

Le nom du journal ou de la publication de toutes les références bibliographiques citées dans les Tables critiques internationales est indiqué au moyen d'un nombre-clé correspondant à la liste donnée ci-dessous. Les nombres qui suivent ce nombre-clé dans un renvoi bibliographique indiquent dans l'ordre suivant: (1) le volume, (2) la page, et (3) les deux derniers chiffres de l'année. Ainsi 64*V*, **31**: 253; 22, indique Verslag Koninklijke Akademie van Wettenschappen te Amsterdam, vol. 31, page 253, 1922. Les numéros des séries ne sont pas donnés. Les nombres-clés se rapportant à des livres ou à des publications non périodiques sont précédés de la lettre *B* et le numéro du volume est donné en chiffres romains. Ainsi, *B*10, **IV**: 191; 18 indique Doelter, Handbuch der Mineralchemie, page 191 du volume 4 de l'édition de 1918. Le nombre-clé *O* est employé pour indiquer "communication privée de."

DAS LITERATURVERZEICHNIS

In allen Literaturstellen, die in I. C. T. verzeichnet sind, ist der Name der Zeitschrift oder der Publikation mit Hilfe einer *Schlüsselnummer*, entsprechend der unten folgenden Liste, angegeben. Die Zahlen, welche diesen Schlüsselnummern bei einem Literaturzitat folgen, bedeuten der Reihe nach: (1) der Band, (2) die Seite und (3), die letzten zwei Zahlen des Jahrganges. So bedeutet z. B. 64*V*, **31**: 253; 22, Verslag Koninklijke Akademie van Wettenschappen te Amsterdam, Band 31, Seite 253, 1922. Serien-Nummern werden nicht angegeben. Der Schlüsselzahl wird ein *B* vorausgesetzt, wenn sie Bücher, oder eine andre nicht periodische Veröffentlichung bezeichnet. Die Bandnummer wird durch römische Ziffern angegeben. Es bedeutet z. B. also *B*10, **IV**: 191; 18, Doelter, Handbuch der Mineralchemie, Seite 191, des 4. Bandes, der Auflage des Jahres 1918. Die Schlüsselzahl Null wird gebraucht, um anzuzeigen, dass es eine 'private Mitteilung' ist.

INDICAZIONI BIBLIOGRAFICHE

In tutte le indicazioni bibliografiche che si incontrano nele "Tabelle Critiche Internazionali" il nome del giornale o della pubblicazione è espresso con un *numero chiave* riportato nell'elenco dato più oltre. I numeri che, nella citazione, vengono dopo il numero chiave sono disposti con l'ordine seguente: (1) il volume, (2) la pagina, e (3) le ultime due cifre del millesimo. Cosi 64*V*, **31**: 253; 22, indica la Verslag Koninklijke Akademie van Wettenschappen te Amsterdam, Vol. 31, pagina 253, 1922. I numeri di serie non vengono dati. Quando un numero chiave è preceduto dalla lettera *B* si riferisce a libri o ad altre pubblicazioni non periodiche, e il numero del volume viene allora scritto in cifre romane. Cosi *B*10, **IV**: 191; 18 indica Doelter, Handbuch der Mineralchemie, pagine 191 del IV° volume dell'edizione 1918. Il numero chiave *O* indica "Comunicazione privata da　　　"

KEY TO THE PERIODICALS

Data regarding the libraries which receive many of these periodicals may be found through the following sources:

United States and Canada: "Periodicals Abstracted by Chemical Abstracts, 1926" (Chemical Abstracts, Ohio State Univ., Columbus, Ohio); "Union List of Serials in the Libraries of the United States and Canada, 1925–   " (H. W. Wilson & Co., New York City); "A Catalogue of Scientific Periodicals in Canadian Libraries, 1924" (McGill Univ., Montreal, Canada).

Great Britain: "A World List of Scientific Periodicals Published in the Years 1900–1921" (Oxford Univ. Press, London, 1925–   )

Holland: "Chemisch Jaarboekje tevens Jaarboekje der Nederlandsche Chemische Vereeniging, vol. 3." (Amsterdam, D. B. Centen, 1920.)

1. Journal of the American Chemical Society.
2. Physical Review.
3. London, Edinburgh and Dublin Philosophical Magazine and Journal of Science.
4. Journal of the Chemical Society, London
5. Proceedings of the Royal Society (London). A Mathematical and Physical Sciences.
6. Annales de chimie et de physique. *See also* Nos 14 and 16
7. Zeitschrift für physikalische Chemie, Stöchiometrie und Verwandtschaftslehre.
8. Annalen der Physik. [Journal der Physik, 1790–1794. Neues Journal der Physik, 1795–1796. Annalen der Physik, 1799–1819; Annalen der Physik und der physikalische Chemie, 1819–1824 (Gilbert). Annalen der Physik und Chemie, 1824–1899 (Poggendorff, Wiedemann). Annalen der Physik, 1900–   (Drude, Wien and Planck).]
9. Zeitschrift für Elektrochemie und angewandte physikalische Chemie.
11. American Chemical Journal. Combined with No. 1 in 1914
12. American Journal of Science.
13. Annalen der Chemie, Justus Liebig's.
14. Annales de chimie.
16. Annales de physique.
18. Archives Néerlandaises des sciences exactes et naturelles. Series III A. Sciences exactes
19. Arkiv for Kemi, Mineralogi och Geologi
21. Astrophysical Journal.
22. Atti della reale accademia nazionale dei Lincei. (Rendiconti classe di scienze fisiche, matematiche e naturali)
24. Atti del reale istituto Veneto di scienze, lettere ed arti.
25. Berichte der deutschen chemischen Gesellschaft
26. Berichte der deutschen physikalischen Gesellschaft. *See also* No. 96.
27. Bulletin de la société chimique de France.
28. Bulletin de la société chimique de Belgique.
29. Bureau of Mines, Bulletins.
31. Bureau of Standards, Scientific Papers.
31A. Bureau of Standards, Bulletin
32. Bureau of Standards, Technologic Papers.
33. Chemical and Metallurgical Engineering.
34. Comptes rendus hebdomadaires des séances de l'académie des sciences, de l'institut de France.
36. Gazetta chimica italiana.
38. Journal of the American Ceramic Society.

41. Journal of the Chemical Society of Japan (Nippon Kwagaku Kwai Shi)
42. Journal de chimie physique.
45. Industrial and Engineering Chemistry
47. Journal of the Institute of Metals, London.
48. Journal of the Optical Society of America and Review of Scientific Instruments.
50. Journal of Physical Chemistry.
51. Journal de physique et le radium. *See also* No. 199.
53. Journal of the Russian Physico-Chemical Society.
54. Journal of the Society of Chemical Industry.
55. Kolloid-Zeitschrift (*Formerly* Zeitschrift für Chemie und Industrie der Kolloide.)
57. Monatshefte für Chemie und verwandte Teile anderer Wissenschaften.
58. Nature, London.
59. Nuovo Cimento.
62. Philosophical Transactions of the Royal Society of London.
63. Physikalische Zeitschrift.
64P. Proceedings of the Royal Academy of Sciences of Amsterdam.
64V. Verslag koninklijke Akademie van Wetenschappen te Amsterdam.
65. Proceedings of the American Academy of Arts and Sciences.
67. Proceedings of the Physical Society of London.
68. Proceedings of the Royal Society of Edinburgh.
69. Proceedings and Transactions of the Royal Society of Canada.
70. Recueil des travaux chimiques des Pays-Bas.
72. Rendiconti reale istituto Lombardo de scienze e lettere.
75. Sitzungsberichte Akademie der Wissenschaften in Wien, mathematisch-naturwissenschaftliche Klasse.
76. Sitzungsberichte der preussischen Akademie der Wissenschaften.
77. Stahl und Eisen.
78. Transactions of the American Electrochemical Society.
80. Transactions of the American Institute of Mining and Metallurgical Engineers
83. Transactions of the Faraday Society.
88. Verhandlungen der physikalischen Gesellschaft zu Berlin. *See also* No 96
89. Wissenschaftliche Abhandlungen der physikalisch-technischen Reichsanstalt.
91. Zeitschrift für analytische Chemie.
92. Zeitschrift für angewandte Chemie.
93. Zeitschrift für anorganische und allgemeine Chemie.
94. Zeitschrift für Krystallographie. (*Name changed in* 1921 *from* Zeitschrift für Kristallographie und Mineralogie.)
95. Zeitschrift für Metallkunde.
96. Zeitschrift für Physik. (Verhandlungen der physikalischen Gesellschaft zu Berlin, 1882–1898; Verhandlungen der deutschen physikalischen Gesellschaft, 1899–1902; Berichte der deutschen physikalischen Gesellschaft, 1903–1919; Zeitschrift für Physik, 1920–   .)
98. Zeitschrift des Vereines deutscher Ingenieure.
101. Elektrotechnische Zeitschrift
105. Journal of the Society of Glass Technology.
112. Dinglers polytechnisches Journal.
115. Engineering.
119. Proceedings of the American Institute of Electrical Engineers.

413

128. Journal of the Washington Academy of Sciences.
132. Anales de la sociedad española de física y química.
133. British Association for the Advancement of Science, Reports.
135. Chemical News and Journal of Industrial Science. (Name changed in 1921 from Chemical News and Journal of Physical Science.)
136. Chemiker Zeitung.
137. Kongelige Danske Videnskabernes Selskab, Mathematisk-fysiske Meddelelser.
138. Societas scientiarum fennica. Commentationes physico-mathematicae.
139. Ferrum.
140. Journal of the Iron and Steel Institute, London.
141. Journal of Biological Chemistry.
143. Journal of the Franklin Institute.
144. Mathematikai és Természettudományi Értesitö, Budapest.
147. Meddelanden från K. Vetenskapakademiens Nobelinstitut.
149. Archives des sciences physiques et naturelles. (Bibliothèque brittanique, 1796-1815; Bibliothèque universelle des sciences, belles-lettres et arts, 1816-1835; Bibliothèque universelle de Genève, 1836-1845; Supplément à la bibliothèque universelle de Genève. Archives des sciences physiques et naturelles, 1846-1847; Bibliothèque universelle de Genève. Archives des sciences physiques et naturelles, 1848-1857; Bibliothèque universelle, revue suisse et étrangère. Archives des sciences physiques et naturelles, 1858-1861, Bibliothèque universelle et revue suisse. Archives des sciences physiques et naturelles, 1862-1877; Bibliothèque universelle. Archives des sciences physiques et naturelles, 1878-   .)
152. Carnegie Institution of Washington Publications.
153. Minutes of Proceedings of the Institution of Civil Engineers.
156. U. S. Geological Survey, Bulletin.
159. Science Reports of the Tôhoku Imperial University.
166. Science.
168. Communications from the Physical Laboratory of the University of Leiden.
173. Analyst, London.
175. Annales academiae scientiarum fennicae.
176. Chemisch Weekblad, Amsterdam.
186. Bulletin de la classe des sciences, académie royale de Belgique
187. Metall und Erz, Zeitschrift für Metalhuttenwesen und Erzbergbau, einschl. Aufbereitung.
188. Nachrichten von der koniglichen Gesellschaft der Wissenschaften zu Göttingen. Geschäftliche Mitteilungen; mathematisch-physikalische Klasse.
189. Centralblatt für Mineralogie, Geologie und Palaontologie
190. Neues Jahrbuch für Mineralogie, Geologie und Palaontologie.
196. Sammlung chemischer und chemisch-technischer Vortrage.
197. Proceedings of the National Academy of Sciences.
198. Revue générale des sciences pures et appliquées
199. Le Radium. (Merged into No. 51 in 1920.)
200. Jahrbuch der Radioaktivität und Elektronik.
201. Proceedings of the Cambridge Philosophical Society.
202. Zeitschrift für physiologische Chemie.
205. Biochemische Zeitschrift.
207. Geologiska Föreningens i Stockholm Förhandlingar.
208. Physica, Nederlandsch Tijdschrift voor Natuurkunde.
209. Japanese Journal of Chemistry.
210. Scientific Papers, Institute of Physical-Chemical Research, Tokyo.
211. Abhandlungen der mathematisch-physischen Klasse der sächsischen Akademie der Wissenschaften zu Leipzig.
212. Transactions of the American Society for Steel Treating.
213. Sitzungsberichte der mathematisch-physikalischen Klasse der Bayerischen Akademie der Wissenschaften zu München

214. Kongelige Danske Videnskabernis Selskab, Skrifter naturvidenskabelig og mathematisk Afdeling.
215. Lunds Universitets Årsskrift.
216. Giornale di chimica industriale ed applicata. (Annali di chimica applicata, 1914; continued as Giornale di chimica applicata; combined with Giornale di chimica industriale, March, 1920, to form Giornale di chimica industriale ed applicata.)
217. U. S. Coast and Geodetic Survey, Special Publications.
218. Naturwissenschaften.
219. Proceedings of the Physico-Mathematical Society of Japan.
220. Jern-Kontorets Annaler, Stockholm.
221. Berichte über die Verhandlungen der sächsischen Akademie der Wissenschaften zu Leipzig. Mathematisch-physische Klasse.
222. Giornale di mineralogia, cristallografia e petrografia.
223. Journal of General Physiology.
224. Kosmos, Stockholm.
226. Mitteilungen aus dem Kaiser-Wilhelm Institut für Eisenforschung zu Düsseldorf.
227. Proceedings of the Society for Experimental Biology and Medicine.
228. Denkschriften der kaiserlichen Akademie der Wissenschaften zu Wien, mathematisch-naturwissenschaftliche Klasse.
229. Journal of Bacteriology.
230. Biochemical Journal.
231. U. S. Public Health Service, Public Health Reports.
232. Soil Science.
233. Pharmaceutisch Weekblad.
234. Journal of the South African Chemical Institute. (Name changed in 1922 from Journal of the South African Association of Analytical Chemists.)
235. Comptes-rendus des travaux du laboratoire Carlsberg.
236. Ergebnisse der Physiologie.
237. Fortschritte der Chemie, Physik und physikalischen Chemie.
238. Travaux et mémoires du bureau international des poids et mesures.
239. Nouveaux mémoires de l'académie royale des sciences, des lettres et des beaux-arts de Belgique, Brussels.
240. Bibliothèque universelle des sciences, belles-lettres et arts. (Continued as No. 149.)
241. Proceedings of the American Philosophical Society.
242. Vierteljahrsschrift der naturforschenden Gesellschaft, Zürich.
243. Zeitschrift für Instrumentenkunde.
244. Journal of the Society of Automotive Engineers.
245. Zeitschrift für das gesamte Schiess- und Sprengstoffwesen.
246. Ice and Refrigeration.
247. Chemist-Analyst.
248. Proceedings of the University of Durham Philosophical Society.
249. Fortschritte auf dem Gebiete der Röntgenstrahlen.
250. Bulletin de la société française de physique.
251. Proceedings of the Royal Society of Victoria, Melbourne.
252. Chemische Umschau auf dem Gebiete der Fette, Oele, Wachse und Harze. (Before 1916 Chemische Revue über die Fett- und Harz Industrie.)
253. Lubrication.
254. Zeitschrift für Beleuchtungswesen, Heizungs- und Lüftungstechnik.
255. Bulletin of the American Institute of Mining and Metallurgical Engineers. (Continued as No. 329.)
256. Comptes rendus de la société scientifique, Warsaw.
266. Indianapolis Medical Journal.
267. Philippine Journal of Science.
268. Terrestrial Magnetism.

IMPORTANT NOTICE

ANNUAL TABLES OF CONSTANTS AND NUMERICAL DATA
CHEMICAL, PHYSICAL AND TECHNOLOGICAL

Published by an International Committee, under the authority of the International
Research Council and the International Union of
Pure and Applied Chemistry

THE data given in International Critical Tables are based upon determinations published previous to January 1924.

In order to keep always up to date with future progress in physical measurements, the purchaser of International Critical Tables should become a regular subscriber to *International Annual Tables*, which will bring to his desk each year a volume in which he will find assembled and classified all of the data which has appeared in the world's literature, for the year covered by the volume.

Thus, *Annual Tables*, Volume VI, will contain all the data published in the literature of 1923-1924; Volume VII will contain the data published in the literature of 1925; Volume VIII, the data to be found in the literature of 1926, etc. . . .

The older volumes of Annual Tables go back to 1910 and contain all of the original determinations published during that period and which served as all or part of the basis of selection of the "best" values recorded in Inter-

national Critical Tables. These older volumes thus constitute a handy and accessible source of reference to the original determinations.

Through special arrangements made with the International Commission in charge of Annual Tables, all purchasers of International Critical Tables will be given special discounts on all purchases of International Annual Tables, as follows:

1. Volumes I, II, III, IV, V (data for 1910-1922), cloth, $52.50 (regular price, $71.00).

2. 25% discount on Volumes VI to X inclusive, covering the period 1923-1928.

3. 25% discount on the index to Volumes I-V, which will be published in 1927.

These discounts apply only on subscriptions addressed to Dr. C. Marie, 9 rue de Bagneux, Paris VI, France.

All subscriptions should be accompanied by draft or money order payable at Paris.

SPECIMEN SECTIONS

Free copies of any of the following specimen sections may be obtained on application to the above address.

Spectroscopy. Electricity, Magnetism, Electrochemistry. Radioactivity. Crystallography, Mineralogy. Biology. Engineering, Metallurgy. Colloids. The following identification form should be filled out and transmitted with the first subscription.

Name . . . .

Address . . . .

.